

A mobile app providing individually-tailored psychoeducation about sleep for older adults with chronic health conditions and low health literacy

Raymond L Ownby¹

Kamilah Thomas-Purcell²

Donrie Purcell³

Joshua Caballero⁴

Sweta Tewary⁵

Rosemary Davenport¹

Michael Simonson⁶

¹Department of Psychiatry and Behavioral Medicine, Nova Southeastern University, Fort Lauderdale, Florida, USA

²Department of Health Science, College of Health Care Sciences, Nova Southeastern University, Fort Lauderdale, Florida, USA

³Department of Psychiatry and Behavioral Medicine, Nova Southeastern University, Fort Lauderdale, Florida, USA. Now at Department of Community Health & Preventive Medicine, Morehouse School of Medicine, Atlanta, Georgia, USA.

⁴Department of Clinical and Administrative Science, College of Pharmacy, Larkin University, Miami, Florida, USA. Now at Department of Clinical and Administrative Pharmacy, College of Pharmacy, University of Georgia, Athens, Georgia, USA.

⁵Department of Geriatric Medicine, Nova Southeastern University, Fort Lauderdale, Florida, USA

⁵*Instructional Technology and Distance Education Program, Fischler College of Education,
Nova Southeastern University, Fort Lauderdale Florida, USA*

A mobile app providing individually tailored psychoeducation about sleep for older adults with chronic health conditions and low health literacy

Abstract

Objective: This paper reports the effect of a mobile app that provides tailored information about sleep to individuals aged 40 and older who have chronic health conditions and low health literacy.

Methods: The sleep module was a part of a multitopic app focused on chronic disease self-management. Participants were randomly assigned to receive sleep psychoeducation at reading levels equivalent to 3rd, 6th or 8th grade. The primary outcome measure was the Pittsburgh Sleep Quality Index (PSQI), which was completed at baseline, after the intervention, and again three months later. Outcomes were assessed using repeated measures mixed effects models.

Results: Most participants were Black, Indigenous, or Other Persons of Color (BIPOC; 87%); they had average reading level at the 7th grade. Health literacy, socioeconomic status, number of health conditions, and the intervention's effect were related to the PSQI. The PSQI score significantly decreased over the course of the three study visits for all groups, consistent with a small to medium effect size ($d = 0.40$). No effect of treatment group was observed.

Conclusion: Results suggest that a brief tailored information intervention may be beneficial for individuals aged 40 and older who have low health literacy and chronic health conditions. Further development of the intervention may enhance its clinical effectiveness.

Registered at clinicaltrials.gov NCT02922439.

Introduction

Sleep problems are common among individuals in the United States and have a significant impact on their mood, quality of life, (National Sleep Foundation, 2023), workplace productivity (Filip et al., 2017), and cognitive functioning (Goel et al., 2009), and risk for chronic diseases (Liu et al., 2013). Despite available treatments, a significant number of individuals do not seek help for sleep disorders (Gordon et al., 2022; Torrens Darder et al., 2021), highlighting the need to prioritize the dissemination of information about the importance of sleep and available treatments (Di et al., 2022).

Among older adults, especially those with chronic medical conditions, sleep disturbances are common and have substantial clinical and public health implications. Onen (2018) found that 83% of older adults report at least one chronic medical condition, and these conditions are associated with higher rates of insomnia, sleep apnea, and other sleep disturbances. Additionally, Foley (2004) noted that conditions such as heart disease, depression, pain, and memory problems are linked to higher rates of insomnia, while obesity, diabetes, and lung disease are linked to sleep apnea, daytime sleepiness, and restless leg syndrome. Nadorff (2018) found that sleep disorders are strongly associated with anxiety, depression, dementia, and suicidal thoughts in older adults.

Sleep problems may be an even greater concern for older adults with chronic health conditions (Liu et al., 2013; Onen & Onen, 2018). The negative impact of sleep problems on daily functioning in older adults has also been reported (Gooneratne & Vitiello, 2014; Manocchia, San, & Ware, 2001), especially among persons from minoritized groups (Meredith et al., 2020). Sleep problems can have a detrimental impact on older adults' health-related quality of life, daily functioning, and health care utilization (Dai et al., 2019; Manocchia, Keller, & Ware, 2001). Insufficient or poor-quality sleep is associated with chronic health conditions and

worse health outcomes, particularly in minoritized and low-income populations (Jackson et al., 2015; Laposky et al., 2016; Williams et al., 2015). Individuals from minoritized racial and ethnic groups and those with lower socioeconomic status (SES) tend to get less sleep and report poorer sleep quality (Jehan et al., 2018; Williams et al., 2015). Sleep problems may contribute to health disparities in conditions such as obesity, cardiovascular disease, and diabetes (Jackson et al., 2015; Laposky et al., 2016; Williams et al., 2015) and various factors, including SES, racism, discrimination, neighborhood segregation, and access to healthcare, can influence sleep health disparities (Billings et al., 2021).

Health literacy, defined as the ability to understand and use health information, plays a crucial role in shaping health outcomes and disparities. Low health literacy has been linked to sleep disturbances (Ghiassi & Partridge, 2011; Hackney et al., 2008) and may be a factor in health disparities (Jackson et al., 2015), as is health literacy itself (Paasche-Orlow & Wolf, 2007, 2010). Persons with lower health literacy tend to be older, less educated, have lower income and report poorer sleep and overall health (Cutilli et al., 2018; Hackney et al., 2008). They may encounter difficulties in understanding and managing sleep disorders or implementing recommendations to improve their sleep health (Hackney et al., 2008). Several studies have demonstrated a link between inadequate health literacy and poor sleep health knowledge or outcomes. Hackney (2008) reviewed research showing patients with low health literacy understand less about sleep disorders and engage in poorer sleep hygiene. Gazmararian (2003) found that patients with chronic diseases and low health literacy have less knowledge about managing their conditions. Morrow (2006) found that adults with chronic heart failure and lower health literacy tended to have worse sleep, cognition, and more health problems, suggesting that

health literacy interventions should consider patients' sleep and cognitive abilities. A complex group of interactions between sleep, chronic disease, and health literacy may thus exist.

Treatment options for sleep disorders in older adults encompass a range of approaches, both pharmacological and non-pharmacological (Bloom et al., 2009). Non-drug therapies particularly cognitive behavioral therapy can be highly effective and have long-lasting benefits (Gooneratne & Vitiello, 2014). Medications may also be used but can cause side effects such as impaired thinking or movement (Gooneratne & Vitiello, 2014). Given the bidirectional relationship between sleep and health in older adults, accurate diagnosis and proper treatment of sleep disorders are crucial.

Evidence suggests that older adults often rely on pharmacologic treatments such as over-the-counter sleep aids or prescription sleep medications (Abraham et al., 2017; Musich et al., 2018), highlighting the need for accessible alternatives such as digital interventions. Although behavioral treatments have demonstrated effectiveness, access to them is limited by the number of trained providers and their distribution (Thomas et al., 2016). One strategy to address these problems is the adaptation of behavioral sleep medicine techniques to digital interventions making them more widely available. Several apps, for example, for cognitive behavioral therapy for insomnia (CBT-I), have shown promise in helping people improve their sleep (Espie et al., 2016; Kuhn et al., 2016; Yu et al., 2018).

Digital therapeutics and mobile health interventions show great potential for expanding healthcare access and reducing health disparities. Rivers (2014) argues that mobile technology and mHealth interventions can play a vital role in addressing health disparities by providing accessible health information and care to underserved groups. Similarly, Gibbons (2011) proposes that Web 2.0 technologies such as social media could be harnessed to provide health

promotion and education to minoritized and medically underserved groups. Several authors have emphasized the need to address sleep disparities and proposed actionable strategies (Billings et al., 2021; Hughes et al., 2022; Jackson et al., 2020; Miller et al., 2023). By applying health equity frameworks, engaging communities, and targeting social determinants of health, digital therapeutics can fulfil their potential to reduce disparities and empower marginalized groups.

This paper presents the results of a study of a multitopic mobile app for chronic disease self-management (CDSM). The app included a module providing individually-tailored psychoeducation on sleep to older adults with chronic health conditions and low health literacy. Participants' response to the app and the relations of sleep to their SES, health literacy, and chronic health conditions, as well as the effects of the intervention, were evaluated.

Method

The app's content was initially developed by examining existing resources related to CDSM and conducting a qualitative study to identify the information needs of older adults with chronic health conditions (Jacobs et al., 2017; Thomas-Purcell et al., 2019). Input from potential users was crucial in shaping the content and format of the intervention. To ensure evidence-based content, a diverse team was assembled that comprised professionals from relevant disciplines including medicine, nursing, psychology, pharmacy, public health, and education.

The app consisted of thematic sections, each presenting content on multiple screens. Guided by cognitive load theory-based instructional design (Paas et al., 2004), each module included an introduction, an assessment of participants' current status, general health-related insights, personalized content, and a summary. Participants could test their understanding through self-check questions. Information was presented through text complemented by images,

diagrams, and narrated animations in line with the principles of multimedia learning (Mayer, 2009).

More detailed information on the app's initial development and testing can be found in a recently-published paper (Patel et al., 2023). The modules were devised with identical content but different reading complexity levels, categorized based on the Fry and Flesch Reading Ease scores (3rd grade, with narration; 6th and 8th grade). This was facilitated through the use of Health Literacy Advisor® software integrated with Microsoft Word®.

The study took place at two locations: Fort Lauderdale, Florida and Atlanta, Georgia. Participants were recruited through various channels, such as past unrelated studies, local healthcare clinics, medical practices, and word of mouth. In Atlanta, collaboration with local churches helped identify and inform potential participants about the study. Data collection followed U.S. National Institutes of Health guidelines, including recording participant information including race, ethnicity, and gender.

Screening began with a brief telephone interview to determine eligibility based on medical history, medication use, and educational background. The Rapid Estimate of Adult Literacy in Medicine was used with a predefined cutoff for health literacy set at or below an 8th-grade level (Han et al., 2017, October). Eligibility criteria included being 40 years or older, having at least one ongoing chronic health condition under treatment, having an education below college graduate, and a score below the cutoff on the health literacy assessment. The college education criterion stemmed from prior research showing that successful completion of a college degree correlated with adequate health literacy (Ownby et al., 2019).

Participant assessments were completed at baseline, post-intervention, and at a follow-up three months later. The FLIGHT/VIDAS health literacy scale (Ownby, Acevedo, Jacobs, et al.,

2014; Ownby et al., 2013; Ownby, Acevedo, Waldrop-Valverde, et al., 2014; Ownby et al., 2021) was chosen due to its comprehensive score range, unlike to other measures with ceiling effects (Pleasant, 2014). Participants' socioeconomic status was evaluated using an index created through a principal components analysis of self-reported education and income; component scores were used for data analysis.

Participants' health status was assessed through a structured health conditions interview based on the Functional Comorbidity Scale (Groll et al., 2005) but expanded to include additional health conditions common in older adults (Centers for Medicare and Medicaid Services, 2012). Self-report measures were administered via audio computer-assisted self-interview software (Bethesda MD: Questionnaire Development System) that read all questions aloud to participants to minimize the effect of reading ability on their responses.

The sleep module

Table 1 provides an outline of the topics covered in the module. After viewing a section on the purpose of the module, participants completed the Insomnia Severity Index or ISI (Morin et al., 2011). Responses to the ISI were used to individually tailor content presented later in the module. For example, participants who reported difficulty in getting to sleep received suggestions to address it. Next, participants received feedback on how depression and stress could potentially impact their sleep based on their responses to the CES-D and PSS. The module then proceeded to discuss sleep stages, the two-process theory (Borbély, 1982), and how chronic health conditions can affect sleep. It continued with a review of sleep hygiene, approaches for keeping a sleep log, and a discussion of sleep restriction.

Insert Table 1 about here

Statistical analyses

Statistical analyses assessed the impact of SES, health literacy, number of health conditions, the intervention and treatment group assignment. Data processing and analysis were performed using SPSS and R (R Core Team, 2022). The intended sample size was determined using the mixed effects model simulation routine in PASS (NCSS, 2018), ensuring a minimum of 30 participants per group for adequate statistical power.

The effects of covariates and the intervention were evaluated using a series of random intercept mixed effects models that progressively increased in complexity (Luke, 2017). These were implemented with the R package *lme4* (Bates et al., 2015). The order in which covariates were included in the models was determined based on their anticipated impact on outcomes and potential for intervention. The baseline model (model 1 in Table 3) included only age, race, and gender (model 1). Subsequently, study site, SES, health literacy, health conditions, intervention effect over time, and treatment group were added. After computing model 1, likelihood ratio tests were conducted to compare the previous model to one with an additional covariate in order to assess its significance. To facilitate interpretation of results, chi-square values obtained from model comparisons were converted to Cohen's *d* (J. Cohen, 1988) using the R package *esc* (Ludecke, 2019).

Human subjects approval.

All procedures were approved by the Institutional Review Boards of Nova Southeastern University (2018-685-NSU) and of Emory University (MODCR001-IRB00087112). Participants

provided verbal consent for initial screening procedures and written informed consent for all other aspects of their participation.

Results

A total of 286 individuals completed the baseline assessments and the sleep module. Of these individuals, 195 had ISI scores of 8 or greater, suggesting insomnia according to the criterion reported by Morin et al. (2011). This report focuses on the data from these participants. Data from these participants are reported here. Of the 195 participants, 91 (46.7%) were men and 104 (53.3%) were women. There were 168 participants who identified as Black, Indigenous, or Other People of Color (BIPOC) accounting for 86% of the sample, while 27 (14.0%) identified as white individuals. The majority (61%) reported annual incomes of less than \$10,000 (US). Following the interpretation suggested by Morin et al. (2011), 87 (45%) had subthreshold, 79 (41%) had moderate, and 29 (15%) had severe insomnia. Complete information on recruitment, inclusion, and randomization, including a CONSORT diagram detailing patient flow through the study, is available online as a preprint (Ownby et al., 2023).

Descriptive statistics for continuous variables are presented in Table 2. The average level of health literacy on the FLIGHT/VIDAS measure was 10.02, indicating a basic level of health literacy that would enable a person to manage simple tasks such as reading straightforward prescription labels (Ownby et al., 2019). However, persons at this level of health literacy may struggle with understanding more complex directions or text that requires inferences. Their average score on the PSS was at a level above the population averages previously reported (S. Cohen & Janicki-Deverts, 2012), and their average score on the CES-D was greater than typical cutoff scores for major depression (Vilagut et al., 2016). Average scores on the PSQI and ISI

were consistent with a substantial sleep disturbance. Participants spent an average of 32 minutes completing the sleep module (SD = 13.7).

Insert Table 2 about here

Chronic health conditions and sleep disturbance

We evaluated which health conditions might be related to sleep disturbance by assessing the presence or absence of a condition related to sleep disturbance defined as having a PSQI score greater than 5 (Buysse et al., 2008) in multiple chi-square analyses uncorrected for multiple comparisons. The relation of sleep disturbance to chronic obstructive pulmonary disease approached statistical significance ($\chi^2 [1] = 3.53, p = 0.06$) as did that for angina ($\chi^2 [1] = 3.57, p = 0.06$). It was related to a diagnosis of hypercholesterolemia ($\chi^2 [1] = 5.10, p = 0.02$), depression ($\chi^2 [1] = 10.60, p = 0.001$), other nervous condition ($\chi^2 [1] = 13.85, p = < 0.001$), and bipolar disorder ($\chi^2 [1] = 6.77, p = 0.009$).

Models for the PSQI

Results of the likelihood ratio tests for sequential assessments of progressively more complex models for the PSQI are presented in Table 3. These models represent, in order, baseline (including age, gender, and race as covariates), site (Fort Lauderdale vs Atlanta), SES (an index of education and income), health literacy, total health conditions, time (baseline, immediate follow-up, and three-month follow-up), and treatment group (3rd, 6th, or 8th grade reading level). It can be seen that multiple factors are significantly associated with sleep quality over time, including SES, health literacy, and total health conditions. The effect of the intervention (assessed change in PSQI over time) was statistically significant and consistent with

a small to medium effect size ($d = 0.40$). Figure 1 displays a plot of the model-estimated means for each group during the three study visits. This model-corrected change illustrated in the figure represents a decline in PSQI of 0.85 points (95%CI [0.11, 1.60]).

Insert Table 3 about here

Insert Figure 1 about here

Discussion

The purpose of this paper was to report on the effects of a brief tailored information intervention for sleep psychoeducation. The intervention specifically targeted individuals aged 40 years and older with chronic health conditions and low health literacy. The sleep module was part of a larger intervention for chronic disease self-management which was delivered in three sessions over two to three weeks. Results of tests of the primary hypotheses of the main study are currently under review and available online as a preprint (Ownby et al., 2023). In addition, we assessed the impact of individual topic modules on relevant measures, and in this instance, we evaluated the modules' impact on sleep using the PSQI. We found significant decreases in participants' PSQI scores over the course of the study. While the magnitude of the change was relatively small, it may still be important considering the brevity of the intervention and the limited resources available to individuals with limited economic resources and low health literacy.

Several studies have examined the impact of CBT-I interventions on sleep in older adults. One study (Riedel et al., 1995) found that watching a 15-minute educational video had a significant positive effect on sleep satisfaction compared to a control condition. Participants in this study were older adults with insomnia with an average age of 67.4 years. Some participants in this study were excluded due to chronic health conditions and psychotropic medication use. In another study Lichstein et al. (2001), individuals aged 60 years or older participated in multisession training in either relaxation or sleep compression strategies. This study showed positive effects on sleep satisfaction, with moderate effect sizes. Participants with high levels of caffeine, nicotine, and alcohol use as well as those with elevated levels of anxiety and depression, were excluded from this study.

Buysse et al. (2011), on the other hand, included individuals who might be considered as having comorbid insomnia related to medical or psychiatric disorders, groups previously excluded in some studies. They found that a brief behavioral intervention delivered in two sessions with two follow-up phone calls had a significant effect compared to an information-only control group. The effect size for this intervention was large ($d = 1.10$). It should be noted that the present study's exclusion criteria were different from those of these studies. Some studies excluded individuals with common comorbidities and substance or medication use, while in our study these individuals participated. Participants' sleep in our study was less rigorously evaluated than in these studies, but at the same time may have been more representative of individuals who might not have participated in those studies.

Other researchers have explored the effectiveness of brief interventions for sleep problems. Reid et al. did not find an effect on the PSQI of one session of sleep hygiene education (2010), but Tucker et al. (2021) reported a large effect size ($d = 0.77$) for an online six-session

educational intervention. In a study that examined the optimal number of sessions in interventions for insomnia, Edinger et al. (Edinger et al., 2007) found that one session was effective in the short term but did not produce lasting effects. Okun and Glidewell (2023) showed that a single four-hour group class was effective in reducing insomnia symptoms in a group whose age averaged 54 years and who had multiple health conditions. These studies suggest that brief interventions can have positive effects on sleep, but longer interventions may have greater and longer-lasting impacts. Despite the smaller effect size observed in our study compared to others, the intervention's online availability and relatively brief nature may make it a useful initial element of a stepped-care approach (Espie, 2009) for sleep problems. Individuals who do not see improvement after the brief intervention can be referred for more intensive treatment.

The limitations of the present analyses should be acknowledged. The primary focus of the study was chronic disease self-management, with the sleep intervention being just one aspect of it. Participants also completed tailored information modules for mood disturbance, pain, and stress which may have influenced their perception of sleep. Additionally, it is important to note that this report relies on a single self-report measure of sleep, leaving unanswered the question of how the intervention might have affected objective sleep measures. Further, our participants were primarily individuals with low incomes and minoritized groups, limiting the generalizability of these findings to others. Finally, although the intervention did have a statistically significant effect on the PSQI, the change represented a small to medium effect size. It is likely that a more intensive intervention could yield more substantial and clinically meaningful improvements in sleep.

In summary, this paper aimed to present the sleep-specific outcomes of an intervention targeting a range of chronic disease-related issues in individuals aged 40 and older with chronic health conditions and low health literacy. Results suggest a modest improvement in participants' sleep over the course of the study and follow-up assessments. These findings suggest that a brief intervention offering personalized sleep information and guidance at appropriate reading levels may be helpful in improving sleep. Further development of the intervention appears warranted.

3385 words

Disclosure statement

Dr. Ownby is an applicant on a US patent application (US 2021/0065908) focused on automated assessment of patient understanding of health information. Dr. Ownby is a stockholder in Enalan Communications, Inc., a company that develops digital therapeutics.

Funding

This study was funded by grants from the US National Heart, Lung, and Blood Institute (R56HL096578) and the US National Institute on Minority Health and Health Disparities (R01 MD010368) to Dr. Ownby.

Data availability

Data from this study are available on request from the first author.

ORCID

Raymond L. Ownby: <https://orcid.org/0000-0001-9433-2820>

Kamilah Thomas-Purcell: <https://orcid.org/0000-0002-7523-128X>

Donrie Purcell: <https://orcid.org/0000-0002-4710-9412>

Joshua Caballero: <https://orcid.org/0000-0003-4406-2425>

Michael Simonson: <https://orcid.org/0000-0003-3478-7879>

References

- Abraham, O., Pu, J., Schleiden, L. J., & Albert, S. M. (2017). Factors contributing to poor satisfaction with sleep and healthcare seeking behavior in older adults. *Sleep Health*, 3(1), 43-48. <https://doi.org/10.1016/j.sleh.2016.11.004>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1:48. <https://doi.org/10.18637/jss.v067.i01>
- Billings, M. E., Cohen, R. T., Baldwin, C. M., Johnson, D. A., Palen, B. N., Parthasarathy, S., Patel, S. R., Russell, M., Tapia, I. E., Williamson, A. A., & Sharma, S. (2021). Disparities in sleep health and potential intervention models: A focused review. *Chest*, 159(3), 1232-1240. <https://doi.org/10.1016/j.chest.2020.09.249>
- Bloom, H. G., Ahmed, I., Alessi, C. A., Ancoli-Israel, S., Buysse, D. J., Kryger, M. H., Phillips, B. A., Thorpy, M. J., Vitiello, M. V., & Zee, P. C. (2009). Evidence-based recommendations for the assessment and management of sleep disorders in older persons. *Journal of the American Geriatrics Society*, 57(5), 761-789-761-789.
- Borbély, A. A. (1982). A two process model of sleep regulation. *Hum Neurobiol*, 1(3), 195-204.
- Buysse, D. J., Germain, A., Moul, D. E., Franzen, P. L., Brar, L. K., Fletcher, M. E., Begley, A., Houck, P. R., Mazumdar, S., Reynolds, C. F., 3rd, & Monk, T. H. (2011). Efficacy of brief behavioral treatment for chronic insomnia in older adults. *Arch Intern Med*, 171(10), 887-895. <https://doi.org/10.1001/archinternmed.2010.535>
- Buysse, D. J., Hall, M. L., Strollo, P. J., Kamarck, T. W., Owens, J., Lee, L., Reis, S. E., & Matthews, K. A. (2008). Relationships between the Pittsburgh Sleep Quality Index (PSQI), Epworth Sleepiness Scale (ESS), and clinical/polysomnographic measures in a community sample. *J Clin Sleep Med*, 4(6), 563-571.
- Centers for Medicare and Medicaid Services. (2012). *Chronic conditions among Medicare beneficiaries, chartbook (2012 ed.)*. Centers for Medicare and Medicaid Services.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences (2nd ed.)*. Routledge.
- Cohen, S., & Janicki-Deverts, D. (2012). Who's stressed? Distributions of psychological stress in the United States in probability samples from 1983, 2006, and 2009. *Journal of Applied Social Psychology*, 42(6), 1320-1334. <https://doi.org/doi:10.1111/j.1559-1816.2012.00900.x>
- Cutilli, C. C., Simko, L. C., Colbert, A. M., & Bennett, I. M. (2018). Health literacy, health disparities, and sources of health information in U.S. older adults. *Orthopaedic Nursing*, 37(1), 54-65-54-65.

- Dai, H., Mei, Z., An, A., & Wu, J. (2019). Association between sleep problems and health-related quality of life in Canadian adults with chronic diseases. *Sleep Med*, *61*, 26-30. <https://doi.org/10.1016/j.sleep.2019.04.015>
- Di, H., Guo, Y., Daghlas, I., Wang, L., Liu, G., Pan, A., Liu, L., & Shan, Z. (2022). Evaluation of sleep habits and disturbances among us adults, 2017-2020. *JAMA Network Open*, *5*(11), e2240788-e2240788. <https://doi.org/10.1001/jamanetworkopen.2022.40788>
- Edinger, J. D., Wohlgemuth, W. K., Radtke, R. A., Coffman, C. J., & Carney, C. E. (2007). Dose-response effects of cognitive-behavioral insomnia therapy: a randomized clinical trial. *Sleep*, *30*(2), 203-212. <https://doi.org/10.1093/sleep/30.2.203>
- Espie, C. A. (2009). “Stepped Care”: A health technology solution for delivering cognitive behavioral therapy as a first line insomnia treatment. *Sleep*, *32*(12), 1549-1558. <https://doi.org/10.1093/sleep/32.12.1549>
- Espie, C. A., Luik, A. I., Cape, J., Drake, C. L., Sriwardena, A. N., Ong, J. C., Gordon, C., Bostock, S., Hames, P., Nisbet, M., Sheaves, B., G Foster, R., Freeman, D., Costa-Font, J., Emsley, R., & Kyle, S. D. (2016). Digital Cognitive Behavioural Therapy for Insomnia versus sleep hygiene education: The impact of improved sleep on functional health, quality of life and psychological well-being. Study protocol for a randomised controlled trial. *Trials*, *17*(1).
- Filip, I., Tidman, M., Saheba, N., Bennett, H., Wick, B., Rouse, N., Patriche, D., & Radfar, A. (2017). Public health burden of sleep disorders: Underreported problem. *Journal of Public Health*, *25*(3), 243-248. <https://doi.org/10.1007/s10389-016-0781-0>
- Foley, D., Ancoli-Israel, S., Britz, P., & Walsh, J. (2004). Sleep disturbances and chronic disease in older adults. *Journal of Psychosomatic Research*, *56*(5), 497–502. <https://doi.org/10.1016/j.jpsychores.2004.02.010>
- Gazmararian, J. A., Williams, M. V., Peel, J., & Baker, D. W. (2003). Health literacy and knowledge of chronic disease. *Patient Education and Counseling*, *51*(3), 267–275.
- Ghiassi, R., & Partridge, M. R. (2011). Health literacy and sleep apnoea. *Thorax*, *66*(2), 180. <https://doi.org/10.1136/thx.2010.149070>
- Gibbons, M. C., Fleisher, L., Slamon, R. E., Bass, S., Kandadai, V., & Beck, J. R. (2011). Exploring the potential of web 2.0 to address health disparities. *Journal of Health Communication*, *16*(sup1), 77–89-77–89.
- Goel, N., Rao, H., Durmer, J. S., & Dinges, D. F. (2009). Neurocognitive consequences of sleep deprivation. *Semin Neurol*, *29*(4), 320-339. <https://doi.org/10.1055/s-0029-1237117>
- Gooneratne, N. S., & Vitiello, M. V. (2014). Sleep in older adults. *Clinics in Geriatric Medicine*, *30*(3), 591–627.

- Gordon, N. P., Yao, J. H., Brickner, L. A., & Lo, J. C. (2022). Prevalence of sleep-related problems and risks in a community-dwelling older adult population: a cross-sectional survey-based study. *BMC Public Health*, 22(1), 2045. <https://doi.org/10.1186/s12889-022-14443-8>
- Groll, D. L., To, T., Bombardier, C., & Wright, J. G. (2005). The development of a comorbidity index with physical function as the outcome. *Journal of clinical epidemiology*, 58(6), 595-602. <http://linkinghub.elsevier.com/retrieve/pii/S0895435605000211?showall=true> (Not in File)
- Hackney, J. E., Weaver, T. E., & Pack, A. I. (2008). Health literacy and sleep disorders: A review. *Sleep Medicine Reviews*, 12(2), 143–151-143–151.
- Han, H. J., Acevedo, A., Waldrop-Valverde, D., & Ownby, R. L. (2017, October, October, 2017). *Evaluation of short forms of the Rapid Estimate of Adult Literacy in Medicine (REALM)*. International Conference on Communication in Healthcare/Health Literacy Annual Research Conference, Baltimore MD.
- Hughes, A. J., Gunn, H., Siengsukon, C., Stearns, M. A., James, E., Donley, T., Grandner, M. A., Thomas, S. J., Hansen, K., & Williams, N. J. (2022). Eliminating sleep health disparities and achieving health equity: Seven areas for action in the behavioral sleep medicine community. *Behavioral Sleep Medicine*, 1-13. <https://doi.org/10.1080/15402002.2022.2149523>
- Jackson, C. L., Redline, S., & Emmons, K. M. (2015). Sleep as a potential fundamental contributor to disparities in cardiovascular health. *Annu Rev Public Health*, 36, 417-440. <https://doi.org/10.1146/annurev-publhealth-031914-122838>
- Jackson, C. L., Walker, J. R., Brown, M. K., Das, R., & Jones, N. L. (2020). A workshop report on the causes and consequences of sleep health disparities. *Sleep*, 43(8). <https://doi.org/10.1093/sleep/zsaa037>
- Jacobs, R. J., Ownby, R. L., Acevedo, A., & Waldrop-Valverde, D. (2017). A qualitative study examining health literacy and chronic illness self-management in Hispanic and non-Hispanic older adults. *J Multidiscip Healthc*, 10, 167-177. <https://doi.org/10.2147/JMDH.S135370>
- Jehan, S., Myers, A. K., Zizi, F., Pandi-Perumal, S. R., Jean-Louis, G., Singh, N., Ray, J., & McFarlane, S. I. (2018). Sleep health disparity: The putative role of race, ethnicity and socioeconomic status. *Sleep Med Disord*, 2(5), 127-133. <https://www.ncbi.nlm.nih.gov/pubmed/31179440>
- Kuhn, E., Weiss, B. J., Taylor, K. L., Hoffman, J. E., Ramsey, K. M., Manber, R., Gehrman, P., Crowley, J. J., Ruzek, J. I., & Trockel, M. (2016). CBT-I Coach: A description and clinician perceptions of a mobile app for cognitive behavioral therapy for insomnia. *Journal of clinical sleep medicine* 12(4), 597-606. <https://doi.org/10.5664/jcsm.5700>

- Laposky, A. D., Van Cauter, E., & Diez-Roux, A. V. (2016). Reducing health disparities: The role of sleep deficiency and sleep disorders. *Sleep Medicine, 18*, 3–6–3–6.
- Lichstein, K. L., Riedel, B. W., Wilson, N. M., Lester, K. W., & Aguillard, R. N. (2001). Relaxation and sleep compression for late-life insomnia: A placebo-controlled trial. *J Consult Clin Psychol, 69*(2), 227-239. <https://doi.org/10.1037//0022-006x.69.2.227>
- Liu, Y., Wheaton, A. G., Chapman, D. P., & Croft, J. B. (2013). Sleep duration and chronic diseases among U.S. adults age 45 years and older: evidence from the 2010 Behavioral Risk Factor Surveillance System. *Sleep, 36*(10), 1421-1427. <https://doi.org/10.5665/sleep.3028>
- Ludecke, D. (2019). esc: Effect size computation for meta analysis (version 0.5.1). <https://doi.org/10.5281/zenodo.1249218>
- Luke, S. G. (2017). Evaluating significance in linear mixed-effects models in R. *Behav Res Methods, 49*(4), 1494-1502. <https://doi.org/10.3758/s13428-016-0809-y>
- Manocchia, M., Keller, S., & Ware, J. E. (2001). Sleep problems, health-related quality of life, work functioning and health care utilization among the chronically ill. *Quality of Life Research, 10*(4), 331-345. <https://doi.org/10.1023/A:1012299519637>
- Manocchia, M., San, K., & Ware, J. E. (2001). Sleep problems, health-related quality of life, work functioning and health care utilization among the chronically ill. *Quality of Life Research, 10*(4), 331-345. <http://www.jstor.org/stable/4037547>
- Mayer, R. E. (2009). *Multimedia learning (2nd ed.)*. Cambridge.
- Meredith, S., Frawley, J., Sibbritt, D., & Adams, J. (2020). Risk factors for developing comorbid sleeping problems: Results of a survey of 1,925 women over 50 with a chronic health condition. *Journal of Aging and Health, 32*(5-6), 472-480. <https://doi.org/10.1177/0898264319832134>
- Miller, S. J., Sly, J. R., Alcaraz, K. I., Ashing, K., Christy, S. M., Gonzalez, B., Lu, Q., Newton, R. L., Redmond, M., Shen, M., Thomas-Purcell, K., Yi, J., Veinot, T., & Meade, C. D. (2023). Equity and behavioral digital health interventions: Strategies to improve benefit and reach. *Transl Behav Med, 13*(6), 400-405. <https://doi.org/10.1093/tbm/ibad010>
- Morin, C. M., Belleville, G., Bélanger, L., & Ivers, H. (2011). The Insomnia Severity Index: Psychometric indicators to detect insomnia cases and evaluate treatment response. *Sleep, 34*(5), 601-608. <https://doi.org/10.1093/sleep/34.5.601>
- Morrow, D., Clark, D., Tu, W., Wu, J., Weiner, M., Steinley, D., & Murray, M. D. (2006). Correlates of health literacy in patients with chronic heart failure. *Gerontologist, 46*(5), 669-676. <https://doi.org/10.1093/geront/46.5.669>
- Musich, S., Wang, S. S., Slindee, L. B., Saphire, L., & Wicker, E. (2018). Characteristics of new-onset and chronic sleep medication users among older adults: A retrospective study

- of a US Medigap plan population using propensity score matching. *Drugs & Aging*, 35(5), 467-476. <https://doi.org/10.1007/s40266-018-0543-5>
- Nadorff, M. R., Drapeau, C. W., & Pigeon, W. R. (2018). Psychiatric illness and sleep in older adults. *Sleep Medicine Clinics*, 13(1), 81–91–81–91.
- National Sleep Foundation. (2023). *National Sleep Foundation's 2023 Sleep in America Poll*. National Sleep Foundation. <https://www.thensf.org/wp-content/uploads/2023/03/NSF-2023-Sleep-in-America-Poll-Report.pdf>
- NCSS, L. (2018). *PASS 16 power analysis and sample size software*. NCSS, LLC. [ncss.com/software/pass](https://www.ncss.com/software/pass)
- Okun, M. L., & Glidewell, R. N. (2023). Improvement of insomnia symptoms following a single 4-hour CBT-I workshop. *Behav Sleep Med*, 21(1), 72-83. <https://doi.org/10.1080/15402002.2022.2036741>
- Onen, S.-H., & Onen, F. (2018). Chronic medical conditions and sleep in the older adult. *Sleep Medicine Clinics*, 13(1), 71–79.
- Ownby, R. L., Acevedo, A., Jacobs, R. J., Caballero, J., & Waldrop-Valverde, D. (2014). Quality of life, health status, and health service utilization related to a new measure of health literacy: FLIGHT/VIDAS. *Patient Educ Couns*, 96(3), 404-410. <https://doi.org/10.1016/j.pec.2014.05.005>. Available online May 14, 2014. (Not in File)
- Ownby, R. L., Acevedo, A., & Waldrop-Valverde, D. (2019). Enhancing the impact of mobile health literacy interventions to reduce health disparities. *Quarterly Review of Distance Education*, 10(1), 15-34.
- Ownby, R. L., Acevedo, A., Waldrop-Valverde, D., Jacobs, R. J., & Caballero, J. (2013). *A new computer-administered measure of health literacy: Validity and relation to quality of life in Spanish and English speakers. Paper presented at the International Conference on Communication in Healthcare, Montreal, October, .*
- Ownby, R. L., Acevedo, A., Waldrop-Valverde, D., Jacobs, R. J., & Caballero, J. (2014). Abilities, skills and knowledge in measures of health literacy. *Patient Educ Couns*, 95(2), 211-217. <https://doi.org/10.1016/j.pec.2014.02.002> (Not in File)
- Ownby, R. L., Acevedo, A., & Waldrop, D. (2021). Abilities, skills and knowledge related to measures of health literacy: A replication of the ASK model. *Ann Behav Med*, 55, S29.
- Ownby, R. L., Waldrop, D., Davenport, R., Simonson, M., Caballero, J., Thomas-Purcell, K., Purcell, D., Ayala, V., Gonzalez, J., Patel, N., & Kondwani, K. (2023). A mobile app for chronic disease self-management for individuals with low health literacy: A multisite randomized controlled clinical trial. *medRxiv*, 2023.2004.2001.23288020. <https://doi.org/10.1101/2023.04.01.23288020>

- Paas, F., Renkl, A., & Sweller, J. (2004). Cognitive Load Theory: Instructional implications of the interaction between information structures and cognitive architecture. *Instructional Science*, 32(1), 1-8. <http://dx.doi.org/10.1023/B:TRUC.0000021806.17516.d0> (Not in File)
- Paasche-Orlow, M. K., & Wolf, M. S. (2007). The causal pathways linking health literacy to health outcomes. *Am. J Health Behav*, 31 Suppl 1, S19-S26. <https://doi.org/10.5555/ajhb.2007.31.supp.S19> [doi] (Not in File)
- Paasche-Orlow, M. K., & Wolf, M. S. (2010). Promoting health literacy research to reduce health disparities. *J Health Commun*, 15 Suppl 2, 34-41. <https://doi.org/926961883> [pii];10.1080/10810730.2010.499994 [doi] (Not in File)
- Patel, N., Waldrop, D., & Ownby, R. L. (2023). Creating a tailored info app to promote self-management skills in persons with chronic health conditions: Development strategies and user experience [preprint] *Distance Learning*, 20(1), 9-18. <https://www.infoagepub.com/products/distance-learning-vol-20-1>
- Pleasant, A. (2014). Advancing health literacy measurement: a pathway to better health and health system performance. *J. Health Commun*, 19(12), 1481-1496. <https://doi.org/10.1080/10810730.2014.954083> [doi] (Not in File)
- R Core Team. (2022). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.r-project.org/about.html>
- Reid, K. J., Baron, K. G., Lu, B., Naylor, E., Wolfe, L. F., & Zee, P. C. (2010). Aerobic exercise improves self-reported sleep and quality of life in older adults with insomnia. *Sleep Medicine*. <https://doi.org/10.1016/j.sleep.2010.04.014>
- Riedel, B. W., Lichstein, K. L., & Dwyer, W. O. (1995). Sleep compression and sleep education for older insomniacs: Self-help versus therapist guidance. *Psychol Aging*, 10(1), 54-63. <https://doi.org/10.1037//0882-7974.10.1.54>
- Rivers, B. M., Bernhardt, J. M., Fleisher, L., & Green, B. L. (2014). Opportunities and challenges of using technology to address health disparities. *Future Oncology*, 10(4), 519-522-519-522.
- Thomas-Purcell, K. B., Jacobs, R. J., Seidman, T. L., Acevedo, A., Waldrop-Valverde, D., & Ownby, R. L. (2019). A mixed analysis approach to elucidate the multiple chronic condition experience of English- and Spanish-speaking older adults. *Clin Interv Aging*, 14, 407-418. <https://doi.org/10.2147/CIA.S193215>
- Thomas, A., Grandner, M., Nowakowski, S., Nesom, G., Corbitt, C., & Perlis, M. L. (2016). Where are the behavioral sleep medicine providers and where are they needed? A geographic assessment. *Behavioral Sleep Medicine*, 14(6), 687-698. <https://doi.org/10.1080/15402002.2016.1173551>

- Torrens Darder, I., Argüelles-Vázquez, R., Lorente-Montalvo, P., Torrens-Darder, M. D. M., & Esteva, M. (2021). Primary care is the frontline for help-seeking insomnia patients. *Eur J Gen Pract*, 27(1), 286-293. <https://doi.org/10.1080/13814788.2021.1960308>
- Tucker, R. M., Contreras, D., Carlson, B. R., Carter, A., & Drake, C. L. (2021). Sleep Education for Elders Program (SLEEP): Promising pilot results of a virtual, health educator-led, community-delivered sleep behavior change intervention. *Nature and Science of Sleep*. <https://doi.org/10.2147/nss.s304035>
- Vilagut, G., Forero, C. G., Barbaglia, G., & Alonso, J. (2016). Screening for depression in the general population with the Center for Epidemiologic Studies Depression (CES-D): A systematic review with meta-analysis. *PLOS ONE*, 11(5), e0155431. <https://doi.org/10.1371/journal.pone.0155431>
- Williams, N. J., Grandner, M. A., Snipes, S. A., Rogers, A., Williams, O., Airhihenbuwa, C., & Jean-Louis, G. (2015). Racial/ethnic disparities in sleep health and health care: Importance of the sociocultural context. *Sleep Health*, 1(1), 28–35-28–35.
- Yu, J. S., Kuhn, E., Miller, K. E., & Taylor, K. (2018). Smartphone apps for insomnia: Examining existing apps' usability and adherence to evidence-based principles for insomnia management. *Translational Behavioral Medicine*, 9(1), 110-119. <https://doi.org/10.1093/tbm/iby014>

Table 1

Content overview of chronic disease self-management sleep module

Topics
Purpose of the module
Association of chronic health conditions and sleep problems
Association of sleep disturbance with depressive symptoms and stress
Assessment of current sleep status via Insomnia Severity Index
Personalized feedback on participants' levels of depressive symptoms and stress from the CES-D and PSS
Sleep and the immune system
What is sleep? Understanding that it's a time when the brain is active and that good sleep is essential for functioning
Basic information about sleep, including sleep stages and importance of each, with narrated animations illustrating the stages and explaining movement from one stage to the next.
How chronic health conditions can affect sleep via factors such as pain and shortness of breath.
The two process model of sleep, linking the model to factors likely to improve or adversely affect sleep
Rules for better sleep: Sleep hygiene recommendations
How to use a sleep diary to see patterns related to medication and substance use, daily activity, and stress.
Feedback on responses to the Insomnia Severity Index with individualized recommendations to address problems reported.
Creating an action plan to improve sleep
Summary

Table 2

Descriptive statistics for continuous variables (n = 195)

	Minimum	Maximum	Mean	SD
Age	40.00	81.00	57.04	7.86
Education	6.00	15.00	11.80	1.81
Health	2.00	18.00	6.79	2.79
SES Index	-2.42	2.48	-0.15	0.85
WJ GE	0.90	18.00	6.99	4.16
FVA	2.00	18.00	10.04	3.95
CESD	5.00	55.00	25.06	8.70
PSS	12.00	37.00	22.80	3.51
PSQI	0.00	21.00	9.74	3.95
ISI	8.00	28.00	15.90	4.96

Note: BMI = body mass index; SES Index = index of socioeconomic status (see text); Health = Total chronic health diagnoses; WJ GE = Woodcock-Johnson Passage Comprehension (reading) Grade Equivalent Score; FVA = FLIGHT/VIDAS health literacy measure; CESD = Center for Epidemiological Studies—Depression Scale; PSS = Perceived Stress Scale, 10-item version; PSQI = Pittsburgh Sleep Quality Index; ISI = Insomnia Severity Index (note that only persons with ISI scores of 8 or greater were included in analyses).

Table 3

Likelihood ratio tests for progressively more complex models in random intercept analysis

	Parameters	Log Likelihood	χ^2	<i>df</i>	<i>p</i>	<i>d</i>	95% CI	Effect
model1	6	-1270						Base
model2	7	-1270	0.62	1	0.43	0.11	-0.17, 0.39	Site
model3	8	-1270	0.01	1	0.93	0.01	-0.27, 0.80	SES
model4	9	-1264	12.01	1	0.001	0.51	0.22, 0.80	HL
model5	10	-1260	6.84	1	0.01	0.38	0.10, 0.67	Health
model6	12	-1257	7.46	2	0.02	0.40	0.11, 0.69	Time
model7	14	-1256	0.56	2	0.76	0.11	-0.17, 0.39	Group

Note: For all models, the Pittsburgh Sleep Quality Index in the dependent variable, measured at three time points. Model 1 is base model that included only age, gender and race; Model 2 added the effect of site (Atlanta vs. Fort Lauderdale) to model 1; model 3 added socioeconomic status (SES) to model 3; model 4 added health literacy to model 3; model 5 added total number of health conditions; model 6 added the time (the effect of the intervention over study visits); and model 7 added the effect of treatment group (whether participants received the intervention at 3rd, 6th, of 8th grade reading difficulties).

Figure Caption

Figure 1. Pittsburgh Sleep Quality Index at Each Visit for Treatment Groups

Figure note: PSQI = Pittsburgh Sleep Quality Index; Evaluations = Study Visit (0 = Baseline; 1= First Follow-Up; 2 = Second Follow-Up; Colors represent model-corrected mean values for each treatment group (3rd, 6th, and 8th grade reading difficulty of intervention materials).

