

Designing multimedia patient education materials for adolescent idiopathic scoliosis: a protocol for a feasibility randomized controlled trial of patient education videos

Short title – Multimedia education materials for adolescent idiopathic scoliosis

Garett Van Oirschot^{1*}

Cailbhe Doherty^{1,2}

¹School of Public Health, Physiotherapy & Sport Science, University College Dublin, Dublin, Ireland.

²Insight SFI Research Centre for Data Analytics, University College Dublin, Ireland.

*Corresponding author e-mail: garett.vanoirschot@ucdconnect.ie

Both authors contributed to the research design, drafting the manuscript and approving the final version. Both authors agree to be accountable for all aspects of the work.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

This trial forms part of a PhD funded under the UCD Ad Astra program and received no additional funding.

The authors certify that they have no affiliations with or financial involvement in any organization or entity with a direct financial interest in the subject matter or materials discussed in the article.

This protocol was updated as version 7 on 27 November 2023

Abstract

Triple-masked three-armed feasibility parallel randomized controlled trial

Multimedia patient education materials are increasingly used in healthcare. While much research focuses on optimising their scientific content, research is equally needed to optimise design and implementation. This study aims to determine the feasibility of a study examining how the implementation of scientific advice on design affects patient outcomes. Participants aged 10-18 with radiographically confirmed adolescent idiopathic scoliosis will be recruited from community settings in Ireland and randomized into usual care or receiving multimedia educational videos with or without evidence-informed design principles. Participants will be masked in the two video intervention arms, as will the therapist sending the educational videos.

Outcomes will include the number of participants recruited and randomized, the number analysed post-intervention and at week eight, and the outcomes for baseline, post-intervention, and week 8. Adverse events will also be reported. This feasibility randomized controlled trial will offer insight into the feasibility of implementing advice from the literature in designing a trial of multimedia patient education materials for a population with adolescent idiopathic scoliosis.

Trial is registered on ClinicalTrials.gov as NCT06090344.

Keywords: *Adolescent Idiopathic Scoliosis, Health Education, Multimedia, Patient Education, Scoliosis*

Introduction

Patient education is a pillar of treatment that is recommended in many aspects of musculoskeletal (MSK) care, [1-10] including adolescent idiopathic scoliosis (AIS). [11, 12] Multimedia education is one option for conveying some of the information that healthcare providers may wish to pass on to a patient. They provide the accessibility, universality and usability that is desired in patient education aids. [13] While much healthcare research focuses on the content of educational materials, there are fewer interventional studies that examine the influence of design. Where such design research does exist, it is often in other medical areas [14-27] and the small amount of design recommendations in MSK healthcare are based mostly on literature arising narrative reviews, [28-30] non-randomized studies, cross-sectional surveys, or case series. [12] Regardless, there is potential for educational materials to fill a gap where healthcare services are known to be under strain, which can occur due to a lack of resources or rural/remote locations. [31]

Under resourcing is found in public healthcare systems for scoliosis, where waitlists can prevail for services [32-34], mirroring the dilemma of many musculoskeletal health services. Such services cannot rely on immediate one-to-one clinical encounters to provide education, whether for financial or geographical reasons. [35] Innovation has been required in healthcare to bring information and guidance to patients in these situations, with digital mediums providing such solutions. Scoliosis patients and their families have well-documented information needs [12, 36, 37] and multimedia education materials offer the advantage of dispersing healthcare information in a manner that is cheap, en masse, [31, 38, 39] and without physical proximity. [40, 41]

Rationale

Multimedia education has been examined extensively in the pedagogical literature, resulting in recommendations on how to best design materials in order to maximise engagement and learning [13, 42] While this information has informed the design of educational materials in healthcare [14-27] this has been less frequent in the MSK literature with only one RCT examining osteoarthritis-related pain, [43] and other MSK conditions being informed by narrative advice only. [28-30] The use of such principles [42] has not been tested in a randomized controlled trial of education for a MSK population such as adolescent idiopathic scoliosis.

Given the uncertainty surrounding the recruitment of this age bracket from the community and the unknown ability to engage with patient education materials over a trial period, a feasibility study is warranted to provide information on the design and implementation of a potential trial.

Aims and Objectives

The aim of the eventual three-armed parallel randomized controlled trial will be to compare the effectiveness of multimedia educational videos with and without evidence-informed design implementation against usual care for patients with adolescent idiopathic scoliosis. It will have an objective to examine if using videos with evidence-informed design results in superior post-treatment, 8 week, and 26 week outcomes for video engagement, knowledge translation and retention, and patient reported outcomes compared to videos formatted to match existing online resources [44] or treatment as usual for this population with AIS. We hypothesise that such an intervention will result in superior outcomes for video engagement, knowledge translation and retention, and for patient reported outcomes.

The more immediate aim of this feasibility RCT will be to assess the feasibility of running this trial with the objective of examining the recruiting, the intervention use, the intervention adherence, and the outcome assessment at baseline and eight weeks. It will be planned and evaluated using the RE-AIM framework [45] in order to optimise reach, effectiveness, adoption, implementation, and maintenance and determine the feasibility of this RCT and any aspects that cannot be pragmatically explored in sufficient depth will be explained to avoid shortcomings in reporting that have been previously observed [46-48].

Materials and methods

This triple-masked three-armed feasibility parallel randomized controlled trial will be conducted in Ireland and is prospectively registered on ClinicalTrials.gov. It was designed in consideration of the CONSORT guidelines for feasibility studies and the TIDieR [49] and RoB-2 [50] appraisal tools. The RE-AIM framework [45] will be used to ensure proper consideration of sustainable adoption and implementation. While it is not expected or intended, any changes to the trial protocol will be reported along with reasons for such changes. The trial is approved by University College Dublin Human Research Ethics Committee (LS-23-15-VanOirschot-Doherty, 3 May 2023).

Participants

The inclusion criteria will be as follows: aged 10-18 years, parent/guardian consent for those under 18 years of age, AIS as confirmed by Cobb angle ≥ 10 deg on plain film radiographs, able to watch and listen to online educational materials as

well as read and complete online surveys. Non-AIS scoliosis including but not limited to neurological conditions will be excluded.

Patient and public Involvement

Two individuals over age 18 who have been through the adolescent care pathway for AIS were consulted when designing the study protocol for feedback on the design and educational content to be used in the interventions. Information was also sought from a physiotherapist who treats scoliosis on a daily basis and collaborates with scoliosis orthopaedic surgeons in Ireland. Their opinion will also be sought about disseminating the results, which parts of the results should be shared, and in which digital/or print format.

Recruitment

Participants will be recruited in Ireland via digital and poster advertising through two private healthcare clinics serving an AIS population, two scoliosis advocacy groups based in Ireland, as well as through social media and word of mouth. The primary author, a PhD student and physiotherapist of 18 years' experience with a post-graduate musculoskeletal and sports masters degree, will conduct the preliminary screening and provide the participant information package to those who express interest. All participants will be screened by phone or by video call prior to inclusion, and all participants not meeting eligibility will have their exclusion reasons recorded but without any identifying information. Advertising and recruitment opened on 28 November 2023, with the study due to commence in January 2024.

Consent process

Informed consent will be sought from all potential participants meeting the inclusion criteria. The primary author will discuss the trial methodology and answer any questions or concerns about the trial before taking consent / assent electronically. Those aged 18 will be able to consent for themselves, while potential participants under the age of 18 will require parent/guardian consent before granting assent themselves. All participant rights will remain protected.

Fidelity

Treatment fidelity for the two video intervention groups will be monitored through YouTube analytics, which will show the lack of views for each participant with their video or with sections of video that are skipped over.

Baseline Assessment

Following informed consent, participants will electronically complete a series of baseline assessments of demographics and outcome measures, with the primary author being available by phone or electronically if assistance is required. A description of the baseline variables plus other study outcomes and their purpose is shown in Table 1.

Table 1. Baseline and outcome variables collected and their purpose, with the primary feasibility outcomes in **bold**

Variable	Measure	Time points (weeks)	Purpose & dimension of RE-AIM framework
Age	Date of birth	0	Describe population, reach
Gender	Male/female	0	Describe population, reach
Date of plain film radiographs confirming scoliosis	Weeks	0	Describe population, reach
Date of last radiograph	Weeks	0	Describe population, reach
Duration since care-seeking	Weeks	0	Describe population, reach
Major Cobb Angle	Degrees	0	Describe population
Minor Cobb Angle (if applicable)	Degrees	0	Describe population
Brace treatment	Yes/No	0	Describe population
Surgical treatment	Yes/No	0	Describe population
Physiotherapy treatment	Yes/No	0	Describe population
Quality of life	SRS-22	0, 8	Secondary outcome
	EQ-5D-Y	0, 8	Secondary outcome
Physical Activity	PAQ-C	0, 8	Secondary outcome
Anxiety	STAI-Ch	0, 8	Secondary outcome
Immediate Knowledge translation	5-item MCQ	1,2,3,4,5,6,8	Secondary outcome
Cognitive Engagement			
<i>One week Knowledge retention</i>	5-item MCQ	2,3,4,5,6,7	Secondary outcome, maintenance
<i>End of study Knowledge retention</i>	30-item MCQ	0, 8	Secondary outcome, maintenance
Affective engagement	YouTube Like/Dislike	1,2,3,4,5,6	Secondary outcome
Intervention adherence / Behavioral Engagement			

Watch Time	YouTube watch time %	1,2,3,4,5,6	Primary outcome, adoption, implementation
Viewings	YouTube # views	1,2,3,4,5,6	Primary outcome, adoption, implementation
Number recruited, rejected, randomized, dropped off, completed	Number	0,1,2,3,4,5,6,7,8	Primary outcome, implementation
Dropout reasons	Participant response	1,2,3,4,5,6,7,8	Primary outcome, implementation
Outcomes completed	%	8	Primary outcome, implementation
Adverse events	Number and type	1,2,3,4,5,6,8	Primary outcome, effectiveness
Satisfaction with care and treatment	Tailored question	8	Primary outcome, effectiveness

Abbreviations: SRS-22r, Scoliosis Research Society 22-revised; EQ-5D-Y, EuroQOL 5 Dimension Youth version; PAQ-C, Physical Activity Questionnaire for Children; STAI-Ch, State Trait Anxiety Index for Children; MCQ, Multiple Choice Questionnaire

Randomization & Masking

Upon consenting to participate in the study, participants will be randomized by a random number generator on a 1:1:1 ratio into one of the three intervention arms by a member of the research team who will be masked to the interventions and follow-up assessment of outcomes. Participants will be told that the trial will compare usual care with two other multimedia video interventions. Participants in the usual care group will not be masked to their allocation but the participants in the two video interventions will not be aware of their allocation to videos with traditional format or augmented format.

For the two video groups, an email will be sent by the supervising author, a registered Physiotherapist and researcher for fifteen years, with a link to the correct video but this author will be masked to which intervention the link pertains to. The primary author will record and analyze all outcomes that will be taken via the SurveyMonkey online platform but will remain masked to the treatment delivered when viewing these outcomes.

Interventions

In the usual care group, participants will continue along their existing healthcare pathway and will not receive any educational intervention beyond the usual education from their healthcare providers, which will not be determined by participation in the study and will not be funded by it.

In the video groups, participants will continue along their usual healthcare pathway as well, however they will also be sent a link to six multimedia videos, each three to four minutes long, using the information and format of a typical online resource, in this case the Frequently Asked Questions section of Scoliosis Research Society website [44] with some additional content from the qualitative literature on information needs in the scoliosis population. [11, 36, 37, 51] The outline of the content is shown in Table 2. Adjustments were also made to ensure spelling and grammar in Irish-version English language and some terms were altered to reflect Irish phrases i.e. consultant instead of specialist. The script was edited again following feedback from the individuals who formed the public and patient involvement described above, and this assisted in making minor corrections and updating statistics such as higher prevalence in girls vs boys, [52-55] as well as elaborating on discussion points about lifestyle factors and back pain. The video channel will be set up as a private channel unique to each participant, so that the

engagement statistics can be tracked for each participant. This will consist of six videos, sent once weekly from Week 1 to Week 6. Participants can skip forward, back and re-watch a released video an unlimited number of times until Week 8.

Table 2. Outline of educational content

Video	Questions to be answered in the educational content
1	What is scoliosis? Is scoliosis more common in boys or girls? What are the other types of scoliosis? Does everyone with scoliosis wear a brace or need surgery? Why do kids get scoliosis? Does bad posture lead to scoliosis? Are there exercises I can do to make my spine straight?
2	Can we screen for scoliosis? Can scoliosis curves get better on their own? How often will I need my scoliosis re-checked? Will the x-rays harm me?
3	What can I do to keep my scoliosis from getting worse? How do you decide what the treatment will be? What is involved in conservative? Can you tell me about bracing? How do you know if a patient needs surgery? Where can I find healthcare specialists who work with scoliosis? Should I try other conservative methods first?
4	Which kind of brace is right for me? What are the different types of braces? Why should I wear a brace? Does it matter how many hours a day I wear the brace? Can I take the brace off for PE at school, sports activities, swimming, etc.? How will I know if the brace is working? I wore a brace for two years and my curve was the same size when I stopped wearing it, why? Did I waste my time? What does a brace look like?
5	If I need scoliosis surgery, what can I expect? Will surgery only stop my curve from growing or will it help straighten it? How long will it take me to recover from surgery? Does that include participation in sports? Will I have my rods removed after my spine is fused? Can the rods break?
6	Should I avoid certain activities because of my scoliosis? Is carrying a heavy back-pack bad? What if I have to wear a brace? How do I cope with having scoliosis?

How can I connect with other kids who have scoliosis?
Will my curves get worse?
Can my scoliosis be cured?
If I have mild scoliosis, do I have to see a spine specialist?
Will I pass my scoliosis on to my future children?

Similarities between the two video intervention groups

The videos will contain identical information and it will be presented in the same sequence in both types of videos. The script will be identical except for specific components that are described below.

Differences between the two video intervention groups

For the video group with augmented (AUGM), the text, graphics, and audio formatted in accordance with recommendations from the Cognitive Theory of Multimedia Learning and other advice from the literature. [13, 28-30, 42]

Materials

The multimedia materials for the TRAD and AUGM video groups will be recorded using an iPhone13 to allow for content to be formatted to a phone screen format in case users view it on mobile devices and edited on Final Cut Pro X [56] with additional audio editing on Garage Band. [57] The final video will upload to a private YouTube channel setup by the primary author and the corresponding link will be sent via email. All participants will require internet access and a device to view and/or hear the educational materials. Participants in the two video intervention groups will be sent a link to a private YouTube channel, unique to each participant so that individual YouTube analytics can be recorded and analyzed. Additional patient-reported outcomes will be collected via SurveyMonkey.

Following completion of the study, the videos will be made available on a YouTube link for all participants. They will also be made public via an open link in the final manuscript of this study, to assist with any future replication and appraisal.

Outcomes

As recommended in the CONSORT guidelines for feasibility studies [58, 59], the feasibility outcomes for this trial will be the number of participants recruited, rejected, dropped out and reasons; adherence to videos via behavioural engagement; number of outcomes completed, and adverse events. Adverse events will be recorded with preparations made for any adverse effects from asking about the participants' mental health status or from being presented with unexpected information about scoliosis. The preparations are outlined Table 3.

Adherence to the RE-AIM framework will also ensure that there is an evaluation of the reach: recording of the percentage of excluded individuals, the percentage of individuals who participate, the characteristics of participants versus non-participants and the recording of qualitative indicators as to why participants were reached and/or recruited. The effectiveness will be measured using the knowledge outcomes and the overall quality of life outcomes in the SRS-22r and the EQ-5D-Y, a moderation analysis across the subgroups, and attrition analysis of the characteristics of these participants and their reasons. Adoption to specific settings is expected to be assessed easily by the online delivery of this intervention. The implementation will be evaluated by recording the number of interventions delivered perfectly, the adaptations made over the course of the study, the total budget, and the consistency of implementation across the eight weeks. Maintenance will be assessed by the measurement of all outcomes at the follow-up assessment in week eight, attrition rate, and analysis of the subgroups for each of these items.

Table 3. Protocol for adverse events

Potential event	Research member responsible	Action to be taken
Participant distress from answering mental health questions	Primary author will be available by phone or electronic means	Provide support and explain the rationale for the questions in the outcome measures and if needed, liaising with a psychologist with experience in treating an AIS population.
Participant distress from receiving unexpected information about AIS	Primary author will be available by phone or electronic means	Answer any AIS-related questions and provide reassurance where needed. Onward referral can be made back to any scoliosis-specialised health services that the participant is already attending, or to a healthcare practitioner known to the primary author who has experience with this population

The final fully-powered RCT, should it proceed based on results of this feasibility study, will assess a primary outcome of post-intervention cognitive engagement, meaning the knowledge translation for each video for weeks 1-6 and retention for all videos at Week 8. Secondary outcomes will be knowledge retention one week later for each of the six videos for weeks 2-7; affective & behavioural engagement from YouTube analytics determined by like/dislike, watch-time, number of views; health related quality of life measured by the SRS-22r and the EQ-5D-Y, anxiety measured by the STAI-Ch, and physical activity measured by the PAQ-C. Secondary outcomes will be measured at baseline and week 8.

The SPIRIT schedule of enrolment, interventions, and assessments is shown in Figure 1.

Figure 1. SPIRIT schedule of enrolment, interventions, and assessments.

Statistical analysis

Ratio and interval variables (YouTube analytics) will be analysed for differences by ANOVA. The remaining ordinal variables will have differences between groups analysed using Friedmans. Pre- and post-test changes will be analysed using Kruskal-Wallis. Comparisons will be made using intention-to-treat principles.

Generalised estimating equations (GEE) will be used to evaluate the relationship between the design characteristics and the measures of engagement. Each of the characteristics described above (Watch time / Total time (%), Retention (%)), Time spent on specific segments, Average view duration (minutes:seconds) will be included as co-variates in separate GEE models for each measure of engagement.

There was an audit of previous year's records to note that were approximately 60 eligible participants over twelve months in one of the private practice recruitment sites, and this will be in addition to social media and advocacy groups advertising (with two advocacy groups having 2542 and 579 followers in Ireland), as well as two other pending consultant clinics. Statistical power for the eventual RCT has been calculated using GPower [60] to detect a minimum change of 1 out of 5 in the primary outcome, knowledge quizzes, with a one way ANOVA to compare three groups with an alpha of 0.05 and determined that a sample size of 92 participants

were needed. This feasibility study will assess the recruitment potential using 10-15 participants per group.

Data will be stored and handled using password-protected institutional storage services, in accordance with institutional policy. The final data will be accessibly by the primary and supervising authors.

Results

This trial will be reported in conformance with the CONSORT guideline for feasibility randomized controlled trials. A CONSORT flow diagram will outline the number of participants in each stage of the study (Figure 2). All anonymised data will be publicly available as per Open Science recommendations on <https://osf.io/rcfsd/>. Results will be communicated to participants directly and through academic publication with no publishing restrictions.

Figure 2. CONSORT flow diagram

Discussion

Potential limitations and attempts to address

SRS-22r does not capture all aspects of physical functioning, such as mobility and self-care. [61] Additionally, SRS-22r was evaluated in the COS study that included all spinal deformities and thus limiting applicability to specific AIS [61] therefor additional quality of life measure will be taken via the EQ-5D-Y. It is acknowledged the EQ-5D-Y may require supplemental assessment tools for school function, [62] and that it's correlation with the SRS-22r has been found to be good for

non-scoliotic populations. [63] Proxy reporting may occur with the EQ-5D-Y as participants complete at home in the presence of parent(s)/guardian(s), but this yields similar outcomes as the 3-layer predecessor. [64]

Future potential

This study will determine the feasibility of implementing a fully powered trial and may offer an option to provide basic information about AIS to patients and the general public. Its utility could be particularly found in addressing those individuals on the waitlists for many services related to AIS. [11, 32, 33] This could have further implications for countries with under resourced healthcare systems or rural and remote regions lacking the access to clinicians that would often provide this specialty advice and education.

Conclusion

This feasibility randomized controlled trial will offer insight into the feasibility of implementing advice from the literature in designing multimedia patient education materials for a population with adolescent idiopathic scoliosis and allow a comparison with materials that do not undergo similar redesign as well as a comparison with usual care.

Acknowledgements

The authors wish to thank Scoliosis Awareness & Support Ireland, the Scoliosis Advocacy Network, McGowan Physio for their offer of assistance in recruiting for this study.

References

1. Bannuru RR, Osani MC, Vaysbrot EE, Arden NK, Bennell K, Bierma-Zeinstra SMA, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage*. 2019;27(11):1578-89. Epub 20190703. doi: 10.1016/j.joca.2019.06.011. PubMed PMID: 31278997.
2. Barrett E, Larkin L, Caulfield S, De Burca N, Flanagan A, Gilsenan C, et al. Physical Therapy Management of Nontraumatic Shoulder Problems Lacks High-Quality Clinical Practice Guidelines: A Systematic Review With Quality Assessment Using the AGREE II Checklist. *J Orthop Sports Phys Ther*. 2021;51(2):63-71. doi: 10.2519/jospt.2021.9397. PubMed PMID: WOS:000615835200006.
3. Collins NJ, Barton CJ, van Middelkoop M, Callaghan MJ, Rathleff MS, Vicenzino BT, et al. 2018 Consensus statement on exercise therapy and physical interventions (orthoses, taping and manual therapy) to treat patellofemoral pain: recommendations from the 5th International Patellofemoral Pain Research Retreat, Gold Coast, Australia, 2017. *Br J Sports Med*. 2018;52(18):1170-8. Epub 20180620. doi: 10.1136/bjsports-2018-099397. PubMed PMID: 29925502.
4. Coombes BK, Bisset L, Vicenzino B. Management of Lateral Elbow Tendinopathy: One Size Does Not Fit All. *J Orthop Sports Phys Ther*. 2015;45(11):938-49. doi: 10.2519/jospt.2015.5841. PubMed PMID: WOS:000364600900015.
5. Day JM, Lucado AM, Uhl TL. A Comprehensive Rehabilitation Program for Treating Lateral Elbow Tendinopathy. *Int J Sports Phys Th*. 2019;14(5):818-34. doi: 10.26603/ijsp20190818. PubMed PMID: WOS:000500486300014.
6. Doiron-Cadrin P, Lafrance S, Saulnier M, Cournoyer E, Roy JS, Dyer JO, et al. Shoulder Rotator Cuff Disorders: A Systematic Review of Clinical Practice Guidelines and Semantic Analyses of Recommendations. *Arch Phys Med Rehab*. 2020;101(7):1233-42. doi: 10.1016/j.apmr.2019.12.017. PubMed PMID: WOS:000543033600016.
7. Fernandes L, Storheim K, Sandvik L, Nordsletten L, Risberg MA. Efficacy of patient education and supervised exercise vs patient education alone in patients with hip osteoarthritis: a single blind randomized clinical trial. *Osteoarthritis Cartilage*. 2010;18(10):1237-43.
8. Morrissey D, Cotchett M, Said J'Bari A, Prior T, Griffiths IB, Rathleff MS, et al. Management of plantar heel pain: a best practice guide informed by a systematic review, expert clinical reasoning and patient values. *Br J Sports Med*. 2021;55(19):1106-18. Epub 20210330. doi: 10.1136/bjsports-2019-101970. PubMed PMID: 33785535; PubMed Central PMCID: PMC8458083.
9. Willy RW, Högglund LT, Barton CJ, Bolgla LA, Scalzitti DA, Logerstedt DS, et al. Patellofemoral Pain. *J Orthop Sports Phys Ther*. 2019;49(9):CPG1-CPG95. doi: 10.2519/jospt.2019.0302. PubMed PMID: 31475628.
10. Zhang W, Moskowitz R, Nuki G, Abramson S, Altman R, Arden N, et al. OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartilage*. 2008;16(2):137-62.
11. Negrini S, Donzelli S, Aulisa AG, Czaprowski D, Schreiber S, de Mauroy JC, et al. 2016 SOSORT guidelines: orthopaedic and rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis Spinal Disord*. 2018;13:3. Epub 20180110. doi: 10.1186/s13013-017-0145-8. PubMed PMID: 29435499; PubMed Central PMCID: PMC5795289.
12. Motyer G, Dooley B, Kiely P, Fitzgerald A. Parents' information needs, treatment concerns, and psychological well-being when their child is diagnosed with adolescent idiopathic scoliosis: A systematic review. *Patient Education and Counseling*. 2021;104(6):1347-55. doi: <https://doi.org/10.1016/j.pec.2020.11.023>.
13. Hoffman AS, Volk RJ, Saarimaki A, Stirling C, Li LC, Härter M, et al. Delivering patient decision aids on the Internet: definitions, theories, current evidence, and emerging

research areas. *BMC Medical Informatics and Decision Making*. 2013;13(2):S13. doi: 10.1186/1472-6947-13-S2-S13.

14. Abujarad F, Peduzzi P, Mun S, Carlson K, Edwards C, Dziura J, et al. Comparing a Multimedia Digital Informed Consent Tool With Traditional Paper-Based Methods: Randomized Controlled Trial. *JMIR Form Res*. 2021;5(10):e20458. Epub 20211019. doi: 10.2196/20458. PubMed PMID: 34665142; PubMed Central PMCID: PMC8564662.
15. Athilingam P, Osorio R, Kaplan H, Oliver D, O'Neachtain T, Rogal P. Embedding Patient Education in Mobile Platform for Patients With Heart Failure: Theory-Based Development and Beta Testing. *Comput Inform Nurs*. 2016;34(2):92-8. doi: 10.1097/cin.0000000000000216. PubMed PMID: 26765655.
16. Choi J. Development and pilot test of pictograph-enhanced breast health-care instructions for community-residing immigrant women. *Int J Nurs Pract*. 2012;18(4):373-8. doi: 10.1111/j.1440-172X.2012.02051.x. PubMed PMID: 22845637.
17. Dash S, Kamath U, Rao G, Prakash J, Mishra S. Audio-visual aid in teaching "fatty liver". *Biochem Mol Biol Educ*. 2016;44(3):241-5. Epub 20151202. doi: 10.1002/bmb.20935. PubMed PMID: 26625860.
18. Ferguson I, Phillips A, Lin M. Continuing Medical Education Speakers with High Evaluation Scores Use more Image-based Slides. *West J Emerg Med*. 2017;18(1):152-8. Epub 20161205. doi: 10.5811/westjem.2016.10.31484. PubMed PMID: 28116029; PubMed Central PMCID: PMC5226752.
19. Grech V. The application of the Mayer multimedia learning theory to medical PowerPoint slide show presentations. *J Vis Commun Med*. 2018;41(1):1745-3062 (Electronic). doi: 10.1080/17453054.2017.1408400.
20. Iorio-Morin C, Brisebois S, Becotte A, Mior F. Improving the pedagogical effectiveness of medical videos. *J Vis Commun Med*. 2017;40(3):96-100. Epub 20170919. doi: 10.1080/17453054.2017.1366826. PubMed PMID: 28925762.
21. Issa N, Mayer R, Schuller M, Wang E, Shapiro M, DaRosa D. Teaching for understanding in medical classrooms using multimedia design principles. *Med Educ*. 2013;47(4):388-96. doi: 10.1111/medu.12127. PubMed PMID: 23488758.
22. Issa N, Schuller M, Santacaterina S, Shapiro M, Wang E, Mayer R, et al. Applying multimedia design principles enhances learning in medical education. *Med Educ*. 2011;45(8):818-26. doi: 10.1111/j.1365-2923.2011.03988.x. PubMed PMID: 21752078.
23. Javaid M, Schellekens H, Cryan J, Toulouse A. eNEUROANAT-CF: a Conceptual Instructional Design Framework for Neuroanatomy e-Learning Tools. *Med Sci Educ*. 2021;31(2):777-85. Epub 20210122. doi: 10.1007/s40670-020-01149-y. PubMed PMID: 34457925; PubMed Central PMCID: PMC8368470.
24. Kayler L, Dolph B, Seibert R, Keller M, Cadzow R, Feeley T. Development of the living donation and kidney transplantation information made easy (KidneyTIME) educational animations. *Clin Transplant*. 2020;34(4):e13830. Epub 20200309. doi: 10.1111/ctr.13830. PubMed PMID: 32072670.
25. Nagmoti J. Departing from PowerPoint default mode: Applying Mayer's multimedia principles for enhanced learning of parasitology. *Indian J Med Microbiol*. 2017;35(2):199-203. doi: 10.4103/ijmm.IJMM_16_251. PubMed PMID: 28681806.
26. Sait S, Tombs M. Teaching Medical Students How to Interpret Chest X-Rays: The Design and Development of an e-Learning Resource. *Adv Med Educ Pract*. 2021;12:123-32. Epub 20210205. doi: 10.2147/amep.S280941. PubMed PMID: 33574725; PubMed Central PMCID: PMC8787294.
27. Wyatt T, Krauskopf P, Gaylord N, Ward A, Huffstutler-Hawkins S, Goodwin L. Cooperative m-learning with nurse practitioner students. *Nurs Educ Perspect*. 2010;31(2):109-13. PubMed PMID: 20455369.
28. Hernandez-Sanchez S, Moreno-Perez V, Garcia-Campos J, Marco-Lledo J, Navarrete-Munoz EM, Lozano-Quijada C. Twelve tips to make successful medical infographics. *Med Teach*. 2021;43(12):1353-9. doi: 10.1080/0142159x.2020.1855323. PubMed PMID: WOS:000600444700001.

29. Scott H, Fawcner S, Oliver C, Murray A. Why healthcare professionals should know a little about infographics. *Br J Sports Med*. 2016;50(18):1104-+. doi: 10.1136/bjsports-2016-096133. PubMed PMID: WOS:000383467400005.
30. Scott H, Fawcner S, Oliver CW, Murray A. How to make an engaging infographic? *Br J Sports Med*. 2017;51(16):1183. doi: 10.1136/bjsports-2016-097023.
31. Sharma S, Blyth FM, Mishra SR, Briggs AM. Health system strengthening is needed to respond to the burden of pain in low- and middle-income countries and to support healthy ageing. *J Glob Health*. 2019;9(2):020317-. doi: 10.7189/jogh.09.020317. PubMed PMID: 33274068.
32. British Scoliosis Society, British Association of Spine Surgeons. Statement from the British Scoliosis Society (BSS) and the British Association of Spine Surgeons (BASS) regarding wait times for adult and paediatric elective spine patients UK: British Scoliosis Society and British Association of Spine Surgeons; 2021 [cited 2022 14 November]. Available from: <https://spinesurgeons.ac.uk/News/10283819>.
33. Children's Hospital Group. Scoliosis Co-Design 10 Point Action Plan 2018/2019. In: Group CsH, editor. Ireland2018.
34. Dabke HV, Jones A, Ahuja S, Howes J, Davies PR, editors. SHOULD PATIENTS WAIT FOR SCOLIOSIS SURGERY? *Orthopaedic Proceedings*; 2006: The Bone & Joint Journal.
35. Sharma S, Traeger AC, Mishra SR, Sharma S, Maher CG. Delivering the right care to people with low back pain in low- and middle-income countries: the case of Nepal. *J Glob Health*. 2019;9(1):010304. doi: 10.7189/jogh.09.010304. PubMed PMID: 30774940; PubMed Central PMCID: PMC6367652.
36. van Schaik P, Flynn D, van Wersch A, Ryan KA, Ferguson V. Factors important in the design of information material for scoliosis. *Int J Rehabil Res*. 2007;30(2):163-5. doi: 10.1097/MRR.0b013e32813a2ec6. PubMed PMID: 17473629.
37. Wellburn S, van Schaik P, Bettany-Saltikov J. The Information Needs of Adolescent Idiopathic Scoliosis Patients and Their Parents in the UK: An Online Survey. *Healthcare (Basel)*. 2019;7(2). Epub 20190620. doi: 10.3390/healthcare7020078. PubMed PMID: 31226769; PubMed Central PMCID: PMC6627603.
38. Hauser-Ulrich S, Kunzli H, Meier-Peterhans D, Kowatsch T. A Smartphone-Based Health Care Chatbot to Promote Self-Management of Chronic Pain (SELMA): Pilot Randomized Controlled Trial. *JMIR Mhealth Uhealth*. 2020;8(4):e15806. Epub 20200403. doi: 10.2196/15806. PubMed PMID: 32242820; PubMed Central PMCID: PMC67165314.
39. O'Keeffe M, O'Sullivan P, Purtill H, Bargary N, O'Sullivan K. Cognitive functional therapy compared with a group-based exercise and education intervention for chronic low back pain: a multicentre randomised controlled trial (RCT). *Br J Sports Med*. 2020;54(13):782-9. doi: 10.1136/bjsports-2019-100780. PubMed PMID: 31630089.
40. Eccleston C, Blyth FM, Dear BF, Fisher EA, Keefe FJ, Lynch ME, et al. Managing patients with chronic pain during the COVID-19 outbreak: considerations for the rapid introduction of remotely supported (eHealth) pain management services. *Pain*. 2020;161(5):889-93. doi: 10.1097/j.pain.0000000000001885. PubMed PMID: WOS:000531374900003.
41. Garcia LM, Birckhead BJ, Krishnamurthy P, Sackman J, Mackey IG, Louis RG, et al. An 8-week self-administered at-home behavioral skills-based virtual reality program for chronic low back pain: Double-blind, randomized, placebo-controlled trial conducted during COVID-19. *J Med Internet Res*. 2021;23(2). doi: 10.2196/26292.
42. Mayer RE. *Multimedia Learning*. 3rd ed. Cambridge: Cambridge University Press; 2020.
43. Rini C, Porter LS, Somers TJ, McKee DC, DeVellis RF, Smith M, et al. Automated Internet-based pain coping skills training to manage osteoarthritis pain: a randomized controlled trial. *Pain*. 2015;156(5):837-48. doi: 10.1097/j.pain.000000000000121. PubMed PMID: 25734997; PubMed Central PMCID: PMC4402249.

44. Scoliosis Research Society. Frequently Asked Questions Video Library 2018 [cited 2023 8 January]. Available from: <https://www.srs.org/patients-and-families/patient-education-videos/frequently-asked-question-videos#Scoliosis>.
45. Glasgow RE, Harden SM, Gaglio B, Rabin B, Smith ML, Porter GC, et al. RE-AIM Planning and Evaluation Framework: Adapting to New Science and Practice With a 20-Year Review. *Front Public Health*. 2019;7:64. Epub 20190329. doi: 10.3389/fpubh.2019.00064. PubMed PMID: 30984733; PubMed Central PMCID: PMC6450067.
46. D'Lima D, Soukup T, Hull L. Evaluating the Application of the RE-AIM Planning and Evaluation Framework: An Updated Systematic Review and Exploration of Pragmatic Application. *Front Public Health*. 2021;9:755738. Epub 20220126. doi: 10.3389/fpubh.2021.755738. PubMed PMID: 35155336; PubMed Central PMCID: PMC68826088.
47. Harden SM, Gaglio B, Shoup JA, Kinney KA, Johnson SB, Brito F, et al. Fidelity to and comparative results across behavioral interventions evaluated through the RE-AIM framework: a systematic review. *Syst Rev*. 2015;4:155. Epub 20151108. doi: 10.1186/s13643-015-0141-0. PubMed PMID: 26547687; PubMed Central PMCID: PMC4637141.
48. Gaglio B, Shoup JA, Glasgow RE. The RE-AIM framework: a systematic review of use over time. *Am J Public Health*. 2013;103(6):e38-46. Epub 20130418. doi: 10.2105/ajph.2013.301299. PubMed PMID: 23597377; PubMed Central PMCID: PMC3698732.
49. Hoffmann T, Glasziou P, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *Br Med J*. 2014;348:g1687. Epub 20140307. doi: 10.1136/bmj.g1687. PubMed PMID: 24609605.
50. Sterne J, Savovic J, Page M, Elbers R, Blencowe N, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *Br Med J*. 2019;366:l4898. Epub 20190828. doi: 10.1136/bmj.l4898. PubMed PMID: 31462531.
51. Green BN, Johnson C, Moreau W. Is physical activity contraindicated for individuals with scoliosis? A systematic literature review. *J Chiropr Med*. 2009;8(1):25-37. doi: 10.1016/j.jcm.2008.11.001. PubMed PMID: 19646383; PubMed Central PMCID: PMC2697577.
52. Catanzariti JF, Rimetz A, Genevieve F, Renaud G, Mounet N. Idiopathic adolescent scoliosis and obesity: prevalence study. *Eur Spine J*. 2023;32(6):2196-202. Epub 20230425. doi: 10.1007/s00586-023-07709-1. PubMed PMID: 37097342.
53. Konieczny MR, Senyurt H, Krauspe R. Epidemiology of adolescent idiopathic scoliosis. *J Child Orthop*. 2013;7(1):3-9. Epub 20121211. doi: 10.1007/s11832-012-0457-4. PubMed PMID: 24432052; PubMed Central PMCID: PMC3566258.
54. Suh SW, Modi HN, Yang JH, Hong JY. Idiopathic scoliosis in Korean schoolchildren: a prospective screening study of over 1 million children. *Eur Spine J*. 2011;20(7):1087-94. Epub 20110128. doi: 10.1007/s00586-011-1695-8. PubMed PMID: 21274729; PubMed Central PMCID: PMC3176687.
55. Wong HK, Hui JH, Rajan U, Chia HP. Idiopathic scoliosis in Singapore schoolchildren: a prevalence study 15 years into the screening program. *Spine (Phila Pa 1976)*. 2005;30(10):1188-96. doi: 10.1097/01.brs.0000162280.95076.bb. PubMed PMID: 15897834.
56. Apple Inc. Apple Final Cut Pro X license and download [electronic resource]. 2023.
57. Apple Inc. Garage Band license and download [electronic resource]. 2023.
58. Abbade LPF, Abbade JF, Thabane L. Introducing the CONSORT extension to pilot trials: enhancing the design, conduct and reporting of pilot or feasibility trials. (1678-9199 (Print)).
59. Eldridge SM, Chan CL, Campbell MJ, Bond CM, Hopewell S, Thabane L, et al. CONSORT 2010 statement: extension to randomised pilot and feasibility trials. (1756-1833 (Electronic)).

60. Faul F, Erdfelder E, Lang A-G, Buchner A. G* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior research methods*. 2007;39(2):175-91.
61. Alamrani S, Rushton A, Gardner A, Falla D, Heneghan NR. Outcome measures evaluating physical functioning and their measurement properties in adolescent idiopathic scoliosis: a protocol for a systematic review. *BMJ Open*. 2020;10(4):e034286. Epub 20200401. doi: 10.1136/bmjopen-2019-034286. PubMed PMID: 32241788; PubMed Central PMCID: PMC7170637.
62. Golicki D, Młyńczak K. Measurement Properties of the EQ-5D-Y: A Systematic Review. *Value Health*. 2022. Epub 20220622. doi: 10.1016/j.jval.2022.05.013. PubMed PMID: 35752534.
63. Diarbakerli E, Grauers A, Gerdhem P. Population-based normative data for the Scoliosis Research Society 22r questionnaire in adolescents and adults, including a comparison with EQ-5D. *Eur Spine J*. 2017;26(6):1631-7. doi: 10.1007/s00586-016-4854-0.
64. Lin J, Wong CKH, Cheung JPY, Cheung PWH, Luo N. Psychometric performance of proxy-reported EQ-5D youth version 5-level (EQ-5D-Y-5L) in comparison with three-level (EQ-5D-Y-3L) in children and adolescents with scoliosis. *The European Journal of Health Economics*. 2022;23(8):1383-95. doi: 10.1007/s10198-022-01435-z.




Figure 2

Figure 1. Schedule of enrolment, interventions, and assessments.

	STUDY PERIOD									
	Enrolment	Allocation	Post-allocation							Close-out
TIMEPOINT	-8 wk	0	1 wk	2 wk	3 wk	4 wk	5 wk	6 wk	7 wk	8 wk
ENROLMENT:										
Eligibility screen	X									
Informed consent	X									
Allocation		X								
INTERVENTIONS:										
<i>Augmented Videos</i>			←————→							
<i>Traditional Videos</i>			←————→							
<i>Usual care</i>			←————→							
ASSESSMENTS:										
<i>Baseline variables</i>		X								
<i>SRS-22r, EQ-5D-Y, STAI-Ch, PAQ-C</i>		X								X
<i>Engagement</i>			X	X	X	X	X	X	X	X
<i>Knowledge (weekly topics)</i>			X	X	X	X	X	X	X	
<i>Knowledge (all topics)</i>		X								X

Note: wk = Week

Figure 1