

Predictors of blood pressure control and antihypertensive drug adherence among hypertensive patients: Hospital-based cross-sectional study

Tamrat P. Elias, Asteray T. Minyilshewa, Mengesha A. Tekle, Tsegaye W. Gebreamlak, Binyam L. Adde

Abstract

Background: Hypertension or elevated blood pressure is a serious medical condition that significantly increases the risk of diseases of the heart, brain, kidneys, and other organs. Antihypertensive drug adherence is a key to controlling blood pressure.

Objective: To assess factors associated with antihypertensive drug adherence and blood pressure control among hypertensive patients in Selected Public Hospitals under Addis Ababa City Administration.

Method: A hospital-based cross-sectional study was conducted among hypertensive patients on follow-up in randomly selected Public Hospitals under the Addis Ababa City Administration from May 1, 2022, to August 31, 2022. The study population was 393 and patients who fulfilled the eligibility criteria were selected by systematic random sampling and the first participant was selected randomly. Data collection was conducted by reviewing the patient's electronic medical records and by interviewing patients with a structured questionnaire. Data was entered into Epi-Info 7.2.1 and exported to SPSS version 25 software for analysis. Logistic regression analysis was done to see the association between the dependent and independent variables.

Result: The rate of antihypertensive drug adherence and blood pressure control was 72.5

% and 23.4% respectively. Participants with uncontrolled blood pressure were found to

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

be 41.7% less adherent than those with controlled blood pressure (AOR= 0.59; 95% CI, 0.36-0.97). Non-adherence to dietary restriction (AOR, 3.31; 95% CI, 1.84–5.96) and chronic kidney disease (AOR=3.85; 95% CI, 1.41-10.52) are associated with good adherence, whereas using single antihypertensive drug (AOR=0.53; 95% CI, 0.30-0.94), and non-adherence to moderate physical exercise (AOR=0.30; 95% CI, 0.20-0.65) were associated with poor adherence to antihypertensive medications. Male sex (AOR=1.95; 95% CI, 1.04–3.28) and blood pressure measurement at home (AOR=0.59; 95% CI, 0.36–0.99) were independent predictors of controlled blood pressure. Drinking alcohol (AOR=1.92; 95% CI, 1.05-3.49) is inversely associated with blood pressure control.

Conclusion: Despite good adherence to antihypertensive medications, blood pressure control was low. This signifies the importance of lifestyle measures beyond pharmacologic intervention.

Key terms: Hypertension, Blood pressure control, Antihypertensive drugs, Adherence.

Introduction

Hypertension or elevated blood pressure is a serious medical condition that can be defined as office systolic blood pressure (SBP) values ≥ 140 mmHg and/or diastolic blood pressure (DBP) values ≥ 90 mmHg, or the level of BP at which the benefits of treatment unequivocally outweigh the risks of treatment [1].

In 2019, an estimated 1.3 billion people Worldwide have high blood pressure, and it is estimated that the number of people with hypertension will increase by 15–20% by 2025, reaching close to 1.5 billion [2,3]. The overall prevalence of hypertension in adults is around 30 - 45% [4]. The estimated pooled prevalence of hypertension is about 30.8% in Africa [5] and 30.0%- 31.1% in Sub-Saharan Africa [6]. In Ethiopia, there are different prevalence reports in different studies and across different regions of the country.

Systematic meta-analyses conducted in 2019 using published and unpublished articles showed that the overall pooled prevalence of hypertension in the country was 21.8% [7].

Hypertension is associated with a substantial financial burden. The global financial burden of high BP in 2001 was estimated to be around US\$370 billion or about 10% of the world's overall healthcare expenditure [8].

Elevated BP is also associated with a large global burden of cardiovascular disease (CVD) and premature death. According to data from different studies, an estimated 7.7–10.4 million annual deaths are attributable to elevated blood pressure levels, 88% of which were in low- and middle-income countries (LMICs) [9, 10, 11].

Randomized clinical trials have demonstrated that BP lowering with commonly used regimens, such as diuretics, angiotensin-converting-enzyme (ACE) inhibitors, angiotensin-receptor blockers (ARB), and calcium channel blockers reduces the risk of CVD and all-cause mortality [12, 13].

Despite these effective interventions, hypertension control remains unacceptably low, particularly in LMICs [14]. Recent global estimates suggest that only 45.6% of people with hypertension were aware of their condition, only 36.9% were receiving treatment and only 13.8% had achieved BP control (defined as systolic BP <140 mmHg and diastolic BP <90 mmHg) [14].

In chronic diseases, including hypertension, medication mainly serves as a preventive measure and not to suppress symptoms, maintaining long-term adherence is particularly difficult, and the risk of treatment discontinuation is very high [15].

Various studies have shown that poor medication adherence is a primary reason for poor BP control [16, 17]. Poor adherence is responsible for unnecessary overprescription of drugs, substantial worsening of diseases, increased hospital admission rates, and longer

hospital stays all leading to a significant medical burden such as reduced optimal clinical benefit and increased risk of cardiovascular events including acute coronary syndromes, stroke, chronic heart failure as well as mortality [18]. In addition to adverse cardiovascular events, it is also associated with reduced quality of life, disability, reduced work productivity, and greater health care costs [19].

The need of conducting this study is to assess the adherence rate and blood pressure control among hypertensive patients in selected public hospitals under Addis Ababa city administration and identify factors for poor adherence and blood pressure control. So that, knowing the magnitude and reasons for the problem, appropriate interventions can be planned to improve adherence and blood pressure control.

Methodology

Study design and setting:

A hospital-based cross-sectional study was conducted at Yekatit 12 Hospital Medical College and Menelik II comprehensive specialized hospital medical referral clinics among hypertensive patients on follow-up from May 1, 2022, to August 31, 2022, and the study areas were selected randomly.

Yekatit 12 Hospital Medical College, former Bethsaida Hospital established in 1915, is located in the Arada Sub-city of Addis Ababa. It serves around 230,000 people annually both in the emergency and outpatient departments. Menelik II Comprehensive Specialized Hospital was established in 1884 by the Russian Red Cross Society as a nursing school. Progressively it was named as Menelik II referral hospital and started working under the city government of Addis Ababa Health Bureau.

Study participants:

All hypertensive patients on follow-up at Yekatit 12 Hospital Medical College and Menelik II Comprehensive Specialized Hospital medical referral clinics were used as a source population and patients on antihypertensive treatment were the study population. All patients ≥ 18 years of age with hypertension and on antihypertensive treatment for at least 6 months were included in the study. The exclusion criteria were refusal to sign informed consent, pregnancy, and known mental illness.

Data collection tools and procedures:

A structured questionnaire and Checklists were developed after a review of relevant works of literature. Several questions that can address the objective of this study were constructed including the independent variables like socio-demographic characteristics (Age, sex, educational status, occupational status, income, marital status, ethnicity, religion, and residence), behavioral characteristics (medication adherence, low salt diet adherence, physical activity status, alcohol status, smoking status), clinical characteristics (duration of hypertension, a family of hypertension, availability of BP cuff at home, BP monitoring at home or elsewhere, comorbidity, number of antihypertensive drugs) and the dependent variables like antihypertensive drug adherence and blood pressure control. The tool was amended after a pre-test was done on 5% of patients having follow-up for hypertension at St. Paul's Hospital Millennium Medical College. The data was collected by general practitioners working in the chronic medical follow-up clinic of the hospital from May 1, 2022, to November 15, 2022.

Sample size and statistical analysis:

The sample size was determined by using the single population proportion formula. From

the study conducted in Dessie Referral Hospital, the overall good adherence rate was 51.9% (P=0.519%) [20]. Since the total population was 4420, a correction formula was used, and a 10% non-response rate was assumed. The final sample size was calculated to be 393. Study participants were selected using systematic random sampling. After the first patient was selected randomly, then every 13th patient who visited the follow-up clinic was selected, until the required sample size was obtained.

After checking the data for consistency, completeness, clarity, and accuracy, it was coded manually, entered into Epi-Info version 7.2.1, and exported to Statistical Package for Social Sciences (SPSS) version 25.0 for analysis. After descriptive statistics, bivariate analysis was done to see the association between the independent and dependent variables. Variables with a P-value of less than 0.25 in the bivariate analysis were further entered into the multivariate logistic regression model for final analysis. Multivariate analysis was done using a backward logistic regression method. P-value less than 0.05 was considered to determine the statistical significance of the association and odds ratio with a 95% confidence interval was used to determine the presence, strength, and direction of association between covariates and the outcome variable.

Operational definitions:

Hypertension: is defined as office SBP values ≥ 140 mmHg and/or DBP values ≥ 90 mmHg.

Controlled Blood pressure: is defined as SBP < 130 and DBP < 80 .

Drug adherence: is defined based on Morisky's Medication Adherence Scale (MMAS-8) [73]

- Patients were considered as adherent to their medication(s) when Morisky's medication adherence scale (MMAS-8) score is > 6

- Non-adherence was considered when the patient's MMAS-8 score was ≤ 6 .

Ethical Consideration

Ethical clearance was obtained from the ethical review committee of Addis Ababa City Administration Health Bureau and the ethical committee of Yekatit 12 Hospital Medical College. Written informed consent was obtained from study participants. Participant identity remains confidential. The answers obtained were documented and analyzed anonymously.

Results:

Socio-demographic Characteristics

A total of 393 participants were included in the study with a 100% response rate. The majority of the respondents were between 46-60 years (42%) and the mean age of patients was 59.2 (± 12.1 SD) years. More than half (57.3%) of study participants were female. About 25% of the respondents completed college/university. Those who live in urban areas constitute the majority of participants (98%) and about 45.8 % of them had a monthly total income greater than 3,000 Ethiopian birr (ETB) per month followed by those who earn 1,000-2,000 ETB (22.4%).

Table 1: Socio-demographic characteristics of the patients (n=393).

Characteristics	Frequency	Percentage
Sex		
Male	168	42.7
Female	225	57.3
Age in years		

18-44	48	12.2
45-60	165	42.0
61-75	146	37.2
>76	34	8.7
Educational level		
Unable to read and write	65	16.5
Read and write	26	6.6
Primary School	67	17.0
Secondary School	68	17.3
High School/Preparatory	68	17.3
College/University	99	25.2
Marital Status		
Single	30	7.6
Married	253	64.4
Divorced	25	6.4
Widowed	85	21.6
Occupation		
Government Employee	58	14.8
Retired	88	22.4
Housewife	122	31.0
Private	85	21.6
No job	40	10.2
Residence		
Urban	385	98
Rural	8	2
Monthly income in ETB		
<1,000	52	13.2
1,000 - 2,000	88	22.4
2,001 – 3,000	73	18.6
>3,000	180	45.8

Behavioral Characteristics

Of the total 393 participants, 4.6% are active cigarette smokers. The prevalence of alcohol consumption was 22.9%. Most of the respondents ate Vegetables (44%) and Cereal Products (42.7%) over the past 6 months on most days of the week. More than half (59.5%) of respondents use additional salt in their diet. A quarter (75.1%) of respondents had moderate physical activity with a median time of 30 minutes per day.

Table 2: Behavioral characteristics of the patients (n=393)

Behavioral character	Frequency	Percentage
Cigarette Smoking		
Yes	18	4.6
No	375	95.4
Alcohol Consumption		
Yes	90	22.9
No	303	77.1
Additional Salt Added on Plate		
Yes	234	59.5
No	159	40.5
Vigorous Physical Exercise		
Yes	17	4.3
No	376	95.7
Moderate Physical Exercise		
Yes	295	75.1
No	98	24.9

Comorbidities

Considering the duration of Hypertension, in almost half (49.6%) of the patients, hypertension was diagnosed within the past 5 years. Diabetes mellitus (49.1%) and heart

failure (19.8%) are the two most common Comorbidities. More than one-third of respondents had a family history of hypertension.

Table 3: Duration of hypertension and comorbidities of the patients (n=393)

Duration of Hypertension	Frequency	Percentage
<5 Years	195	49.6
5-10 Years	120	30.5
>10 Years	78	19.8
Comorbidities		
Cardiovascular Diseases	78	19.8%
Diabetes Mellitus	193	49.1%
Chronic kidney disease	23	5.9%
Asthma	14	3.6
Family History of Hypertension		
Yes	149	37.9
No	244	62.1

Antihypertensive Drug Utilization

CCBs (67.2%), ACEIs (58.8%), and Thiazide diuretics (26.5%) are the three commonly utilized antihypertensive drugs. ACEIs and CCBSs are the most common combination treatment accounting for about 33.8%. Monotherapy is utilized in 37.2% of patients, whereas polytherapy is utilized in 62.8% of study participants. Adverse drug reactions were reported in five (1.3%) patients and each of two is due to CCBs and ACEIs and for one patient no specific drug was identified. Most participants (95.4%) bought antihypertensive medication from hospitals and got physician advice about the benefits of antihypertensive drugs.

Table 4: Antihypertensive drug utilization of the patients (n=393)

Antihypertensive drugs	Frequency	Percentage
CCBs	264	67.2%

ACEIs	231	58.8%
Thiazide Diuretics	104	26.5%
Beta Blockers	56	14.2%
Loop Diuretics	49	12.5%
ARBs	8	2%
MRAs	18	4.6%
ACEIs + CCBs	133	33.8%
ACEIs + Thiazide diuretics	47	12%
CCBs + Thiazide diuretics	55	14%
ACEIs +CCBs + Thiazide diuretics	23	5.9%
Monotherapy	146	37.2%
Polytherapy	247	62.8%
Drug Availability in Hospital		
Yes	375	95.4
No	18	4.6
Physician Advice About the Benefits of Antihypertensive drugs		
Yes	334	85.0
No	59	15.0
Average Cost of Antihypertensive Drugs	600.53 birr	

The respondent's level of adherence to antihypertensive drugs was assessed using the eight-item Morisky Medication Adherence Scale. It was found that 72.5 % of the respondents were adherent to medication, whereas 27.5 % were not.

More than half (52.4%) of patients measure Blood Pressure at home. The rate of blood pressure control before six months, three months of Study, and at the time of Study was 20.4%, 27.2%, and 23.4% respectively and the rate of recent blood pressure control in Yekatit 12 Hospital Medical College and Menelik II Referral Hospital was 28.8% and 11.5% respectively.

Factors associated with antihypertensive drug adherence

A multiple logistic regression model was used to assess the association between adherence status and various factors. All statistically significant variables from univariate analysis at a level of 0.25 were included in the multiple logistic regression analysis to control confounding effects. The thirteen factors that were included in the model were sex, age, level of education, monthly income, number of antihypertensive medications, additional salt on a plate, vigorous physical exercise, moderate physical exercise, duration of hypertension, cardiovascular diseases, chronic kidney disease, blood pressure measurement at home, and blood pressure control status.

From the above table, those participants who added additional salt on the plate were 3.3 times more adherent than those who didn't add additional salt on a plate (odds ratio [OR], 3.31; 95% CI, 1.84– 5.96), those participants who took single antihypertensive medications are 47% less likely to be adherent than those who take two or more antihypertensive medications (odds ratio [OR], 0.53; 95% CI, 0.30-0.94), participants who didn't do moderate physical exercise were 70% less adherent than those who do moderate physical exercise (odds ratio [OR], 0.3; 95% CI, 0.20-0.65), and participants without chronic kidney disease are 3.8 times more likely to be non-adherent than those with chronic kidney disease (odds ratio [OR], 3.85; 95% CI, 1.41-10.52).

From Bivariate regression analysis, those patients with uncontrolled blood pressure are 41.7% less adherent than those with controlled blood pressure ($p=0.041$, $p < 0.05$) (odds ratio [OR], 0.59; 95% CI, 0.36–0.97).

Table 5: Factors associated with antihypertensive drug adherence

Variable	Drug Adherence		COR (95%CI)	P value	AOR (95%CI)	P value
	Yes	No				
Sex						
Male	127	41	0.76(0.48-1.19)	0.23	0.74(0.376-1.45)	0.38

Female	158	67	1			
Age						
25-44	29	19	1.2(0.48-2.9)	0.69		
45-60	124	41	0.6(0.27-1.3)	0.21	1.04(0.37-2.94)	0.92
61-75	110	36	0.6(0.27-1.3)	0.20	1.44(0.55-3.77)	0.45
76-90	22	12	1			
Level of education						
Unable to read and write	44	21	2.14(1.03-4.45)	0.04	1.02(0.35-2.95)	0.95
Read and write	15	11	3.30(1.30-8.37)	0.01	0.68(0.20-2.28)	0.53
Primary School	51	16	1.41(0.66-3.01)	0.37		
Secondary School	47	21	2.01(0.97-4.15)	0.05	0.67(.276-1.63)	0.38
High School/Preparatory	47	21	2.01(0.97-4.15)	0.05	0.42(.18-1.00)	0.05
College/University	81	18	1			
Occupation						
Government Employee	43	15	0.81(0.33-1.99)	0.65		
Retired	71	17	0.55(0.23-1.31)	0.18	1.76(.60-5.16)	0.30
Housewife	79	43	1.27(0.58-2.74)	0.54		
Private	64	21	0.76(0.33-1.76)	0.53		
No job	28	12	1			
Monthly income						
<1000	32	20	2.05(1.06-3.96)	0.03	0.55(0.23-1.29)	0.17
1000-2000	59	29	1.61(0.92-2.83)	0.09	0.63(0.31-1.30)	0.21
2001-3000	56	17	0.99(0.52-1.89)	0.99		
>3000	138	42	1			
Added salt to plate						
Yes	146	88	0.23(0.13-0.4)	0.00*	3.31(1.84-5.966)	0.00**
No	139	20				
Vigorous physical exercise						
Yes	9	8	0.40(0.15-1.08)	0.07	2.31(0.66-8.01)	0.18
No	276	100	1			
Moderate physical exercise						
Yes	230	65	0.76(1.70-4.49)	0.00*	0.30(0.20-0.65)	0.01**
No	55	43	1			
Duration of Hypertension						
<5 Years	135	60	1.06(0.59-1.88)	0.83		
5-10 Years	95	25	0.62(0.32-1.21)	0.16	1.17(0.54-2.53)	0.68
>10 Years	55	23	1			
Cardiovascular Disease						
Yes	49	29	0.56(0.33-0.97)	0.03*	1.40(0.73-2.67)	0.30
No	236	79	1			
CKD						
Yes	9	14	0.21(0.08-0.51)	0.00*	3.85(1.41-10.52)	0.08**

No	276	94	1			
Health education						
Yes	44	23	1.48(0.84-2.59)	0.17	0.82(0.42-1.59)	0.56
No	241	85	1			
Number of Antihypertensive drugs						
Single	118	28	0.49(0.30-0.80)	0.00	0.53(0.30-0.94)	0.03**
Two and above	167	80	1			
BP measurement at home						
Yes	161	45	1.81(1.16-2.84)	0.00	0.84(0.48-1.46)	0.54
No	124	63				
Blood Pressure Control						
Controlled	59	33	0.59(0.36-0.97)	0.04	1.58(0.87-2.85)	0.12
Not controlled	226	75				

Factors associated with blood pressure control

The association between blood pressure control with patient-related, medication-related, and Institution-related factors was assessed using bivariate and multivariate logistic regression. Multiple logistic regression models are used to assess the adjusted association of predictors with adherence status. All statistically significant variables from bivariate analysis at a level of 0.25 were included in the multiple logistic regression analysis to control confounding effects. The ten factors that were included in the model were sex, cigarette smoking, alcohol drinking, diet, type of oil used, duration of hypertension, cardiovascular diseases, health education about hypertension and its treatment, drug adherence, and blood pressure measurement at home. Drinking alcohol and male sex and blood pressure measurement at home are independent predictors of controlled blood pressure.

As shown in Table 6, male participants had 1.95 times more controlled blood pressure than female participants (odds ratio [AOR], 1.95; 95% CI, 1.04–3.28), those participants who didn't drink alcohol had 1.92 times more controlled blood pressure than who drink alcohol (odds ratio [AOR], 1.92; 95% CI, 1.05-3.49) and those participants who didn't

measure blood pressure at home had 41% less controlled blood pressure than those who measure blood pressure at home (odds ratio [AOR], 0.59; 95% CI, 0.36–0.99).

Table 6: Factors associated with blood pressure control

Variable	BP control		COR (95% CI)	P- value	AOR (95% CI)	P-value
	Yes	No				
Sex						
Male	27	141	0.3(0.28-0.71)	0.004*	1.854(1.04-3.28)	0.035**
Female	65	160	1			
Cigarette smoking	.		.			
Yes	91	284	6.66(0.75-58.8)	0.08*	0.15(0.01-1.28)	0.08
No	1	17	1			
Alcohol Drinking						
Yes	63	240	0.44(0.23-0.83)	0.01*	1.92(1.05-3.49)	0.031**
No	29	61	1			
Diet						
Meat	12	40	1.16(0.47-2.90)	0.73		
Vegetables	48	125	1.75(0.96-3.21)	0.06*	0.69(0.40-1.21)	0.20
Cereal Products	32	136	1			
Type of oil						
Vegetable Oil	87	276	3.22(0.55-18.67)	0.19*	0.41(0.08-1.94)	0.26
Butter	1	4	2.74(0.14-50.73)	0.49		
Sesame/Nug Oil	2	5	2.30(0.17-30.57)	0.52		
Others	2	16	1			
Duration of Hypertension						
<5 Years	49	146	0.66(0.32-1.37)	0.27		
5-10 Years	21	99	0.49(0.22-1.09)	0.08*	1.90(0.91-3.96)	0.08
>10 Years	22	56	1			
Cardiovascular Disease						
Yes	68	247	0.52(0.26-1.00)	0.05*	1.77(0.96-3.24)	0.06
No	24	54				
Health education						
Yes	21	46	1.68(0.87-3.25)	0.12*	0.67(0.36-1.25)	0.21
No	71	255	1			
Drug adherence			.			
Yes	59	226	0.56(0.30-1.03)	0.06*	1.4(0.48-1.46)	0.21
No	33	75	1			
BP measurement at home						
Yes	56	131	1.86(1.05-3.30)	0.03*	0.59(0.36-0.99)	0.04**
No	36	170	1			

Discussion

The study showed that the majority of study participants were adherent to antihypertensive medications. But, despite this good adherence, less than one-third of the participants had controlled blood pressure. The addition of salt on the plate, Moderate physical exertion, chronic kidney disease, and the number of antihypertensive drugs were strongly associated with adherence whereas male sex, alcohol drinking, and blood pressure measurement at home were significantly associated with good blood pressure control. Antihypertensive drug adherence was significantly associated with controlled blood pressure.

In this study, the rate of antihypertensive drug adherence was 72.5%, which is higher than other studies conducted in Dessie Referral Hospital (51.9%) [20], Jimma University Medical Center (61.8%) [21], Southwest Ethiopia (60.5%) [22], Nedjo General Hospital (31.4%) [23], and the national medication adherence rate among hypertensive patients (65.1%) [24]. However, the rate of medication adherence was comparable with a study conducted in Sudan, which is 70.5% [25] and lower compared with a study from Tanzania, where the adherence rate was 76.9%, and a study in Japan reported an adherence rate of 91.5% [26].

The variation from other countries could be due to socioeconomic factors and population characteristics and from previous studies in our country could be that most of the studies were done in rural parts where most of the residents didn't attend formal education. This study is conducted in an urban setup where the society has a better awareness of drug adherence. The other reason could be the difference in the assessment tools.

The rate of blood pressure control was 23.4%, which is lower than the studies conducted in Southwest Ethiopia (50.3%) [22], Jimma University Medical Center (42.8%) [27], University of Gondar Referral Hospital (50.4%) [28], Dessie city (55.8%) [29], Eastern

Sudan (45.3%) [25], and Cameroon (36.8%) [30]. However, this study has Comparable results with the US study (24%) [31] which used a similar operational definition of controlled hypertension.

The variation might be due to the difference in the operational definition of controlled blood pressure in previous studies, which was SBP less than 140 and DBP less than 90 as compared to the operational definition in this study, which was SBP less than 130, and DBP less than 80.

A statistically significant association was not found between adherence to antihypertensive medication with patient-related factors including age, sex, level of education, marital status, residence, income, and occupation which is similar to the study conducted at Jimma University Medical Center [27], Eastern Sudan [25], and Cameroon [30].

A statistically significant association was found between adherence to antihypertensive medication and adding additional salt to the plate. Those participants who added additional salt to the plate were more adherent ($P < 0.01$). In this study, the rate of dietary salt restriction was 40% which is higher than reported from two meta-analysis and systematic review studies which was 34% [32] and 37.3% [33]. However, no data report showed an association with medication adherence.

Participants who take single antihypertensive medications were found to be less adherent than those who take two or more antihypertensive medications. Similar findings were reported in Nigeria in which patients taking three or more antihypertensive medications are more adherent with a rate of 73.1% [34] and a study from Lebanon demonstrated taking two or more drugs was a predictor of high adherence ($P = 0.02$) [35]. Another study from Pakistan also reported that patients on monotherapy had a mean adherence of 79% compared to 90% for those on three drugs or more antihypertensive medications

[36]. However, a study conducted at Jimma University Medical Center showed that two or more antihypertensive medication use was associated with poor adherence [21] and a cross-sectional Study conducted in Saudi Arabia showed being on multiple medications was associated with medication non-adherence ($P < 0.01$) [37].

Chronic kidney disease was associated with good medication adherence in our study. Opposite results were reported from various studies in which the presence of chronic kidney disease was associated with poor drug adherence and the adherence rate also declined with worsening CKD [38, 39, 40]. Of note, a Brazilian study has shown that drug adherence tends to improve as renal function deteriorates [41] and another systematic review and meta-analysis showed a 2.54 times increase in the odds of being adherent among patients who had co-morbidities as compared with those without comorbidities [24].

This could be related to physicians and patients becoming more concerned by the quality of BP control as the disease progresses as well as they perceive the combined danger of multiple comorbidities and make them to be more adherent than those without comorbidities.

In this study, male respondents had a higher rate of blood pressure control than females. However, there was an opposite finding from a study conducted in Kenya in which the male sex is an independent predictor of poor blood pressure control [42], a study from the US also reported that male sex was independently associated with uncontrolled blood pressure [43].

Participants who did not drink alcohol had good blood pressure control in this study. REasons for Geographic and Racial Differences in Stroke (REGARDS) Study also showed that heavy alcohol consumption is associated with poor BP control ($P = 0.02$) [40].

Population-based studies conducted in the UK showed higher alcohol use was associated with poor blood pressure control [44].

Participants who measured blood pressure at home had more controlled blood pressure. Similar findings were reported from a systematic review and meta-analysis by Cappuccio FP, blood pressure was lower in people with hypertension who had home blood pressure monitoring than in those who had standard blood pressure monitoring in the healthcare system [45].

Blood pressure control is found to be associated with adherence. Similar findings were reported from several studies demonstrating the strong association between antihypertensive drug adherence and blood pressure control [24, 46-49].

The limitation of this study was the use of indirect methods to assess adherence which predisposes to recall bias due to the self-reporting used for measuring adherence.

Conclusion

Anti-hypertensive drug adherence is the most important management component for hypertensive patients to achieve adequate blood pressure control which is important in reducing hypertension-associated adverse cardiovascular morbidity and mortality. It was found that 72.4% were adherent to antihypertensive medications despite this only 23.4% of patients had controlled blood pressure. This signifies the importance of lifestyle modifications beyond pharmacologic interventions.

References

1. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, Clement DL, Coca A, De Simone G, Dominiczak A, Kahan T. 2018 ESC/ESH Guidelines for the management of arterial hypertension: The Task Force for the management of arterial

- hypertension of the European Society of Cardiology (ESC) and the European Society of Hypertension (ESH). *European Heart Journal*. 2018 Sep 1; 39(33):3021-104.
2. Hypertension. Key facts. 25 August 2021 [website] (<https://www.who.int/news room /fact sheets/ detail/hypertension>, accessed 27 November 2021).
 3. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *The Lancet*. 2005 Jan 15; 365(9455):217-23.
 4. Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, Bahonar A, Chifamba J, Dagenais G, Diaz R, Kazmi K. Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in high-, middle-, and low-income countries. *Jama*. 2013 Sep 4;310(9):959-68.
 5. Adeloje D, Basquill C. Estimating the prevalence and awareness rates of hypertension in Africa: a systematic analysis. *PloS one*. 2014 Aug 4;9(8):e104300.
 6. Ataklte F, Erqou S, Kaptoge S, Taye B, Echouffo-Tcheugui JB, Kengne AP. Burden of undiagnosed hypertension in sub-Saharan Africa: a systematic review and meta-analysis. *Hypertension*. 2015 Feb; 65(2):291-8.
 7. Tiruneh SA, Bukayaw YA, Yigizaw ST, Angaw DA. Prevalence of hypertension and its determinants in Ethiopia: A systematic review and meta-analysis. *PloS one*. 2020 Dec 31;15(12):e0244642.
 8. Gaziano TA, Bitton A, Anand S, Weinstein MC. The global cost of nonoptimal blood pressure. *Journal of hypertension*. 2009 Jul 1;27(7):1472-7.
 9. Danaei G, Lu Y, Singh GM, Carnahan E, Stevens GA, Cowan MJ, Farzadfar F, Lin JK, Finucane MM, Rao M, Khang YH. Cardiovascular disease, chronic kidney disease, and diabetes mortality burden of cardiometabolic risk factors from 1980 to 2010: a comparative risk assessment. *Lancet Diabetes & Endocrinology*. 2014.

10. Stanaway JD, Afshin A, Gakidou E, Lim SS, Abate D, Abate KH, Abbafati C, Abbasi N, Abbastabar H, Abd-Allah F, Abdela J. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*. 2018 Nov 10;392(10159):1923-94.
11. Zhou B, Perel P, Mensah GA, Ezzati M. Global epidemiology, health burden and effective interventions for elevated blood pressure and hypertension. *Nature Reviews Cardiology*. 2021 May 28:1-8.
12. Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of different blood pressure-lowering regimens on major cardiovascular events: results of prospectively designed overviews of randomized trials. *The Lancet*. 2003 Nov 8;362(9395):1527-35.
13. Ettehad D, Emdin CA, Kiran A, Anderson SG, Callender T, Emberson J, Chalmers J, Rodgers A, Rahimi K. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *The Lancet*. 2016 Mar 5;387(10022):957-67.
14. Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, Chen J, He J. Global disparities of hypertension prevalence and control: a systematic analysis of population-based studies from 90 countries. *Circulation*. 2016 Aug 9; 134(6):441-50.
15. Kjeldsen SE. Hypertension and cardiovascular risk: General aspects. *Pharmacological research*. 2018 Mar 1; 129:95-9.
16. Charles H, Good CB, Hanusa BH, Chang CC, Whittle J. Racial differences in adherence to cardiac medications. *Journal of the National Medical Association*. 2003 Jan;95(1):17.
17. Redmond N, Baer HJ, Hicks LS. Health behaviors and racial disparity in blood pressure control in the national health and nutrition examination survey. *Hypertension*. 2011 Mar;57(3):383-9.

18. Mazzaglia G, Ambrosioni E, Alacqua M, Filippi A, Sessa E, Immordino V, Borghi C, Brignoli O, Caputi AP, Cricelli C, Mantovani LG. Adherence to antihypertensive medications and cardiovascular morbidity among newly diagnosed hypertensive patients. *Circulation*. 2009 Oct 20; 120(16):1598-605.
19. Black HR, Elliott W. Hypertension: A Companion to Braunwald's Heart Disease E-Book. Elsevier Health Sciences; 2012 Aug 1.
20. Andualem A, Liknaw T, Edmealem A, Gedefaw M. Adherence to antihypertensive medications among adult hypertensive patients attending chronic follow-up units of Dessie Referral Hospital, Northeastern Ethiopia: A cross-sectional study. *Medicine*. 2021 Aug 6;100(31).
21. Asgedom SW, Atey TM, Desse TA. Antihypertensive medication adherence and associated factors among adult hypertensive patients at Jimma University Specialized Hospital, southwest Ethiopia. *BMC research notes*. 2018 Dec; 11(1):1-8.
22. Asgedom SW, Gudina EK, Desse TA. Assessment of blood pressure control among hypertensive patients in Southwest Ethiopia. *PloS one*. 2016 Nov 23; 11(11):e0166432.
23. Berisa HD, Dedefo MG. Non-adherence related factors to antihypertensive medications among hypertensive patients on follow up at Nedjo General Hospital in West Ethiopia. *The Open Public Health Journal*. 2018 Feb 28;11(1).
24. Desta M, Yibeltal D, Memiah P, Ayenew T, Mulugeta H, Gedefaw M, Kidanie BB, Birhanu MY, Tessema Z, Mengist B, Alene M. Antihypertensive medications adherence and associated uncontrolled blood pressure among hypertensive patients in Ethiopia: systematic review and meta-analysis. *International Journal of Africa Nursing Sciences*. 2022 Mar 15:100404. 53.

25. Omar SM, Elnour O, Adam GK, Osman OE, Adam I. Assessment of blood pressure control in adult hypertensive patients in eastern Sudan. *BMC cardiovascular disorders*. 2018 Dec;18(1):1-6.
26. Ibaraki A, Goto W, Iura R, Tominaga M, Tsuchihashi T. Current prescription status of antihypertensive drugs with special reference to the use of diuretics in Japan. *Hypertension Research*. 2017 Feb;40(2):203-6.
27. Kebede B, Chelkeba L, Dessie B. Rate of blood pressure control and its determinants among adult hypertensive patients at Jimma University Medical Center, Ethiopia: Prospective cohort study. *SAGE open medicine*. 2021 Mar; 9:20503121211006000.
28. Animut Y, Assefa AT, Lemma DG. Blood pressure control status and associated factors among adult hypertensive patients on outpatient follow-up at University of Gondar Referral Hospital, northwest Ethiopia: a retrospective follow-up study. *Integrated blood pressure control*. 2018;11:37.
29. Fentaw Z, Adamu K, Wedajo S. Blood pressure control status of patients with hypertension on treatment in Dessie City Northeast Ethiopia. *BMC Public Health*. 2022 Dec;22(1):1-7
30. Menanga A, Edie S, Nkoke C, Boombhi J, Musa AJ, Mfeukeu LK, Kingue S. Factors associated with blood pressure control amongst adults with hypertension in Yaounde, Cameroon: a cross-sectional study. *Cardiovascular diagnosis and therapy*. 2016 Oct;6(5):439.
31. Centers for Disease Control and Prevention (CDC). Hypertension Cascade: Hypertension Prevalence. Treatment and Control Estimates Among US Adults Aged. 2017;18:2013-6.
32. de Terline DM, Kramoh KE, Diop IB, Nhavoto C, Balde DM, Ferreira B, Houenassi MD, Hounsou D, Ikama MS, Kane A, Kimbally-Kaki SG. Poor adherence to medication and salt restriction as a barrier to reaching blood pressure control in patients with

- hypertension: Cross-sectional study from 12 sub-Saharan countries. *Archives of cardiovascular diseases*. 2020 Jun 1;113(6-7):433-42.
33. Edmealem A, Ademe S, Gedamu S. Adherence to Self-Care among Patients with Hypertension in Ethiopia: A Systematic Review and Meta-Analysis. *International Journal of Hypertension*. 2022 Jul 16;2022.
 34. Usman MN, Umar MD, Idris FA, Abdullahi Y. Medication adherence and its associated factors among hypertensive patients in a tertiary health facility in Minna, North central Nigeria. *Archives of Clinical Hypertension*. 2019 May 16;5(1):003-7.
 35. Yassine M, Al-Hajje A, Awada S, Rachidi S, Zein S, Bawab W, Zeid MB, El Hajj M, Salameh P. Evaluation of medication adherence in Lebanese hypertensive patients. *Journal of epidemiology and global health*. 2016 Sep 1;6(3):157-67.
 36. Hashmi SK, Afridi MB, Abbas K, Sajwani RA, Saleheen D, Frossard PM, Ishaq M, Ambreen A, Ahmad U. Factors associated with adherence to anti-hypertensive treatment in Pakistan. *PloS one*. 2007 Mar 14;2(3):e280
 37. Algabbani AM. Treatment adherence among patients with hypertension: findings from a cross-sectional study. *Clinical Hypertension*. 2020 Dec;26(1):1-9.
 38. Schmitt KE, Edie CF, Laflam P, Simbartl LA, Thakar CV. Adherence to antihypertensive agents and blood pressure control in chronic kidney disease. *American journal of nephrology*. 2010;32(6):541-8.
 39. Burnier M, Pruijm M, Wuerzner G, Santschi V. Drug adherence in chronic kidney diseases and dialysis. *Nephrology Dialysis Transplantation*. 2015 Jan 1;30(1):39-44.
 40. Judd SE, McClure LA, Howard VJ, Lackland DT, Halanych JH, Kabagambe EK. Heavy drinking is associated with poor blood pressure control in the REasons for Geographic and Racial Differences in Stroke (REGARDS) study. *International journal of environmental research and public health*. 2011 May;8(5):1601-12

41. Magacho EJC, Ribeiro LC, Chaoubah A, et al. Adherence to drug therapy in kidney disease, Braz J Med Biol Res, 2011, vol. 44 (pg. 258-262)
42. Kubo MN, Kayima JK, Were AJ, McLigeyo SO, Ogola EN. Factors associated with uncontrolled hypertension among renal transplant recipients attending nephrology clinics in Nairobi, Kenya. Journal of transplantation. 2015 Jul 14;2015.],
43. Hyman DJ, Pavlik VN. Characteristics of patients with uncontrolled hypertension in the United States. New England Journal of Medicine. 2001 Aug 16;345(7):479-86.]. This difference might be due to different characteristics of the study population.
44. Tapela N, Collister J, Clifton L, Turnbull I, Rahimi K, Hunter DJ. Prevalence and determinants of hypertension control among almost 100,000 treated adults in the UK. Open heart. 2021 Feb 1;8(1):e001461
45. Cappuccio FP, Kerry SM, Forbes L, Donald A. Blood pressure control by home monitoring: meta-analysis of randomized trials. Bmj. 2004 Jul 15;329(7458):145.
46. Yue Z, Bin W, Weilin Q, Aifang Y. Effect of medication adherence on blood pressure control and risk factors for antihypertensive medication adherence. Journal of evaluation in clinical practice. 2015 Feb;21(1):166-72.
47. Khayyat SM, HyatAlhazmi RS, Mohamed MM, Abdul Hadi M. Predictors of medication adherence and blood pressure control among Saudi hypertensive patients attending primary care clinics: a cross-sectional study. PloS one. 2017 Jan 30;12(1):e0171255.],
48. Hasen G, Abdo A. Medication Adherence and Associated Factors in Management of Hypertension in Shashemene Referral Hospital, Ethiopia. Journal of Hypertension and Management. 2020;3.
49. Yue Z, Bin W, Weilin Q, Aifang Y. Effect of medication adherence on blood pressure control and risk factors for antihypertensive medication adherence. Journal of evaluation in clinical practice. 2015 Feb;21(1):166-72.