
EVALUATING AND MITIGATING LIMITATIONS OF LARGE LANGUAGE MODELS IN CLINICAL DECISION MAKING

Paul Hager^{1,2,*}, Friederike Jungmann^{1,2,*}, Kunal Bhagat³, Inga Hubrecht⁴, Manuel Knauer⁴, Jakob Vielhauer⁵, Robbie Holland⁷, Rickmer Braren², Marcus Makowski², Georgios Kaisis^{1,6,7}, Daniel Rueckert^{1,7}

ABSTRACT

1 Clinical decision making is one of the most impactful parts of a physician's responsibilities and stands
2 to benefit greatly from AI solutions and large language models (LLMs) in particular. However, while
3 LLMs have achieved excellent performance on medical licensing exams, these tests fail to assess
4 many skills that are necessary for deployment in a realistic clinical decision making environment,
5 including gathering information, adhering to established guidelines, and integrating into clinical
6 workflows. To understand how useful LLMs are in real-world settings, we must evaluate them *in the*
7 *wild*, i.e. on real-world data under realistic conditions. Here we have created a curated dataset based
8 on the MIMIC-IV database spanning 2400 real patient cases and four common abdominal pathologies
9 as well as a framework to simulate a realistic clinical setting. We show that current state-of-the-art
10 LLMs do not accurately diagnose patients across all pathologies (performing significantly worse
11 than physicians on average), follow neither diagnostic nor treatment guidelines, and cannot interpret
12 laboratory results, thus posing a serious risk to the health of patients. Furthermore, we move beyond
13 diagnostic accuracy and demonstrate that they cannot be easily integrated into existing workflows
14 because they often fail to follow instructions and are sensitive to both the quantity and order of
15 information. Overall, our analysis reveals that LLMs are currently not ready for clinical deployment
16 while providing a dataset and framework to guide future studies.

17 1 Main

18 Large language models (LLMs) have the potential to revolutionize our medical system[54]. They can already streamline
19 report generation and summarization[61, 59, 8, 37], answer biomedical questions with[60, 5, 59] and without[50, 49, 38]
20 images, and could soon effectively interpret multimodal data for precision medicine in the clinic[6]. Importantly,
21 as humans primarily interact with the world through language, LLMs are poised to be the point of access to the
22 multimodal medical AI solutions of the future[36]. Until now, however, the diagnostic capabilities of models have
23 been tested in structurally simple medical contexts, such as canonical vignettes of hypothetical patients or clinical case
24 challenges. In both scenarios, all the required diagnostic information is provided upfront and there is a single answer to
25 be selected from a list of options. This type of question dominates the medical licensing exams that have been used to
26 test LLMs such as the United States Medical Licensing Exam (USMLE)[27, 26], Applied Knowledge Test (AKT) of
27 the Membership of the Royal College of General Practitioners (RCGP)[53], and All India Institute of Medical Sciences
28 (AIIMS) & National Eligibility cum Entrance Test Postgraduate (NEETS PG) entrance exam[42]. LLMs have proven
29 their ability to excel in such scenarios, scoring well above passing on medical licensing exams[31, 21, 49, 50, 38, 56, 39]
30 and rivaling clinician performance on clinical case challenges [34, 28, 10, 18].

31 However, while these medical licensing exams and clinical case challenges are suitable for testing the general medical
32 knowledge of the test-taker, they are far removed from the daily task of clinical decision making. For medical
33 practitioners, clinical decision making is one of the most important and complex everyday responsibilities. It is a
34 multi-step process that requires gathering and synthesizing data from diverse sources and continuously evaluating
35 the facts to reach an evidence-based decision on patient diagnosis and treatment[7, 55]. To reach a precise diagnosis,
36 physicians must gather the necessary information based on the availability of diagnostic resources and adhere to
37 established guidelines. Furthermore, they must carefully consider the patient-specific symptoms to plan the optimal
38 treatment. As this process is very labor-intensive, there exists great potential in harnessing AI, such as LLMs, to
39 alleviate much of the workload, ultimately aiming to autonomously, efficiently, and safely reach a final diagnosis which
40 can then be confirmed by physicians[48, 3]. Thus, to understand how useful LLMs would be in such a real-world

¹ Lab for AI in Medicine, TUM, Munich, Germany ² Institute for Diagnostic and Interventional Radiology, Klinikum rechts der Isar, TUM, Munich, Germany ³ Department of Medicine, ChristianaCare Health System, Wilmington, DE, USA ⁴ Department of Medicine III, Klinikum rechts der Isar, TUM, Munich, Germany ⁵ Department of Medicine II, University Hospital of the LMU Munich, Munich, Germany ⁶ Helmholtz Munich ⁷ Imperial College London * Equal contribution. E-mail: paul.hager@tum.de




Figure 1: (a.) To properly evaluate LLMs for clinical decision making in realistic conditions, we created a curated dataset from real-world patient cases derived from the MIMIC-IV database, which contains comprehensive electronic health record data recorded during hospital admissions. (b.) Our evaluation framework reflects a realistic clinical setting and thoroughly evaluates LLMs across multiple criteria, including diagnostic accuracy, adherence to diagnostic and treatment guidelines, consistency in following instructions, ability to interpret laboratory results, and robustness to changes in instruction, information quantity, and information order. Abbreviations: ICD: International Classification of Diseases. CT: Computed Tomography. US: Ultrasound. MRCP: Magnetic resonance cholangiopancreatography.

41 setting, they must be evaluated on real-world data and under realistic conditions. However, the only analysis that tested
42 an LLM throughout the clinical workflow, used curated lists of possible answers and examined only 36 hypothetical
43 clinical vignettes[46]. Furthermore, any model that is used in such a high-stakes clinical context must not only be
44 highly accurate, but also adhere to diagnostic and treatment guidelines, be robust, and follow instructions, all of which
45 have not been tested in previous medical evaluations.

46 Here, we present a curated dataset based on the Medical Information Mart for Intensive Care (MIMIC-IV) database
47 spanning 2400 real patient cases and four common abdominal pathologies (appendicitis, pancreatitis, cholecystitis,
48 diverticulitis) as well as a comprehensive evaluation framework around our dataset to simulate a realistic clinical setting.
49 We provide LLMs with a patient’s history of present illness and ask them to iteratively gather and synthesize additional
50 information such as physical examinations, laboratory results, and imaging reports until they are confident enough to
51 provide a diagnosis and treatment plan. Our new dataset, task, and analysis represent the first large-scale evaluation of
52 LLMs on every-day clinical decision making tasks in a realistic, open-ended environment. Unlike previous works, we
53 test the autonomous information gathering and open-ended diagnostic capabilities of models, representing an essential
54 step towards evaluating their suitability as clinical decision makers.

55 To understand how useful LLMs would be in the clinic today, we compare the diagnostic accuracy of the models on our
56 dataset with that of clinicians. Furthermore, we propose and evaluate a new range of characteristics beyond diagnostic
57 accuracy, such as adherence to diagnostic and treatment guidelines, correct interpretation of laboratory test results,
58 instruction following capabilities, and robustness to changes in instructions, information order, and information quantity.
59 Finally, we show that summarizing progress and filtering laboratory results for only abnormal results addresses some of
60 the current limitations of models. We make our evaluation framework and dataset freely and openly available to guide
61 future studies considering the use of LLMs in clinical practice.

62 **2 Results**

63 **2.1 Creating the MIMIC-CDM dataset and evaluation framework**

64 Our curated dataset, MIMIC Clinical Decision Making (MIMIC-CDM), is created using the well-established MIMIC-IV
65 database, which is managed by the Massachusetts Institute of Technology (MIT). MIMIC-IV contains de-identified
66 records of patient measurements, diagnoses, procedures, treatments, and free-text clinical notes such as discharge
67 summaries and radiologists reports from patients admitted to the Beth Israel Deaconess Medical Center in Boston,
68 USA, from 2008 to 2019[22]. Figure 1a and Materials Section 6 list the steps involved in creating the MIMIC-CDM
69 dataset and its makeup. Our dataset contains electronic health record data from 2400 unique patients presenting with
70 acute abdominal pain to the emergency department and whose primary diagnosis was one of the following pathologies:
71 appendicitis, cholecystitis, diverticulitis, or pancreatitis. We chose these target pathologies as they represent clinically
72 important diagnoses of a common chief complaint, abdominal pain, which accounts for 10% of all emergency department
73 visits [11, 15]. Furthermore, different treatment strategies, ranging from antibiotics to surgery, are necessary depending
74 on the severity of the condition. Thus, a thorough understanding of the specifics of a patient’s case is required to
75 recommend optimal treatment. Importantly, a good differentiation between the four pathologies can be achieved using
76 standard diagnostic tests, all of which are present in MIMIC-CDM.

77 To reflect a realistic clinical setting that allows LLMs to autonomously engage in every step of the clinical decision
78 making process we have created a comprehensive evaluation framework around our dataset. Using our framework and
79 dataset, we present LLMs with a patient’s history of present illness and task them to gather and synthesize information
80 such as physical examinations, specific laboratory results, and distinct imaging reports to arrive at a diagnosis and
81 treatment plan, as shown in Figure 1b. We then take the complete interaction and evaluate the LLM for diagnostic
82 accuracy as well as adherence to diagnostic and treatment guidelines, and instruction following capabilities. For
83 comparisons with practicing clinicians and further tests concerning robustness, we evaluate the diagnostic accuracy
84 of LLMs when provided with all necessary information for a diagnosis; a dataset we call MIMIC Clinical Decision
85 Making with Full Information (MIMIC-CDM-FI). In this dataset, we include the history of present illness, physical
86 examination, relevant laboratory results, and all abdominal imaging, before directly asking for a diagnosis.

87 In our study, we tested the leading open-access LLM developed by Meta, Llama 2[58], and its derivatives. We focus
88 on the largest model with 70 billion (B) parameters, as it performed best on benchmarks including medical questions.
89 We test both generalist versions such as Llama 2 Chat (70B)[58], Open Assistant (OASST) (70B)[30], and WizardLM
90 (70B)[63], as well as medical-domain aligned models such as Clinical Camel (70B)[56], and Meditron (70B)[12].
91 Further information on the models and our selection criteria can be found in Materials Section 6.3. Data taken from the
92 MIMIC database is currently prohibited from being used with external APIs, such as that of OpenAI or Google, due to
93 data privacy concerns and data usage agreements, so neither ChatGPT nor GPT-4 could be tested. Furthermore, multiple

94 requests for access to Google’s MedPaLM models were denied. We note that Llama 2, Clinical Camel, and Meditron
 95 have been shown to match and even exceed ChatGPT performance on medical licensing exams and biomedical question
 96 answering tests[56, 12].

97 2.2 LLMs diagnose significantly worse than clinicians

98 To ensure patient safety, LLMs must diagnose at least as well as clinicians. Thus, we compared the diagnostic accuracy
 99 of the models on a subset of MIMIC-CDM-FI to four practicing hospitalists: two from the Klinikum Rechts der Isar
 100 Hospital of the Technical University of Munich, Germany (with two and three years of experience), one from the
 101 Ludwig Maximilian University Hospital in Munich, Germany, (four years of experience) and one from the Christiana
 102 Care hospital in Delaware, United States of America (29 years of experience). All four of the hospitalists are internal
 103 medicine physicians with emergency department experience. Each hospitalist was instructed to provide the primary
 104 pathology afflicting the patient and was given the same 100 patients in a random order to diagnose. 20 patients of each
 105 target pathology (appendicitis, cholecystitis, diverticulitis, pancreatitis) were included, with an additional five patients
 106 each of four other abdominal pathologies: gastritis, urinary tract infection, esophageal reflux, and hernia. Each LLM
 107 model was evaluated 10 times, using different random seeds, over the subset of 80 patients to increase statistical power.
 108 All statistical tests were corrected for multiple comparisons.




Figure 2: On a subset (n=80) of MIMIC-CDM-FI, LLMs perform significantly worse than doctors on average and especially on diseases such as cholecystitis and diverticulitis. The exact diagnostic accuracy is shown above each bar. Vertical lines indicate range of scores.

109 We find that current LLMs perform significantly worse than clinicians as each model’s mean performance, averaged
 110 over all four pathologies, is significantly lower than the mean of all clinicians (Doctors vs. Llama 2 Chat, $p=8.06e-10$;
 111 Doctors vs. OASST, $p=1.10e-4$; Doctors vs. WizardLM, $p=6.18e-6$; Doctors vs. ClinicalCamel, $p=1.89e-4$; Doctors vs
 112 Meditron, $p=6.40e-5$) (Fig. 2). The diagnostic accuracy of the clinicians also varied, with the German hospitalists in
 113 residency (Mean = $87.50\% \pm 3.68\%$) performing slightly worse than the more senior US hospitalist (Mean = 92.50%),
 114 which can be attributed to differences in both experience and language, considering all text was in English. Most
 115 models were able to match clinician performance on the simplest diagnosis, appendicitis, where 3 out of 4 clinicians
 116 also correctly diagnosed 20 out of 20 patients. While the Meditron model matched or exceeded the other models at
 117 diagnosing appendicitis, diverticulitis, and pancreatitis patients, it failed for cholecystitis, diagnosing only one patient
 118 with ‘cholecystitis’ and the others primarily with ‘gallstones’ without specifying their location or inflammatory effects.
 119 This mirrors the general performance of the models, which may perform well on certain pathologies but currently lack
 120 the diagnostic range of human hospitalists. In a standard clinical scenario, where every diagnosis is a possibility, models
 121 must perform consistently across all pathologies of a single initial complaint, such as abdominal pain, to be useful.

122 No specialist medical models performs significantly better on average than any generalist models (Clinical Camel vs.
 123 Llama 2 Chat, $p=0.13$; Clinical Camel vs. OASST, $p=1.13$, Clinical Camel vs. WizardLM, $p=0.47$; Meditron vs. Llama
 124 2 Chat, $p=1.11$; Meditron vs. OASST, $p=0.61$; Meditron vs. WizardLM, $p=1.04$) (Fig. 2). Furthermore, as the medical
 125 LLMs are not instruction-fine-tuned (i.e. trained to understand and undertake new tasks), they are unable to complete

126 the full clinical decision making task where they must first gather information and then come to a diagnosis. As this is
127 the primary use-case of a clinical decision making model, we exclude them from all further analysis and only examine
128 the Llama 2 Chat, OASST and WizardLM models for the rest of this work.

129 In our simulated clinical environment, which uses the MIMIC-CDM dataset, the LLM must specify all information
130 it wishes to gather to accurately diagnose a patient. We observe a general decrease in performance, compared to
131 MIMIC-CDM-FI (Supplementary Fig. 13), across the pathologies (Fig. 3). The mean diagnostic averages fall to
132 45.5% (vs. 58.8% on MIMIC-CDM-FI) for Llama 2 Chat, 54.9% (vs. 67.8%) for OASST, and 53.9% (vs. 65.1%) for
133 WizardLM. Mirroring MIMIC-CDM-FI, all models performed best in diagnosing appendicitis (Llama 2 Chat: 74.6%,
134 OASST: 82.0%, WizardLM: 78.4%) which is most likely due to the fact that appendicitis patients have consistent
135 key symptoms with 791 of 957 radiologist reports (82.7%) directly stating that the appendix is dilated, enlarged, or
136 fluid-filled, and typically lack other intra-abdominal pathology descriptions which distract from the acute diagnosis.




Figure 3: When tasked with gathering all information required for clinical decision making themselves, LLMs perform best when diagnosing appendicitis but perform poorly on the other three pathologies of cholecystitis, diverticulitis, and pancreatitis. In such a realistic clinical scenario, model performance decreased compared to the retrospective diagnosis with all information provided (MIMIC-CDM-FI). The exact diagnostic accuracy is shown above each bar.

137 In summary, LLMs do not reach the diagnostic accuracy of clinicians across all pathologies, especially when they must
138 gather all information themselves. Thus, without extensive physician supervision, they would reduce the quality of care
139 that patients receive and are currently unfit for the task of clinical decision making.

140 2.3 LLMs are hasty and unsafe clinical decision makers

141 In addition to poor diagnostic accuracy, LLMs often fail to order the exams required by diagnostic guidelines, do not
142 follow treatment guidelines, and are incapable of interpreting lab results, making them a risk to patient safety.

143 To help clinicians consistently and safely reach a final diagnosis, diagnostic guidelines are published by associations of
144 medical experts. The guidelines help guarantee a consistent standard of care by establishing a clinical consensus of
145 which tests should be used to effectively diagnose a pathology, based on large clinical trials and clinician experience.
146 The current clinical guidelines used for this study were: appendicitis[16], cholecystitis[44], diverticulitis [25], and
147 pancreatitis[32].

148 All guidelines recommended physical examinations as an essential part of the diagnostic process, preferably as the first
149 action. This is because they immediately provide valuable information as to the severity of the patient's condition and a
150 base for subsequent requests for laboratory tests and imaging. We find that only Llama 2 Chat consistently asks for
151 physical examination results, either as the first action (97.1%) or at all (98.1%) (Supplementary Fig. 11). The other
152 two models requested less examinations (OASST: 79.8% & 87.7%; WizardLM: 53.1% & 63.9%), thereby omitting an
153 essential piece of information.

154 Hospitalists routinely order laboratory tests to measure biological markers and evaluate organ function, allowing them
155 to track changes in a patient's health and detect underlying biochemical shifts due to disease. Based on the diagnostic
156 guidelines, we defined categories of necessary laboratory tests for each pathology, including signs of inflammation,




Figure 4: LLMs often do not order the necessary laboratory tests required to establish a diagnosis, despite each necessary test being ordered independently by all doctors in the MIMIC-CDM dataset. The tests, defined by current diagnostic guidelines, help differentiate abdominal pathologies, as results can indicate which organ is currently pathologically afflicted or functioning normally.

157 functional fitness of the liver and gallbladder, pancreas enzymes, and the severity of a patient’s pancreatitis. The exact
 158 tests included in each category can be found in Supplementary Section D. We find that no model consistently orders
 159 all necessary tests, despite each test being independently requested by all doctors in the MIMIC-CDM dataset (Fig.
 160 4). While OASST performs better than the other two models, reaching up to 93.3% and 87.2% in the inflammation
 161 category for appendicitis and diverticulitis, it often does not order the necessary tests for a diagnosis of pancreatitis
 162 (pancreas enzymes: 56.5%, severity: 76.2%), partially explaining why its diagnostic performance on pancreatitis was
 163 only 44.1% (Fig. 3). Not ordering all necessary tests makes it difficult to differentiate between abdominal pathologies,
 164 as laboratory results can provide an indication which organ is currently afflicted or functioning normally.

165 While it is important to order the correct laboratory tests, it is even more essential to correctly interpret them. The
 166 laboratory tests in MIMIC-IV have reference ranges included when applicable, so we tested the interpretation capabilities
 167 of the models by providing the test result with the accompanying reference range and asking them to classify each result
 168 as either below, within, or above the provided range. Any human with numerical literacy should be able to achieve
 169 perfect accuracy on such a task, however all LLMs perform very poorly, especially in the critical categories of low
 170 test results (Chat: 26.5%, OASST: 70.2%, WizardLM: 45.8%) and high test results (Chat: 50.1%, OASST: 77.2%,
 171 WizardLM: 24.1%) (Supplementary Fig. 15). Such a basic incomprehension of laboratory test results is a great risk to
 172 patient safety and must be resolved before LLMs become clinically useful.

173 While diagnostic guidelines also have recommendations for which imaging modality is best suited for establishing
 174 a diagnosis, the modality used in practice can vary based on current availability and the particularities of a patient’s
 175 case. We find that models sometimes match the modalities requested by the doctors in the dataset, but often come to
 176 a diagnosis without requesting an abdominal imaging scan (Supplementary Fig. 12). We include the first abdominal
 177 imaging modality recorded for each patient in MIMIC-CDM for comparison. As we later show that imaging is the
 178 most useful diagnostic tool for the LLMs for all pathologies except pancreatitis (Fig. 8), occasional failure to request
 179 imaging could be partly responsible for their low diagnostic accuracy.

180 Taken together, the lack of consistency of the LLMs in ordering all of the required tests for a diagnosis based on current
 181 guidelines indicates a tendency to diagnose before understanding or considering all of the facts of the patient’s case.
 182 This could pose a risk to patients’ health as they would be diagnosed based on insufficient information.

183 In addition to not following diagnostic guidelines, LLMs generally fail to adhere to treatment guidelines (Fig. 5).
 184 To evaluate their ability to recommend appropriate treatments, we used the aforementioned guidelines to extract the
 185 possible treatments for each pathology and then to classify each treatment as either essential (e.g. antibiotics, support)
 186 or case specific (e.g. appendectomy, cholecystectomy, drainage). For each patient, we then determined if the case
 187 specific treatment was appropriate by matching against the actual operations performed on that patient, read from the
 188 MIMIC-CDM dataset. We evaluate a model’s treatment recommendation only when it correctly diagnosed a patient
 189 since an inaccurate diagnosis likely leads to inappropriate treatment. Fig. 5 shows the total counts of each treatment in
 190 MIMIC-CDM under each treatment name and the counts of each models’ correct diagnoses under each bar. We find
 191 that the LLMs consistently do not recommend appropriate and sufficient treatment, especially for patients with more
 192 severe forms of the pathologies. While they are consistent in recommending some treatments such as appendectomy
 193 for appendicitis (Chat: 97.5%, OASST: 97.5%, WizardLM: 96.9%) and antibiotics for diverticulitis (Chat: 88.6%,
 194 OASST: 97.4%, WizardLM: 86.0%), they rarely recommend other treatments when appropriate such as colectomy for
 195 diverticulitis patients with perforated colons (Chat: 22.2%, OASST: 0.0%, WizardLM: 10.0%) or drainage of infected
 196 pancreatic necrosis (Chat: 0.0%, OASST: 0.0%, WizardLM: 0.0%) and abscesses near the diverticula (Chat: 16.7%,
 197 OASST: 22.2%, WizardLM: 0.0%). Furthermore, they drastically under-treat appendicitis with regards to the necessity
 198 of antibiotics (Chat: 8.5%, OASST: 26.0%, WizardLM: 6.0%) and providing support (Chat: 4.3%, OASST: 9.9%,
 199 WizardLM: 2.7%), diverticulitis with the need for a colonoscopy in the future to check for colon cancer (Chat: 4.5%,
 200 OASST: 18.8%, WizardLM: 9.6%), and pancreatitis with sufficient support (where we expect mentions of fluids, pain
 201 management, and monitoring) (Chat: 39.8%, OASST: 71.3%, WizardLM: 42.9%). In summary, following the treatment
 202 recommendations of the models would negatively impact the health of patients, particularly those with more advanced
 203 stages of disease where indications for emergency operations were ignored.




Figure 5: We find that LLMs do not consistently recommend essential and patient-specific treatment, especially for patients with more severe forms of the pathologies. We only score models on the subset of patients that they correctly diagnosed and that actually received a specific treatment. For example, of the 957 appendicitis patients, 808 received an appendectomy (indicated below the treatment name). Of those 808 patients, Llama 2 Chat correctly diagnosed 603 (indicated below the Llama 2 Chat bar). Of those 603 patients, Llama 2 Chat correctly recommended an appendectomy 97.5% of the time.

204 The tendency of LLMs to not gather the information required by diagnostic guidelines before returning a final diagnosis
205 and treatment plan is particularly problematic considering their low overall diagnostic and treatment accuracy. Such
206 hasty decision making combined with their poor diagnoses and treatment recommendations pose a serious risk to the
207 health of patients without extensive clinician supervision and control.

208 2.4 LLMs require extensive clinician supervision to effectively integrate into clinical workflows

209 In addition to consistently and safely arriving at the correct diagnosis and treatment plan, models must integrate into
210 established clinical workflows to be useful. Central to this is the ability to follow instructions and generate answers so
211 they can be easily processed and used by other parts of the clinic without physician supervision. During the clinical
212 decision making process, we provide clear instructions to the models as to how they should provide their requests and
213 diagnosis, as well as which tools are available to them (see Materials Section 6.2). For example, diagnostic tools must
214 be written in the ‘Action:’ field and desired tests must be specified in the ‘Action Input:’ field, and not in the middle of
215 a paragraph surrounded by other text. This is essential to ensure that the desired tests can be consistently extracted
216 so no manual clinician supervision and interpretation is required. Through extensive comparisons of LLM outputs
217 with dictionaries of known exams and their synonyms, we go to great lengths to understand what tests are requested,
218 even if the models do not follow our schema, recording every time they fail to follow instructions. We investigate the
219 capabilities of models to follow our instructions at three timepoints during our analysis: when providing the next action
220 to take, when requesting a tool, and when providing a diagnosis.

221 All models struggle to follow the provided instructions (Fig. 6), making errors every two to four patients when providing
222 actions (Chat: 1.91, OASST: 3.90, WizardLM: 2.13) and hallucinating non-existent tools every two (Chat: 2.20 and
223 WizardLM: 2.41) to five patients (OASST: 5.48). When providing diagnoses on the MIMIC-CDM dataset, errors are
224 made every three to five patients (Chat: 3.18, OASST: 4.55, WizardLM: 3.64) while on the MIMIC-CDM-FI dataset
225 there is a greater discrepancy in the number of patients diagnosed without formatting error (Chat: 1.12, OASST: 6.60,
226 WizardLM: 28.21). While many of these errors are easily caught (Supplementary Table 6), the error rate is so high that
227 extensive manual controls would be necessary to ensure model output is being correctly interpreted. Such poor abilities
228 to follow instructions greatly reduces their usefulness in clinical systems as they would require large amounts of manual
229 supervision to ensure proper performance.




Figure 6: LLMs struggle to follow instructions, often introducing errors when providing the next action to take and hallucinating non-existent tools, up to once every two patients. Formatting errors while providing the diagnosis also regularly occur. In the clinic, extensive manual supervision would be required to ensure proper performance.

230 Another key component that must be fulfilled before we consider integrating such models into real world workflows
231 is robustness. Models must not be sensitive to small changes in user instructions as their performance will then vary
232 greatly based on who is interacting with them. On the MIMIC-CDM-FI dataset, we find that changes in instructions
233 (Supplementary B) can lead to large changes (both positive and negative) in diagnostic accuracy (Fig. 7). For example,
234 large changes are seen when removing system and user instructions (up to +5.1% for Chat on cholecystitis, down to
235 -16.0% for Chat on pancreatitis), or when removing all medical terminology from the system instruction (up to +6.2
236 for WizardLM on diverticulitis, down to -3.5% for OASST on pancreatitis). Additionally, we see that even minor

237 changes in instructions can greatly change diagnostic accuracy such as asking for the ‘Main Diagnosis:’ (up to +7.0%
 238 for Chat on diverticulitis, down to -10.6% for WizardLM on cholecystitis) or ‘Primary Diagnosis:’ (up +8.7% for Chat
 239 on Pancreatitis, down to -5.2% for WizardLM on cholecystitis) instead of ‘Final Diagnosis:’. Models should be able to
 240 provide the most appropriate diagnosis given the situation, in this case the reason for the patient’s abdominal pain, and
 241 not be sensitive to minute changes in phrasing so as not to require extensive clinician training before use.




Figure 7: Often small changes in instructions, such as changing final diagnosis to main diagnosis or primary diagnosis, greatly affects the performance of the LLMs on the MIMIC-CDM-FI dataset. This would vary the quality of responses received depending on who is using the model.

242 Furthermore, LLMs must perform better or equally well when provided with more information. Large amounts of
 243 data are gathered during a patient’s hospital stay, with the patients in our dataset averaging 150 laboratory tests and 3
 244 imaging exams per admission. Models must be able to focus on the key facts of the case to make a diagnosis. We show
 245 that models perform worse when all diagnostic exams are provided, typically attaining their best performance when only
 246 a single exam is provided in addition to the history of present illness (Fig 8). Removing information greatly increases
 247 diagnostic accuracy, with cholecystitis diagnosis improving by 18.5% for the Chat and 16.5% for the WizardLM
 248 models when only providing radiologist reports, and pancreatitis diagnosis improving by 21.6% (Chat), 9.5% (OASST),
 249 and 8.6% (WizardLM) when only providing laboratory results. This greatly reduces the usefulness of such models
 250 as they cannot simply be given all relevant information and be trusted to arrive at their best diagnosis. To optimize
 251 model performance, clinicians would have to decide which diagnosis is most likely to effectively filter the information
 252 presented. While such filtering by clinicians would increase their mean scores to a theoretical best of Chat: 72.2%,
 253 OASST: 70.9%, WizardLM: 71.8%, it would remove any benefit of deploying such a clinical decision making model.

254 Models must also provide the same diagnosis, irrespective of the order in which the information is presented. We test
 255 the diagnostic consistency of the models on the MIMIC-CDM-FI dataset by switching the order of the information from
 256 the canonical physical examination, then laboratory tests, then imaging, to all possible permutations thereof (history of
 257 present illness is always included first). We show that all models have large ranges of performance, up to 18.0% (Chat -
 258 Pancreatitis), 7.9% (OASST - Cholecystitis), and 5.4% (WizardLM - Cholecystitis) (Fig. 9 and Supplementary Fig. 16).
 259 Importantly, we find that the order of information that delivers the best performance for each model is different for
 260 each pathology (Supplementary Section I). This again reduces the benefits of deploying the models as clinicians must
 261 constantly consider and monitor in which order they provide the models with information, in a disease specific manner,
 262 to not degrade performance.

263 In summary, extensive clinician supervision and prior evaluation of the most probable diagnosis would be required
 264 to ensure proper functioning of LLMs because they do not reliably follow instructions, perform better with a disease
 265 dependant order of information and degrade in performance when given relevant information. Furthermore, their
 266 sensitivity to small changes in instructions that seem inconsequential to humans would require extensive clinician
 267 training to ensure good performance.




Figure 8: For almost all diseases, providing all information does not lead to the best performance on the MIMIC-CDM-FI dataset. This suggests that LLMs cannot focus on the key facts and degrade in performance when too much information is provided. This is a problem in the clinic where an abundance of information is typically gathered to wholistically understand the patients health and being able to focus on key facts is an essential skill. The theoretical best shows the mean performance if a clinician were to select the best diagnostic test for each pathology.




Figure 9: By mixing the order in which information is presented to LLMs, their performance changes despite the information included staying the same. This places an unnecessary burden upon clinicians who would need to make preliminary diagnoses to decide the order in which they feed the models with information for best performance.

268 2.5 Summarization and filtering for abnormal laboratory results partially mitigates limitations of current 269 LLMs

270 To help address some of the limitations found in this analysis, we explore simple modifications that can be done without
271 retraining the model. One major limitation is that LLMs are currently limited in the amount of text they can read, with
272 all models tested in this study having a limit of 4096 tokens or approximately 2400 words. Due to this, if the history
273 of present illness or radiologist reports are disproportionately long, the limit of the model is sometimes reached. To
274 alleviate this, we implement an automatic summarization protocol that monitors the length of the text that the model
275 receives. Once it approaches its maximum token amount, we ask the model to summarize the test results it has received,
276 combining the healthy observations and emphasizing the pathological indications (Supplementary B.2). Removing such
277 a summarization protocol resulted in marginal but consistent losses on the mean of -1.3% (Chat), -0.8% (OASST), and

278 -0.5% (WizardLM), and particularly hurt the diagnosis of diverticulitis -4.7% (Chat), -2.7% (OASST), -3.5% (Wizard)
 279 (Supplementary Fig. 17).

280 Due to the inability of LLMs to reliably interpret laboratory results (Supplementary Fig. 15), even when provided
 281 with reference ranges, and their issues understanding larger quantities of information (Fig. 8), we find that filtering
 282 the laboratory results and removing all normal test results generally improved performance on the MIMIC-CDM-FI
 283 Dataset (Fig. 10). This increases the amount of other information that can be included before reaching the input length
 284 limit of the LLMs and reduces the amount of healthy information, which tends to confuse the tested models. Removing
 285 this information increased Llama 2 Chat and WizardLM mean performance by 6.0% and 3.9% respectively, while
 286 slightly changing mean OASST diagnostic accuracy by -0.9%. On the MIMIC-CDM-FI dataset, cholecystitis patients
 287 benefited the most from such a filter, with all models increasing in diagnostic accuracy (Chat: +16.8%, OASST: +3.5%,
 288 WizardLM: +14.8%). Pancreatitis patients in MIMIC-CDM-FI on the other-hand saw a light decrease in performance
 289 for the OASST (-6.1%) and WizardLM (-4.1%) models, with a slight increase for Llama 2 Chat (+2.0%). As many of
 290 our other analyses examine the general behaviour of laboratory tests and their impact on model performance, we do
 291 not use this fix for any other sections of this work. While this filtering improves the performance of the LLMs as they
 292 function today, ideally a model would perform best with all available information. Healthy laboratory test results are an
 293 important source of information for clinicians and should not degrade model performance.




Figure 10: Filtering to include only abnormal laboratory results using the laboratory reference ranges provided in MIMIC-IV database generally improves model performance, especially for the cholecystitis pathology. This allows the model to focus more on abnormal, pathological signals.

294 3 Discussion

295 The strong performance of LLMs on medical licensing exams has led to increased interest in using them in clinical
 296 decision making scenarios involving real patients. However, medical licensing exams do not test the capabilities
 297 required for real world clinical decision making. We have, for the first time, evaluated leading open access large
 298 language models on making open-ended clinical decisions with thousands of real world patient cases to assess their
 299 potential benefits and possible harms. By not only comparing their diagnostic performance against clinicians, but also
 300 testing their information gathering abilities, adherence to guidelines, and instruction following capabilities as well as
 301 their robustness to changes in prompts, information order, and information quantity, we move for the first time beyond
 302 simple evaluations of diagnostic accuracy and establish a range of characteristics that are necessary for safe and robust
 303 integration into the clinic. In this work, we have shown that current leading LLMs are unsuitable for clinical decision
 304 making on all of these accounts.

305 Among the models tested in this study, we find that OASST performed best overall as it had decent diagnostic accuracy,
306 generally requested appropriate laboratory exams, and was most robust to changes in information quantity. Llama 2 Chat
307 had the worst overall diagnostic accuracy, often refused to follow instructions, and was heavily influenced by the order
308 and quantity of information, but was the only model to consistently ask for a physical examination. WizardLM was
309 the most robust to changes in the order of diagnostic exams and followed instructions well when returning diagnoses,
310 but was the worst at following diagnostic guidelines, failing to consistently order physical examinations and necessary
311 laboratory tests. Despite the performance of OASST being generally better than Chat and WizardLM across the diverse
312 set of analyses included in this study, it is still not currently suitable for clinical use due to its inferior performance
313 compared to clinicians, broad failure to order correct treatments, and general lack of robustness. While one of the
314 medical domain models, Clinical Camel, achieved the highest diagnostic accuracy (mean=73% vs OASST mean=68%,
315 Supplementary Fig. 13), its inability to participate in the iterative clinical decision making task precluded it from
316 evaluations of its robustness and consistency which we believe to be essential to ensure safe deployment in the clinic.
317 Other LLMs such as ChatGPT, GPT-4, Med-PALM, and Med-PALM 2 could not be tested due to the data privacy and
318 usage agreements of MIMIC-IV, highlighting the risk of using corporate models in a sensitive area such as medicine,
319 where patient privacy, transparency, and reliability are essential[57, 24].

320 The biggest barrier to using current LLMs in clinical practice is that no model consistently reached the diagnostic
321 accuracy of clinicians across all pathologies, with a further decrease in accuracy when having to gathering diagnostic
322 information themselves. A major weakness of these models is that their diagnostic accuracy is disease dependent,
323 with markedly higher accuracy on appendicitis compared to cholecystitis, diverticulitis, and pancreatitis. A clinical
324 model must diagnose well on all major differential diagnoses of a chief complaint, such as abdominal pain, to be useful.
325 Additionally, LLMs are unable to classify a lab result as normal or abnormal, even when provided with its reference
326 range. This is underscored by the fact that presenting the model with only abnormal laboratory results generally
327 improved diagnostic performance.

328 We further found that the models do not follow diagnostic guidelines, which is particularly problematic considering
329 their low overall diagnostic accuracy, indicating a tendency to diagnose before fully understanding a patient's case.
330 Insufficient diagnostic information also negatively affected the treatment recommendations of LLMs, where we showed
331 that models do not follow all established treatment guidelines, especially for severe cases. The hasty decision making of
332 the models combined with their low diagnostic performance and poor treatment recommendations pose a serious risk to
333 the health of patients without extensive clinician supervision and control.

334 Beyond diagnostic accuracy, we extensively test models on their reliability and robustness which are essential character-
335 istics to ensure consistent and safe patient care. We found that models struggle to follow instructions, often hallucinating
336 non-existent tools and requiring continuous manual supervision to ensure proper performance. Models are also sensitive
337 to seemingly inconsequential changes in instruction phrasing, requiring clinicians to carefully monitor the language
338 they use to interact with the models to not degrade performance. Contrary to expectation, LLMs diagnose best when
339 only a single diagnostic exam is provided rather than when given all relevant diagnostic information, demonstrating an
340 inability to extract the most important diagnostic signal from the evidence. Counterintuitively, we found models to
341 be sensitive to the order in which information is presented, resulting in large changes in diagnostic accuracy despite
342 identical diagnostic information. Importantly, all of these weaknesses are disease-specific within each model, meaning
343 that a different instruction, diagnostic test, and order of tests achieved best results for each pathology. As physicians
344 would have to perform preliminary diagnostic evaluations in an attempt to maximize model performance according to
345 their suspected diagnosis, all benefits of a clinical decision making system would be lost.

346 Many of the current limitations of LLMs exposed in our study have been shown concurrently in domains outside of
347 medicine. It has been shown that LLM performance on tasks can vary by between 8% and 50% just by optimizing the
348 instructions[64]. The sensitivity of LLMs to the order of presented information has been well documented on multiple
349 choice questions [66, 43] and information retrieval[33]. The difficulty LLMs have in interpreting numbers[51] and
350 solving simple arithmetic[17] is an active research topic[23, 52]. Even the largest models currently available, PaLM 2
351 and GPT4, perform poorly on instruction following tests[67]. Our analysis demonstrates how these current limitations
352 of LLMs become harmful in medical contexts where robustness and consistency are essential. We argue that these
353 understudied aspects of model performance should become normal parts of medical model evaluations and that all of
354 these issues must be addressed before LLMs can be considered for clinical decision making.

355 4 Limitations

356 While we have been able to demonstrate the limitations of current leading large language models for clinical decision
357 making, we consider the following limitations of our study.

358 First, as we are using a dataset of real-world clinical data, we rely on the accuracy of the discharge diagnosis written by
359 the attending clinicians. Since a diagnosis using our dataset must be made with available information and without direct
360 access to the patient, the task of coming to the exact diagnosis of the attending physician is a difficult one. We can
361 only provide the information that was gathered during the patient's hospital stay, so model requests for information
362 not in the MIMIC-IV database must be denied. However, as the MIMIC-IV database contains all data gathered during
363 a patient's hospital stay, we can assume that all information required for a diagnosis and treatment plan is contained
364 within our data considering the attending physicians successfully diagnosed and treated all patients. Furthermore, being
365 flexible enough to handle acute restrictions, such as unavailable imaging modalities or laboratory tests, and still come to
366 a correct diagnosis is a desirable ability for any real-world clinical AI application. Due to this difficulty, we were lenient
367 in our evaluation of the diagnoses, accepting alternative names for the pathologies, as long as they were medically
368 correct (see Supplementary Table 5).

369 Another issue concerns the handling of data gathered multiple times during a patient's hospital stay, such as laboratory
370 data. As we test LLMs in an emergency-department-centered clinical decision making scenario, we only provide the
371 models with the first collected value for each laboratory test as they are likely most indicative for diagnostic purposes
372 rather than treatment monitoring. Thus, we currently do not capture the changes in laboratory test values if the patient's
373 condition deteriorates over their hospital stay. This could be remedied by examining all time points and determining the
374 most abnormal test result to be returned or by allowing multiple requests for laboratory tests to return successive test
375 results. However, both of these approaches would widen the temporal gap of provided test results, possibly providing
376 conflicting diagnostic signals. Considering LLMs also have poor temporal reasoning capabilities[62], simply including
377 the timestamp would most likely not be an adequate solution.

378 Our comparison between models and clinicians included only three doctors in residency from Germany and one senior
379 hospitalist from the United States. Increasing the diversity and number of clinicians as well as the number of patient
380 cases evaluated would give a more nuanced view of model performance compared to practicing hospitalists. Future
381 models could possibly soon reach or even outperform clinicians in residency and thus provide a low-cost, interactive
382 second opinion to consult, as is already the case for AI models in other areas such as mammography screening [35].

383 Lastly, we only examined the initial complaint of abdominal pain and four related diagnostic endpoints in our study.
384 While these pathologies are clinically important and well suited for our analysis, it will be important to test future
385 models on both additional diagnostic endpoints and a broader range of initial complaints. A clinical decision making
386 model must show strong performance across all possible pathologies of a particular initial complaint to guarantee
387 adequate patient care without extensive preliminary diagnoses by clinicians. As each initial complaint has its own
388 set of relevant diagnostic exams and is often best investigated with complaint-dependant exams, model performance
389 could vary greatly. We believe the MIMIC-IV database to be a rich resource to help create such additional datasets
390 considering its unfiltered inclusion of all patients that presented to the emergency room.

391 **5 Conclusion**

392 In conclusion, our study presents the first analysis of the capabilities of current state-of-the-art large language models
393 on real-world data in a realistic clinical decision making scenario. Our main finding is that current models do not
394 achieve satisfactory diagnostic accuracy, performing significantly worse than trained physicians, and do not follow
395 treatment guidelines, thus posing a serious risk to the health of patients. This is exacerbated by the fact that they do
396 not request the necessary exams for a safe differential diagnosis, as required by diagnostic guidelines, indicating a
397 tendency to diagnose before fully understanding a patient's case. Furthermore, we show LLMs are distracted by relevant
398 diagnostic information, are sensitive to the order of diagnostic tests, and struggle to follow instructions, prohibiting
399 their autonomous deployment in the clinic and requiring extensive clinician supervision.

400 By sourcing our dataset from real clinical cases and closely aligning our evaluation criteria with official diagnostic and
401 treatment guidelines, we present the first analysis to help physicians understand how well LLMs would perform in the
402 clinic today. While our findings cast doubt on the suitability of LLMs for clinical decision making as they currently
403 exist, we believe there lies great potential in their use after the issues raised are resolved. By making our dataset
404 and framework freely available we hope to guide the development of the next generation of clinical AI models and
405 contribute towards a future where physicians can feel confident in using safe and robust models to improve patient
406 outcomes.

407 6 Methods

408 6.1 MIMIC-CDM Dataset

409 We created our curated dataset of 2400 patients which we call MIMIC-Clinical Decision Making (MIMIC-CDM),
410 using the MIMIC-IV Database [22]. The MIMIC-IV Database is a comprehensive, publicly available database of the
411 de-identified electronic health records of almost 300,000 patients that presented to the Beth Israel Deaconess Medical
412 Center (BIDMC) in Boston, Massachusetts from 2008 to 2019. It contains real patient cases from the hospital and
413 includes all recorded measurements such as laboratory and microbiology test results, diagnoses, procedures, treatments
414 and free-text clinical notes such as discharge summaries and radiologist reports.

415 In this work, we focus on four target pathologies for which we filter: appendicitis, cholecystitis, diverticulitis, and
416 pancreatitis. As we are only testing for these pathologies, we must ensure that they are the primary diagnosis and reason
417 for the patient presenting to the emergency department and not merely a secondary diagnosis during a longer and more
418 serious hospital admission. Thus, we first filtered patients for our targets using the diagnosis table which contains all
419 recorded diagnostic International Statistical Classification of Diseases and Related Health Problems (ICD) codes. Then,
420 we manually checked the discharge diagnosis of each patient's discharge summary and only included those patients
421 whose very first primary diagnosis was one of our pathologies. If any other diagnosis was written in the discharge
422 diagnosis before one of our targets, the patient's case was removed from the dataset. If a patient was diagnosed with
423 more than one of the four pathologies included in our analysis, the patient was removed from the dataset.

424 After filtering for the appropriate pathologies, we split the discharge summary into its individual sections, extracting the
425 history of present illness and physical examination. First, we removed all patients where the pathology was mentioned in
426 the history of present illness as these admissions were mostly transfers where the diagnosis had already been established
427 and the hospital admission data was thus missing the initial emergency-department test results relevant for diagnostic
428 purposes. Furthermore, we removed all patients where no physical examination was included as this is a crucial source
429 of information according to the diagnostic guidelines of each pathology.

430 We gathered all laboratory and microbiology tests recorded during a patient's hospital admission and those up to one
431 day before admission. We included tests up to one day before admission as the MIMIC-IV documentation states that
432 there are millions of laboratory tests that are not associated with any hospital admission but can be joined to patient
433 stays using the patient's id, recorded test time, and hospital admission time. The tests before the official start of the
434 admission often had highly relevant values for diagnostic purposes and were thus included, though only if they were
435 not associated with any other hospital admission. If a laboratory test was recorded multiple times, we included only
436 the first entry in our dataset to simulate a therapy-naive diagnostic clinical-decision-making scenario. Furthermore,
437 we saved all reference ranges of the laboratory tests provided by MIMIC and established a comprehensive dictionary
438 mapping possible synonyms and abbreviations of tests to their original entry to be able to interpret all requests of the
439 LLMs for test results. This dictionary of synonyms was constantly expanded during initial testing of the models until
440 no unrecognized tests were recorded. The dictionary also contains common laboratory test panel names that map to a
441 list of the individual tests of that panel, such as complete blood count, basic metabolic panel, liver function panel, renal
442 function panel, and urinalysis, among others.

443 Similar to the laboratory data, many radiology reports were not associated with any hospital admission but their
444 timestamp was a few hours before the recorded start of the hospital admission. These often contained diagnosis-relevant
445 information and so we used the same 24 hour inclusion criteria as for the laboratory results and again allowed only those
446 exams not associated with any other hospital admission. Next, we established a list of uniquely identifying keywords
447 for each anatomical region and imaging modality. We used this list of keywords to determine the region and modality
448 of each included report from its MIMIC-IV provided exam name. Mappings were made for special exams such as CTU
449 to CT and MRCP to MRI, to provide them if, for example, a CT scan or MRI was requested, due to their low frequency.
450 We also used this list when interpreting the model requests for imaging information during evaluation. We manually
451 checked and adjusted the keywords until all reports in MIMIC-IV were correctly classified. Radiology reports were
452 split into report sections and only the *findings* section was included. This was done as many other sections such as
453 *conclusions* or *impressions* contained the diagnosis of the radiologist, which would have made the task trivial.

454 Finally, all procedures and operations performed during a patient's hospital stay were saved to understand which
455 patient-specific treatments were undertaken. The procedures in the MIMIC-IV procedures table saved as ICD9 and
456 ICD10 codes were extracted and combined with the free-text procedures section from each patient's discharge summary.
457 The free-text extraction from the discharge summary was required as many essential procedures, including surgeries,
458 were often not included in the procedures table.

459 A final round of data cleaning replaced any remaining mentions of the primary diagnosis with three underscores '___',
460 which is used by MIMIC-IV to censor data such as a patient's name or age. To increase data quality we excluded

461 patients that had no associated laboratory tests or for which no abdominal imaging was recorded. The final dataset,
462 MIMIC-CDM, contains 2400 unique patients presenting to the emergency department with one of the four target
463 pathologies (957 appendicitis, 648 cholecystitis, 257 diverticulitis, 538 pancreatitis) and whose makeup is detailed in
464 Figure 1a. The dataset contains physical examinations for all patients (2400), 138,788 laboratory results from 480
465 unique laboratory tests and 4403 microbiology results from 74 unique tests. Furthermore, MIMIC-CDM contains 5959
466 radiology reports, including 1836 abdominal CTs, 1728 chest x-rays, 1325 abdominal ultrasounds, 342 abdominal
467 x-rays and 227 MRCPs. Finally, there were 395 unique procedures recorded over all patients, with a total of 2917 ICD
468 procedures plus the 2400 free text procedures specified in the discharge summaries. Supplementary Table 2 shows the
469 age, sex and race statistics of the patients in the dataset split up by pathology. As the reports provided were de-identified,
470 the models did not have access to any of these characteristics during evaluation.

471 A second version of the dataset, which we call MIMIC-CDM-Full Information (MIMIC-CDM-FI), combines all the
472 information required for diagnosing each case and presents it all at once. Here we include the history of present illness,
473 physical examination, all abdominal imaging and all laboratory data helpful for both reaching the correct diagnosis and
474 ruling out differential diagnoses. To determine which laboratory data to include, we used the diagnostic guidelines of
475 each disease: appendicitis[16], cholecystitis[44], diverticulitis[25], and pancreatitis[32]. The specific tests included in
476 each category can be found in Supplementary Table 4. The information is presented in the order: history of present
477 illness, physical examination, laboratory results, imaging. The imaging is ordered by chart time from earliest to latest.

478 A subset of 80 representative patients of the MIMIC-CDM-FI dataset was randomly selected to be used for comparison
479 with physicians. The subset is evenly split between the four target pathologies with 20 patients randomly selected from
480 each pathology and matching the makeup of the full dataset, as shown in Supplementary Table 3. For the physicians,
481 the data was prepared as a PDF and the information was provided exactly in the same order and quantity as for the
482 models. Reference ranges were included when provided by MIMIC-IV. The abbreviations in the history of present
483 illness and physical examination were replaced with unabbreviated text for the German doctors, as they were unfamiliar
484 with US-specific abbreviations. The models performed worse with unabbreviated text (Supplementary Fig. 14). The
485 laboratory data were provided as a table in the PDF to increase readability. To mitigate the risk of physicians recognizing
486 the pattern of four distinct target pathologies, a further five patients each of patients presenting with gastritis, urinary
487 tract infaction, esophageal reflux and hernia were included. Thus, the final dataset used in the reader study spanned 100
488 patients, 80 of which are part of MIMIC-CDM.

489 6.2 Evaluation Framework

490 To realistically test the capabilities of LLMs on the task of clinical decision making, we simulated a clinical environment
491 in which a patient presents to the emergency department with acute abdominal pain and information must be iteratively
492 gathered before a final diagnosis is made. The LLM is tasked with the Clinical Decision Making (CDM) task
493 (Supplementary Materials B.1) which instructs it to consider a patient's symptoms and gather information to come
494 to a diagnosis and treatment plan while also explaining the two formats it should answer with. Both formats ask the
495 LLM to first think (i.e. consider the evidence, which has been shown to improve the quality of reasoning and future
496 actions[65]), and then either request more information or provide a diagnosis and treatment plan. If it chooses to
497 request more information, it must state 'Action:' followed by either 'Physical Examination', 'Laboratory Test', or
498 'Imaging'. Additionally, it must provide an 'Action Input' which specifies what information is desired from the action
499 (i.e. 'Complete Blood Count' or 'Abdominal Ultrasound'). The 'Action Input' field is ignored if a physical examination
500 is requested. The second format is to be used when the model decides enough information has been gathered for a
501 diagnosis, and asks the model to consider the evidence one last time and then provide a final diagnosis and treatment
502 plan.

503 The model is initially presented with these instructions and the history of present illness of the patient and then prompted
504 to record its 'Thoughts:', thus beginning the clinical decision making task. Outputs are generated until either a stop
505 token is reached or the model generates the 'Observation(s):' phrase, indicating that it has reached the end of its action
506 request and would potentially start hallucinating the result of its request. We stop model text generation here and then
507 examine the response of the model, extracting which action was desired and what the input to that action is. If the model
508 does not follow the instructions and, for example, writes 'Perform a physical examination' instead of 'Action: Physical
509 Examination', we still provide the appropriate information but record every instance of it not following instructions for
510 our evaluations (see Figure 6). We call these errors 'Next Action Errors'.

511 If the requested information is available for that patient case, we return it and prompt the model again to consider
512 the evidence. If the information is not available, we inform the model and ask for an alternative action. We return
513 only the laboratory tests and radiologist reports that were specifically requested. Laboratory tests are compared to the
514 previously mentioned dictionary of available tests to return the best match. If no match is found, 'N/A' is returned.
515 Requests for imaging have the exam modality and anatomical region extracted using the aforementioned keyword lists

(see Materials Section 6.1) and used to match against those saved for that patient in MIMIC-CDM. If multiple reports exist for a modality and region combination, we return the first report chronologically. The next request for an imaging examination of that modality and region will return the next report chronologically. Once there are no reports left to return, we inform the model that we can no longer provide reports of that modality and region combination. If a physical examination was requested, we return the entire physical examination regardless of any specifications made in the ‘action input’ field. We do this because we consider it best practice to perform a complete physical examination of a patient rather than only partially and reliably separating a physical examination report into its parts is difficult due to their free-form and heavily abbreviated style. If an invalid tool is requested (‘hallucinated’), we state that the tool does not exist and remind it which tools are available, or that it should make a diagnosis and provide a treatment plan. These occasions are also recorded as tool hallucinations for our evaluations (see Figure 6). An example exchange between an LLM and our framework can be seen in Figure 1b and Supplementary Section C.

We repeat this process, prompting the LLM to think and in turn receiving requests for information. Once the model decides that it has gathered sufficient information, it outputs its final diagnosis and treatment plan, ending the clinical decision making task for that patient. The final diagnosis is then evaluated to see if it contains the recorded pathology of the patient. In addition to a direct match of the pathology name (i.e. appendicitis, cholecystitis, diverticulitis, pancreatitis), we allow for a range of alternative phrasings as long as they are medically correct (see Supplementary Table 5). If multiple diagnoses are given, we only examine the first diagnosis mentioned. This is how we calculate the diagnostic accuracy for all analyses. A new instance of the task is then started for the next patient.

As LLMs can only take a limited amount of words as input, we monitor the length of the conversation and if we approach the input limit of the model, we ask it to summarize the information it has received so far to reduce the length of the conversation (Supplementary Materials B.2). We first summarize each gathered piece of information individually, leaving the initial history of present illness and instructions untouched. As LLMs have no memory and interpret each request independently, we replace the original pieces of information with the summaries. If we have summarized all steps of the interaction and still approach the limit of the model, we force a generation of diagnosis and treatment plan.

For the MIMIC-CDM-FI dataset we instruct the model to consider the facts of the case and then provide a diagnosis and only a diagnosis (Supplementary Materials B.3). As explained in Materials Section 6.1, the MIMIC-CDM-FI dataset includes the history of present illness, physical examination, all relevant laboratory results and every radiologist report where the abdominal region was inspected. Radiologist reports of other regions were not included due to the input length limits of the models. If including all of this information exceeds the input length of the model, we ask the LLM to summarize each radiologist report individually. If the input length is still exceeded, we ask the LLM to summarize all imaging information at once. In the rare cases where the input length continues to be exceeded, we remove words from the final imaging summary until there is enough space for a diagnosis (i.e. 25 tokens or 20 words).

548 6.3 Models

549 An overview of the models included and considered is given in Table 1.

550 When deciding which models to test, we started by only considering models with a context length of 4096 tokens due
551 to the large amounts of text contained within the MIMIC-CDM clinical cases. The context length defines how many
552 combined tokens a large language model can read and write. For example, if a model has a context length of 2048 and
553 receives an input of 2000 tokens, it can only generate 48 new tokens. A minimum length of 4096 is required, as the
554 average number of tokens of relevant information per patient case in MIMIC-CDM-FI is 2080 tokens with a maximum
555 count of 15,023 tokens. If one considers the extra tokens that are required for the information gathering back-and-forth
556 using MIMIC-CDM data, this quickly exceeds the limit of 2048 tokens of smaller models (context length windows
557 almost always differ in powers of 2).

558 Next, we considered which open access models performed best on medical reasoning tasks. To gauge model strength,
559 we used the MedQA[27] dataset as it is comprised of 12,723 questions from the USMLE and is thus a good gauge of
560 general medical knowledge contained within the model. As of the time of writing, Llama 2 is the leading open-access
561 model on the MedQA (USMLE) dataset, with the 70B model achieving a score of 58.4[12] exceeding that of GPT3.5
562 which scored only 53.6[38].

563 To effectively complete the clinical decision making task without specific fine-tuning to the task and format, the model
564 must be instruction fine-tuned. Instruction fine-tuning involves training a model to adapt to a wide range of new
565 tasks so that it can, with minimal instruction or example, complete an unseen task, like our clinical decision making
566 objective. The most popular and performant instruction fine-tuned versions of Llama 2 are Llama 2 Chat, fine-tuned by
567 Meta themselves; WizardLM, fine-tuned by Microsoft using evolutionary algorithm (Evo-Instruct) generated training
568 data; and OASST, fine-tuned using a crowd-sourced collection of 161,443 messages. Currently the only two existing
569 medically fine-tuned versions of Llama 2 with a context length of 4096 and 70B parameters are Clinical Camel and

570 Meditron. Neither has been extensively instruction fine-tuned and thus they both generated nonsensical and repetitive
571 responses to the clinical decision making objective using MIMIC-CDM data.

572 Currently, the most popular and leading closed-source LLMs for medical question answering are Chat-GPT (MedQA:
573 ~53.6[38]), GPT-4 (MedQA: 90.2[39]), Med-PaLM (MedQA: 67.2[50]), and Med-PaLM 2 (MedQA: 86.5[50]). As
574 previously stated, due to the signed data usage agreements of the MIMIC-IV database, the data cannot be sent to
575 external servers[19], precluding its use with closed-source models that are only accessible through an API and whose
576 models cannot be downloaded.

577 Furthermore, Chat-GPT is fine-tuned primarily through user conversations with the model, and since it is impossible
578 to know if portions of the MIMIC-IV database have already been used for queries by users less aware of the data
579 usage agreement[19], the data could already have been seen by the model during training, invalidating any results it
580 produces. Little to no information is known about the training data of GPT4, giving rise to analogous concerns about
581 its performance. While the exact pre-training data of Llama 2 is also not known, Meta has stated that it only used
582 'public available online data', which strongly mitigates the risk of MIMIC-IV data having being used. Med-PaLM and
583 Med-PaLM 2 achieve strong scores on MedQA but the exact data used for training are unknown, the models are only
584 accessible through an API, and access to the models is currently unavailable for all researchers. Repeated requests for
585 access were denied.

586 We strongly agree with current sentiment that open source models must drive progress in the field of medical AI due to
587 patient privacy and safety concerns, corporate lack of transparency, and the danger of unreliable external providers[57].
588 It is a serious risk to patient safety if key medical infrastructure is based on external company APIs and models whose
589 performance could change erratically with updates and which could generally be deactivated for any reason.

590 For each model tested, we downloaded and used the GPTQ quantized version from huggingface, the central repository
591 for all LLM models. GPTQ quantization reduces the numerical precision of the weights while monitoring the generated
592 output to reduce the GPU memory requirements of a model while not significantly degrading performance[20]. The
593 GPTQ parameters of the downloaded models were 4 bits, 32 group size, act order true, 0.1 damp% and 4096 sequence
594 length. This gives the highest possible inference quality while reducing model size to around 40 GB which can fit onto
595 a single A40 GPU. This reflects an economically realistic scenario of a single high-end GPU being used to host the
596 model to run the clinical decision making task. A fixed seed of 2023 and greedy decoding were used for all experiments
597 making all results deterministic and reproducible, except for the evaluation on the 80 patient subset for comparison with
598 clinicians where 10 different seeds were used for increased statistical power.

599 **6.4 Data and Code Availability**

600 The dataset is available to all researchers who create an account on <https://physionet.org/> and follow the steps to gain ac-
601 cess to the MIMIC-IV database (<https://physionet.org/content/mimiciv/2.2/>). Access is given after completing the "CITI
602 Data or Specimens Only Research" training course. The data use agreement of physionet for "credentialed health data"
603 must also be signed. The dataset can then be recreated using the code found at: [https://github.com/paulhager/MIMIC-
604 Clinical-Decision-Making-Dataset](https://github.com/paulhager/MIMIC-Clinical-Decision-Making-Dataset). The code to create the dataset uses python v3.10 and pandas v2.1.3.

605 The publication of the dataset on the Physionet website for those with access to MIMIC is currently under review. Once
606 it has been accepted, anyone credentialed to access MIMIC will be able to download the data directly.

607 The evaluation framework used for this study can be found at: [https://github.com/paulhager/MIMIC-Clinical-Decision-
608 Making-Framework](https://github.com/paulhager/MIMIC-Clinical-Decision-Making-Framework). The analysis framework to evaluate all results, generate all plots and do all statistical analysis can
609 be found at: <https://github.com/paulhager/MIMIC-Clinical-Decision-Making-Analysis>. All code uses python v3.10,
610 pytorch v2.1.1, transformers v4.35.2, spacy v3.4.4, langchain v0.0.339, optimum v1.14, thefuzz v0.20, exllamav2
611 v0.0.8, nltk v3.8.1, negspacy v1.0.4, scispacy v0.5.2

612 **6.5 Statistics**

613 All statistical tests were conducted using the Python programming language, version 3.10 and using the SciPy library.
614 Comparisons of means were tested for statistical significance using two-sided Student's *t*-tests with unequal variances
615 (tested through Bartlett's tests). To account for multiple comparisons, *p*-values were Bonferroni corrected with a
616 multiplier of 5 for the comparison of the doctors against the models and 3 for the comparison of the specialist and
617 generalist models.

Model	Base	Parameters	Training Dataset	Downloadable
Llama 2 Chat[58]	Llama 2[58]	70B	Public data*	✓
OASST[30]	Llama 2[58]	70B	Public data* Open Source Data	✓
WizardLM[63]	Llama 2[58]	70B	Public data* Evol-Instruct Generated[63]	✓
Clinical Camel[56]	Llama 2[58]	70B	Public data* ShareGPT PubMed Articles (pre 2021)[56] MedQA[27]	✓
Meditron[12]	Llama 2[58]	70B	Public data* Clinical Guidelines Public PubMed Abstracts[12] Public PubMed Papers[12] RedPajama[14]	✓
Chat-GPT[41]	GPT3.5[9]	???	User Conversations† Common Crawl[47] WebText2[45] Books1[29] Books2[29] Wikipedia	✗
GPT4[40]	???	???	???	✗
Med-PaLM[49]	Flan-PaLM[13]	540B	Webpages† Wikipedia† Social Media† Github† News Articles† Books† 473 Instruction fine-tuning Datasets[13] HealthSearchQA[49] MedicationQA[2] LiveQA[1]	✗
Med-PaLM 2[50]	PaLM 2[4]	340B	Web Documents† Books† Code† Mathematics† Conversational Data† MedQA[27] HealthSearchQA[49] MedicationQA[2] LiveQA[1]	✗

Table 1: An overview of the considered LLMs and their properties. Due to the data usage agreement of MIMIC-IV, only open access models that can be downloaded can be used with the data. Thus, only LLMs based on Llama 2 were used in this study.

* Meta defines ‘public data’ as a ‘mix of data from publicly available sources’.

† No further information provided.

??? indicates no information has been made public.

References

- [1] A. B. Abacha, E. Agichtein, Y. Pinter, and D. Demner-Fushman. Overview of the medical question answering task at trec 2017 liveqa. In *TREC*, pages 1–12, 2017.
- [2] A. B. Abacha, Y. Mrabet, M. Sharp, T. R. Goodwin, S. E. Shooshan, and D. Demner-Fushman. Bridging the gap between consumers’ medication questions and trusted answers. In *MedInfo*, pages 25–29, 2019.
- [3] L. Adlung, Y. Cohen, U. Mor, and E. Elinav. Machine learning in clinical decision making. *Med*, 2(6):642–665, 2021.
- [4] R. Anil, A. M. Dai, O. Firat, M. Johnson, D. Lepikhin, A. Passos, S. Shakeri, E. Taropa, P. Bailey, Z. Chen, et al. Palm 2 technical report. *arXiv preprint arXiv:2305.10403*, 2023.
- [5] Y. Bazi, M. M. A. Rahhal, L. Bashmal, and M. Zuair. Vision–language model for visual question answering in medical imagery. *Bioengineering*, 10(3):380, 2023.
- [6] A. Belyaeva, J. Cosentino, F. Hormozdiari, C. Y. McLean, and N. A. Furlotte. Multimodal llms for health grounded in individual-specific data. *arXiv preprint arXiv:2307.09018*, 2023.
- [7] S. Berman. Clinical decision making. In L. Bajaj, S. J. Hambidge, G. Kerby, and A.-C. Nyquist, editors, *Berman’s Pediatric Decision Making (Fifth Edition)*, pages 1–6. Mosby, fifth edition edition.
- [8] S. Biswas. Chatgpt and the future of medical writing, 2023.
- [9] T. Brown, B. Mann, N. Ryder, M. Subbiah, J. D. Kaplan, P. Dhariwal, A. Neelakantan, P. Shyam, G. Sastry, A. Askell, et al. Language models are few-shot learners. *Advances in neural information processing systems*, 33:1877–1901, 2020.
- [10] T. Buckley, J. A. Diao, A. Rodman, and A. K. Manrai. Accuracy of a vision-language model on challenging medical cases. *arXiv preprint arXiv:2311.05591*, 2023.
- [11] G. Cervellin, R. Mora, A. Ticinesi, T. Meschi, I. Comelli, F. Catena, and G. Lippi. Epidemiology and outcomes of acute abdominal pain in a large urban emergency department: retrospective analysis of 5,340 cases. 4(19):362–362.
- [12] Z. Chen, A. H. Cano, A. Romanou, A. Bonnet, K. Matoba, F. Salvi, M. Pagliardini, S. Fan, A. Köpf, A. Mhatahshami, A. Sallinen, A. Sakhaeirad, V. Swamy, I. Krawczuk, D. Bayazit, A. Marmet, S. Montariol, M.-A. Hartley, M. Jaggi, and A. Bosselut. Meditron-70b: Scaling medical pretraining for large language models, 2023.
- [13] H. W. Chung, L. Hou, S. Longpre, B. Zoph, Y. Tay, W. Fedus, Y. Li, X. Wang, M. Dehghani, S. Brahma, et al. Scaling instruction-finetuned language models. *arXiv preprint arXiv:2210.11416*, 2022.
- [14] T. Computer. Redpajama: an open dataset for training large language models, 2023.
- [15] S. Di Saverio, M. Podda, B. De Simone, M. Ceresoli, G. Augustin, A. Gori, M. Boermeester, M. Sartelli, F. Coccolini, A. Tarasconi, N. de’ Angelis, D. G. Weber, M. Tolonen, A. Birindelli, W. Biffi, E. E. Moore, M. Kelly, K. Soreide, J. Kashuk, R. Ten Broek, C. A. Gomes, M. Sugrue, R. J. Davies, D. Damaskos, A. Leppäniemi, A. Kirkpatrick, A. B. Peitzman, G. P. Fraga, R. V. Maier, R. Coimbra, M. Chiarugi, G. Sganga, A. Pisanu, G. L. de’ Angelis, E. Tan, H. Van Goor, F. Pata, I. Di Carlo, O. Chiara, A. Litvin, F. C. Campanile, B. Sakakushev, G. Tomadze, Z. Demetrashvili, R. Latifi, F. Abu-Zidan, O. Romeo, H. Segovia-Lohse, G. Baiocchi, D. Costa, S. Rizoli, Z. J. Balogh, C. Bendinelli, T. Scalea, R. Ivatury, G. Velmahos, R. Andersson, Y. Kluger, L. Ansaloni, and F. Catena. Diagnosis and treatment of acute appendicitis: 2020 update of the wses jerusalem guidelines. 15(1).
- [16] S. Di Saverio, M. Podda, B. De Simone, M. Ceresoli, G. Augustin, A. Gori, M. Boermeester, M. Sartelli, F. Coccolini, A. Tarasconi, et al. Diagnosis and treatment of acute appendicitis: 2020 update of the wses jerusalem guidelines. *World journal of emergency surgery*, 15:1–42, 2020.
- [17] N. Dziri, X. Lu, M. Sclar, X. L. Li, L. Jian, B. Y. Lin, P. West, C. Bhagavatula, R. L. Bras, J. D. Hwang, et al. Faith and fate: Limits of transformers on compositionality. *arXiv preprint arXiv:2305.18654*, 2023.
- [18] A. V. Eriksen, S. Möller, and J. Ryg. Use of gpt-4 to diagnose complex clinical cases. *NEJM AI*, 2023.
- [19] M. L. for Computational Physiology. Responsible use of mimic data with online services like gpt, 2023. Accessed on 16.01.2024.
- [20] E. Frantar, S. Ashkboos, T. Hoefler, and D. Alistarh. Gptq: Accurate post-training quantization for generative pre-trained transformers. *arXiv preprint arXiv:2210.17323*, 2022.
- [21] A. Gilson, C. W. Safranek, T. Huang, V. Socrates, L. Chi, R. A. Taylor, D. Chartash, et al. How does chatgpt perform on the united states medical licensing examination? the implications of large language models for medical education and knowledge assessment. *JMIR Medical Education*, 9(1):e45312, 2023.

-
- [22] A. L. Goldberger, L. A. Amaral, L. Glass, J. M. Hausdorff, P. C. Ivanov, R. G. Mark, J. E. Mietus, G. B. Moody, C.-K. Peng, and H. E. Stanley. Physiobank, physiobank, and physionet: components of a new research resource for complex physiologic signals. *circulation*, 101(23):e215–e220, 2000.
- [23] S. Golkar, M. Pettee, M. Eickenberg, A. Bietti, M. Cranmer, G. Krawezik, F. Lanusse, M. McCabe, R. Ohana, L. Parker, et al. xval: A continuous number encoding for large language models. *arXiv preprint arXiv:2310.02989*, 2023.
- [24] B. Haibe-Kains, G. A. Adam, A. Hosny, F. Khodakarami, M. A. Q. C. M. S. B. of Directors Shradha Thakkar 35 Kusko Rebecca 36 Sansone Susanna-Assunta 37 Tong Weida 35 Wolfinger Russ D. 38 Mason Christopher E. 39 Jones Wendell 40 Dopazo Joaquin 41 Furlanello Cesare 42, L. Waldron, B. Wang, C. McIntosh, A. Goldenberg, A. Kundaje, et al. Transparency and reproducibility in artificial intelligence. *Nature*, 586(7829):E14–E16, 2020.
- [25] J. Hall, K. Hardiman, S. Lee, A. Lightner, L. Stocchi, I. M. Paquette, S. R. Steele, D. L. Feingold, et al. The american society of colon and rectal surgeons clinical practice guidelines for the treatment of left-sided colonic diverticulitis. *Diseases of the Colon & Rectum*, 63(6):728–747, 2020.
- [26] D. Hendrycks, C. Burns, S. Basart, A. Zou, M. Mazeika, D. Song, and J. Steinhardt. Measuring massive multitask language understanding. *arXiv preprint arXiv:2009.03300*, 2020.
- [27] D. Jin, E. Pan, N. Oufattole, W.-H. Weng, H. Fang, and P. Szolovits. What disease does this patient have? a large-scale open domain question answering dataset from medical exams. *Applied Sciences*, 11(14):6421, 2021.
- [28] Z. Kanjee, B. Crowe, and A. Rodman. Accuracy of a generative artificial intelligence model in a complex diagnostic challenge. *JAMA*, 2023.
- [29] J. Kaplan, S. McCandlish, T. Henighan, T. B. Brown, B. Chess, R. Child, S. Gray, A. Radford, J. Wu, and D. Amodei. Scaling laws for neural language models. *arXiv preprint arXiv:2001.08361*, 2020.
- [30] A. Köpf, Y. Kilcher, D. von Rütte, S. Anagnostidis, Z.-R. Tam, K. Stevens, A. Barhoum, N. M. Duc, O. Stanley, R. Nagyfi, et al. Openassistant conversations—democratizing large language model alignment. *arXiv preprint arXiv:2304.07327*, 2023.
- [31] T. H. Kung, M. Cheatham, A. Medenilla, C. Sillos, L. De Leon, C. Elepaño, M. Madriaga, R. Aggabao, G. Diaz-Candido, J. Maningo, et al. Performance of chatgpt on usmle: Potential for ai-assisted medical education using large language models. *PLoS digital health*, 2(2):e0000198, 2023.
- [32] A. Leppäniemi, M. Tolonen, A. Tarasconi, H. Segovia-Lohse, E. Gamberini, A. W. Kirkpatrick, C. G. Ball, N. Parry, M. Sartelli, D. Wolbrink, et al. 2019 wses guidelines for the management of severe acute pancreatitis. *World journal of emergency surgery*, 14(1):1–20, 2019.
- [33] N. F. Liu, K. Lin, J. Hewitt, A. Paranjape, M. Bevilacqua, F. Petroni, and P. Liang. Lost in the middle: How language models use long contexts. *arXiv preprint arXiv:2307.03172*, 2023.
- [34] D. McDuff, M. Schaekermann, T. Tu, A. Palepu, A. Wang, J. Garrison, K. Singhal, Y. Sharma, S. Azizi, K. Kulka-rni, et al. Towards accurate differential diagnosis with large language models. *arXiv preprint arXiv:2312.00164*, 2023.
- [35] S. M. McKinney, M. Sieniek, V. Godbole, J. Godwin, N. Antropova, H. Ashrafiyan, T. Back, M. Chesus, G. S. Corrado, A. Darzi, et al. International evaluation of an ai system for breast cancer screening. *Nature*, 577(7788):89–94, 2020.
- [36] M. Moor, O. Banerjee, Z. S. H. Abad, H. M. Krumholz, J. Leskovec, E. J. Topol, and P. Rajpurkar. Foundation models for generalist medical artificial intelligence. *Nature*, 616(7956):259–265, 2023.
- [37] A. Nicolson, J. Dowling, and B. Koopman. Improving chest x-ray report generation by leveraging warm-starting. *arXiv preprint arXiv:2201.09405*, 2022.
- [38] H. Nori, N. King, S. M. McKinney, D. Carignan, and E. Horvitz. Capabilities of gpt-4 on medical challenge problems. *arXiv preprint arXiv:2303.13375*, 2023.
- [39] H. Nori, Y. T. Lee, S. Zhang, D. Carignan, R. Edgar, N. Fusi, N. King, J. Larson, Y. Li, W. Liu, et al. Can generalist foundation models outcompete special-purpose tuning? case study in medicine. *arXiv preprint arXiv:2311.16452*, 2023.
- [40] R. OpenAI. Gpt-4 technical report. *arXiv*, pages 2303–08774, 2023.
- [41] L. Ouyang, J. Wu, X. Jiang, D. Almeida, C. Wainwright, P. Mishkin, C. Zhang, S. Agarwal, K. Slama, A. Ray, et al. Training language models to follow instructions with human feedback. *Advances in Neural Information Processing Systems*, 35:27730–27744, 2022.

-
- [42] A. Pal, L. K. Umapathi, and M. Sankarasubbu. Medmcqa: A large-scale multi-subject multi-choice dataset for medical domain question answering. In *Conference on Health, Inference, and Learning*, pages 248–260. PMLR, 2022.
- [43] P. Pezeshkpour and E. Hruschka. Large language models sensitivity to the order of options in multiple-choice questions. *arXiv preprint arXiv:2308.11483*, 2023.
- [44] M. Pisano, N. Allievi, K. Gurusamy, G. Borzellino, S. Cimbanassi, D. Boerna, F. Coccolini, A. Tufo, M. Di Martino, J. Leung, et al. 2020 world society of emergency surgery updated guidelines for the diagnosis and treatment of acute calculus cholecystitis. *World journal of emergency surgery*, 15(1):1–26, 2020.
- [45] A. Radford, J. Wu, R. Child, D. Luan, D. Amodei, I. Sutskever, et al. Language models are unsupervised multitask learners. *OpenAI blog*, 1(8):9, 2019.
- [46] A. Rao, M. Pang, J. Kim, M. Kamineni, W. Lie, A. K. Prasad, A. Landman, K. Dreyer, and M. D. Succi. Assessing the utility of chatgpt throughout the entire clinical workflow: Development and usability study. *Journal of Medical Internet Research*, 25:e48659, 2023.
- [47] A. Roberts, C. Raffel, K. Lee, M. Matena, N. Shazeer, P. J. Liu, S. Narang, W. Li, and Y. Zhou. Exploring the limits of transfer learning with a unified text-to-text transformer. 2019.
- [48] E. H. Shortliffe and M. J. Sepúlveda. Clinical Decision Support in the Era of Artificial Intelligence. *JAMA*, 320(21):2199–2200, 12 2018.
- [49] K. Singhal, S. Azizi, T. Tu, S. S. Mahdavi, J. Wei, H. W. Chung, N. Scales, A. Tanwani, H. Cole-Lewis, S. Pfohl, et al. Large language models encode clinical knowledge. *Nature*, pages 1–9, 2023.
- [50] K. Singhal, T. Tu, J. Gottweis, R. Sayres, E. Wulczyn, L. Hou, K. Clark, S. Pfohl, H. Cole-Lewis, D. Neal, et al. Towards expert-level medical question answering with large language models. *arXiv preprint arXiv:2305.09617*, 2023.
- [51] A. Testolin. Can neural networks do arithmetic? a survey on the elementary numerical skills of state-of-the-art deep learning models. *arXiv preprint arXiv:2303.07735*, 2023.
- [52] A. Thawani, J. Pujara, F. Ilievski, and P. Szekely. Representing numbers in nlp: a survey and a vision. In *Proceedings of the 2021 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies*, pages 644–656, 2021.
- [53] A. J. Thirunavukarasu, R. Hassan, S. Mahmood, R. Sanghera, K. Barzangi, M. El Mukashfi, and S. Shah. Trialling a large language model (chatgpt) in general practice with the applied knowledge test: observational study demonstrating opportunities and limitations in primary care. *JMIR Medical Education*, 9(1):e46599, 2023.
- [54] A. J. Thirunavukarasu, D. S. J. Ting, K. Elangovan, L. Gutierrez, T. F. Tan, and D. S. W. Ting. Large language models in medicine. *Nature Medicine*, pages 1–11, 2023.
- [55] J. Tiffen, S. J. Corbridge, and L. Slimmer. Enhancing clinical decision making: Development of a contiguous definition and conceptual framework. *Journal of Professional Nursing*, 30(5):399–405, 2014.
- [56] A. Toma, P. R. Lawler, J. Ba, R. G. Krishnan, B. B. Rubin, and B. Wang. Clinical camel: An open-source expert-level medical language model with dialogue-based knowledge encoding. *arXiv preprint arXiv:2305.12031*, 2023.
- [57] A. Toma, S. Senkaiahliyan, P. R. Lawler, B. Rubin, and B. Wang. Generative ai could revolutionize health care—but not if control is ceded to big tech. *Nature*, 624(7990):36–38, 2023.
- [58] H. Touvron, L. Martin, K. Stone, P. Albert, A. Almahairi, Y. Babaei, N. Bashlykov, S. Batra, P. Bhargava, S. Bhosale, et al. Llama 2: Open foundation and fine-tuned chat models. *arXiv preprint arXiv:2307.09288*, 2023.
- [59] T. Tu, S. Azizi, D. Driess, M. Schaekermann, M. Amin, P.-C. Chang, A. Carroll, C. Lau, R. Tanno, I. Ktena, et al. Towards generalist biomedical ai. *arXiv preprint arXiv:2307.14334*, 2023.
- [60] T. van Sonsbeek, M. M. Derakhshani, I. Najdenkoska, C. G. Snoek, and M. Worring. Open-ended medical visual question answering through prefix tuning of language models. *arXiv preprint arXiv:2303.05977*, 2023.
- [61] D. Van Veen, C. Van Uden, M. Attias, A. Pareek, C. Bluethgen, M. Polacin, W. Chiu, J.-B. Delbrouck, J. M. Z. Chaves, C. P. Langlotz, et al. Radadapt: Radiology report summarization via lightweight domain adaptation of large language models. *arXiv preprint arXiv:2305.01146*, 2023.
- [62] Y. Wang and Y. Zhao. Tram: Benchmarking temporal reasoning for large language models. *arXiv preprint arXiv:2310.00835*, 2023.
- [63] C. Xu, Q. Sun, K. Zheng, X. Geng, P. Zhao, J. Feng, C. Tao, and D. Jiang. Wizardlm: Empowering large language models to follow complex instructions. *arXiv preprint arXiv:2304.12244*, 2023.

- [64] C. Yang, X. Wang, Y. Lu, H. Liu, Q. V. Le, D. Zhou, and X. Chen. Large language models as optimizers. *arXiv preprint arXiv:2309.03409*, 2023.
- [65] S. Yao, J. Zhao, D. Yu, N. Du, I. Shafran, K. Narasimhan, and Y. Cao. React: Synergizing reasoning and acting in language models. *arXiv preprint arXiv:2210.03629*, 2022.
- [66] C. Zheng, H. Zhou, F. Meng, J. Zhou, and M. Huang. On large language models' selection bias in multi-choice questions. *arXiv preprint arXiv:2309.03882*, 2023.
- [67] J. Zhou, T. Lu, S. Mishra, S. Brahma, S. Basu, Y. Luan, D. Zhou, and L. Hou. Instruction-following evaluation for large language models. *arXiv preprint arXiv:2311.07911*, 2023.

A Dataset Statistics

Disease	Median Age	Age Range	Gender Distribution	Race Distribution
Appendicitis	36.0	18-95	F (48.8%), M (51.2%)	WHITE (64.9%) BLACK (10.3%) HISPANIC (8.3%) ASIAN (8.0%) OTHER (8.5%)
Cholecystitis	59.0	18-95	F (55.8%), M (44.2%)	WHITE (63.8%) BLACK (15.1%) HISPANIC (9.2%) ASIAN (5.4%) OTHER (6.5%)
Pancreatitis	59.0	18-93	F (55.1%), M (44.9%)	WHITE (65.1%) BLACK (15.3%) HISPANIC (8.9%) ASIAN (2.6%) OTHER (8.1%)
Diverticulitis	60.0	18-91	F (58.9%), M (41.1%)	WHITE (72.1%) BLACK (10.5%) HISPANIC (5.8%) ASIAN (4.7%) OTHER (7.0%)

Table 2: Demographic Statistics of Patients with Different Diseases

Disease	Median Age	Age Range	Gender Distribution	Race Distribution
Appendicitis	38.5	20-75	F (45.0%), M (55.0%)	WHITE (65.0%) BLACK (15.0%) HISPANIC (5.0%) ASIAN (5.0%) OTHER (10.0%)
Cholecystitis	66.5	31-85	F (45.0%), M (55.0%)	WHITE (65.0%) BLACK (10.0%) HISPANIC (5.0%) ASIAN (5.0%) OTHER (15.0%)
Pancreatitis	52.5	20-82	F (60.0%), M (40.0%)	WHITE (50.0%) BLACK (10.0%) HISPANIC (15.0%) ASIAN (0.0%) OTHER (25.0%)
Diverticulitis	56.0	31-85	F (45.0%), M (55.0%)	WHITE (75.0%) BLACK (15.0%) HISPANIC (0.0%) ASIAN (5.0%) OTHER (5.0%)

Table 3: Demographic Statistics of Physician Comparison Subset of 80 patients

B Prompts

B.1 CDM Template

`{system_tag_start}`You are a medical artificial intelligence assistant. You give helpful, detailed and factually correct answers to the doctors questions to help him in his clinical duties. Your goal is to correctly diagnose the patient and provide treatment advice. You will consider information about a patient and provide a final diagnosis.

You can only respond with a single complete

Thought:

Action:

Action Input:

format OR a single

Thought:

Final Diagnosis:

Treatment:

format. Keep all reasoning in the Thought section. The Action, Action Input, Final Diagnosis, and Treatment sections should be direct and to the point. The results of the action will be returned directly after the Action Input field in the "Observation:" field.

Format 1:

Thought: (reflect on your progress and decide what to do next)

Action: (the action name, should be one of `[{tool_names}]`)

Action Input: (the input string to the action)

Observation: (the observation from the action will be returned here)

OR

Format 2:

Thought: (reflect on the gathered information and explain the reasoning for the final diagnosis)

Final Diagnosis: (the final diagnosis to the original case)

Treatment: (the treatment for the given diagnosis)

The tools you can use are:

Physical Examination: Perform physical examination of patient and receive the observations.

Laboratory Tests: Run specific laboratory tests and receive their values. The specific tests must be specified in the 'Action Input' field.

Imaging: Do specific imaging scans and receive the radiologist report.

Scan region AND modality must be specified in the 'Action Input'

field.`{add_tool_descr}{system_tag_end}{user_tag_start}{examples}`Consider the following case and come to a final diagnosis and treatment by thinking, planning, and using the aforementioned tools and format.

Patient History:

`{input}{user_tag_end}{ai_tag_start}`Thought:`{agent_scratchpad}`

B.2 CDM Observation Summarize Template

`{system_tag_start}`You are a medical artificial intelligence assistant. Your goal is to effectively, efficiently and accurately reduce text without inventing information. You want to return verbatim observations that are abnormal and of interest to a possible diagnosis of the patient. Normal observations can be combined. Do not invent information. Use medical abbreviations when possible to save characters. Put the most important information first.`{system_tag_end}{user_tag_start}`Please summarize the following result:
`{observation}{user_tag_end}{ai_tag_start}`
Summary:

B.3 CDM-FI Template

`{system_tag_start}`You are a medical artificial intelligence assistant. You directly diagnose patients based on the provided information to assist a doctor in his clinical duties. Your goal is to correctly diagnose the patient. Based on the provided information you will provide a final diagnosis of the most severe pathology. Don't write any further information. Give only a single diagnosis.`{system_tag_end}{fewshot_examples}{user_tag_start}`Provide the most likely final diagnosis of the following patient.

`{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}`Final Diagnosis:

B.4 Reference Range Test Zeroshot Template

`{system_tag_start}`You are a technical AI assistant working in a laboratory that handles tests for a hospital. You are good at interpreting numbers. You are responsible for reviewing the results of lab tests and determining whether they are Low, Normal, or High. You will be given the test, its value and then the reference range for that test, which will be written as "Reference Range [Lower Reference Range - Upper Reference Range]". You will write just one word, indicating if the test results are Low, Normal, or High. Do not write anything other than your one word

answer.{system_tag_end}{user_tag_start}{lab_test_string_rr}{user_tag_end}{ai_tag_start}Test Result:

B.5 CDM-FI No Final Template

{system_tag_start}You are a medical artificial intelligence assistant. You directly diagnose patients based on the provided information to assist a doctor in his clinical duties. Your goal is to correctly diagnose the patient. Based on the provided information you will provide the diagnosis. Don't write any further information. Give only a single diagnosis.{system_tag_end}{fewshot_examples}{user_tag_start}Provide the diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Diagnosis:""

B.6 CDM-FI Main Diagnosis Template

{system_tag_start}You are a medical artificial intelligence assistant. You directly diagnose patients based on the provided information to assist a doctor in his clinical duties. Your goal is to correctly diagnose the patient. Based on the provided information you will provide the main diagnosis. Don't write any further information. Give only a single diagnosis.{system_tag_end}{fewshot_examples}{user_tag_start}Provide the main diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Main Diagnosis:""

B.7 CDM-FI Primary Diagnosis Template

{system_tag_start}You are a medical artificial intelligence assistant. You directly diagnose patients based on the provided information to assist a doctor in his clinical duties. Your goal is to correctly diagnose the patient. Based on the provided information you will provide the primary diagnosis. Don't write any further information. Give only a single diagnosis.{system_tag_end}{fewshot_examples}{user_tag_start}Provide the primary diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Primary Diagnosis:""

B.8 CDM-FI No System Template

{system_tag_start}{system_tag_end}{fewshot_examples}{user_tag_start}Provide the most likely final diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Final Diagnosis:""

B.9 CDM-FI No User Template

{system_tag_start}You are a medical artificial intelligence assistant. You directly diagnose patients based on the provided information to assist a doctor in his clinical duties. Your goal is to correctly diagnose the patient. Based on the provided information you will provide a final diagnosis of the most severe pathology. Don't write any further information. Give only a single diagnosis.{system_tag_end}{fewshot_examples}{user_tag_start}{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Final Diagnosis:""

B.10 CDM-FI No Medical Template

{system_tag_start}You are an artificial intelligence assistant. You answer questions to the best of your abilities. Think hard about the following problem and then provide an answer.{system_tag_end}{fewshot_examples}{user_tag_start}Provide the most likely final diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Final Diagnosis:""

B.11 CDM-FI Serious Final Template

{system_tag_start}You are a medical artificial intelligence assistant. You directly diagnose patients based on the provided information to assist a doctor in his clinical duties. Your goal is to correctly diagnose the patient. Based on the provided information you will provide a final diagnosis of the most severe pathology. Don't write any further information. Give only a single diagnosis.{system_tag_end}{fewshot_examples}{user_tag_start}Provide the most serious final diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Final Diagnosis:""

B.12 CDM-FI Minimal System Template

{system_tag_start}You are a medical artificial intelligence assistant. You diagnose patients based on the provided information to assist a doctor in his clinical duties.{system_tag_end}{fewshot_examples}{user_tag_start}Provide the most likely final diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Final Diagnosis:""

B.13 CDM-FI No System No User Template

{system_tag_start}{system_tag_end}{fewshot_examples}{user_tag_start}{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}Final Diagnosis:""

B.14 CDM-FI No Diagnosis Prompt Template

{system_tag_start}You are a medical artificial intelligence assistant. You directly diagnose patients based on the provided information to assist a doctor in his clinical duties. Your goal is to correctly diagnose the patient. Based on the provided information you will provide a final diagnosis of the most severe pathology. Don't write any further information. Give only a single diagnosis.{system_tag_end}{fewshot_examples}{user_tag_start}Provide the most likely final diagnosis of the following patient.

{input}{diagnostic_criteria}{user_tag_end}{ai_tag_start}""

C Example Exchange using Synthetic Data

In this section we show an example exchange between our simulated clinical environment provided by our evaluation framework using the MIMIC-CDM dataset and the WizardLM model. The first prompt is the system prompt given by the framework to the model explaining the task and its options. The rest of the exchange pulls (synthetic) data from MIMIC-CDM and records the LLM's response. Seen in this exchange is an example of the model requesting an imaging modality that is not available in the dataset and also hallucinating a surgery tool that does not exist. Upon reminding it of the available tools it correctly diagnoses the patient. This example exchange uses synthetic data to further protect patient privacy.

Evaluation Framework using MIMIC-CDM

You are a medical artificial intelligence assistant. You give helpful, detailed and factually correct answers to the doctors questions to help him in his clinical duties. Your goal is to correctly diagnose the patient and provide treatment advice. You will consider information about a patient and provide a final diagnosis.

You can only respond with a single complete

Thought:

Action:

Action Input:

format OR a single

Thought:

Final Diagnosis:

Treatment:

format. Keep all reasoning in the Thought section. The Action, Action Input, Final Diagnosis, and Treatment sections should be direct and to the point. The results of the action will be returned directly after the Action Input field in the "Observation:" field.

Format 1:

Thought: (reflect on your progress and decide what to do next)

Action: (the action name, should be one of [Physical Examination, Laboratory Tests, Imaging])

Action Input: (the input string to the action)

Observation: (the observation from the action will be returned here)

OR

Format 2:

Thought: (reflect on the gathered information and explain the reasoning for the final diagnosis)

Final Diagnosis: (the final diagnosis to the original case)

Treatment: (the treatment for the given diagnosis)

The tools you can use are:

Physical Examination: Perform physical examination of patient and receive the observations.

Laboratory Tests: Run specific laboratory tests and receive their values. The specific tests must be specified in the 'Action Input' field.

Imaging: Do specific imaging scans and receive the radiologist report. Scan region AND modality must be specified in the 'Action Input' field. Consider the following case and come to a final diagnosis and treatment by thinking, planning, and using the aforementioned tools and format.

Evaluation Framework using MIMIC-CDM

Patient History:

___ presenting with 3 days since acute onset of pain in RLQ. Pain started ___ afternoon but worsened significantly overnight. Intense episodes woke her from sleep. Patient proceeds to experience nausea and chills. No vomiting. Patient endorses anorexia. Past Medical History: Anorexia, Anxiety. Family History: Non-contributory.

LLM

Thought: Given the acute onset of right lower quadrant pain, along with anorexia and right-sided flank pain, there could be several possibilities such as appendicitis, kidney stones, or inflammatory bowel disease like Crohn's disease. However, since the patient does not have any past medical history or family history of these conditions, it might be less likely. Therefore, we need to perform further investigations to determine the exact cause of the pain.

Action: Physical Examination

Action Input: N/A

Evaluation Framework using MIMIC-CDM

Observation: Physical Examination:

Temp: 97.8 HR: 44 BP: 104/69 RR: 17 100% Ra Gen: NAD HEENT: No scleral icterus, mucus membranes moist CV: RRR no m,r,g RESP: CTABL Abd: soft, tender to palpation inRLQ, non distended, no rebound or guarding Ext: wwp4, palpable distal pulses

LLM

Thought: After performing the physical examination, the vital signs appear normal except for a slightly elevated blood pressure. Findings of tenderness to palpation in the right lower quadrant of the abdomen supports the forementioned possibilities of the patient's complaints, such as appendicitis, kidney stones, urinary tract infection, ovarian cysts or inflammatory bowel disease. We need to investigate further to rule out potential causes.

Action: Laboratory Tests

Action Input: Complete Blood Count (CBC), Liver Function Tests (LFTs), Renal panel, Urinalysis

Evaluation Framework using MIMIC-CDM

Observation: Laboratory Tests:

(Blood) Red Blood Cells: 4.39 m/uL
(Blood) White Blood Cells : 16.8 K/uL
(Blood) Hemoglobin: 12.4 g/dL
(Blood) Hematocrit: 38.8 %
(Blood) MCV: 85.4 fL
(Blood) MCH: 27.3 pg
(Blood) MCHC: 33.4 %
(Blood) Platelet Count: 226.8 K/uL
(Blood) Lymphocytes: 22.0 %
(Blood) Basophils: 0.5 %
(Blood) Eosinophils: 1.9 %
(Blood) Monocytes: 5.5 %
(Blood) Neutrophils: 69.8 %
(Blood) RDW: 13.0 %
(Blood) Alanine Aminotransferase (ALT): 18.1 IU/L
(Blood) Aspartate Aminotransferase (AST): 29.1 IU/L
(Blood) Alkaline Phosphatase: 68.8 IU/L
(Blood) Bilirubin, Total: 0.3 mg/dL
(Blood) PT: 10.5 sec
(Blood) INR(PT): 0.9
(Blood) Albumin: 5.2 g/dL
(Blood) Creatinine: 0.6 mg/dl
(Urine) Ketones: NEG
(Urine) Urine Color: Yellow.
(Urine) Urine Appearance: Clear.
(Urine) Specific Gravity: 1.0
(Urine) Protein: NEG.
(Urine) pH: 5.8 units
(Urine) Bilirubin: NEG.
(Urine) Glucose: NEG.
(Urine) Urobilinogen: NEG.
(Urine) Ketone: TR.
(Urine) Nitrite: NEG.
(Urine) Leukocytes: NEG.

LLM

Thought: The laboratory test results show normal values for most parameters analyzed. However, the patients has elevated white blood cells as a marker for inflammation. Based on the diagnostic test results so far, the patient is suspected to have an inflammation in the abdomen, such as appendicitis or inflammatory bowel disease.

Action: Imaging

Action Input: Ultrasound of the Right Lower Quadrant

Evaluation Framework using MIMIC-CDM

Observation: Imaging:

Abdomen Ultrasound: Not available. Try a different imaging modality.

LLM

Thought: Since ultrasound is not suitable for this case, we should proceed with another imaging modality. Given the possibility of hepatobiliary involvement, a computed tomography (CT) scan would be more appropriate to visualize the internal organs better.

Action: Imaging

Action Input: Computed Tomography (CT) scan of the Abdom

Evaluation Framework using MIMIC-CDM

Observation: Imaging:

Abdomen CT: EXAMINATION:

CT ABD AND PELVIS WITH CONTRAST:

TECHNIQUE:

MDCT imaging, extending from the lung bases to the pubic symphysis, was performed post-administration of 130cc intravenous contrast. Axial scans were analyzed alongside coronal and sagittal reformatted images.

FINDINGS:

CHEST:

Lung bases are clear without pleural or pericardial effusions.

ABDOMEN:

Liver and gallbladder present normally; portal and hepatic veins are patent. Pancreas and spleen are normal; bilateral adrenal glands are unremarkable. Kidneys show symmetrical enhancement without hydronephrosis.

Stomach and small bowel maintain normal caliber. A hyperdense, dilated structure, likely the appendix, arises from the cecal base, measuring up to 10 mm and exhibiting mucosal hyperenhancement compared to adjacent bowel. Predominant features include hyperenhancement and wall thickening, with less emphasis on fluid content. Inflammatory changes are evident at the cecal base, around the appendix, especially at its indistinct tip.

Adjacent to the ascending colon's mesentery, two enlarged lymph nodes, 1.5 cm and 1.9 cm in size, show brisk enhancement. The smaller node contains central low-density material. Both nodes are surrounded by marked fat stranding and suspected necrotic or suppurative changes.

The ascending, transverse, and descending colon appear normal. No mesenteric free air is observed.

PELVIS:

Urinary bladder and adnexa are unremarkable. Uterus contains an intrauterine device in standard position. A small volume of simple free fluid is present in the pelvis, likely physiologic.

VESSELS:

Aorta and its major branches are normal in caliber and patent.

OSSEOUS STRUCTURES:

No focal lytic or sclerotic lesions suggestive of malignancy are identified.

LLM

Thought: The CT scan supports the diagnosis of acute appendicitis due to the enlarged and inflamed appendix observed. I will recommend surgical intervention for removal of the affected organ.

Action: Surgery

Action Input: Appendectomy

Evaluation Framework using MIMIC-CDM

Observation: Provide a diagnosis and treatment OR a valid tool. That is not a valid tool, try one of [Physical Examination, Laboratory Tests, Imaging].

LLM

Thought:

The CT scan revealed an inflamed and enlarged appendix, indicating acute appendicitis as the most likely cause of the patient's pain. Additionally, there are two enlarged and briskly enhancing lymph nodes in the mesentery, which could suggest a secondary infectious process or inflammatory condition. The presence of a small amount of simple free fluid in the pelvis might be due to the inflammatory response.

Final Diagnosis: Acute Appendicitis

Treatment: Surgical intervention - Appendectomy

D Laboratory Test Categories

Pathologies	Category	Items
Appendicitis Cholecystitis Diverticulitis Pancreatitis	Inflammation	White Blood Cells WBC Count C-Reactive Protein High-Sensitivity CRP
Cholecystitis	Gallbladder	Bilirubin Gamma Glutamyltransferase
Cholecystitis	Liver	Alanine Aminotransferase (ALT) Aspartate Aminotransferase (AST)
Pancreatitis	Pancreas	Amylase Lipase
Pancreatitis	Pancreatitis Severity	Hematocrit Urea Nitrogen Triglycerides Calcium Potassium

Table 4: The individual laboratory test category items defined using the guidelines for appendicitis[16], cholecystitis[44], diverticulitis [25], and pancreatitis[32]. At least one of these tests must have been ordered either individually or as part of a panel for the category to be counted.

E Diagnosis Definitions

Pathology	Location	Modifiers
appendicitis	appendi	gangren infect inflam abscess rupture necros perf
cholecystitis	gallbladder	gangren infect inflam abscess necros perf
cholecystitis	cholangitis	cholangitis
cholecystitis	acute gallbladder	disease attack
cholecystitis	acute biliary	colic
diverticulitis	diverticul	inflam infect abscess perf rupture
diverticulitis	acute colonic	perfor
diverticulitis	sigmoid	perfor colitis
pancreatitis	pancrea	gangren infect inflam abscess necros

Table 5: For each pathology, alternative diagnostic phrasings were accepted. If within one diagnostic phrase (delineated through e.g. periods, commas, ‘vs’, or other separators) the location and modifier occurred without negation, the diagnosis was marked as correct. For example, for cholecystitis a diagnosis of ‘Perforated gallbladder’ or ‘Gallbladder infection’ were both accepted. A diagnosis of ‘Gallbladder disease vs perforated appendix’ would not be accepted. Shortened substrings of the locations and modifiers were used to allow for alternative endings to match such as both ‘An *infect* of the gallbladder’ and ‘*Infected* gallbladder’, or ‘*Periappendicular* abscess’ and ‘Perforated *appendix*’. Manual controls were done to verify the specificity and sensitivity of the definitions.

F Adherence to Diagnostic Guidelines




Figure 11: Llama 2 Chat was the only LLM that consistently requested physical examinations. (Late) Physical Examinations counted the physical examination if it wasn't the first information requested.




Figure 12: The first imaging modality requested by the LLMs and the attending doctors in the MIMIC dataset are shown. LLMs sometimes follow diagnostic guidelines concerning imaging but often diagnose without requesting any imaging at all. As we show that imaging is the most useful diagnostic tool for all LLMs for each pathology except pancreatitis, this could be partly responsible for their low diagnostic accuracy. The legend specifies the colors of the imaging modalities and the patterns of the models.

G FI Performance




Figure 13: LLMs perform better when all information is given, especially on pathologies with strong indications such as appendicitis (dilated appendix described in radiologist report) and pancreatitis (elevated pancreatic enzymes listed in laboratory test results).

H LLMs Diagnostic Accuracy Without Medical Abbreviations




Figure 14: Diagnostic accuracy on the clinician subset of MIMIC-CDM-FI stays the same when medical abbreviations are written out.

I LLMs are sensitive to information order

Changing the order of the presented information changes diagnostic accuracy. Crucially, the best order is disease specific, meaning a clinician must deliver a preliminary diagnosis to ensure proper model performance, eliminating many of the benefits of an AI clinical decision making system. Best order for each pathology is in bold.

I.1 Llama 2 Chat

Experiment	Imaging, Lab Physical	Imaging, Physical Lab	Lab, Imaging Physical	Lab, Physical Imaging	Physical, Imaging Lab	Physical, Lab Imaging
Pathology						
Appendicitis	95.72	95.72	95.61	94.88	95.40	94.98
Cholecystitis	39.04	40.43	46.76	48.30	41.05	64.35
Diverticulitis	45.91	46.30	41.63	42.41	47.08	49.03
Pancreatitis	60.41	65.99	53.16	50.37	63.01	48.14
Mean	60.27	62.11	59.29	58.99	61.64	64.13

I.2 OASST

Experiment Pathology	Imaging, Lab Physical	Imaging, Physical Lab	Lab, Imaging Physical	Lab, Physical Imaging	Physical, Imaging Lab	Physical, Lab Imaging
Appendicitis	97.18	97.60	96.97	96.97	97.60	97.39
Cholecystitis	69.29	67.28	64.51	61.57	65.59	64.66
Diverticulitis	48.64	48.64	41.63	47.08	46.69	47.86
Pancreatitis	64.13	64.13	65.80	65.06	65.43	60.22
Mean	69.81	69.41	67.23	67.67	68.83	67.53

I.3 WizardLM

Experiment Pathology	Imaging, Lab Physical	Imaging, Physical Lab	Lab, Imaging Physical	Lab, Physical Imaging	Physical, Imaging Lab	Physical, Lab Imaging
Appendicitis	96.97	96.76	96.55	96.34	97.18	96.66
Cholecystitis	40.74	37.19	41.82	37.96	36.42	51.39
Diverticulitis	51.36	52.53	54.09	53.70	53.70	52.92
Pancreatitis	78.81	77.88	75.84	75.28	78.07	69.14
Mean	66.97	66.09	67.07	65.82	66.34	67.53

J LLMs Cannot Interpret Numbers




Figure 15: When providing a laboratory test result and its reference range, LLMs are incapable of consistently interpreting the result as normal, low or high.

K LLMs Struggle to Follow Instructions

Error Name	Error Example	Corrected Example
Next Action Error	Order labs	Action: Laboratory Tests
	Perform a physical examination	Action: Physical Examination
	Run a CT scan of the abdomen	Action: Imaging Action Input: Abdominal CT
Tool Hallucination	Action: Consult specialist	
	Action: Consider the evidence	
Diagnosis Error	Final Diagnosis: Based on the patient's symptoms, laboratory results, and imaging studies, the final diagnosis is: Appendicitis	Final Diagnosis: Appendicitis
	Final Diagnosis: Acute Gallstone Pancreatitis. The patient's symptoms of abdominal pain, nausea, vomiting, and fever, along with the laboratory results of elevated serum amylase and lipase levels, suggest a diagnosis of acute pancreatitis.	Final Diagnosis: Acute Gallstone Pancreatitis
	Final Diagnosis: Appendicitis, abdominal pain, and elevated white blood cells	Final Diagnosis: Appendicitis

Table 6: Examples of the types of errors commonly made by models when providing actions and diagnoses. The corrected example in the desired format is also provided. Note that tool hallucination examples are simply not valid actions and so there are no corrected examples provided.

L LLMs are Sensitive to the Order of Information




Figure 16: By mixing the order in which information is presented to LLMs, their diagnostic accuracy changes despite the information included staying the same. This places an unnecessary burden upon clinicians who would need to consider and monitor the order in which they feed the models with information. The value above each whisker shows the difference between the best performing and worst performing order.

M Summarizing Progress Improves CDM Diagnostic Accuracy




Figure 17: When an LLM approaches its input limit, we ask it to summarize the information gathered thus far to allow for more context. Increased input sizes allows it more opportunities to ask for information and increases the chances of requesting information that is important for the diagnosis.