1 2	Yellow fever in Ghana: Predicting emergence and ecology from historical outbreaks
3	Seth D. Judson ^{1*} , Ernest Kenu ² , Trevon Fuller ³ , Franklin Asiedu-Bekoe ⁴ , Alberta Biritwum-
4	Nyarko ⁵ , Lee F. Schroeder ⁶ , David W. Dowdy ⁷
5	¹ Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland,
6	USA
7	² Department of Epidemiology, University of Ghana School of Public Health, Accra, Ghana
8	³ Institute of the Environment and Sustainability, University of California Los Angeles, USA
9	⁴ Disease Surveillance Department, Ghana Health Service, Accra, Ghana
10	⁵ Policy, Planning, Monitoring & Evaluation Division, Ghana Health Service, Accra, Ghana
11	⁶ Department of Pathology, University of Michigan, Ann Arbor, Michigan, USA
12	⁷ Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore,
13	Maryland, USA
14	
15	*Corresponding author:
16	Seth D. Judson MD ORCID: 0000-0001-9911-7839
17	Email: sjudson4@jh.edu
18	
19	
20	
21	
22	

23 Abstract

24 Understanding the epidemiology and ecology of yellow fever in endemic regions is 25 critical for preventing future outbreaks. Ghana is a high-risk country for yellow fever. In this 26 study we estimate the epidemiology, ecological cycles, and areas at risk for yellow fever in 27 Ghana based on historical outbreaks. We identify 2371 cases and 887 deaths (case fatality rate 28 37.4%) from yellow fever reported in Ghana from 1910 to 2022. Since implementation of routine 29 childhood vaccination in 1992, the estimated mean annual number of cases decreased by 81% 30 and the geographic distribution of yellow fever cases also changed. While there have been 31 multiple large historical outbreaks of yellow fever in Ghana from the urban cycle, recent 32 outbreaks have originated among unvaccinated nomadic groups in rural areas with the 33 sylvatic/savanna cycles. Using machine learning and an ecological niche modeling framework, 34 we predict areas in Ghana that are similar to where prior yellow fever outbreaks have originated 35 based on temperature, precipitation, landcover, elevation, and human population density. We 36 find differences in predictions depending on the ecological cycles of outbreaks. Ultimately, these 37 findings and methods could be used to inform further subnational risk assessments for yellow 38 fever in Ghana and other high-risk countries.

39

40 Author Summary

41 Yellow fever is a viral hemorrhagic fever transmitted by mosquitoes in Africa and South 42 America through different ecological transmission cycles. While West Africa has had the most 43 cases of yellow fever, less is known about the epidemiology and ecology of yellow fever among 44 countries in this region. Ghana has had multiple yellow fever outbreaks, including a recent 45 outbreak in 2021-2022. In this study we estimate cases and deaths due to yellow fever in Ghana,

compare the ecological cycles of outbreaks, and predict future areas at risk based on prior yellow
fever cases and environmental conditions. We find that the populations at risk for yellow fever in
Ghana have changed over the past century and that different ecological factors influence the risk
of future emergence. Understanding these changes and the nuances of yellow fever epidemiology
and ecology within countries will be important for future outbreak preparedness.

51

52 Introduction

53 Yellow fever (YF) outbreaks continue to occur in Africa and South America despite an 54 effective vaccine having been created nearly a century ago. Yellow fever is a viral hemorrhagic 55 fever caused by yellow fever virus (YFV), a mosquito-borne flavivirus. In Africa, YFV 56 circulates through three transmission cycles. In the sylvatic cycle, forest-dwelling *Aedes* spp. 57 mosquitoes infect non-human primates (NHPs) and intermittently transmit YFV to humans after feeding on infected NHPs [1]. In the savanna or intermediate cycle, humans are infected by 58 59 mosquitoes in forest border areas, and there is human-to-human and NHP-to-human transmission 60 via mosquito vectors [1]. In the urban cycle, there is only transmission among humans from the 61 anthropophilic mosquito Aedes aegypti, which breeds in water-containing vessels in urban sites 62 [1]. While all three transmission cycles cause human cases of YF in Africa, it remains unknown how each cycle contributes to the burden of YF, limiting targeted interventions and predictive 63 models [2,3]. 64

Among infectious diseases in the WHO Africa Region (AFRO), YF caused the fifth most
outbreaks in 2020 [4]. And while there are more cases of YF in West Africa compared to
anywhere else in the world, a recent systematic review found only 12 studies of YF incidence
and mortality within African countries [5]. Compounding the limited knowledge of the historical

burden of YF, there are gaps in predicting future emergence of YF within African countries, and
it is unknown how each of the three transmission cycles (sylvatic, savanna, and urban) contribute
to YF outbreaks.

72 There is a contrasting situation in South America, where all outbreaks of YF in Brazil 73 have been attributed to the sylvatic cycle since 1942 and there are known seasonal and climatic 74 drivers of outbreaks [6]. In addition, some species of New World NHPs develop lethal infections 75 due to YFV, and the finding of moribund or dead NHPs can indicate an epizootic outbreak, 76 signaling impending sylvatic YF outbreaks among humans [1]. In contrast, most YFV infections 77 among Old World NHPs in Africa are believed to be inapparent or subclinical, and therefore 78 outbreaks from sylvatic and savanna transmission occur without alerting indicators in Africa. 79 Initial sylvatic or savanna outbreaks could lead to epidemics if the urban cycle subsequently 80 becomes established. Therefore, understanding the historical epidemiology and ecology of YF is crucial to predicting future emergence among YF-endemic African nations. 81 82 There is also concern for YF outbreaks within African countries to cause epidemics. In 83 2016 a YF outbreak in Angola spread to neighboring countries, and the epidemic response 84 required more than 28 million YF vaccines, exhausting the global YF vaccine supply [7]. In 85 response to this YF epidemic, the WHO, GAVI, and UNICEF developed the Eliminate Yellow Fever Epidemics (EYE) strategy with the aim of eliminating yellow fever epidemics by 2026 [7]. 86 87 The core objectives of the EYE strategy are to (1) protect at-risk populations, (2) prevent 88 international spread, and (3) contain outbreaks rapidly [8]. Central to achieving these objectives is understanding the regions and populations at greatest risk for YF outbreaks. Ghana is among 89 90 the 27 countries identified by EYE as being high-risk for YF and has had multiple YF outbreaks.

91 Understanding the contexts of YF outbreaks in Ghana and identifying areas at future risk 92 could inform resource allocation for vaccination campaigns, vector control, and diagnostic 93 testing. Modeling tools could aid risk assessment for YF by predicting areas at risk based on 94 prior outbreaks. For zoonotic and vector-borne diseases like YF, the locations of cases, vectors, 95 or hosts can be used to create spatial models which can be translated into risk maps [9]. For 96 example, researchers have predicted the local risk for sylvatic YF in Brazil from models based 97 on human and NHP cases as well as ecological factors [10,11]. Through combining epidemiological data from historical YF outbreaks with local 98 99 ecological knowledge, we aim to predict the habitats and factors associated with YF in Ghana. 100 Because there are limited data on the multiple *Aedes* spp. and NHPs that may propagate YFV circulation in Ghana, we use a machine learning approach incorporating confirmed human YF 101 102 cases as inputs and multiple abiotic and biotic covariates as explanatory variables. Ultimately, 103 our overall goals are to (1) review the historical epidemiology of YF outbreaks in Ghana, (2) 104 assess contributions of different YF ecological cycles to these outbreaks, and (3) identify areas at 105 risk for YF emergence based on this historical and ecological understanding. 106

- 107 Methods
- 108 Setting

109 The Republic of Ghana is a country in West Africa on the Gulf of Guinea, bordered by 110 Côte d'Ivoire, Burkina Faso, and Togo. Ghana consisted of 10 regions until December 2018 111 when the Brong-Ahafo, Northern, Volta, and Western regions were split to make a total of 16 112 regions. For the purposes of this study, we used the current geographic boundaries of the 16 113 regions and 260 districts in Ghana. For outbreaks prior to 2019 with available geographic

114	information, we re-classified these locations to align with the current terminology for regions and
115	districts. We also compared outbreak locations by the five recently proposed agro-climatic zones
116	for Ghana, which are named from north to south: Sudan Savannah, Guinea Savannah, Transition
117	Zone, Forest, and Coastal [12]. These zones encompass geographic regions with similar
118	temperature and precipitation patterns [12].
119	The Government of Ghana introduced the YF vaccine as part of the Expanded Program
120	on Immunization (EPI) in 1992, making it a routine immunization for children in Ghana.
121	Although the WHO estimates that 95% of the population in Ghana has been vaccinated against
122	YF [13], Ghana had a recent YF outbreak from October 2021 to February 2022 with 70
123	confirmed cases and 35 deaths [14]. The origin and epicenter of the outbreak was in the West
124	Gonja district of the Savannah region, among unvaccinated nomadic populations, from which the
125	outbreak spread to other regions [15,16]. We reviewed the epidemiology of this latest outbreak
126	and prior outbreaks using the methods described below.
127	
128	Identifying YF Outbreaks in Ghana
129	To identify YF outbreaks in Ghana and the locations of corresponding cases, we searched
130	PubMed, EBSCO, Google Scholar, and Scopus for literature using the search terms "yellow
131	fever" AND "Ghana." We also searched the ProMed archives using the same search terms. To
132	find WHO YF reports, we used the WHO's Institutional Repository for Information Sharing
133	(IRIS) and queried the Medical Subject Heading (MeSH) "yellow fever", which included the
134	Disease Outbreak News (DON) and Weekly Epidemiology Record (WER). This also included a
135	2000 WHO report which included YF case reporting among countries from 1950-1998 [17]. We
136	also reviewed additional WHO documents, including a 2005 report on YF in Ghana [18] which

137	was referenced in a review of arboviruses in Ghana [19]. Finally, we reviewed publicly available
138	online reports from the Ghana Health Service (GHS) and the Ghana Weekly Epidemiological
139	Report. The literature search was conducted up to October 20th 2023. We collated all of the data
140	from the aforementioned sources into a dataset of reported annual YF cases, deaths, and case
141	fatality rates (CFR) for individuals in Ghana, as well as a dataset of outbreaks in Ghana
142	(locations, month of outbreak onset, and reactive vaccination campaigns) [20].
143	
144	Estimating Annual YF Cases and Outbreak Characteristics
145	For the period from 1910-1950, we extracted YF cases, deaths, and outbreak
146	characteristics primarily from a single source (Scott 1965) [21]. For 1950-2022, we compared
147	YF cases and deaths between multiple sources, including the primary literature and WHO
148	reports. If there was a discrepancy in annual cases/deaths, we used the total cases/deaths reported
149	in the primary literature, given delays between reporting cases and confirmatory testing.
150	We excluded cases from 2003 which were reported only in the WHO 2005 report [18]
151	because we could not find these cases referenced in another source. We were also unable to find
152	deaths reported for 31 cases in 2012 [22]. If the source distinguished between reported and
153	confirmed cases, as well as diagnostics used, we also included this information in our
154	dataset. For our calculations of disease burden and CFR, we used overall reported cases and
155	deaths.
156	
157	Categorizing YF Outbreaks According to Likely Transmission Cycles
158	For outbreaks with sufficient epidemiological information (including location details,
159	diagnostic method, and vector surveillance), we categorized the outbreaks as likely occurring due

160 to the urban, sylvatic, or savanna cycles [20]. If vector surveillance identified sufficient A. 161 aegypti based on the indices described below, or if the outbreak was concentrated around urban 162 dwellings or structures (i.e. wells or water storage containers) then these outbreaks were 163 attributed to the urban cycle. Outbreaks where vector surveillance revealed insufficient A. 164 *aegypti* and/or cases originated in forest or savanna habitat were attributed to the sylvatic or 165 savanna cycle. Given limitations in distinguishing between the sylvatic and savanna cycles, these 166 were considered together for our modeling analyses. 167 For categorizing urban YF, we considered the following vector indices: (1) house index (% of houses with at least one positive breeding place). (2) container index (% containers with A. 168 169 *aegypti* larvae), and (3) Breteau index (# of positive larval breeding places per 100 houses) [23]. There is a high risk for urban YF when the house index > 35, container index > 20, and Breteau 170 171 index > 50 [23]. If the Breteau index is between 5 and 50, then there is considered to be 172 sufficient A. aegypti to cause an outbreak. There is unlikely to be urban transmission of 173 YFV when the house index <4, container index <3, and Breteau index <5 [23]. 174 175 Georeferencing YF Occurrences 176 To obtain occurrence data for our models, we identified the locations of laboratory 177 confirmed human YF cases in Ghana. A panel of YF experts had previously determined that one 178 of the most important risk factors for YF in the African region was confirmed YF cases since

179 1960 [24]. Therefore, we included only confirmed cases since 1960 for our models. We

180 georeferenced each case to the town of residence; if this was unknown then we used the location

181 of the healthcare facility where the patient was evaluated. To improve precision, we excluded

cases that were only referenced at the region or district level. Our methods for georeferencing arefurther described in S1 text.

184 To mitigate sampling bias, we treated multiple cases at the same location in a year as a 185 single occurrence. We also excluded locations that where within 1 km of another occurrence, 186 given our spatial resolution of covariates at 30 arc seconds ($\sim 1 \text{ km}^2$). We categorized the 187 transmission cycles of occurrences as described above. 188 Through this process we identified 21 occurrences for YF in Ghana from 1960-2022. We 189 categorized six of these occurrences as being due to the urban cycle given positive vector indices 190 for A. aegypti. Additional details on the occurrences are available in the online dataset and S1 191 text. To compare the regional distribution of YF cases as well as districts where YF outbreaks 192 have originated, we created maps in ArcGIS Pro (https://www.esri.com).

193

194 Modeling Habitat Suitability for YF Emergence

195 To identify suitable habitats for YF emergence, we built on a previously described 196 approach that is optimized for small sample sizes to identify habitat suitability for pathogen 197 spillover [25]. We also used a similar methodology and covariates as those that were recently 198 used to model YF emergence in Brazil [11]. We used Maximum Entropy Species Distribution 199 Modeling (Maxent) version 3.3.3 [26], a machine learning algorithm commonly used in species 200 distribution modeling and ecological niche modeling, to predict the distribution of a species 201 based on presence locations as input and multiple ecological covariates. We chose Maxent given 202 that it has outperformed other modeling algorithms for small sample sizes with presence-only 203 data, including in environments in Africa [27]. Our aim was to model the environments suitable 204 for YF emergence in Ghana, as well as to explore the relative importance of different abiotic,

biotic, and human covariates. Therefore, instead of making the stringent assumptions to interpret
Maxent output as an ecological niche, we instead built our models to indicate habitat suitability
[28].

208 We modeled YF habitat suitability as a function of multiple covariates (Table 1). These

209 covariates were identified as factors that could influence the ecological cycles of YF and

210 included (1) abiotic covariates: 19 bioclimatic variables and elevation (WorldClim 2.1,

211 https://www.worldclim.org) [29], (2) biotic covariates: landcover (MODIS MCD12Q1,

212 https://www.earthdata.nasa.gov) [30] and NHP species richness

213 (https://sedac.ciesin.columbia.edu/data/set/species-global-mammal-richness-2015) [31], and (3)

214 human population density (https://sedac.ciesin.columbia.edu/data/set/gpw-v4-population-

215 density-adjusted-to-2015-unwpp-country-totals-rev11) [32].

216 To identify potential NHPs that could propagate the sylvatic/savanna cycles, we

217 referenced the IUCN Red List (https://www.iucnredlist.org/) and identified 17 species of NHPs

in Ghana. Of these species, six had epidemiological and laboratory evidence of YF infection at

219 the species level (*Cercopithecus mona*, *Colobus vellerosus*, *Erythrocebus patas*, *Pan troglodytes*,

220 Papio anubis, Perodicticus potto), and three had evidence at the genus level (Cercopithecus

221 lowei, Cercopithecus petaurista, and Cercopithecus roloway) [24]. Eight of these nine species

are members of the Cercopithecidae family. The exception was *P. troglodytes*, which is not a

common species in Ghana and is unlikely to have a significant role in YF ecology [24].

224 Therefore, we evaluated species richness of Old World monkeys (Cercopithecidae) as a

covariate. Further details about the covariates are available in S1 text.

226

Covariate	Details	Resolution	Source	
19 bioclimatic	Gridded climate data 1970- 30 arc seconds/1		Worldolim 2.1	
variables	2000	km ²		
	Derived from the Shuttle	30 arc seconds/1	Worldclim 2.1	
Elevation	Radar Tomography	km ²		
	(SRTM) elevation data	KIII		
Species richness of	Gridded values of the		Global Mammal	
Old World	number of species of	30 arc seconds/1	Richness Grids, 2015	
monkeys	Cercopithecidae from the	km ²	Release (2013)	
(Cercopithecidae)	IUCN Red list		(SEDAC)	
	Moderate Resolution			
Londoovor	Imaging Spectroradiometer	500 m resolution	MODIS MCD12Q1	
Lanucover	(MODIS) Land Cover	re-scaled to 1 km	NASA	
	from 2021-2023			
Unman nonviotion	2020 LIN adjusted	20 are accords/1	UN WPP-Adjusted	
density	2020 ON adjusted	June 2	population density	
density	population density	KIN ²	v4.11 (SEDAC)	

Table 1. Explanatory covariates for YF habitat suitability models

228

We created one set of models with all YF occurrences undifferentiated by ecological cycle, to predict areas with habitat suitability for YF irrespective of cycle. We created another set of models which excluded the six urban occurrences that had confirmed positive *A. aegypti* vector indices. For our model parameters, we used a leave-one out cross-validation (LOOCV) method that is optimized for small sample sizes [25,33]. To make our models comparable to recent Maxent models for YF from Brazil, we used similar Maxent parameters and methods for selecting covariates [11].

We ran our initial models in Maxent with all covariates, using a jackknife procedure to evaluate relative covariate importance. For our initial models, we used the default Maxent settings and additionally selected random seed, replicated run type as cross validate, replicates equal to the sample size, and set the maximum iterations to 100,000. To identify collinearity among covariates, we used a correlation matrix in ArcGIS Pro. We removed covariates that had >70% correlation as well as those with less than 5% percent contribution and permutation

importance. We used both of these metrics and the jackknife test to compare the relative 242 243 importance of covariates. Percent contribution was deemed less informative than the other 244 metrics since it is determined heuristically and depends on the sequence of covariates [34]. 245 Following this exclusion of potentially less-relevant covariates, we re-ran the models 246 using the same LOOCV approach and settings. We used the cumulative output of the Maxent 247 models as a representation of habitat suitability, where the value of each cell is the sum of all raw 248 values less than or equal to the value for that location and rescaled from 0 to 100 [28]. The 249 cumulative output can be used to create binary maps and determine omission rates for the 250 models. For the binary maps and omission rates we used the minimum training presence 251 cumulative threshold, which corresponds to the minimum habitat suitability in the training data. 252 To compare the relative performance of the Maxent models we used the area under the receiver 253 operating characteristic curve (AUC), which ranges from 0 to 1 and reflects the probability that 254 randomly chosen presence locations are ranked higher than background points [28]. 255 256 **Results** 257 We identified 2371 cases and 887 deaths due to YF in Ghana from 1910 to 2022. The 258 overall mean CFR was 37.4%. The first definitive outbreak of YF in Ghana occurred in 1910. 259 From 1910-1960 there were 569 cases and 335 deaths (CFR 58.9%). In 1945 YF vaccination 260 became required for foreigners in Ghana, and the first reactive mass vaccination campaign for 261 YF in Ghana occurred in 1951. We identified 12 reactive vaccination campaigns from 1951-2022 262 (Fig S1). 263 We successfully identified the regions in which 1722/1802 (95.6%) of reported cases 264 from 1960 to 2022 occurred (Fig 1). From 1960-1992 there were 1511 cases and 478 deaths

265 (CFR 31.6%) that occurred in 11 of Ghana's 16 regions. The greatest number of YF cases were

266	reported in Upper West (n=384), Volta (n=340), Eastern (n=250), Upper East (n=163), Savannah

- 267 (n=124), and Bono (n=105) regions. Following implementation of routine childhood YF
- vaccination in 1992, the mean annual number of YF cases fell by 81%. From 1992-2022 there
- were 291 cases and 70 deaths (CFR 24.1%) with the highest number of cases in the Upper West
- 270 (n = 135), Savannah (n = 45), and Upper East (n=33) regions and few cases in the southern
- 271 regions. The districts where YF outbreaks were suspected to originate in Ghana based on
- 272 primary/index cases are shown in Fig 2.
- 273

274 Fig 1. Yellow Fever Cases Reported by Region in Ghana

275 Reported YF cases that could be located on a regional level in Ghana are shown for the periods
276 before and after routine childhood YF vaccination, 1960-1992 and 1993-2022. The regions of

historical cases have been updated to match the current terminology for the 16 regions in Ghana.

278

279 Fig 2. Yellow Fever Outbreak Origins by District in Ghana

280 The districts where reported YF outbreaks were suspected to originate in Ghana based on

primary/index cases are shown for the periods before and after routine childhood YF vaccination,

282 1960-1992 and 1993-2022. Also shown is Mole National Park and the West Gonja District

where the 2021 and 1983 outbreaks were suspected to originate. For the purposes of this figure,

284 outbreaks are defined as more than one case of YF.

285

We identified certain outbreaks and cases that could be categorized as likely due to urban, savanna, and sylvatic cycles in Ghana (Table 2). Of eleven outbreaks with available epidemiological data that occurred between 1910 and 1979, we estimated that nine likely

- corresponded to the urban cycle. One outbreak (1983) had evidence of starting in the
- sylvatic/savanna cycles and then spreading through the urban cycle, and at least two recent
- 291 outbreaks appear to have originated via the sylvatic/savanna cycles. In addition to these
- 292 outbreaks there have been sporadic individual cases likely from the sylvatic/savanna cycles.
- 293

Table 2. Select YF outbreaks and cases in Ghana with epidemiologic details from 1910-2022*

Year	Region	Initial location	Agro-climatic zone	Suspected Cycle	Details	Ref
1910	Western	Sekondi	Coastal	urban	cases in confined area near commercial waterfront/dwellings	[21]
1926	Eastern	Asamankese	Forest	urban	outbreak confined within town, "aedes index" 89%	[21]
1927	Greater Accra	Accra	Coastal	urban	outbreak around particular dwellings	[21]
1937	Greater Accra	Accra	Coastal	urban	localized outbreak around a well with <i>A. aegypti</i>	[21]
1955	Bono East	Kintampo	Transition	sylvatic/savanna	no <i>A. aegypti</i> found, <i>A. africanus</i> found to be widespread	[35]
1969	Northern	Pong- Tamale	Guinea Savannah	sylvatic/savanna	only 2/246 houses with <i>A. aegypti</i> larvae	[36]
1969	Upper East	Bolgatanga	Sudan Savannah	urban	house index 11%	[36]
1970	Eastern	Akim- Manso, Asikasu, Akwatia	Forest	urban	<i>A. aegypti</i> found breeding in all towns, house index as high as 50%	[36]
1977	Upper West	Jirapa	Sudan Savannah	urban	house index 9.1%, container index 5.9%, Breteau 14%	[23]
1978	Eastern	Maase	Forest	urban	house index 36.4%, container index 38%, Bretau index 96%	[23]
1978	Volta	Hohoe, Kpandu	Forest/Guinea Savannah	urban	house index 4-16%, container index 3- 7%, Bretau index 4- 16%	[23]
1983	Savannah	Damongo	Guinea Savannah	sylvatic/savanna to urban	laboratory confirmed deaths of baboons at Mole reserve, subsequent urban spread	[37]
2006	Bono East	Kountaya village	Guinea Savannah	sylvatic/savanna	single case in village near forest	[38]
2011	Upper	Kassena-	Sudan	sylvatic/savanna	index case went to a	[39]

	East	Nankana	Savannah		farm in a forest	
		West			border	
2021	Savannah	West Gonja	Guinea Savannah	sylvatic/savanna	initial cases in savanna habitat	[15]
*Additional categorized outbreaks and cases in online dataset [20]						

294

295 The covariates selected for the final models and their relative contributions are shown in 296 Table 3. The average habitat suitability models for YF in Ghana are shown in Fig 3. The 297 minimum training presence threshold was used to translate these models into the binary maps in 298 Fig 4. The overall YF model (AUC 0.846) included landcover, precipitation of the driest quarter 299 (BIO17), human population density, annual mean temperature (BIO1), and elevation. The model 300 that excluded confirmed urban YF occurrences (AUC 0.880) also included landcover and 301 precipitation of driest quarter (BIO17), as well as precipitation of wettest quarter (BIO16) and 302 mean temperature of the driest quarter (BIO9). The most important type of landcover based on 303 the response curves for the models was dense herbaceous vegetation. Human population density 304 had the highest permutation importance and jackknife variable importance for the overall YF 305 model. Precipitation of the driest quarter (BIO17) had the highest permutation importance and 306 jackknife variable importance for the model excluding urban YF. The overall YF model 307 predicted 20/21 (95%) of occurrences tested at the minimum training presence threshold. The 308 model excluding urban YF predicted 14/15 (93%) of occurrences at the same threshold. The only 309 location that was not predicted by both models at the minimum training presence threshold was a 310 single case of likely sylvatic YF in Sene district of the Bono East region. Both models identified 311 regions and districts with recent YF outbreaks as having high suitability including Upper West, 312 Upper East, North East, Savannah, Northern, Bono, and Bono East regions. The overall YF 313 model identified all regions in Ghana as having areas of suitable habitat for YF.

314 Fig 3. Predicted Habitat Suitability for YF in Ghana

315	The left panel shows the average Maxent cumulative output for models including 21 YF
316	occurrences in Ghana and the following covariates: human population density, elevation,
317	landcover, precipitation of driest quarter (BIO17), and annual mean temperature (BIO 1). The
318	right panel shows the average output of the models excluding 6 confirmed urban YF occurrences
319	and includes the following covariates: precipitation of driest quarter (BIO17), precipitation of
320	wettest quarter (BIO16), landcover, and mean temperature of driest quarter (BIO9).
321	
322	Fig 4. Predicted Habitat Suitability for YF in Ghana based on the Minimum Training
323	Presence Threshold
323 324	Presence Threshold These binary habitat suitability maps represent areas greater or equal to the minimum training
323 324 325	Presence Threshold These binary habitat suitability maps represent areas greater or equal to the minimum training presence threshold (MTP) for each group of YF models. The MTP is defined as the lowest
323 324 325 326	Presence ThresholdThese binary habitat suitability maps represent areas greater or equal to the minimum trainingpresence threshold (MTP) for each group of YF models. The MTP is defined as the lowestpredicted habitat suitability value for the occurrences in the training data. Therefore, the
323 324 325 326 327	Presence ThresholdThese binary habitat suitability maps represent areas greater or equal to the minimum trainingpresence threshold (MTP) for each group of YF models. The MTP is defined as the lowestpredicted habitat suitability value for the occurrences in the training data. Therefore, thehighlighted areas above the MTP represent locations with ecological conditions at least as
323 324 325 326 327 328	Presence ThresholdThese binary habitat suitability maps represent areas greater or equal to the minimum trainingpresence threshold (MTP) for each group of YF models. The MTP is defined as the lowestpredicted habitat suitability value for the occurrences in the training data. Therefore, thehighlighted areas above the MTP represent locations with ecological conditions at least asfavorable as the location with the least suitable known YF occurrence in Ghana.
323 324 325 326 327 328 329	Presence Threshold These binary habitat suitability maps represent areas greater or equal to the minimum training presence threshold (MTP) for each group of YF models. The MTP is defined as the lowest predicted habitat suitability value for the occurrences in the training data. Therefore, the highlighted areas above the MTP represent locations with ecological conditions at least as favorable as the location with the least suitable known YF occurrence in Ghana.

Table 3	Contribution	of final	covariates to	YF	habitat	suitability	models
						~	

Model	Covariate	Percent Contribution*	Permutation Importance [†]		
	BIO17 (Precipitation of Driest Quarter)	29.4	63.8		
FI	BIO16 (Precipitation of Wettest Quarter)	11.2	26.5		
Excluding Urban VE	Landcover	50.5	9.3		
UIDAII IF	BIO9 (Mean Temperature of Driest Quarter)	8.9	0.3		
	Human Population Density	46.3	35.1		
	Elevation	15	23		
Overall YF	Landcover	25.5	14.8		
	BIO17 (Precipitation of Driest Quarter)	6.1	14.5		
	BIO1 (Annual Mean Temperature)	7.1	12.6		
* Total model gain attributed to particular covariate, determined heuristically					

[†]The importance of each covariate calculated from randomly changing values of a covariate and calculating decrease in model performance. The average performance decrease for each covariate is normalized, resulting in percentages.

331

332 Discussion

333 Our analysis estimates the historical epidemiology and ecology of YF in Ghana as well as 334 areas at potential risk for future emergence of YF. We found that the geographic distribution and 335 disease burden of YF in Ghana have changed since the introduction of routine childhood 336 vaccination in 1992. While many historical outbreaks have occurred through the urban cycle, 337 recent outbreaks appear to have originated from the sylvatic or savanna cycles. Therefore, there 338 appears to be a higher risk for YF outbreaks originating among unvaccinated populations in 339 environments where the appropriate vectors, NHPs, and human hosts are present to maintain the 340 sylvatic or savanna cycles. Using machine learning we identified these potential suitable habitats 341 on a more granular scale in Ghana as well as factors that should be considered when predicting 342 YF based on different transmission cycles.

The overall mean CFR we calculated for YF in Ghana (37.4%) was similar to the mean 343 344 CFR from a recent meta-analysis of global YF cases (39%) and severe cases in African countries 345 (36%) [5]. Given that there is limited surveillance and diagnostic capacity for YF in Ghana, our 346 data likely underestimate of the true burden of YF (while overestimating the true CFR) given 347 that mild or subclinical infections may not be detected. The YF cases in our study more likely 348 reflect severe YF, and it is estimated that $\sim 15\%$ of YF cases are from severe YF. If only severe 349 YF cases were captured in our dataset, this would suggest that over 15,000 cases of YF have 350 occurred in Ghana since 1910 – a number that may be even higher given historical and current 351 limitations in diagnosing YF. Both the number of cases and the case-fatality of YF have declined 352 in Ghana since 1992, likely in part due to introduction of routine childhood YF vaccination in

1992. Nevertheless, YF outbreaks continue to occur, highlighting the need for ongoing riskprediction and surveillance.

355 The geographic distribution, epidemiology, and ecology of YF outbreaks in Ghana has 356 changed substantially over the past century. During the first half of the 20th century, the majority 357 of YF outbreaks were detected along coastal regions and periodically in the north, with large 358 outbreaks occurring roughly every 10 years [21]. Relatively higher population density and the 359 presence of susceptible foreign populations likely contributed to these urban YF outbreaks. 360 Detection bias also likely contributed to these observations given differences in accessibility to 361 diagnostics and reporting. The requirement for YF immunization among foreigners in 1945, local 362 reactive vaccination campaigns, and increasing diagnostic recognition likely contributed to the 363 changing distribution of YF outbreaks with greater recognition of outbreaks in more remote 364 northern regions. In the context of increasing vaccination coverage for YF in Ghana, recent 365 outbreaks have originated among unvaccinated populations in rural areas and border districts in 366 the Guinea Savannah and Sudan Savannah agro-climatic zones. These areas may have more 367 nomadic pastoralist communities and as well as immigrants from neighboring countries, 368 potentially contributing to lower vaccination coverage. Recent YF outbreaks appear to reflect the 369 sylvatic and savanna cycles, which could transition into the urban cycle if there are sufficient 370 susceptible hosts. We found at least one large historical outbreak that began in the 371 sylvatic/savanna cycle and then spread to multiple other regions through the urban cycle. Climate 372 change, urbanization, and changing human population movements could fuel the spread of 373 sylvatic/savanna YF to urban outbreaks.

The Upper West, Savannah, and Upper East regions have experienced the majority of YF cases in outbreaks since routine immunization. Our models identified these regions as well as

376 neighboring regions (Upper East, North East, Northern, Bono, and Bono East) as being highly 377 suitable for YF emergence. The West Gonja district where the 2021 YF outbreak originated is 378 located near Mole National Park (Figure 2), which is the largest protected area in Ghana and is 379 likely suitable habitat for the NHPs and mosquitoes that could maintain the sylvatic and savanna 380 cycles of YF. Mole National Park is also a tourist site where nomadic populations may work and 381 also has surrounding pastoralist communities [40]. The 1983 YF outbreak was also believed to 382 have originated from this area given the finding of laboratory confirmed deaths of baboons with 383 YF [37]. Reviewing the locations of historical YF emergence, we found additional locations with 384 recurring cases, which could also be hotspots for YF. At least four outbreaks appear to have 385 originated in the Jirapa district in the Upper West and three in Bawku Municipal district in the Upper East. During the 2021 outbreak there were two cases from Tinga village in the Bole 386 387 region, and given the distance from the epicenter of the outbreak it is possible that this could 388 have been a separate YF emergence event. This village also had clinically diagnosed cases of YF 389 in 1983. While there have been isolated cases of sylvatic/savanna YF since routine 390 immunization, with index cases exposed to forest-savanna ecotones, these cases did not lead to 391 sustained outbreaks. In contrast, a significant unvaccinated population likely contributed to the 392 2021-2022 outbreak. Therefore, identifying both susceptible populations as well as the habitat 393 for YF emergence will be crucial for predicting areas at future risk.

We used confirmed human cases and machine learning to identify likely suitable habitats for the emergence of YF in Ghana. The models do not represent the distribution or ecological niche of YFV, but instead represent habitats with similar conditions to where prior YF cases occurred. Given that YF can start in the sylvatic/savanna cycles and then spread through urban cycle, we also created a set of models excluding confirmed urban YF cases to provide a better

399 prediction of habitats where outbreaks may originate. This model is a proxy for the habitats 400 similar to where savanna/sylvatic cases have occurred, which appears to be the likely origin of 401 recent and future YF outbreaks in Ghana. Because we used the location of residence or 402 healthcare facility for our occurrences, the habitats identified by this model do not reflect where 403 YFV circulates in the savanna/sylvatic cycle, but instead habitats similar to where human cases 404 from these cycles have been detected.

405 To our knowledge, this is the first time that the different ecological cycles of YF have 406 been integrated into spatial models in Africa. Overall these models illustrate the important 407 differences in habitats and locations where YF may occur in Ghana. For example, while 408 landcover, precipitation, and temperature are important covariates in both sets of models, human 409 population density was the most important covariate for models that included outbreaks from the 410 urban cycle– reflecting the fact that the urban cycle requires a higher density of susceptible 411 human hosts compared to the sylvatic/savanna cycles. In contrast, precipitation of the driest 412 quarter was the most important covariate for the model that excluded urban YF. Predictive 413 models that do not distinguish between the ecological cycles of YF will risk misidentifying areas 414 at risk for emergence. For instance, our models excluding urban cases reflect habitat similar to 415 where the recent YF outbreaks have originated in Ghana and could be a better predictor for 416 where future outbreaks may originate. In contrast, the overall model is more appropriate for 417 predicting areas where YF may spread through the urban cycle. Therefore, it is important that 418 researchers think carefully about the goals and inferences of their models based on YF ecology 419 and epidemiology.

420 There are multiple limitations to our analysis. We used reported YF cases from the421 literature and online reports, and there were relatively few laboratory confirmed historical cases.

422 For our habitat suitability models, we used strict inclusion criteria of confirmed cases since 1960 423 with georeferenced locations, limiting our analysis to 21 total occurrences. We used AUC values 424 to compare the relative performance between models, but AUC values can be inflated in 425 presence-only models such as Maxent. We attempted to minimize this inflation and reduce 426 overfitting by using the LOOCV approach and a small number of covariates. Another limitation 427 was sparse epidemiologic and entomologic surveillance data for most outbreaks. Vector indices 428 were only available for some of the occurrences, so we had to make assumptions about certain 429 outbreaks that lacked entomologic data, and we only excluded confirmed urban YF locations in 430 our second set of models. Similarly, it is possible that both urban and savanna/sylvatic cycles 431 may occur together during outbreaks. For example, while the 2021 YF outbreak may have originated in the savanna/sylvatic cycle, it could have spread through dwellings via the urban 432 433 cycle. We categorized this outbreak and similar outbreaks as originating from the 434 savanna/sylvatic cycle since our goal was to model habitat for YF emergence. The alternative 435 explanation, that this YF outbreak originated from the urban cycle of another outbreak, seemed 436 less likely given lack of nearby preceding outbreaks. Overall, additional vector surveillance data 437 are needed for further clarity. Lastly, the covariates we chose for our models were intended to 438 reflect habitat suitability via presence of the appropriate climatic conditions, environments, and 439 hosts. To make models more reflective of relative risk for future outbreaks, they would need to 440 include additional explanatory variables, most importantly YF vaccination coverage. 441 Despite these limitations, our study has important implications for preparing for future 442 outbreaks of YF in Ghana. For example, our models could be used to inform further risk

444 committee of YF experts developed a WHO protocol for national risk assessment for YF [24].

assessment which could guide diagnostic testing, vector control, and vaccination campaigns. A

443

445	However, the strategies in this protocol are resource intensive and involve sampling humans,
446	vectors, and NHPs for YF. Our methods for estimating habitat suitability based on historical
447	cases could be used to prioritize field surveillance for YF with the WHO risk assessment
448	protocol [24]. Our findings also have important implications for models and risk maps for YF.
449	While predictive models for YF risk have been made on a global scale [41–43], these models
450	either contain sparse data from Ghana or do not predict subnational risk for YF. Additionally,
451	these existing models do not distinguish between the different ecological cycles of YF in Africa,
452	which may influence prediction and interpretation. Therefore, additional subnational data and
453	local ecological knowledge are needed to create future risk maps for YF and other arboviruses if
454	they are to be informative for national policymakers [9].

455 While the habitats suitable for YF emergence appear to cluster in the northern regions of 456 Ghana, currently all YF diagnostic testing is done at the National Public Health Reference 457 Laboratory in Korle Bu in the south of the country. Expanding YF diagnostic testing to the Zonal 458 Public Health Laboratory in Tamale, which is close to recent YF outbreaks and predicted habitat 459 suitability, could improve early detection of outbreaks. While the burden of YF has decreased 460 since implementation of routine childhood vaccination in 1992, the recent outbreak has also 461 demonstrated the risk for future outbreaks among unvaccinated groups. Identifying unvaccinated 462 nomadic populations in habitats that are suitable for YF emergence will be an important step for 463 preventing future outbreaks. Overall, despite a century of research on YF, critical questions 464 remain regarding the ecology, epidemiology, and emergence of YF in African countries. 465 Addressing these research gaps and using this knowledge to inform public health interventions will be essential to improving health equity and preventing future YF epidemics. 466

467

468 Funding

- 469 This work was supported by the National Institute of Health T32 AI007291-32 to SDJ as well as
- 470 R01AI136977 to DWD, LFS, FA, and EK. The content is solely the responsibility of the authors
- 471 and does not necessarily represent the official views of the National Institutes of Health.
- 472

473 Acknowledgements

- 474 The authors would like to thank Dr. Eric Agboli for sharing the 2005 WHO YF in Ghana report
- and Dr. Sarah Louise Poynton for editing advice. They would also like to thank their colleagues
- 476 at the Ghana Health Service, University of Ghana, and Noguchi Memorial Institute for Medical
- 477 Research.
- 478

479 Author contributions

- 480 Conceptualization, Data curation, Analysis, Writing-original Draft: SDJ
- 481 Methodology: SDJ, TF
- 482 Writing-reviewing and editing: SDJ, DWD, LFS, TF, FA, EK, AB
- 483 Funding: DWD, LFS, FA, EK

484

485 **References**

- Monath TP, Vasconcelos PFC. Yellow fever. Journal of Clinical Virology. 2015;64: 160–
 173. doi:10.1016/J.JCV.2014.08.030
- Gaythorpe KAM, Jean K, Cibrelus L, Garske T. Quantifying model evidence for yellow
 fever transmission routes in Africa. PLoS Comput Biol. 2019;15: e1007355.
 doi:10.1371/journal.pcbi.1007355
- Jean K, Hamlet A, Benzler J, Cibrelus L, Gaythorpe KAM, Sall A, et al. Eliminating yellow
 fever epidemics in Africa: Vaccine demand forecast and impact modelling. PLOS Neglected
 Tropical Diseases. 2020;14: e0008304. doi:10.1371/JOURNAL.PNTD.0008304
- 494 4. Regional Committee for Africa. Progress report on the implementation of the regional
 495 strategy for health security and emergencies 2016–2020: information document. World

496 Health Organization. Regional Office for Africa; 2019. Report No.: AFR/RC69/INF.DOC/1. 497 Available: https://apps.who.int/iris/handle/10665/331442 498 5. Nwaiwu AU, Musekiwa A, Tamuzi JL, Sambala EZ, Nyasulu PS. The incidence and 499 mortality of yellow fever in Africa: a systematic review and meta-analysis. BMC Infectious 500 Diseases. 2021;21: 1-11. doi:10.1186/S12879-021-06728-X/FIGURES/4 501 6. Hamlet A, Ramos DG, Gaythorpe KAM, Romano APM, Garske T, Ferguson NM. 502 Seasonality of agricultural exposure as an important predictor of seasonal yellow fever 503 spillover in Brazil. Nature communications. 2021;12. doi:10.1038/S41467-021-23926-Y 504 7. World Health Organization. A global strategy to eliminate yellow fever epidemics (EYE) 505 2017–2026. Geneva: World Health Organization; 2018. Available: 506 https://apps.who.int/iris/handle/10665/272408 507 8. Eliminate yellow fever epidemics (EYE) strategy 2017-2026. [cited 5 Mar 2023]. Available: 508 https://www.who.int/initiatives/eve-strategy 509 9. Judson SD, LeBreton M, Fuller T, Hoffman RM, Njabo K, Brewer TF, et al. Translating 510 Predictions of Zoonotic Viruses for Policymakers. EcoHealth. 2017. doi:10.1007/s10393-511 017-1304-3 512 10. Aliaga-Samanez A, Real R, Segura M, Marfil-Daza C, Olivero J. Yellow fever surveillance 513 suggests zoonotic and anthroponotic emergent potential. Communications Biology 2022 5:1. 514 2022;5: 1-12. doi:10.1038/s42003-022-03492-9 515 11. de Thoisy B, Silva NIO, Sacchetto L, Trindade G de S, Drumond BP. Spatial epidemiology 516 of yellow fever: Identification of determinants of the 2016-2018 epidemics and at-risk areas 517 in Brazil. PLOS Neglected Tropical Diseases. 2020;14: e0008691. 518 doi:10.1371/JOURNAL.PNTD.0008691 519 12. Yamba EI, Aryee JNA, Quansah E, Davies P, Wemegah CS, Osei MA, et al. Revisiting the 520 agro-climatic zones of Ghana: A re-classification in conformity with climate change and 521 variability. PLOS Climate. 2023;2: e0000023. doi:10.1371/journal.pclm.0000023 522 13. Yellow Fever (YF) vaccination coverage. [cited 28 Sep 2022]. Available: 523 https://immunizationdata.who.int/pages/coverage/yfv.html?CODE=GHA&YEAR= 524 14. Bonney JHK, Sanders T, Pratt D, Agbodzi B, Laryea D, Agyeman NKF, et al. Molecular 525 Characterization of Circulating Yellow Fever Viruses from Outbreak in Ghana, 2021-2022 -526 Volume 29, Number 9—September 2023 - Emerging Infectious Diseases journal - CDC. 527 [cited 29 Aug 2023]. doi:10.3201/eid2909.221671 528 15. World Health Organization. Disease Outbreak News; Yellow Fever - Ghana. 1 Dec 2021 529 [cited 1 Aug 2023]. Available: https://www.who.int/emergencies/disease-outbreak-530 news/item/yellow-fever---ghana

16. Word Health Organization. Yellow Fever - West and Central Africa. In: Disease Outbreak
News [Internet]. 23 Dec 2021 [cited 17 Jan 2024]. Available:

- https://www.who.int/emergencies/disease-outbreak-news/item/yellow-fever---west-and central-africa
- 535 17. WHO report on global surveillance of epidemic-prone infectious diseases. [cited 2 Dec
- 536 2023]. Available: https://www.who.int/publications-detail-redirect/WHO-CDS-CSR-ISR 537 2000.1
- 538 18. WHO. Yellow Fever Cases Reported in Ghana, 1950–2004. World Health Organization;
 2005.
- 540 19. Agboli E, Tomazatos A, Maiga-Ascofaré O, May J, Lühken R, Schmidt-Chanasit J, et al.
 541 Arbovirus Epidemiology: The Mystery of Unnoticed Epidemics in Ghana, West Africa.
 542 2022. doi:10.3390/microorganisms10101914
- 543 20. Yellow Fever in Ghana: Predicting Emergence and Ecology from Historical Outbreaks
 544 (Datasets). figshare; 2023. doi:10.6084/m9.figshare.24747165
- 545 21. Scott D. Epidemic Disease in Ghana 1901-1960. London: Oxford Univesity Press; 1965.
- 546 22. WHO. Yellow fever in Africa and South America, 2011-2012 = Fièvre jaune en Afrique et
 547 en Amérique du Sud, 2011-2012. Weekly Epidemiological Record = Relevé
 548 épidémiologique hebdomadaire. 2013;88: 285–296.
- 549 23. Agadzi VK, Boatin BA, Appawu MA, Mingle JA, Addy PA. Yellow fever in Ghana, 1977550 80. Bulletin of the World Health Organization. 1984;62: 577.
- 24. Risk assessment on yellow fever virus circulation in endemic countries. [cited 7 Nov 2023].
 Available: https://www.who.int/publications-detail-redirect/WHO-HSE-PED-CED-2014-2
- Judson SD, Fischer R, Judson A, Munster VJ. Ecological Contexts of Index Cases and
 Spillover Events of Different Ebolaviruses. PLOS Pathogens. 2016;12: e1005780.
 doi:10.1371/journal.ppat.1005780
- 556 26. Phillips SJ, Anderson RP, Schapire RE. Maximum entropy modeling of species geographic
 557 distributions. Ecological Modelling. 2006;190: 231–259.
 558 doi:10.1016/j.ecolmodel.2005.03.026
- 27. van Proosdij ASJ, Sosef MSM, Wieringa JJ, Raes N. Minimum required number of specimen
 records to develop accurate species distribution models. Ecography. 2016;39: 542–552.
 doi:10.1111/ecog.01509
- 562 28. Merow C, Smith MJ, Silander Jr JA. A practical guide to MaxEnt for modeling species'
 563 distributions: what it does, and why inputs and settings matter. Ecography. 2013;36: 1058–
 564 1069. doi:10.1111/j.1600-0587.2013.07872.x

- 29. Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution interpolated
 climate surfaces for global land areas. International Journal of Climatology. 2005;25: 1965–
 1978. doi:10.1002/joc.1276
- 568 30. Friedl MA, Sulla-Menashe D, Tan B, Schneider A, Ramankutty N, Sibley A, et al. MODIS
 569 Collection 5 global land cover: Algorithm refinements and characterization of new datasets.
 570 Remote Sensing of Environment. 2010;114: 168–182. doi:10.1016/j.rse.2009.08.016
- 31. International Union for Conservation of Nature IUCN, Center for International Earth
 Science Information Network CIESIN Columbia University. Gridded Species
 Distribution: Global Mammal Richness Grids, 2015 Release. Palisades, New York: NASA
 Socioeconomic Data and Applications Center (SEDAC); 2015. Available:
 https://doi.org/10.7927/H4N014G5
- 32. Center for International Earth Science Information Network CIESIN Columbia
 University. Gridded Population of the World, Version 4 (GPWv4): Population Density
 Adjusted to Match 2015 Revision UN WPP Country Totals, Revision 11. Palisades, New
 York: NASA Socioeconomic Data and Applications Center (SEDAC); 2018. Available:
 https://doi.org/10.7927/H4F47M65
- 33. Pearson RG, Raxworthy CJ, Nakamura M, Townsend Peterson a. Predicting species
 distributions from small numbers of occurrence records: A test case using cryptic geckos in
 Madagascar. Journal of Biogeography. 2007;34: 102–117. doi:10.1111/j.13652699.2006.01594.x
- 585 34. Phillips S. A Brief Tutorial on Maxent. AT&T Research. 2005;190: 231–259.
- 35. Boorman JPT, Porterfield JS. A small outbreak of yellow fever in the Gold Coast.
 Transactions of The Royal Society of Tropical Medicine and Hygiene. 1957;51: 439–449.
 doi:10.1016/0035-9203(57)90079-2
- 36. Beausoleil EG, Mukhesee AB, Grant FC, Herron CA. Surveillance of yellow fever in Ghana,
 1960 et 1970. ORSTOM Ser Entomol Med Parasitol. 1972;10: 99–101.
- 591 37. P.A.K. ADDY KM and VKA. Recent Yellow Fever Epidemics in Ghana (1969-1983).
 592 EAST AFRICAN MEDICAL JOURNAL. 1986;63.
- 38. World Health Organization. Weekly Epidemiological Record, 2008, vol. 83, 08 [full issue].
 Weekly Epidemiological Record = Relevé épidémiologique hebdomadaire. 2008;83: 69–76.
- 39. WHO. Yellow fever in Ghana. In: Disease Outbreak News [Internet]. 3 Feb 2012 [cited 3
 Dec 2023]. Available: https://www.who.int/emergencies/disease-outbreaknews/item/2012_02_03b-en
- 40. Inusah A-W, Collins G, Dzomeku P, Head Id M, Ziblim S-D. Knowledge, attitudes and
 practice towards yellow fever among nomadic populations: A cross-sectional study in yellow
 fever outbreak communities in Ghana. PLOS Global Public Health. 2023;3: e0000733.
 doi:10.1371/JOURNAL.PGPH.0000733

- 41. Shearer FM, Longbottom J, Browne AJ, Pigott DM, Brady OJ, Kraemer MUG, et al.
 Existing and potential infection risk zones of yellow fever worldwide: a modelling analysis.
- 604The Lancet Global Health. 2018;6: e270–e278. doi:10.1016/S2214-109X(18)30024-X
- 42. Jentes ES, Poumerol G, Gershman MD, Hill DR, Lemarchand J, Lewis RF, et al. The revised
 global yellow fever risk map and recommendations for vaccination, 2010: consensus of the
 Informal WHO Working Group on Geographic Risk for Yellow Fever. The Lancet
- 608 Infectious Diseases. 2011;11: 622–632. doi:10.1016/S1473-3099(11)70147-5
- 609 43. Gaythorpe KAM, Hamlet A, Jean K, Ramos DG, Cibrelus L, Garske T, et al. The global
 610 burden of yellow fever. eLife. 2021;10. doi:10.7554/ELIFE.64670

612 Supporting Information

613

614 S1 Fig. Yellow Fever Annual Cases and Deaths in Ghana 1910-2022

- 615 The reported annual number of YF cases and deaths in Ghana since the first detected outbreak in
- 616 1910 until 2022 are shown. Also depicted are the years of reactive YF vaccination campaigns
- 617 and immunization policies.
- 618







Overall YF Model

Model excluding urban YF



