

1 **Title:**

2 Knowledge, Attitudes, and Practices Regarding Urinary Tract Infections Among Women in the United
3 Arab Emirates

4 **Authors:**

5 Shaikha Salah Alhaj^{1*}, Sajad Allami¹, Amjad Mohamadiyeh¹, Ammar Agha¹, Abdul Kareem Abu Ali¹,
6 Joudi M Bassam Habbal¹, Balsam Qubais Saeed²

7 **Affiliations:**

8 ¹ College of Medicine, University of Sharjah, Sharjah, UAE

9 ² Clinical Sciences Department, College of Medicine, University of Sharjah, Sharjah, UAE

10 **Corresponding Author:**

11 * Shaikha Salah Alhaj

12 College of Medicine, University of Sharjah, Sharjah, UAE

13 Email: shawa5i1@hotmail.com

14 **Acknowledgments:**

15 None to Declare

16 **Abstract:**

17 **Background:**

18 Urinary tract infections (UTIs), which are infections of the kidneys, ureters, bladder, or urethra, are a
19 worldwide public health concern. As compared to men, women are more prone to UTIs. There have been
20 several studies that explore the knowledge, attitudes, and practices of women regarding UTIs in different
21 countries, but no such study has been conducted in the UAE; therefore, we conducted this study in the
22 UAE setting.

23 **Methods:**

24 This study was conducted using an online survey created on Microsoft Forms. The minimum sample size
25 required for our study was 385. This study was conducted after obtaining ethical approval from the
26 Research Ethics Committee at the University of Sharjah (Reference Number: REC-23-09-19-01-F). A
27 personally designed questionnaire consisting of 21 items, derived from previous research was used to
28 record data. The data was analyzed using SPSS.

29 **Results:**

30 A total of 475 females were included in the study. Most respondents were aged 18-30 years (47.4%). Our
31 study found that a majority of the participants (69.7%) correctly identified bacteria as the most common
32 cause of UTIs. With regards to practices undertaken during UTIs, among the participants with a history
33 of UTI episodes, 32.6% waited 24-48 hours before seeking medical attention at a hospital or clinic, while
34 10% did not visit a hospital at all. Distinct trends were found when comparing demographic factors with
35 knowledge levels. Most notably, the age group of 18-30 years showed the highest percentage of high-
36 knowledge individuals (49%) compared to other age groups ($p < 0.05$). Education level was significantly
37 ($p = 0.003$) associated with UTI knowledge. Going to the hospital/clinic was reported by 41% with high
38 knowledge but only by 20% of those with poor knowledge. Moreover, a higher proportion of individuals
39 with high knowledge sought medical attention immediately within 24 hours (47%).

40 **Conclusion:**

41 Most of the participants possessed adequate knowledge regarding UTIs. Higher knowledge levels were
42 associated with more proactive & appropriate health behaviors, such as seeking medical attention
43 promptly & drinking more water.

44 **Introduction:**

45 Urinary tract infections (UTIs), which are infections of the kidneys, ureters, bladder, or urethra, are a
46 worldwide public health concern [1]. According to a study, over 404.6 million people all around the
47 globe had UTIs in 2019, and nearly 236,786 died as a result of UTIs the same year [2]. As compared to
48 men, women are more prone to UTIs [1]. One of the reasons attributable to this fact is the presence of
49 shorter urethra in women, which makes it easier for bacteria or other microbes to reach the bladder or
50 urinary tract and cause infection [1, 3]. It is estimated that approximately 60% of females experience at
51 least one UTI in their lifetime [3], which is more likely to occur in menopausal or pregnant women [3-4].
52 The urinary system is designed to keep the foreign bacteria out, but sometimes, failure of host defenses
53 to remove these bacteria can result in UTIs. *Escherichia coli* is the most associated bacteria with UTIs.
54 Other causative pathogens include *Staphylococcus saprophyticus*, *Proteus*, *Klebsiella*, and *Pseudomonas*
55 *aeruginosa* [5-6]. Additionally, certain risk factors predispose women to UTIs, which include pregnancy,
56 sexual activity (new sexual partner), menopause, use of certain birth controls, catheters, or a suppressed
57 immune system [5-6]. UTIs can be both symptomatic and asymptomatic. Symptomatic UTIs are
58 characterized by painful and frequent urination, hematuria, suprapubic pain, and fever [4-5]. On the other
59 hand, asymptomatic UTIs do not have any apparent symptoms but are characterized by the presence of
60 bacteria in urine [5, 7]. However, both symptomatic and asymptomatic infections should be treated [7] to
61 avoid further complications such as recurrent infections, kidney damage, bladder dysfunction, or preterm
62 labor [6, 8].

63 Despite being preventable, UTIs are quite prevalent among women in the UAE [4]. They can be
64 prevented by staying well-hydrated, maintaining hygiene, urinating after sexual activity, and raising
65 awareness among girls and women [1, 6]. There have been several studies that explore the knowledge,
66 attitudes, and practices of women regarding UTIs in different countries [9-10], but no such study has
67 been conducted in the UAE; therefore, in this study, we aim to explore the knowledge, attitudes, and
68 practices of women in the UAE regarding UTIs. The results of this study will aid us in determining the

69 level of awareness among them, which can contribute to targeted awareness campaigns and educational
70 initiatives related to women's health.

71

72 **Methods & Materials:**

73 The Strengthening the Report of the Observational Studies in Epidemiology (STOBE) guidelines were
74 used throughout conducting this study [11].

75

76 **Study Design:**

77 This study was a cross-sectional survey to assess the knowledge, attitude, and practices (KAP) regarding
78 Urinary Tract Infections (UTI) among women in the United Arab Emirates (UAE).

79

80 **Study Settings and Selection of Participants:**

81 This study was conducted using an online survey created on Microsoft Forms. The minimum sample size
82 required for our study was 385, collected using the Raosoft Sample Size Calculator with a confidence
83 interval of 95%, a margin of error of 5%, and a population size of 3.2 million. The data for the
84 population size of women in the UAE was used according to the latest population statistics [12].
85 However, we calculated data from 475 participants. The inclusion criteria for participants included: i)
86 females of any age; ii) residents of the UAE. The exclusion criteria included: i) males; ii) not a resident
87 of the UAE.

88

89 **Ethics Statement:**

90 Data was collected from participants after obtaining informed consent. The online consent was taken
91 prior to filling out the questionnaire. A brief summary of the questionnaire ahead was provided along
92 with the purpose of filling out this form. It was clarified that participation is anonymous and voluntary.
93 Only those who consented to participate could access the form ahead. In the case of minors, it was

94 mandatory that the consent form was filled out in the presence of parents or guardians. Participants were
95 ensured that their information would be kept confidential. This study was conducted after obtaining
96 ethical approval from the Research Ethics Committee at the University of Sharjah (Reference Number:
97 REC-23-09-19-01-F).

98

99 **Data Collection and Statistical Analysis:**

100 A personally designed questionnaire consisting of 21 items, derived from previous research and expert
101 opinion [9-10], was used to record data. It was divided into four sections: i) demographics details; ii)
102 questions to determine knowledge of the respondents; iii) questions to determine the attitude of the
103 respondents; iv) questions to determine practices of the respondents. Both English and Arabic versions of
104 the questionnaire were used to collect data. Data collection was started on 24th October 2023 and we
105 completed collecting data on 1st December 2023.

106 The data was collected and coded. It was then analyzed using IBM Statistical Package for Social
107 Sciences (SPSS) for Windows, Version 21 (IBM Corp, Armonk, NY, USA). A descriptive analysis was
108 run. A knowledge score was calculated, where a 0-1 score was categorized as poor knowledge, a 2-3
109 score was categorized as moderate knowledge, and a 4-6 score was categorized as high knowledge. To
110 study the association of factors with knowledge, a chi-square test was run, and a $p < 0.05$ was used as the
111 level of significance.

112

113 **Results:**

114 **1. Characteristics of the Study Population:**

115 A total of 475 females were included in the study. Most respondents were aged 18-30 years (47.4%).
116 The participants were equally distributed between married (48.8%) and single (48.2%), with a small
117 minority divided between divorced and widowed. A significant majority of the participants held a
118 bachelor's degree (66.9%), followed by those with a high school diploma (18.1%). Most participants

119 lived in Sharjah and Dubai (73.1%), while the rest were distributed among the remaining emirates. 35
120 participants (7.3%) had either diabetes or hypertension. A summary of the demographic characteristics
121 of the participants is shown in *Table 1*.

122

123 **Table 1: Demographic Characteristics of the Participants**

124

Variable	N	%
Age		
<18	16	3.4
18-30	225	47.4
31-40	113	23.8
41 and above	121	25.5
Marital Status		
Divorced	6	1.3
Married	232	48.8
Single	229	48.2
Widow	8	1.7
Educational Level		
Bachelor's degree	318	66.9
Post Graduate Studies	66	13.9
High school	86	18.1
Primary/middle school	5	1.1
Emirate		
Abu Dhabi	88	18.5
Ajman	31	6.5
Dubai	122	25.7
Fujairah	4	0.8
Ras Al Khaimah	3	0.6
Sharjah	225	47.4
Um al Quwain	2	0.4

125

126 **2. Knowledge of the Participants Regarding Urinary Tract Infections:**

127 Participants who responded 'No' to the question “Have you heard of the term urinary tract infection
128 (UTI)?” were categorized as possessing a minimal level of knowledge (18.7%), and the questionnaire
129 was subsequently concluded for them at that point. The most prevalent understanding of UTIs among
130 these respondents is inflammation of the kidney, bladder, or urethra (53.5% chose “Can be in all the

131 above”), followed by 'Inflammation of the urethra' only (23.2%). A majority (69.7%) correctly identify
 132 bacteria as the most common cause of UTIs. Additionally, most participants were able to accurately
 133 identify the symptoms, habits that increase the risk, and preventive measures of UTIs.

134
 135 **Table 2: Knowledge of the Participants Regarding Urinary Tract Infections**
 136

Variable	N	%
Have they heard of the term UTI?		
No	89	18.7
Yes	386	81.3
Definition of UTI		
Inflammation of the bladder	17	4.4
Inflammation of the kidney	2	0.5
Inflammation of the urethra	110	28.5
None of the above	3	0.8
All of the above*	254	65.8
Most common cause of UTI		
Bacteria*	269	69.7
Bad hygiene	37	9.6
Fungi	58	15.0
Viruses	22	5.7
Which symptom/s do you think occur with UTI?		
Pain during urination*	351	19.5
Change in the urine color/appearance*	302	16.8
Abdominal pain*	205	11.4
Back pain*	170	9.5
Increase in the frequency of urination*	251	14.0
Leg pain	31	1.7
Fever*	199	11.1
Constipation	22	1.2
A sudden desire to go to bathroom to urinate*	266	14.8
Which habit/s do you think increases the chances of having UTI?		
Cleaning the genitalia from back to front*	238	27.5
Drinking a small amount of water*	334	38.6
Delaying urination*	267	30.8
Urinating after sexual intercourse	10	1.2
Cleaning the genitalia from front to back	9	1.0
Drinking large amounts of water	8	0.9
Which factor/s do you think prevent UTI?		
Cleaning the genitalia from back to front	20	2.3
Drinking a small amount of water	3	0.3
Delaying urination	5	0.6

Urinating after sexual intercourse*	207	23.8
Cleaning the genitalia from front to back*	278	31.9
Drinking large amounts of water*	358	41.1

137 * represents the correct answers

138

139 3. Attitude of the Participants Regarding Urinary Tract Infections:

140 Most respondents (69.6%) believe that going to the hospital/clinic is the appropriate way to deal with a
 141 UTI. Drinking more water is also a common response, with 19.2% of respondents suggesting this
 142 method. 46.1% of respondents perceive UTI as a serious condition/disease. Some of the most commonly
 143 believed complications of UTIs included “damage to the kidneys” and “decrease in quality of life”
 144 (39.9% and 23.6%, respectively).

145

146 **Table 3: Attitude of the Participants Regarding Urinary Tract Infections**

147

Variable	N	%
In your opinion, what is the appropriate way of dealing with UTI?		
Drink more water	74	19.2
Go to the Hospital/Clinic	269	69.6
I don't know	9	2.3
Rest at home	1	0.3
Shower more frequently	6	1.6
Take analgesics (painkillers)	1	0.3
Take antibiotics directly	26	6.7
Do you think UTIs are common?		
Yes	363	94
No	23	6
Which of the following statements do you think is true regarding UTI?		
I don't know	34	8.9
UTIs affect both genders equally	52	13.5
UTIs affect females more	295	76.3
UTIs affect males more	5	1.3
Which complications do you expect from having a UTI?		
It can damage the kidneys	289	39.9
It can decrease the quality of life	171	23.6
It can lead to recurrent UTIs	142	19.6
It can affect pregnancies	85	11.8
It can lead to death	29	4
It can cause weight loss	8	1.1

Do you feel UTI is a serious condition/disease?		
I don't know	70	18.1
Yes	178	46.1
No	138	35.8

148

149 **4. Practices of the Participants Regarding Urinary Tract Infections:**

150 With regards to practices undertaken during UTIs, among the participants with a history of UTI
 151 episodes, 56 (32.6%) waited 24-48 hours before seeking medical attention at a hospital or clinic, while
 152 17 (10%) did not visit a hospital at all. The most common initial actions taken by these participants
 153 included visiting a hospital or clinic and increasing water intake (48% and 38.5%, respectively).

154

155 **Table 4: Practices of the Participants Regarding Urinary Tract Infections**

156

Variable	N	%
Have you ever had a UTI before?		
Yes	172	44.6
No	214	55.4
What did you do when you felt symptoms of UTI?		
Drank more water	114	38.5
Go to the hospital/clinic	142	48.0
Had rest at home	6	2.0
Took antibiotics directly	8	2.7
Took more showers	5	1.7
Took analgesics (painkillers)	19	6.4
I don't remember	2	0.7
If yes, how long did you wait before visiting the hospital/clinic?		
< 24 hours	49	28.4
24-48 hours	56	32.6
2-5 days	32	18.6
More than 5 days	18	10.4
I did not go to the hospital	17	10
Which symptoms do you have?		
Pain during urination	129	23.5
Change in the urine color & appearance	83	15.1
Back pain	55	10.0
Increase in the frequency of urination	94	17.2
A sudden desire to go to bathroom to urinate	87	15.9
Fever	33	6.0
Abdominal pain	67	12.2

How much water do you drink per day?		
< 0.5 Litres	34	8.8
0.5 Litres to 1 Litre	159	41.2
1 Litre to 2 Litres	142	36.8
More than 2 Litres	51	13.2

157

158 **5. Outcomes of Consultation:**

159 Regarding participants' expectation of consultation outcomes, participants were required to rank four
160 possible outcomes of a visit to the GP in order of importance. The majority of the participants (44%)
161 ranked Confirmation of UTI Diagnosis as the most important factor, with a mean ranking of 1.71.
162 However, the least prioritized outcome was Obtaining an Antibiotic Prescription (3%), with a mean
163 ranking of 3.04.

164

165 **Table 5: Ranking of the Outcomes of Consultation**

Outcome of Consultation	N	% prioritized as the most important	Mean Score
Confirmation of UTI diagnosis	75	44	1.71
Pain relief	48	28	2.49
Obtaining advice on when to contact a GP	43	25	2.74
Obtaining an antibiotic prescription	5	3	3.04

166

167 GP = General Practitioner

168

169 **6. Association of Level of Knowledge Regarding UTI with Demographics:**

170 Distinct trends were found when comparing demographic factors with knowledge levels. Most notably,
171 the age group of 18-30 years showed the highest percentage of high-knowledge individuals (49%)
172 compared to other age groups, while individuals below 18 years predominantly (75%) had poor
173 knowledge about UTIs ($p = <0.05$). Education level was significantly ($p = 0.003$) associated with UTI
174 knowledge as Individuals with higher educational qualifications, such as a bachelor's degree or
175 postgraduate studies, displayed a lower percentage of poor knowledge about UTIs (34% and 21%,
176 respectively). Moreover, the highest percentage (41%) of high-knowledge individuals was found in the

177 postgraduate group. Marital status also exhibited a significant ($p = <0.05$) association with knowledge,
 178 with single individuals demonstrating high levels of knowledge (47%) compared to divorced or married
 179 counterparts. While geographical differences in knowledge levels were noted across various Emirates,
 180 these were not statistically significant ($p = 0.174$).

181
 182

183 **Table 6: Association of Level of Knowledge of Participants Regarding UTI with**
 184 **Demographic Characteristics:**

Variable	Poor Knowledge		Moderate Knowledge		High Knowledge		p-value
	N	n	%	n	%	n	
Age							<0.05
<18	16	12	75	1	6	3	19
18-30	225	63	28	52	23	110	49
31-40	113	43	38	44	39	26	23
41 and above	121	39	32	47	39	35	29
Marital Status							<0.05
Divorced	8	4	50	4	50	0	0
Married	232	78	34	88	38	66	28
Single	229	73	32	49	21	107	47
Widow	6	2	33	3	50	1	17
Educational Level							0.003
Bachelor's degree	318	107	34	97	30	114	36
Post Graduate Studies	66	14	21	25	38	66	28
High school	86	33	38	20	23	33	38
Primary/middle school	3	1	33	2	67	0	0
Emirate							0.174
Abu Dhabi	88	35	40	25	28	28	32
Ajman	31	5	16	12	39	14	45
Dubai	122	45	37	31	25	46	38
Fujairah	4	1	25	2	50	1	25
Ras Al Khaimah	3	3	100	0	0	0	0
Sharjah	225	67	30	74	33	84	37
Um al Quwain	2	1	50	0	0	1	50
Total	475	157	33	144	30	174	37

185
 186

187

188 **7. Association of Level of Knowledge with Attitudes of Participants Regarding UTI:**

189 Individuals with a high level of knowledge about UTIs were significantly ($p = <0.05$) associated with a
190 greater likelihood of recognizing the seriousness of UTIs (63% of the participants who believed that
191 UTIs are serious and were categorized as having high knowledge also) and their potential complications
192 such as damage to the kidneys and leading to recurrent UTIs.

193 The belief that UTIs are more common in females was prevalent across all knowledge levels but was
194 most pronounced (52%) in the high-knowledge group.

195

196 **Table 7: Association of Level of Knowledge with Attitudes of Participants Regarding UTI:**

197

Variable	Poor Knowledge			Moderate Knowledge		High Knowledge	
	N	n	%	n	%	n	%
In your opinion, what is the appropriate way of dealing with UTI?							
Drink more water	74	24	32	27	36	23	31
Go to the Hospital/Clinic	269	32	12	101	38	136	51
I don't know	9	2	22	3	33	4	44
Rest at home	1	0	0	0	0	1	100
Shower more frequently	6	1	17	2	33	3	50
Take analgesics (painkillers)	1	0	0	0	0	1	100
Take antibiotics directly	26	9	35	11	42	6	23
Do you think UTIs are common?							
Yes	363	64	18	130	36	169	47
No	23	4	17	14	61	5	22
Which of the following statements do you think is true regarding UTI?							
I don't know	34	13	38	16	47	5	15
UTIs affect both genders equally	52	16	31	19	37	17	33
UTIs affect females more	295	39	13	104	35	152	52
UTIs affect males more	5	0	0	5	100	0	0
Which complications do you expect from having a UTI?							
It can damage the kidneys	289	37	13	107	37	145	50
It can decrease the quality of life	171	22	13	50	29	99	58
It can lead to recurrent UTIs	142	5	4	33	23	104	73
It can affect pregnancies	85	4	5	18	21	63	74
It can lead to death	29	0	0	2	7	27	93
It can cause weight loss	8	0	0	3	38	5	63
Do you feel UTI is a serious condition/disease?							
I don't know	70	14	20	43	61	13	19
Yes	178	23	13	42	24	113	63
No	138	31	3	59	5	48	4
Total	386	68	18	144	37	174	45

198

199 **8. Association of Level of Knowledge with Practices of Participants Regarding UTI:**

200 Higher knowledge levels were associated with more proactive and appropriate health behaviors, such as
 201 seeking medical attention promptly and drinking more water. Going to the hospital/clinic was reported
 202 41% with high knowledge but only by 20% of those with poor knowledge. Moreover, a higher
 203 proportion of individuals with high knowledge sought medical attention immediately within 24 hours
 204 (47%). Drinking more water was a common action across all knowledge levels, but more prevalent in

205 the high-knowledge group (41%). Individuals with higher knowledge tended to drink more water, with
206 50% of those consuming 1-2 liters having high knowledge and 35% consuming more than 2 liters daily.

207

208 **Table 8: Association of Level of Knowledge with Practices of Participants Regarding UTI:**

209

Variable	Poor Knowledge			Moderate Knowledge		High Knowledge	
	N	n	%	n	%	n	%
Have you ever had a UTI before?							
Yes	386	68	18	144	37	174	45
No	89	89	100	0	0	0	0
Total	475	157	33	144	30	174	37
What did you do when you felt symptoms of UTI?							
Drank more water	114	25	22	42	37	47	41
Go to the hospital/clinic	142	29	20	55	39	58	41
Had rest at home	6	0	0	1	17	5	83
Took antibiotics directly	8	0	0	2	25	6	75
Took more showers	5	1	20	3	60	1	20
Took analgesics (painkillers)	19	5	26	7	37	7	37
I don't remember	2	0	0	1	50	1	50
Total	170	37	22	64	38	69	41
If yes, how long did you wait before visiting the hospital/clinic?							
< 24 hours	49	8	16	18	37	23	47
24-48 hours	56	14	25	23	41	19	34
2-5 days	32	8	25	12	38	12	38
More than 5 days	18	5	28	8	44	5	28
I did not go to the hospital	17	4	24	3	18	10	59
Total	172	39	23	64	37	69	40
Which symptoms do you have?							
Pain during urination	129	24	19	48	37	57	44
Change in the urine color & appearance	83	13	16	33	40	37	45
Back pain	55	10	18	22	40	23	42
Increase in the frequency of urination	94	12	13	36	38	46	49
A sudden desire to go to bathroom to urinate	87	16	18	31	36	40	46
Fever	33	3	9	15	45	15	45
Abdominal pain	67	15	22	23	34	29	43
Total	167	36	22	62	37	69	41
How much water do you drink per day?							
< 0.5 Liters	34	8	24	16	47	10	29
0.5 Liters to 1 Liter	159	24	15	60	38	75	47
1 Liter to 2 Liters	142	24	17	47	33	71	50
More than 2 Liters	51	12	24	21	41	18	35
Total	386	68	18	144	37	174	45

210

211 **Discussion:**

212 Urinary tract infections (UTI) are one of the most common, challenging, and misunderstood diseases that
213 women experience as outpatients [13]. UTIs are a prevalent health issue among females globally [1],
214 with a notable prevalence in the United Arab Emirates (UAE), which is increasing at an exponential rate
215 [14]. The issue of delayed detection of UTIs is very prevalent. It leads to both excessive therapy and
216 inadequate treatment [13]. If not managed properly and timely or left untreated, UTIs may cause serious
217 complications like renal impairment, kidney scarring, sepsis, and pyelonephritis [5]. Women's practices,
218 attitudes, and understanding of urinary tract infections are significant variables that can affect how this
219 illness is managed and prevented. While a substantial number of studies on this subject have been carried
220 out globally [9-10, 13], there remains no data specific to the UAE context. This study aims to bridge this
221 gap by exploring the knowledge, attitudes, and practices of women in the United Arab Emirates about
222 urinary tract infections (UTIs).

223 Our research revealed several noteworthy findings. Most of the participants (80%) were aware of the
224 term UTI and half of them were able to identify the correct definition. The vast majority were able to
225 identify the most common cause of UTIs and the most prevalent symptoms. Additionally, most of them
226 were also able to report the most appropriate ways of dealing with UTI. Therefore, the results of our
227 study concluded that most of the respondents exhibited an acceptable level of awareness regarding UTIs.
228 Our results regarding the knowledge of the participants are consistent with the other studies in the
229 surrounding region like in Saudi Arabia [10, 15], and were also consistent with a study conducted in the
230 Netherlands [9]. On the other hand, studies conducted in Bangalore [16] and Egypt [17] suggested that
231 the participants mostly had poor knowledge and attitude, which may suggest that lower levels of
232 knowledge are associated with developing countries. Similarly, while assessing the attitudes and
233 practices of the respondents, we found out that a significant fraction of our study population understood
234 the seriousness of the condition (46%) and took appropriate measures for the management of UTIs, such
235 as seeking medical help (48%) or increasing water intake (38.5%). When asked about the outcomes of
236 General Practitioner (GP) consultation, the participants in our study showed results similar to those of the

237 study group in the Netherlands from Cox SML et al. [9]. In both studies, most participants chose
238 confirmation of UTI diagnosis as the main outcome behind the GP visit, which shows that patients value
239 getting the definitive diagnostic information during their consultation. Moreover, in our study, only 3%
240 of the participants prioritized obtaining an antibiotic prescription compared to 14.3% in the Netherlands
241 study [9]. These findings may be explained by the wider accessibility of antibiotics without a
242 prescription in the UAE. Uropathogenic bacteria are notable for their frequent resistance to many drugs
243 [18]. It is necessary to monitor the antibiotic resistance patterns in UTIs [19].

244 Regarding antibiotic usage, participants were asked about the appropriate way of dealing with a urinary
245 tract infection and only 6.7% of the participants chose to take antibiotics directly, and only 2.7% took
246 antibiotics directly when they felt UTI symptoms. This may suggest increased awareness of appropriate
247 antibiotic use and a desire for more conservative treatment approaches in the UAE population. Moreover,
248 this trend aligns with global health concerns regarding antibiotic resistance, signifying a positive step
249 towards responsible antibiotic utilization within the UAE community.

250 Furthermore, we studied the association of knowledge level and demographics, attitudes, and practices
251 and concluded that the individuals who possessed a high level of knowledge were 18-30 years old and
252 were well-educated. A similar study was conducted in Malaysia, which stated a possibility that educated
253 respondents most likely would have studied about UTI or UTI-related topics in their syllabus, which
254 could be a possible reason for their high level of knowledge [20]. These individuals also understood the
255 seriousness of UTIs and preferred to seek medical help for treatment. Significant associations were found
256 between knowledge and age, educational level, and marital status; however, no significant association
257 was found between the level of knowledge and Emirate which is in contrast to studies conducted in
258 Saudi Arabia [10, 15].

259 However, certain strengths and limitations of our study need to be acknowledged. This is the first study
260 that assessed the knowledge, attitudes, and practices of women regarding UTIs in the UAE. Enhancing
261 patients' understanding, beliefs, and actions about prevalent illnesses plays a crucial part in health

262 education [21]. The results of this study can guide the tailored educational campaigns for women's health
263 in the UAE. The study's limitations include the possible occurrence of recall bias due to the dependence
264 on self-reported data, as well as the possibility of participants' replies being influenced by social
265 desirability or response bias.

266 In conclusion, our study found that most of the respondents had adequate knowledge about UTIs. The
267 results of this study conclude that most of the women in the UAE exhibited substantial awareness and
268 behavioral patterns regarding UTIs. Higher knowledge levels were associated with more proactive &
269 appropriate health behaviors, such as seeking medical attention promptly & drinking more water.

270

271

272

273

274 **References:**

- 275 1. Clinic C. Why UTIs happen differently in men and women [Internet]. Cleveland Clinic. 2023
276 [cited 2023 Dec 8]. Available from: [https://health.clevelandclinic.org/said-said-worse-urinary-tract-](https://health.clevelandclinic.org/said-said-worse-urinary-tract-infections-infographic)
277 [infections-infographic](https://health.clevelandclinic.org/said-said-worse-urinary-tract-infections-infographic)
- 278 2. Zeng Z, Zhan J, Zhang K, Chen H, Cheng S. Global, regional, and national burden of urinary
279 tract infections from 1990 to 2019: an analysis of the global burden of disease study 2019. World
280 Journal of Urology. 2022 Mar;40(3):755-63. <https://doi.org/10.1007/s00345-021-03913-0>
- 281 3. Weber B. What is the connection between menopause and UTIs? [Internet]. 2022 [cited 2023 Dec
282 8]. Available from: <https://www.medicalnewstoday.com/articles/menopause-and-uti>
- 283 4. Dube R, Al-Zuheiri ST, Syed M, Harilal L, Zuhaira DA, Kar SS. Prevalence, Clinico-
284 Bacteriological Profile, and Antibiotic Resistance of Symptomatic Urinary Tract Infections in
285 Pregnant Women. Antibiotics. 2022 Dec 25;12(1):33. <https://doi.org/10.3390/antibiotics12010033>
- 286 5. Kaur R, Kaur R. Symptoms, risk factors, diagnosis and treatment of urinary tract infections.
287 Postgraduate Medical Journal. 2021 Dec;97(1154):803-12. [https://doi.org/10.1136/postgradmedj-](https://doi.org/10.1136/postgradmedj-2020-139090)
288 [2020-139090](https://doi.org/10.1136/postgradmedj-2020-139090)
- 289 6. Urinary tract infection (UTI) - Symptoms and causes - Mayo Clinic [Internet]. Mayo Clinic. 2022
290 [cited 2023 Dec 9]. Available from: [https://www.mayoclinic.org/diseases-conditions/urinary-tract-](https://www.mayoclinic.org/diseases-conditions/urinary-tract-infection/symptoms-causes/syc-20353447)
291 [infection/symptoms-causes/syc-20353447](https://www.mayoclinic.org/diseases-conditions/urinary-tract-infection/symptoms-causes/syc-20353447)
- 292 7. Givler DN. Asymptomatic bacteriuria [Internet]. StatPearls - NCBI Bookshelf. 2023 [cited 2023
293 Dec 9]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK441848/>
- 294 8. Kassel G. When should you go to the hospital for a urinary tract infection (UTI)? [Internet].
295 Healthline. 2023 [cited 2023 Dec 9]. Available from: [https://www.healthline.com/health/uti-when-to-](https://www.healthline.com/health/uti-when-to-go-to-hospital#complications)
296 [go-to-hospital#complications](https://www.healthline.com/health/uti-when-to-go-to-hospital#complications)
- 297 9. Cox SM, van Hoof MW, Lo-A-Foe K, Dinant GJ, Oudhuis GJ, Savelkoul P, Cals JW, de Bont
298 EG. Cross-sectional internet survey exploring women's knowledge, attitudes and practice regarding

- 299 urinary tract infection-related symptoms in the Netherlands. *BMJ open*. 2022 May 1;12(5):e059978
300 <http://doi.org/10.1136/bmjopen-2021-059978>
- 301 10. Almaghlouth AK, Alkhalaf RA, Alshamrani AA, Alibrahim JA, Alhulibi BS, Al-Yousef AY,
302 Alamer AK, Alsuabie SM, Almuhanha SM, Alshehri AD, ALSHAMRANI AA. Awareness,
303 Knowledge, and Attitude Towards Urinary Tract Infections: An Appraisal From Saudi Arabia.
304 *Cureus*. 2023 Nov 24;15(11). <https://doi.org/10.7759/cureus.49352>
- 305 11. Vandembroucke JP, Elm EV, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, Poole C,
306 Schlesselman JJ, Egger M, Strobe Initiative. Strengthening the Reporting of Observational Studies in
307 Epidemiology (STROBE): explanation and elaboration. *Annals of internal medicine*. 2007 Oct
308 16;147(8):W-163. <https://doi.org/10.7326/0003-4819-147-8-200710160-00010-w1>
- 309 12. Blogger G. United Arab Emirates (UAE) Population Statistics 2023 | GMI [Internet]. Official
310 GMI Blog. 2023 [cited 2023 Dec 8]. Available from: [https://www.globalmediainsight.com/blog/uae-](https://www.globalmediainsight.com/blog/uae-population-statistics)
311 [population-statistics](https://www.globalmediainsight.com/blog/uae-population-statistics)
- 312 13. Ata A, Hussain W, Malik J, Abbasi S, Mehmood A, Khan AH, Parvez A, Abdullah A, Abbasi
313 AS, Shah A. Knowledge Attitude and Practices Regarding Urinary Tract Infection Among Pakistani
314 Diabetic Females. *Pakistan Journal of Medical Research*. 2023 Nov 23;62(3):128-34.
315 <https://www.pjmr.org.pk/index.php/pjmr/article/view/495>
- 316 14. Zhu C, Wang DQ, Zi H, Huang Q, Gu JM, Li LY, Guo XP, Li F, Fang C, Li XD, Zeng XT.
317 Epidemiological trends of urinary tract infections, urolithiasis and benign prostatic hyperplasia in 203
318 countries and territories from 1990 to 2019. *Military Medical Research*. 2021 Dec;8:1-2.
319 <https://doi.org/10.1186/s40779-021-00359-8>
- 320 15. Alshahrani M, Alzahrani AB, Alzahrani AA, Alqhtani AM, Alwabel HH, Asiri KM, Mohammed
321 Y, Abumelha MD, Alshahrani RS, Alshahrani AA, Alhussain RM. Knowledge, Attitude and Practice
322 of Urinary Tract Infection among Female in Aseer Region. *Bahrain Medical Bulletin*. 2022 Mar
323 1;44(1). https://www.bahrainmedicalbulletin.com/MARCH_2022/BMB-22-153.pdf

- 324 16. Muchukota S, Muchukota B, Mekkanti MR, Mounika P, Mathappan R. A cross sectional study
325 on knowledge, attitude & practice towards personal hygiene in women with urinary tract infection and
326 providing patient counselling in esi hospital bangalore. UPI Journal of Pharmaceutical, Medical and
327 Health Sciences. 2020:5-9. <https://doi.org/10.37022/jpmhs.v3i2.20>
- 328 17. El Lawindi, Mona & Sayed, Hanan & El shafei, Arwa & Hayek, Noha & Nour, Hani. Assessment
329 of Urinary Tract Infections' Risk Factors and Knowledge among Attendees of Theodor Bilharz
330 Research Institute, Giza, Egypt.. International Public Health Forum. 2014;1(1):25-9. (PDF)
331 [Assessment of Urinary Tract Infections' Risk Factors and Knowledge among Attendees of Theodor](#)
332 [Bilharz Research Institute, Giza, Egypt. \(researchgate.net\)](#)
- 333 18. Hrbacek J, Cermak P, Zachoval R. Current antibiotic resistance patterns of rare uropathogens:
334 survey from Central European Urology Department 2011–2019. BMC urology. 2021 Dec;21:1-8.
335 <https://doi.org/10.1186/s12894-021-00821-8>
- 336 19. Huang L, Huang C, Yan Y, Sun L, Li H. Urinary tract infection etiological profiles and antibiotic
337 resistance patterns varied among different age categories: a retrospective study from a tertiary general
338 hospital during a 12-year period. Frontiers in Microbiology. 2022 Jan 27;12:813145.
339 <https://doi.org/10.3389/fmicb.2021.813145>
- 340 20. Selamat NW, Martinez KP, Subrain G, Auamnoy T. THE CORRELATION BETWEEN
341 AWARENESS AND ATTITUDE DOMAINS ON URINARY TRACT INFECTION (UTI) AMONG
342 BURAPHA UNIVERSITY STUDENTS IN CHONBURI, THAILAND. Malaysian Journal of Public
343 Health Medicine. 2021 Apr 24;21(1):21-8. <https://doi.org/10.37268/mjphm/vol.21/no.1/art.444>
- 344 21. Kruszewski B. What is Patient Education? | Lecturio Medical [Internet]. Lecturio. 2023 [cited
345 2024 Jan 20]. Available from: [https://www.lecturio.com/blog/medical/what-is-patient-education-and-](https://www.lecturio.com/blog/medical/what-is-patient-education-and-why-is-it-important/)
346 [why-is-it-important/](https://www.lecturio.com/blog/medical/what-is-patient-education-and-why-is-it-important/)