- 1 TITLE: Systematic review of contemporary improvement interventions on care coordination,
- 2 discharge support and transitional care from the patient experience perspective
- 3

# 4 Authors:

5 Tiago S Jesus, PhD, OTD<sup>1,2</sup>; Brocha Z. Stern, PhD, OTR<sup>3</sup>; Dongwook Lee, OTD, OTR<sup>4</sup>; Manrui

6 Zhang, PhD, MPH<sup>2</sup>; Jan Struhar, DPT<sup>5</sup>; Allen W. Heinemann, PhD<sup>6,7</sup>; Neil Jordan, PhD<sup>8,9</sup>; Anne

7 Deutsch, PhD, RN, CRRN<sup>6,7,10</sup>

8	1.	Division of Occupational Therapy, School of Health and Rehabilitation Sciences,
9		College of Medicine, The Ohio State University, Columbus, OH, USA.
10	2.	Center for Education in Health Science, Institute for Public Health and Medicine,
11		Northwestern University Feinberg School of Medicine, Chicago, IL, USA.
12	3.	Institute for Healthcare Delivery Science, Department of Population Health Science
13		and Policy, Icahn School of Medicine at Mount Sinai, New York, NY, USA
14	4.	Center for Child Development & Research, Sensory EL, ROK; Dept. of Physical
15		Medicine and Rehabilitation Medicine, Korehab Clinic, UAE
16	5.	Nerve, Muscle and Bone Innovation Center & Oncology Innovation Center, Shirley
17		Ryan AbilityLab, Chicago, IL, USA.
18	6.	Center for Rehabilitation Outcomes Research, Shirley Ryan AbilityLab, Chicago, IL,
19		USA.
20	7.	Dept. of Physical Medicine and Rehabilitation Medicine, Northwestern University
21		Feinberg School of Medicine, Chicago, IL, USA.
22	8.	Dept. of Psychiatry and Behavioral Sciences, Dept. of Preventive Medicine,
23		Northwestern University Feinberg School of Medicine, Chicago, IL, USA.
24	9.	Center of Innovation for Complex Chronic Healthcare, Hines VA Hospital, Hines, IL,
25		USA
26	10	Center for Health Care Outcomes, RTI International, Chicago, IL, USA.
27		
28	Word cou	nt: 3477; Number figures: 1; Number tables: 4

30 Corresponding author: Tiago Jesus

#### 31 **Contact Information**

- 32 Tiago S Jesus, PhD, OTD (He/Him/His)
- 33 The Ohio State University; College of Medicine, School of Health & Rehabilitation Sciences,
- 34 Division of Occupational Therapy; <u>tiago.jesus@osumc.edu</u>
- 35 453 West 10th Ave. Columbus, OH 43210
- 36 Main Phone: 614-292-1706; Fax: 614-292-0210
- 37

## 38 Author contributions:

- 39 Conceptualization; TSJ, AH, AD; Data curation: TSJ, BZS, MZ, DL, JS; Methodology: TSJ,
- 40 AH, BZS, NJ, AD Project administration: TSJ; Supervision: AH, AD; Roles/Writing original

41 draft: TSJ; and Writing - review & editing: TSJ, AH, BZS, NJ, AD.

42

## 43 **Declarations:**

44 Tiago S. Jesus completed part of this work and Manrui Zhang the totally of this work under a

45 grant from the National Institute on Disability, Independent Living, and Rehabilitation Research

- 46 (NIDILRR; 90ARHF0003). NIDILRR is a center within the Administration for Community
- 47 Living (ACL), U.S. Department of Health and Human Services (HHS). The contents of this
- 48 publication do not necessarily represent the policy of NIDILRR, ACL, or HHS, and the reader
- 49 should not assume endorsement by the U.S. federal government. Dr Jesus initiated this work
- 50 with the Northwestern University's affiliation (under the grant support mentioned) and the work
- 51 was completed with The Ohio State University's affiliation.

52

# 53 **Conflicts of Interest:**

54 All authors declare no conflicts of interest.

55	TITLE: Improving care coordination and discharge support evaluated from the patient
56	experience perspective: systematic review of the contemporary evidence
57	
58	Abstract
59	Aim: To synthesize the impact of improvement interventions related to care coordination and
60	discharge support on patient experience measures.
61	Method: Systematic review. Searches were completed in six scientific databases, five specialty
62	journals and through snowballing. Eligibility included studies in English (2015-2022) on
63	improving care coordination, discharge support, or transitional care assessed by standardized
64	patient experience measures as a primary outcome. Two independent reviewers made
65	eligibility decisions and performed quality appraisals.
66	Results: Of 1087 papers initially screened, 15 were finally included. Seven studies (three
67	randomized controlled trials [RCTs]) focused on care coordination activities and used enhanced
68	supports (e.g., improvement coaching; tailoring for vulnerable populations) for Patient-
69	Centered Medical Homes or other primary care sites. Effectiveness was mixed or neutral
70	relative to standard supports or models of care. Eight studies (three RCTs) focused on enhanced
71	discharge support, including patient education (e.g., "teach back" method) and telephone
72	follow-up or transitional support; mixed or neutral results on the patient experience were also
73	found and with more substantive risks of bias.

74	Conclusion: Enhanced supports for improving care coordination, discharge education, and post-
75	discharge follow-up had mixed or neutral effectiveness for improving the patient experience
76	with care, compared to standard or simpler improvement approaches. Studies on the
77	improvement of patient experiences, especially for enhanced patient discharge, need further
78	strengthening such as including the patient perspective for its co-development.
79	
80	Key words: Patient Experience; Patient Satisfaction; Person-centered Care; Quality

80 **Key words**: Patient Experience; Patient Satisfaction; Person-centered Care; C

81 Improvement; Discharge Planning; Care Coordination

82

## 83 Background:

The patient experience with care is an integral component of healthcare quality and a construct 84 85 used to assess the *person-centeredness* of healthcare delivery, informing quality-improvement (QI) activities.<sup>1-3</sup> Specifically, patient experience refers to how patients have experienced 86 87 valuable aspects of healthcare delivery, such as physician communication, involvement in 88 decision-making, getting timely appointments, or supports to care coordination and discharge planning in the transitions across settings to address care fragmentation issues.<sup>1</sup> 89 90 As synthesized in several systematic reviews, better patient experiences with care have been 91 associated with improved treatment adherence and to improved patient and health system outcomes (e.g., reduced healthcare utilization or malpractice litigation).<sup>4-7</sup> For example, a 92 93 recent systematic review found that better scores on standardized patient experience

measures were associated with greater self-reported physical and mental health, lower 94 95 frequency and length of hospitalizations, and fewer emergency room visits.<sup>4</sup> Furthermore, improving the patient experience of care has been recognized as a QI aim in itself, under the 96 goal of improving the person-centeredness of care.<sup>89</sup> 97 Many health systems increasingly reimburse providers for the value of care, including patient 98 experience scores. The US Agency for Healthcare Research and Quality (AHRQ) has developed 99 the Consumer Assessment of Healthcare Providers and Systems (CAHPS) surveys and quality 100 101 measures, which have been widely used to track, publicly report, and compare providers' patient experience performance.<sup>10</sup> Within selected pay-for-performance programs, the US 102 103 Centers for Medicare & Medicaid Services uses CAHPS scores, reflecting patient experience of care, when computing payment incentives.<sup>11</sup> Several other countries require providers to 104 monitor and improve patient experience.<sup>12 13</sup> Overall, healthcare systems, organizations, and 105 106 frontline practitioners are increasingly subject to quality mandates, value-based incentives, or 107 market pressures (e.g., customer loyalty) to develop systematic activities to improve their patient experience performance,<sup>14</sup> including for care coordination and discharge support.<sup>15 16</sup> 108 Reducing care fragmentation, improving continuity of care, and preventing medication errors or 109 rehospitalizations can be achieved by interventions such as supported discharge, case 110 111 management programs, transitional care services, use of care coordinators as well as models of 112 care such as patient-centered medical homes (PCMHs). Systematic reviews have addressed the 113 effectiveness of these health service interventions, and many of these have been shown to be 114 effective for a set of health, quality of life, and health system outcomes (e.g., reduced rehospitalizations).<sup>17-20</sup> For example, interventions to facilitate the transition from hospital to 115

116	home when started in the hospital and continued in the community through telephone or other
117	telehealth follow-up or scheduled home visits can be effective in reducing hospital readmission,
118	especially when tailored to each patient. <sup>18 19 21</sup>
119	However, we found no systematic review focused on the impact that care coordination,
120	discharge support, or transitional care activities have on the patient's experience with care. A
121	recent overview of systematic reviews for care coordination interventions found patient
122	experiences were rarely addressed or synthesized as outcomes of these interventions. <sup>22</sup> Here,
123	we aim to synthesis the contemporary literature (2015-2022) on the impact of care
124	coordination, discharge support, and transitional care activities on patient experience as
125	assessed by formal patient-reported experience measures.
126	
127	Methods
128	
129	<b>Design:</b> We registered (PROSPERO: CRD42022358337) a systematic review protocol that
130	focused on synthesizing the English-language contemporary evidence (2015-2022) addressing
131	health service interventions to improve the patient-reported experience with care. Here, we
132	focus on the impact of interventions related to enhanced care coordination, discharge support,
133	or transitional care activities. We use the Preferred Reporting Items for Systematic Reviews and

134 Meta-Analyses (PIRSMA) guidance for the review report.

135 Eligibility:

We included controlled trials, longitudinal observational studies with controls, and pre-post 136 137 designs with >30 participants. We excluded qualitative, cross-national studies, or any study not assessing the impact on patient-reported experience measures. Studies needed to have full 138 139 texts available, focus on improving the patient experience (i.e., one of no more than two 140 primary outcomes measures) and report inferential statistics (e.g., p-values) about the impact 141 of the intervention on a patient experience measure. For patient experience outcomes, we included studies that used standardized, quantitative 142 143 patient experience assessments (e.g., validated surveys, their items or composite domains, or 144 surveys that were externally collected and routinely used across providers, including for valuebased reimbursement). We excluded studies that assessed patient experiences only with non-145 146 standardized, gualitative, or other non-validated instruments. Measures that were not patient-147 reported (e.g., observational) were excluded. We did not synthesize the impact of the 148 interventions on measures other than patient-reported experience with care. 149 For participants, we included health systems, organizations, providers, networks, settings, or service units, including any health professionals or staff. We excluded health service 150 151 interventions exclusively delivered by students or clinicians-in-training. Those providing patient 152 experience feedback could be the patient, family/informal caregivers, or proxy respondents. 153 For context, we had no geographic restriction, but we only included English-language articles published in 2015 or thereafter, to reflect contemporary interventions responsive to the more 154 155 recent focus, mandate, or incentives for improving the patient experience of care. For example,

value-based reimbursements that included patient experience scores did not emerge in the US
 until 2013.<sup>23</sup>

- 158 For this review, we only included health-service improvement interventions focused on *care*
- 159 *coordination* (i.e., systematic organizational, service line, or service unit strategies to address
- 160 fragmentation of care delivery and enhance continuity of care<sup>22</sup>), discharge support (i.e.,
- 161 patient education in preparation for discharge and/or post-discharge education or follow-up
- support either in the environment or by telephone or other telehealth/web-based
- 163 mechanisms<sup>18</sup>), and *transitional care* (i.e., subset of intermediate, time-limited set of actions to
- 164 ensure the coordination or continuity of care as patients transfer between different settings or
- levels of care<sup>24</sup>), which often involves a single point of contact to optimize service access,
- 166 communication, and coordination.<sup>25</sup>

#### 167 **Search:**

168 We searched six scientific databases (PubMed/MEDLINE, CINAHL, EconLit, PsycINFO, DOAJ, and

169 Scopus) using a combination of free-text words with indexed terms that reflect our eligibility

170 criteria. We restricted the search for English language and publications after January 2015.

171 **Supplementary Appendix 1** shows the detailed searches for each database.

We also conducted targeted searches within the *Patient Experience Journal, Journal of Patient Experience, Medical Care,* and *Health Expectations. NEJM Catalyst Innovations in Care Delivery* was searched since January 2020, when the journal became peer reviewed. These were journals with frequent papers identified in the initial database searches. Finally, we conducted snowballing strategies (e.g., citation-tracking, similar publications in PubMed) using the final

- 177 included articles and reference lists of similar systematic reviews, including those identified
- through the searches.

#### 179 Selection:

- 180 Two independent reviewers (TJ first reviewer; DL, JS, or MZ for the second reviewer role)
- 181 conducted the title-and-abstract screenings and then the full text reviews against the eligibility
- 182 criteria. Reviewer consensus was achieved on the final inclusion within a round of discussion,
- 183 with no need for third reviewer involvement.

#### 184 Data Extraction

- 185 The research team built a data extraction form that they used to chart data such as study
- 186 characteristics and context, patient experience measures and metrics, interventions, analytic
- 187 approaches, and results pertaining to patient-reported experience with care. The data were
- 188 charted initially by the first author (TJ) and verified by a second author (BS), with subsequent
- rounds for any flagged extractions until they reached consensus. Supplementary Appendix 2
- 190 provides detailed, consensus-based data extractions for each included study.

#### 191 Quality Assessment

The quality assessment was performed independently by TJ and BS. The Cochrane-suggested risk of bias criteria for Effective Practice and Organization of Care (EPOC) reviews were used for the controlled trials, while the National Heart, Lung, and Blood Institute's quality assessment tools were used for the pre-post studies and observational studies that assessed or compared the effects of health service interventions.

As planned in the review protocol, we did not grade the strength of the evidence across the 197 198 research studies and their design. Covering a range of health service or QI interventions, contexts, assessments, and study designs, our review is configurative (not aggregative) in 199 nature. We tabulated the consensus-based appraised risks of bias for each study's context and 200 201 design, based on the respective checklist-based methodological assessment; those detailed assessments are in Supplementary Appendix 3. 202 Analytic plan 203 We tabulated and synthesized the results per two major inclusion categories: 1) care 204 205 coordination and 2) discharge support and follow up. The latter included transitional care because the included studies that had this component were in relation to discharge support. 206 Within each category, we tabulated the methods and appraised the risk of bias for each study, 207 208 and then synthesized the interventions and their findings, ordered by study design (starting with RCTs) and publication date within the same design. Given the configurative nature of the 209

210 review and the heterogeneity of interventions and study contexts, we did not conduct a meta-

analysis.

212

213

#### Results

214

215 **Figure 1** provides the PRISMA flowchart.

216



# Figure 1: PRISMA flowchart of the systematic review.

219

Among the 1087 unique papers screened, 138 full texts were assessed for eligibility and 15

221 were included.<sup>15 16 26-38</sup> The absence of impact assessment with inferential statistics was the

222 main reason for exclusion at the full-text assessment. Among papers included, seven focused

223	on enhanced care coordination supports (e.g., over and beyond models of care based on a
224	PCMH), whereas eight focused on enhanced discharge supports, specifically for enhanced
225	patient education, reengineered discharge, telephone follow-up including for transitional care,
226	or a mix of those interventions.
227	
228	a) Enhanced care coordination supports – primary care level ( $n=7$ )
229	
230	Table 1 summarizes each study's methods (e.g., design, context) and risks of bias;         Table 2
231	summarizes the intervention and study findings.
232	Context-wise, all studies were multi-site PCMHs or other primary care practices, all from a
233	network or integrated health system in the US: Veterans' Health Administration (VHA), <sup>32-34</sup> a
234	statewide PCMH network, <sup>35</sup> Federally Qualified Health Centers, <sup>36</sup> a Safety Net Medical Homes'
235	Initiative, <sup>38</sup> or academically-affiliated primary care practices. <sup>37</sup>
236	Three studies were RCTs for facilitating site-based improvement projects. A recent RCT at the
237	VHA, with the fewest appraised risks of bias (Table 1), used QI facilitation (e.g., coaching) on the
238	site's use of a care-coordination improvement toolkit – beyond its internal dissemination
239	(control); both groups improved, with no significant benefit for the more resource-intensive
240	arm. <sup>34</sup>
241	Another RCT focused on the impact of augmented care coordination for patients with high risk
242	of hospitalization (beyond standard PCMH; control group) and found improvements in 2 of 6
243	care coordination items. <sup>33</sup> The remaining RCT encouraged the selection of high-value care-

	Design & Analysis	Context / Setting	Risk	Risk of Bias – level (within each study design) & Described risks			
Noël et al, 2022	RCT (cluster-randomized) of QI approaches. Multivariable difference-in-differences with adjustment for clustering and patient characteristics.	Patient-Centered Medical Homes, primary care; 12 sites (6 intervention, 6 control) within the Veterans Health Administration system, USA	RC T	Different pre and post groups. Non-coached clinics had significantly lower baseline values for the outcome, thus greater improvement margins.			
Zulm an et al., 2019	Secondary survey of a RCT (patient-level randomization) for quality improvement. Logistic regression, using intention-to-treat analyses. Sensitivity analyses with adjustment.	Patient-Centered Medical Homes, primary care; 5 sites within the Veterans Health Administration system, USA	RC T	No baseline outcomes. Controls were from the same sites, with potential for contamination. Not all patients randomized to the intervention received the intervention. Patients may not distinguish the intervention from the previously-provided support (control). Self-selected sites participated in the program, with 5% greater experience scores than the national average for several health care experience survey questions. No correction for multiple comparisons.			
Dorr et al., 2016	Pragmatic RCT (cluster-randomized) for quality improvement. Adjusted difference-in-differences analysis and generalized estimating equation approach.	Patient-Centered Medical Homes, primary care; 8 sites (4 intervention, 4 control), Oregon, USA	RC T	Use of P value threshold of $P < .10$ and all significant changes identified ranged between $P > 0.5$ and $P < .10$ . No correction for multiple comparisons. Different pre and post groups. Small number of clusters. Selection and respondent bias unable to be adjusted for, e.g., those motivated to take part in the experiment and respond in the pre- and post-periods.			
Nem bhar d et al., 2020	Clustered, non-randomized multi-site controlled before-and-after study. Difference-in- differences analysis.	Patient-Centered Medical Homes (Federally Qualified Health Centers), primary care; 12 sites (6 intervention, 6 control) within a primary care network, USA	NR - CT	No randomization. Centers part of one network and from one state. Selection bias for participant sites.			
Jones et al., 2019	Comparative, retrospective cohort study (one intervention group; two comparators). Multivariable logistic regression models (sensitivity analyses).	Patient-Centered Medical Homes, primary care; 510 sites (25 with intervention, 485 control) within the Veterans Health Administration system, USA	Ob s-C	Observational, retrospective study. No baseline outcome measured. Outcome measure does not address homeless-specific concerns and service needs. Limited response rate. Site-specific inclusion criteria for the intervention. No correction for multiple comparisons.			
Tung et al., 2018	Comparative repeated cross-sectional design with serial random samples. Adjusted difference- in-differences analysis and cross-sectional adjusted post-intervention generalized estimating equations.	Safety Net Medical Home Initiative's primary care practices; 24 sites randomly selected from 65 intervention sites (13 sites included in analyses), USA	Ob s-C	Observational study. Baseline patient experience assessed 13 months after intervention began. All analyses at clinic level with inability to follow individual patients over time. Potential selection bias: only 13 of the 24 clinics that completed baseline surveys were included in the analyses as they also completed the post-intervention ones. No correction for multiple comparisons.			
Nguy en et al., 2020	Comparative retrospective study. Adjusted differences-in-differences and regression models (inverse probability weighting in primary analysis; used sensitivity analyses).	Academically-affiliated primary care practices; 50 sites (13 intervention, 37 comparison) affiliated with Harvard Medical School, USA	Ob s-C	Observational study with external comparators with unknown practice patterns. There may have been similar practice changes in the comparison sites in line with general health care reform initiatives. Although adjusted models were used, the study does not control for variation in the size, structure, and composition of the care teams. No correction for multiple comparisons.			

Table 1: Summary of study methods and risk-of-bias assessment: Enhanced care coordination supported for Patient-Centered Medical Homes or other primary care practices.

Legend: RCT: Randomized Controlled Trial; NR-CT: Non-Randomized Controlled Trial; Obs-C: Observational, comparative study. QI: Quality Improvement

			HCSHS		SHEP <sup>a</sup> PACIC <sup>b</sup>		1.0	2	CG- CAHPS								
	Intervention type Enhanced Care Coordination Supports:	Design	Score	Top-Box	Categorical	Score	Top Box	Score	Score	Score	Results (Descriptive Synthesis)						
Noël et al, 2022	Enhanced QI facilitation & coaching (beyond QI toolkit dissemination)	RCT	+ x								Both the intervention and control groups had significant pre-post care improvements in						
	Centralized, guided coaching of the model. Coaching including a site vis use of a care coordination improver dissemination alone (control)	partici it, dista nent to	pant ' ince o oolkit	VA pr coach (inte	rimary ning, an rventi	-care nd cro on) v	clini oss-si s. int	cs un ite lea ernal	der a PC arning or toolkit	difference over time (95% CI ( $-$ 0.47, 0.50)) $-$ i.e., no benefit for adding the more resource-intensive facilitating coaching component.							
Zulman	Augmented care coordination activities for high-risk patients (beyond standard PCMH)	RCT	x	+ x		+	+ x	x			Care coordination outcomes: Intervention had better results than the controls for 2/6 SHEPS items: talked about health goals (73.1% vs. 68.4%; 95% CI= 1.00–1.59) and talk about barriers to taking care of health (60.4% vs. 54.8%; 95% CI= 1.02–1.56). PACIC chronic illness care scale: Intervention group had higher mean scores (2.91 vs. 2.75.						
et al., 2019	Augmented care coordination activities for patients at high risk for hospitalization. Activities include home visits, health coaching, accompaniment to specialists, and intensive social work. These activities were delivered in addition to the standard PCMHs' support (control).									P= .022). Trust outcomes: Intervention better than controls (60.5% vs. 53.1%; 95% CI=1.10–1.66). Satisfaction outcomes: Intervention better than controls for satisfaction with primary care services (36.5% vs. 31.7%; 95% CI= 1.05–1.47) but not for other items or total score. Perceived access outcomes: no significant differences.							
Dorr et al., 2016	Encouraging selection of high-value care coordination QI goals from a 12-item list (beyond standard QI) Encouragement of PCMH sites (stat high value elements in addition to t had IT based milectone conorting of	RCT e of Or he stan	egon	) for s QI fa	selecti	ng Qi ion (c	goal	s froi	+ x m a list o candard (	f 12 วุเ	The intervention performed better in 2 of 11 composite items, including "Follow-up on test results" ( $p$ =.091) and "Patients' rating of the provider" ( $p$ =.091). The control group performed better in "Access to care" ( $p$ =.093).						
Nembh ard et al., 2020	nau n-based milestone reporting, n practice facilitator. Nurses take on added coordination & navigating support for high-risk patients, i.e., socially vulnerable (beyond PCMH).	NR- CT							goais, an		Difference-in-differences analysis showed no significant difference in patient care experiences between the groups ( $p$ = 0.07).						

	Nursing staff taking on additional coordination and navigation support roles for high- risk, socially vulnerable patients. Nurses also led weekly panel management meetings with enrollees' primary care and behavioral health providers. New role supported by training, a "playbook", and new electronic dashboard. Feedback reports provided to nurses on their performance.													
Jones et al., 2019	Tailoring PCMH model to homeless patients (beyond standard model)Homeless-tailored medical home wi care - controls) to enhance access, a placement. For instance, providers a further coordinate care with other s	Obs -C ith tailo address are train services	red r socia ned to	esource al detern o deliver , mental	s (abov ninants care fo health	e stan , and f or the , addio	dard facilit hom ction	VA prim cate hous eless and , and soc	ary sing d ial).	In adjusted analyses, compared to same-facility controls, intervention group had better experiences in access ( $p$ <.001), communication ( $p$ <.01), office staff helpfulness/courtes ( $p$ <.01), provider ratings ( $p$ <.05), and comprehensiveness ( $p$ <.05). When compared to facilities without intervention, significant differences were found for communication and self-management support ( $p$ <.05). Care coordination and shared decision-making were not significantly different for the intervention group versus either comparator.				
Tung et al., 2018	Practice transformation guided and tailored to the safety net context Practice transformation guided to home principles tailored to the s Regional Coordinating Center wi same intervention, the study cor transformation (high-capability s (controls).	Obs -C by eigh safety-l ith prace mpares setting	nt <i>cha</i> net s ctice s higi s) vs	ange co etting, coache h-impro	+ x suppo s. Whi veme prove	t, base rted le le all nt me ment	ed or ocall sites edica	n medica y by a s had the l home ings	al	In the adjusted models, groups did not have significantly different total PACIC scores (95% <i>CI</i> : -1.1 to 16.5). However, at completion of the intervention, a 10-point higher PCMH capability score was associated with 8.9 points higher total PACIC score (95% <i>CI</i> : 3.1–14.7). Greater patient experience scores in favor of the high capability group were also observed in 4/5 subdomains (patient activation, delivery system design, contextual care, and follow-up/coordination).				
Nguyen et al., 2020	Learning collaborative facilitating a primary-care practice transformation Learning collaborative for a primary site, phased, using lump sum payme included regular care team huddles, monthly webinars to discuss QI strat	Obs -C care tr ents) to , coachi tegies.	ansfc ward ng, tr	ormatior a team- riannual	initiat based 1.5-day	ive (m care. T y learr	ulti-y The co ning s	rear, mul ollaborat essions,	+ X ti- ive and	Relative to comparison practices, the communication score in intervention practices increased by 1.47 percentage points on a 100-point scale ( $P = .02$ ) between pre and post periods. No significant immediate improvements in the 5 other composite measures of patient experience of care.				

Table 2: Enhanced care coordination for Patient-Centered Medical Homes (PCMHs) or other primary care practices evaluated from the patient 246

247 experience perspective as a primary outcome. Keys: Results per measure & metric: + significant positive impact; x non-significant impact.

248 249 Legend: HCSHS: Health Care (System) Hassles Scale; SHEP: VA's Survey of Healthcare Experiences of Patients (adapted from CAHPS); PACIC: Patient Assessment of Chronic Illness Care; CSI: Consumer

Satisfaction Index; CG-CAHPS: Consumer Assessment of Healthcare Providers and System Clinician & Group Survey; PES: Patient Experience Survey; RCT: Randomized Controlled Trial; NR-CT: Non-

250 Randomized Controlled Trials; PPT: Pre-Post Test study; Obs-C: Observational, comparative study. QI: Quality Improvement. VA: Veterans Administration. IT: Information Technology. a Jones et al. 251 specifically reported use of the PCMH-SHEPS. <sup>b</sup> Tung et al. reported use of a modified version of the PACIC.

253	coordination QI goals (beyond standard QI facilitation: control group) and found mixed
254	results. <sup>35</sup> These RCTs had some appraised risks of bias (e.g., potential contamination; <sup>33</sup> selection
255	bias <sup>35</sup> ).
256	Among other study types, one non-randomized trial focused on nurses taking on additional care
257	coordination and navigation support for socially vulnerable patients, beyond a PCMH model
258	(control), and found close but no significant differences between groups ( $p$ = 0.07). <sup>36</sup> Finally,
259	three observational studies (tailoring PCMH model to homeless patients; <sup>32</sup> practice
260	transformation tailored to the safety nets; <sup>38</sup> learning collaborative <sup>37</sup> ) showed mixed results and
261	had substantive risks of bias (Table 1).
262	
263	b) Enhanced discharge or follow-up supports - after hospital or surgical care ( $n=8$ )
264	Table 3 summarizes the methods and Table 4 the interventions and results for the studies on
265	enhanced discharge support, including patient education or follow-up. All studies were based in
266	the US.
267	Among the three RCTs, two focused on patient discharge education using the <i>teach back</i>
268	method, after a health literacy assessment <sup>26</sup> or as part of a multi-modal language-concordant
269	discharge support and telephone follow-up. <sup>27</sup> None of the study interventions led to better
269 270	discharge support and telephone follow-up. <sup>27</sup> None of the study interventions led to better experience results than standard care. The other RCT was on a web-based, surgical patient

272 overall experience score.<sup>28</sup> However, several risks of bias were appraised in all the RCTs such

	Design & Analysis	Setting	Risk o	of Bias – level (within each study design) & Described risks
van Eck et al., 2018	Single-site RCT (patient-level randomization) with posttest measure only. T test and analysis of variance on covariates.	Private practice orthopedic clinic, outpatient and ambulatory surgery; Kerlan Jobe Orthopaedic Clinic, California, USA	RCT	Single site with 2 surgeons. Patients not blinded to group allocation. No baseline outcomes. No correction for multiple comparisons. Limited follow-up time (only two weeks of post-assessment).
Griffey et al., 2015	Single-center RCT (patient-level randomization) with intention-to- treat analysis. Multivariable ordinal logistic regression.	Hospital, emergency department; Barnes Jewish Hospital, St. Louis, Missouri, USA	RCT	No baseline outcomes assessed. Single site. Convenience sampling prior to randomization. About 40% loss to follow-up. No random sequence generation for group assignment or allocation concealment. Contamination may have occurred. No correction for multiple comparisons.
Chan et al., 2015	RCT (patient-level randomization stratified by language). χ2 test on the top-box scores. Sensitivity and subgroup analyses.	Safety-net hospital, various units (internal medicine, family medicine, cardiology, neurology); San Francisco General Hospital and Trauma Center, California, USA	RCT	No baseline outcomes assessed. Single site. Only one month of post-intervention data. Medical teams not blinded to presence of intervention, and possible contamination across nurses within the hospital. No correction for multiple comparisons.
Centrella -Nigro & Alexande r, 2017	Nonrandomized trial, two units of one hospital. Independent T tests.	Community Magnet-designated hospital, medical units (intervention unit mainly neurological patients, control unit mainly genitourinary and infectious diseases); Northern New Jersey, USA	NR- CT	No randomization. Clinical differences in units. Difference in baseline outcomes not adjusted for. Baseline characteristics not assessed or adjusted for. Potential for contamination because all units were in a single hospital. No correction for multiple comparisons. Lack of transparent reporting of participant numbers pre- and post-intervention. No combined analysis of between-group and between- time.
Cancino et al., 2017	Nonrandomized posttest design, two control groups with standard discharge (one in same unit and one in different unit). $\chi^2$ test on the top- box scores.	Safety-net hospital, adult inpatient family medicine (adult general medical units as one comparator); Boston Medical Center, Massachusetts, USA	NR- CT	No randomization. No baseline characteristics or outcomes data. No adjustment for covariates or correction for multiple comparisons. Low response rate. Possibility for contamination as some controls were within the same unit. Nurses selected patients (potential selection bias). Results not transferable to sites with higher baseline scores.
Schreiter et al., 2021	Historical cohort of matched controls at patient level. Wilcoxon rank-sum for scores and Fisher exact test for % of top-box scores.	Academic hospital, transition from surgery; University of Wisconsin Hospital, Wisconsin, USA	Obs -C	No randomization. Historical controls even though matched for some clinical indicators. Single site. No baseline outcomes assessed. Unadjusted analyses for outcome of patient experience. No correction for multiple comparisons.
Thum et al., 2022	Pre- and post-test, one organization with multiple units. χ2 on top-box scores and Kruskal–Wallis for the percentile rank change.	Academic tertiary care center and its affiliated community hospital; Thomas Jefferson University Hospital and affiliate, USA	PPT	No control group. Eligibility criteria not clearly described. Single pretest (although long preintervention period). No establishment of stable baseline through multiple measurement time points. No report on the numbers of surveys used for the analysis, only those mailed. No correction for multiple comparisons.
March et al., 2022	Pre- and post- test, single-center retrospective review of a pilot program. χ2 test on the top-box score after T test on baseline characteristics.	Hospital, pharmacy; Methodist University Hospital, Memphis, Tennessee, USA	PPT	Retrospective, single-site pre-post-test analysis not adjusted for covariates. No establishment of stable baseline through multiple measurement time points. No correction for multiple comparisons. Intervention not delivered to all the eligible patients (business hours). No report on the number of patient experience surveys collected. Risks of selected findings: two different designs for two outcomes.

**Table 3:** Summary of study methods and risk-of-bias assessment: discharge education and follow-up or transitional support from an inpatient or

surgical setting. Legend: RCT: Randomized Controlled Trial; NR-CT: Non-Randomized Controlled Trial; PPT: Pre-Post Test study; Obs-C: Observational, comparative study.

				_									
			OAS - CAHPS	OAS - CAHPS CAHPS		CTM-3		НСАНРЅ					
	Intervention Type Discharge education & post- discharge supports:	Design	Score	Score	Score	Top-Box	Top-Box	Percentile	Top-Box	Results - Descriptive Synthesis			
van Eck et al., 2018	Web-based, Surgical Patient Education Tool for pre-operative & follow-up instructions The intervention group received access to routine perioperative instructions. The too surgery, including preoperative instruction precautions. A random subset of intervent the surgery.	RCT an interac I included s, day-of-s ion patien	+ x instruc surgery its also a	b-bas tions expe acces	sed edu 14 day ectations ssed link	cation s befor s, and cs to vi	tool in re and a postope deos de	additi ofter the erative emons	on to ne e trating	Statistically significant higher OAS CAHPS total patient experience score for the intervention vs. control (97 $\pm$ 5 vs. 94 $\pm$ 8; <i>P</i> = .019). At the domain level, groups only significantly differed for "recovery" (92 $\pm$ 13 for intervention vs. 82 $\pm$ 23 for control; <i>P</i> = .001). No significant difference in total patient satisfaction score based on addition of video.			
Griffey et al., 2015	Teach back for discharge instructions, after health literacy assessment Teach back, after a rapid health literacy ass trained medical student Research Assistant education. Control group had usual dischar	RCT sessment, t (RA) alor rge instruc	appliec ngside c	x I to d harge	lischarge e nurses	e instr s who	uctions have ha	. Deliv d <i>teac</i>	ered by th back	No significant differences in the multivariable models for any the 4 items on the standard patient-reported experience measure ( $P$ values from 0.19 to 0.81), even though the intervention group had higher comprehension of medication ( $P$ <.02), self-care ( $P$ <.03), and follow-up instructions ( $P$ <.0001).			
Chan et al., 2015	Language-concordant discharge support, follow up and transitional support, based on the ReEngineered Discharge, including <i>teach back</i> & motivational interviewing Language-concordant, nurse-led, hospital- transition intervention (in addition to usua concordant inpatient nurse visits, motivatio booklet in the patients' native language - c (RED)), and phone follow-up that included, for participants or transportation plans for arranging follow-up services, if necessary.	RCT based, tail I discharg onal inter onceptual for exam schedule	lored dii e planni viewing Ily like t ple, a ni d appoii	schai ing as , tead hat u urse ntme	x rge supp s contro ch back, ised in F organizi ints, inc	x port, fo I grou discha ReEngi ing pos Iuding	x bllow-up p). Lang arge ma neered st-disch instruc	o, and guage- aterials Discha arge s tions f	care s (a arge ervices or	follow-up instructions ( <i>P</i> <.0001). No statistically significant differences between groups in CTM-3 scores (80.5 % vs. 78.5 %; p= 0.18) or HCAHPS discharge communication domain score (74.8 % vs. 68.7 %; <i>p</i> = 0.11), medicine communication (44.5 % vs. 53.1 %; <i>p</i> = 0.13), or nurse communication (67.9 % vs. 64.9 %; <i>p</i> = 0.43) scores. When stratified by language, no significant differences were seen.			
Centre Ila-	Teach back for discharge instructions, after 1-hour training of nurses	NR-CT		+ x						Significant improvement in top box percentages for 1 of 7 tested items for the intervention group, on information about new			

Nigro & Alexan der, 2017	<i>Teach-back</i> implementation, after 1-hour, by nurse supervisors for assessing fidelity.	multi-moc	medicine: 53.2% vs. 78.5%, $p = .025$ . That item had a non-significant change for the control group: $p = .932$											
Cancin o et al., 2017	Use of the ReEngineered Discharge (RED) Toolkit for selected high- complexity cases Implementation of the ReEngineered Disch discharge preparation processes. A trained selected based on high-complexity dischar	NR-CT harge (RED I nurse del ge plannin	) Toolki ivered t ig or his	t: 12- he R tory (	-item pa ED to al of readu	atient bout tl missio	educati hree pa ns.	ion & tients	+ x a day,	Better top box percentages on the item about instructions: 61% vs. 35% same-unit controls ( $p$ <.001) vs 41% other-units controls (both $p$ <.001). No significant difference on the item about feeling ready to be discharged: 45% vs. 35% same-unit controls ( $P$ = .15) vs. 51% other-units' controls ( $p$ = .25).				
Schrei ter et al., 2021	Transitional follow-up support (telephone) after hospital discharge with possible remediation plans and resources Telephone-based, continued follow-up car medication reconciliation, patient educatio providers and ensuring that contact nurse including remediation care (e.g., same-day direct ward admission) if required.	Obs-C e delivered on about k informatic r clinic app	d by nur ey symp on and fo ointmer	rses a otom: ollow nt, vis	after ho s that w v-up app sit to th	spital /arran pointn e eme	dischar; t contac nents ai ergency	ge for: ct with re in pl room,	+ x ace, or	Intervention got greater percentages of top-box scores than controls in 5 of 11 items: asking about having the needed help (100% vs. 93%, p<.01), educational materials (68% vs. 55% $p$ <.01), understanding of responsibilities (69% vs. 59%, $p$ =.02), instructions on whom to call with post-discharge questions (76% vs. 69%; $p$ =.04), and global experience (57% vs. 46%, $p$ =.02). But the readiness for and discharge satisfaction were similar.				
Thum et al. <i>,</i> 2022	Teach back and new discharge summary, with workflow redesign Nurses were trained in <i>teach back</i> interver the role, and a new discharge summary wa record.	PPT ntion, their as created	r workflo – linked	ow w I to a	vas rede hard st	esigned cop in t	+ d to acc the elec	+ x ommo ctronic	date health	Improved top-box percentages: Care Transitions, $52.4\%-54.5\%$ (p<.001); Discharge Information, $87.4\%-90.1\%$ (p<.001). Improved percentile rank $45.2-74.3$ (p=.0202) for Discharge Information. The change in the percentiles for Care Transitions did not reach statistical significance: $56.8-65.8$ (p=.0591).				
March et al., 2022	Pharmacist-led discharge education & follow-up on medication reconciliation       PPT       +         Pharmacist-led medication reconciliation and education, sensitive to health literacy levels, prior and post discharge following alerts from the Electronic Medical Recording system.       +									Significant improvement in the top-box scores (52.6% vs. 67.3%; $p$ = <.001) in the composite of medication-related HCAHPS results and its specific items: "tell you what the medicine was for" (67.7% vs. 81.9%; $p$ = .018), "describe possible medicine side effects" (37.7% vs. 58.9%; $p$ = 0.004), and "understood the purpose of taking medications" (52.3% vs. 63.7%; $p$ = .035).				

278

**Table 4:** Discharge education and post-discharge follow-up (e.g., transitional care supports) after inpatient or surgical care evaluated from the

280 patient experience perspective as a primary outcome.

study; Obs-C: Observational, comparative study.

Legend: OAS-CAHPS: Outpatient and Ambulatory Surgery Consumer Assessment of Healthcare Providers and System; CAHPS: Consumer Assessment of Healthcare Providers and System; CTM-3: Care
 Transitions Measure; HCAHPS: Hospital Consumer Assessment of Healthcare Providers and Systems; RCT: Randomized Controlled Trial; NR-CT: Non-Randomized Controlled Trials; PPT: Pre-Post Test

single-center study,<sup>26-28</sup> limited follow-up,<sup>27 28</sup> or no correction for multiple comparisons (Table
3).<sup>26-28</sup>

287 Two non-randomized trials were found: one on the implementation of *ReEngineered Discharge* 288 for three patients a day that nurses selected as complex cases,<sup>16</sup> the other on *teach back* 289 discharge education by nurses.<sup>31</sup> Both achieved improvements in selected patient experience items (e.g., information provision). In turn, an observational study with historical controls 290 focused on a telephone follow-up after hospital discharge with possible remediation plans or 291 292 rescue resources (e.g., same-day clinic appointment);<sup>15</sup> it achieved higher percentages of top-box 293 scores (i.e., highest score on the given patient experience items) compared to controls for 5 of 294 the 11 patient experience items. Several risks of bias applied to the three studies (Table 3), 295 including risks of contamination and unadjusted analyses.

Finally, two pre-post studies were included, one using *teach back* and other discharge redesigns,<sup>29</sup> the other a pharmacist-led discharge education and follow-up intervention.<sup>30</sup> These provided either positive or mixed results (improvement in top box percentages but not in the percentile ranks with peer providers). In addition to the uncontrolled study design, these studies had multiple substantive risks of bias (Table 3), limiting the ability to draw conclusions.

301

#### 302 Discussion

303 This knowledge synthesis addresses a gap in the literature regarding the impact of care

304 coordination or discharge support improvement on standardized patient experience measures.

305 Health-service improvement activities, including enhanced supports or tailoring for vulnerable

populations, achieved mixed to null improvements in the patient experience with the care
coordination or discharge processes, compared to standard models of care or simpler qualityimprovement supports. This was found by seven studies (including three multi-site RCTs) on
enhanced QI support provided to PCHMs or other primary care practices. It was also found by
eight studies on enhanced discharge support such as patient education, reengineered
discharge, and/or telephone follow-up or transitional support. These latter studies, as a body of
literature, had a greater risk of bias.

313 Several risks of bias were identified in the reviewed literature across the study designs, limiting 314 the internal and external validity of the findings. The studies on enhanced discharge support 315 were typically single center. Even the multi-site studies on enhanced care coordination were 316 performed within a network or organization. Hence, it is unknown how the findings apply 317 elsewhere. For instance, the study with the lowest appraised risks of bias found that the more 318 resource-intensive type of QI support provided to PCMH units was not superior to the dissemination of the QI tool alone for driving improvements in patient experience of care 319 coordination activities. But this finding directly applies to the VHA. The VHA has been subject to 320 systematic QI programs, studies, or activities, <sup>39 40</sup> which may have contributed to building the 321 capacity for effective QI activities across their units as merely supported by an online QI tool, 322 even without a centralized and more costly QI facilitation. The same may not apply to other 323 324 healthcare organizations and development circumstances. Because context is a vital variable in QI science,<sup>41</sup> future studies may consider generating evidence with validity across organizations 325 and networks and their varied sets of circumstances. QI collaboratives or hubs, including those 326

externally funded, might equitably foster the capacity to develop and assess QI activities across
 service delivery organizations, say within a jurisdiction within the US.<sup>42</sup>

329 Varied study designs were included here. While RCTs can produce the most solid evidence 330 when well conducted, a wider range of designs were included to reflect real-world improvement activities and evaluations, including pragmatic constraints.<sup>43</sup> Study limitations 331 within each design were reflected in the appraised risks of bias, especially among the single-332 center discharge support studies. Moreover, studies' delivery of the tested intervention was 333 334 sometimes pragmatically constrained, too. For instance, a pharmacist-led patient education intervention was performed for patients discharged only during particular hours.<sup>30</sup> Similarly, a 335 336 reengineered discharge support program was capped at a number of patients per day as selected by nurses as "complex" cases.<sup>16</sup> These limitations may affect the fidelity of the 337 338 improvement interventions and accuracy of the results and may often reflect budget or other 339 limitations (e.g., human resources, in-house improvement or research expertise) for the conduct or evaluation of QI activities.<sup>43</sup> Percentile ranks (i.e., provider ranks in patient 340 experience performance relative to peers) were used as an additional metric only in one 341 342 included paper (with a pre-post design), showing mixed results while the percentage of top-box scores had positive results.<sup>29</sup> Reporting results in more than one metric of a patient experience 343 measure can be important, especially for percentile ranks in pre-post studies to partly 344 345 overcome the lack of a control group or the use of historical ones. Overall, the inclusion of varied study designs, interventions, and patient experience measures and metrics may inform 346 further improvement activities and its study, at varying levels of available resources. 347

Across study designs, some common intervention approaches were found, such as enhanced 348 349 supports for QI facilitation on care coordination<sup>34 35</sup> or tailored approaches for at-risk populations (e.g., socially vulnerable, homeless),<sup>32 33 36 38</sup> often showing mixed patient 350 experience results relative to standard supports. In turn, discharge support approaches that 351 352 used a reengineered discharge or teach back strategies were also common but with less favorable results when compared to studies and reviews on other outcomes, such as reducing 353 354 readmissions.<sup>44-46</sup> For example, one recent systematic review of *teach back* strategies applied to discharge education identified a 45% reduction in 30-day readmission.<sup>46</sup> Here, studies using the 355 teach back method had mixed results overall<sup>26 27 29 31</sup> but neutral results in the RCTs,<sup>26 27</sup> either 356 as the main strategy<sup>26</sup> or as part of a multi-modal strategy that also included a reengineered 357 358 discharge, motivational interviewing, follow-up and transitional support, and overall language-359 concordant supports.<sup>27</sup> When found to be effective compared to a control intervention, the 360 teach back method was used in isolation and specifically improved the patient experience with receiving information about new medicines.<sup>31</sup> For transitional care post discharge, which was 361 rarely addressed in the body of literature reviewed, the multi-modal intervention that included 362 this component found neutral results,<sup>27</sup> whereas the intervention specifically focused on this 363 component found positive results in five patient experience items.<sup>30</sup> Further research could 364 365 compare the cost-effectiveness of multi-modal and single-method improvement approaches as 366 well as determine whether multi-modal interventions, due to their complexity, are delivered with fidelity to the main active ingredients of the intervention.<sup>47 48</sup> 367

To be impactful on the patient experience, it may be key for studies to assess how patientcentered (e.g., how attentive, respectful, responsive, with genuine interest) was the delivery of

370	a teach back, telephonic transitional support, and other supportive patient-provider
371	communications intervention. This might include how tailored the communication was to the
372	individual patient <sup>18</sup> to avoid the risks of being perceived as just another <i>box-ticking</i> exercise. <sup>49</sup>
373	Finally, in alignment with growing focus on coproduction and on experience-based codesign for
374	health-service improvement activities, <sup>50 51</sup> the patient experience with care can be further
375	engaged all the way through from the intervention codesign to its evaluation. The literature
376	reviewed here had no focus on engaging the patient perspective in the codesign of the
377	improvement interventions.
378	All the included studies were conducted in the US, which may result from multiple factors. First,
379	the CAHPS program in the US provides established, widely used standardized patient
380	experience measures, which was a criterion for inclusion. Second, providers' performance on
381	CAHPS measures is publicly reported and included in some value-based purchasing programs, <sup>11</sup>
382	which may have driven the interest of providers to develop and report these improvement.
383	Third, CAHPS has been a primary quality target of a "person-centered" medical home, with the
384	US National Committee for Quality Assurance recommending administration of the CAHPS
385	PCMH survey for PCMH transformation, and a related study found that CAHPS surveys were
386	considered actionable for PCMH transformation. <sup>52</sup> Fourth, the VHA in the US has its own
387	patient experience measurement system, which was used a standardized assessment measure
388	in studies externally reported and included here. <sup>33 34</sup> Fifth, improvement activities otherwise
389	fitting the study eligibility criteria from other countries may have been conducted but not
390	reported in English or through peer-reviewed journals.

This review has several limitations. First, it excluded articles published in languages other than 391 392 English or in the grey literature, which affects representativeness but reflects peer-reviewed knowledge readily available to an international audience. Second, representativeness is also 393 affected by the exclusion of articles published before 2015, which allowed for a synthesis of 394 395 contemporary approaches in an ever-changing healthcare delivery landscape. Third, for the data extraction, the second reviewer performed only confirmation tasks while two independent 396 397 reviewer roles were used for all other tasks. Fourth, we did not perform a formal grading of the 398 evidence or of the risks of bias within or across study types, within heterogenic health service 399 delivery contexts, interventions, and study methods. Fifth, we did not include studies that used patient experience as a secondary study outcome. While including these studies may have 400 401 added to the pool of evidence, it would have risked including studies in which improving the patient experience of care was not a primary intention. 402

403

#### 404 Conclusion:

Enhanced support for improving care coordination and discharge support had mixed or neutral results for improving the patient experience with care beyond standard care or simpler supports. Substantial risk of bias and lack of comparison between improvement approaches impede firmer conclusions, especially for the studies on enhanced discharge support, which were also typically single center. The multi-site studies on enhanced care coordination were also limited by being performed within an integrated delivery system, network, or organization, limiting the generalizability of the results across organizational and service delivery contexts.

- 412 Improving the person-centeredness of care may require strengthening the capacity to fund,
- 413 design, conduct, and evaluate improvement studies focused on patient experience across
- 414 providers and their networks. Improving the patient experience with care coordination and
- discharge may require the strengthening of the improvement studies, and further engaging the
- 416 patient perspective from the intervention development to its evaluation.

417

# 419 References

421	1. Agency for Healthcare Research and Quality. What Is Patient Experience? Rockville, MD [Available
422	from: <a href="https://www.ahrq.gov/cahps/about-cahps/patient-experience/index.html">https://www.ahrq.gov/cahps/about-cahps/patient-experience/index.html</a> accessed
423	December 2022.
424	2. Berwick DM. A user's manual for the IOM's 'Quality Chasm' report. Health affairs (Project Hope)
425	2002;21(3):80-90. doi: 10.1377/hlthaff.21.3.80 [published Online First: 2002/05/25]
426	3. Nundy S, Cooper LA, Mate KS. The Quintuple Aim for Health Care Improvement: A New Imperative to
427	Advance Health Equity. <i>Jama</i> 2022;327(6):521-22. doi: 10.1001/jama.2021.25181 [published
428	Online First: 2022/01/22]
429	4. Navarro S, Ochoa CY, Chan E, et al. Will Improvements in Patient Experience With Care Impact Clinical
430	and Quality of Care Outcomes?: A Systematic Review. Medical care 2021;59(9):843-56. doi:
431	10.1097/mlr.0000000000001598 [published Online First: 2021/06/25]
432	5. Goldfarb MJ, Bibas L, Bartlett V, et al. Outcomes of Patient- and Family-Centered Care Interventions in
433	the ICU: A Systematic Review and Meta-Analysis. Critical care medicine 2017;45(10):1751-61.
434	doi: 10.1097/ccm.0000000000002624 [published Online First: 2017/07/28]
435	6. Park M, Giap TT, Lee M, et al. Patient- and family-centered care interventions for improving the
436	quality of health care: A review of systematic reviews. International journal of nursing studies
437	2018;87:69-83. doi: 10.1016/j.ijnurstu.2018.07.006 [published Online First: 2018/07/30]
438	7. Rathert C, Wyrwich MD, Boren SA. Patient-centered care and outcomes: a systematic review of the
439	literature. <i>Med Care Res Rev</i> 2013;70(4):351-79. doi: 10.1177/1077558712465774 [published
440	Online First: 20121120]
441	8. Institute of Medicine. Crossing the Quality Chasm: A New Health System for the 21st Century
442	Washington, DC, 2001.
443	9. National Quality Forum. The Care We Need: Driving better outcomes for people and communities:
444	NQF, 2020.
445	10. Agency for Healthcare Research and Quality. The CAHPS Program Rockville, MD [Available from:
446	https://www.ahrq.gov/cahps/about-cahps/cahps-program/index.html accessed December 22.
447	11. Elliott MN, Beckett MK, Lehrman WG, et al. Understanding The Role Played By Medicare's Patient
448	Experience Points System In Hospital Reimbursement. Health Affairs 2016;35(9):1673-80. doi:
449	10.1377/hlthaff.2015.0691
450	12. Jamieson Gilmore K, Corazza I, Coletta L, Allin S. The uses of Patient Reported Experience Measures
451	in health systems: A systematic narrative review. Health policy (Amsterdam, Netherlands)
452	2023;128:1-10. doi: 10.1016/j.healthpol.2022.07.008 [published Online First: 2022/08/08]
453	13. Mihaljevic AL, Doerr-Harim C, Kalkum E, Strunk G. Measuring patient centeredness with German
454	language Patient-Reported Experience Measures (PREM)-A systematic review and qualitative
455	analysis according to COSMIN. <i>PLoS One</i> 2022;17(11):e0264045. doi:
456	10.1371/journal.pone.0264045 [published Online First: 2022/11/30]
457	14. Quigley DD, Elliott MN, Slaughter ME, et al. Shadow Coaching Improves Patient Experience With
458	Care, But Gains Erode Later. Medical care 2021 doi: 10.1097/mlr.0000000000001629 [published
459	Online First: 2021/08/14]
460	15. Schreiter NA, Fisher A, Barrett JR, et al. A telephone-based surgical transitional care program with
461	improved patient satisfaction scores and fiscal neutrality. <i>Surgery</i> 2021;169(2):347-55. doi:
462	10.1016/j.surg.2020.09.015 [published Online First: 2020/10/24]

463 16. Cancino RS, Manasseh C, Kwong L, et al. Project RED impacts patient experience. J Patient Exp 464 2017;4(4):185-90. doi: 10.1177/2374373517714454 [published Online First: 20170616] 465 17. John JR. Jani H. Peters K. et al. The Effectiveness of Patient-Centred Medical Home-Based Models of 466 Care versus Standard Primary Care in Chronic Disease Management: A Systematic Review and 467 Meta-Analysis of Randomised and Non-Randomised Controlled Trials. Int J Environ Res Public 468 Health 2020;17(18) doi: 10.3390/ijerph17186886 [published Online First: 20200921] 469 18. Gonçalves-Bradley DC, Lannin NA, Clemson L, et al. Discharge planning from hospital. Cochrane 470 Database Syst Rev 2022;2(2):Cd000313. doi: 10.1002/14651858.CD000313.pub6 [published 471 Online First: 20220224] 472 19. Coffey A, Leahy-Warren P, Savage E, et al. Interventions to Promote Early Discharge and Avoid 473 Inappropriate Hospital (Re)Admission: A Systematic Review. Int J Environ Res Public Health 474 2019;16(14) doi: 10.3390/ijerph16142457 [published Online First: 20190710] 475 20. Oyesanya TO, Loflin C, Byom L, et al. Transitions of care interventions to improve quality of life 476 among patients hospitalized with acute conditions: a systematic literature review. Health Qual 477 Life Outcomes 2021;19(1):36. doi: 10.1186/s12955-021-01672-5 [published Online First: 478 20210129] 479 21. Braet A, Weltens C, Sermeus W. Effectiveness of discharge interventions from hospital to home on 480 hospital readmissions: a systematic review. JBI Database System Rev Implement Rep 481 2016;14(2):106-73. doi: 10.11124/jbisrir-2016-2381 482 22. Duan-Porter W, Ullman K, Majeski B, et al. Care Coordination Models and Tools-Systematic Review 483 and Key Informant Interviews. J Gen Intern Med 2022;37(6):1367-79. doi: 10.1007/s11606-021-484 07158-w [published Online First: 20211026] 485 23. Centers for Medicare & Medication Services. HCAHPS: Patients' Perspectives of Care Survey 2021 [Available from: HCAHPS: Patients' Perspectives of Care Survey. 486 487 24. Coleman EA, Boult C. Improving the quality of transitional care for persons with complex care needs. 488 J Am Geriatr Soc 2003;51(4):556-7. doi: 10.1046/j.1532-5415.2003.51186.x 489 25. Sezgin D, O'Caoimh R, O'Donovan MR, et al. Defining the characteristics of intermediate care models 490 including transitional care: an international Delphi study. Aging Clin Exp Res 2020;32(11):2399-491 410. doi: 10.1007/s40520-020-01579-z [published Online First: 20200519] 492 26. Griffey RT, Shin N, Jones S, et al. The impact of teach-back on comprehension of discharge 493 instructions and satisfaction among emergency patients with limited health literacy: A 494 randomized, controlled study. J Commun Healthc 2015;8(1):10-21. doi: 495 10.1179/1753807615y.0000000001 496 27. Chan B, Goldman LE, Sarkar U, et al. The effect of a care transition intervention on the patient 497 experience of older multi-lingual adults in the safety net: results of a randomized controlled 498 trial. J Gen Intern Med 2015;30(12):1788-94. doi: 10.1007/s11606-015-3362-y 499 28. van Eck CF, Toor A, Banffy MB, Gambardella RA. Web-based education prior to outpatient 500 orthopaedic surgery enhances early patient satisfaction scores: a prospective randomized 501 controlled study. Orthop J Sports Med 2018;6(1):2325967117751418. doi: 502 10.1177/2325967117751418 [published Online First: 20180126] 503 29. Thum A, Ackermann L, Edger MB, Riggio J. Improving the discharge experience of hospital patients 504 through standard tools and methods of education. J Healthc Qual 2022;44(2):113-21. doi: 505 10.1097/jhg.000000000000325 506 30. March KL, Peters MJ, Finch CK, et al. Pharmacist transition-of-care services improve patient 507 satisfaction and decrease hospital readmissions. J Pharm Pract 2022;35(1):86-93. doi: 508 10.1177/0897190020958264 [published Online First: 20200918] 509 31. Centrella-Nigro AM, Alexander C. Using the teach-back method in patient education to improve 510 patient satisfaction. J Contin Educ Nurs 2017;48(1):47-52. doi: 10.3928/00220124-20170110-10

32. Jones AL, Hausmann LRM, Kertesz SG, et al. Providing positive primary care experiences for 511 512 homeless veterans through tailored medical homes: The Veterans Health Administration's 513 homeless patient aligned care teams. Med Care 2019;57(4):270-78. doi: 514 10.1097/mlr.0000000000001070 33. Zulman DM, Chang ET, Wong A, et al. Effects of intensive primary care on high-need patient 515 516 experiences: survey findings from a Veterans Affairs randomized quality improvement trial. J 517 Gen Intern Med 2019;34(Suppl 1):75-81. doi: 10.1007/s11606-019-04965-0 34. Noël PH, Barnard JM, Leng M, et al. The Coordination Toolkit and Coaching project: cluster-518 519 randomized quality improvement initiative to improve patient experience of care coordination. J 520 Gen Intern Med 2022;37(1):95-103. doi: 10.1007/s11606-021-06926-y [published Online First: 521 20210609] 522 35. Dorr DA, Anastas T, Ramsey K, et al. Effect of a pragmatic, cluster-randomized controlled trial on patient experience with care: The Transforming Outcomes for Patients through Medical Home 523 524 Evaluation and Redesign (TOPMED) study. Med Care 2016;54(8):745-51. doi: 525 10.1097/mlr.000000000000552 526 36. Nembhard IM, Buta E, Lee YSH, et al. A quasi-experiment assessing the six-months effects of a nurse 527 care coordination program on patient care experiences and clinician teamwork in community 528 health centers. BMC Health Serv Res 2020;20(1):137. doi: 10.1186/s12913-020-4986-0 529 [published Online First: 20200224] 530 37. Nguyen KH, Chien AT, Meyers DJ, et al. Team-based primary care practice transformation initiative 531 and changes in patient experience and recommended cancer screening rates. Inquiry 532 2020;57:46958020952911. doi: 10.1177/0046958020952911 533 38. Tung EL, Gao Y, Peek ME, et al. Patient experience of chronic illness care and medical home 534 transformation in safety net clinics. Health Serv Res 2018;53(1):469-88. doi: 10.1111/1475-535 6773.12608 [published Online First: 20170530] 536 39. Braganza MZ, Kilbourne AM. The Quality Enhancement Research Initiative (QUERI) Impact 537 Framework: Measuring the Real-world Impact of Implementation Science. Journal of general 538 internal medicine 2021;36(2):396-403. doi: 10.1007/s11606-020-06143-z [published Online First: 539 20200901] 540 40. Hagedorn H, Hogan M, Smith JL, et al. Lessons learned about implementing research evidence into 541 clinical practice. Experiences from VA QUERI. Journal of general internal medicine 2006;21 Suppl 542 2(Suppl 2):S21-4. doi: 10.1111/j.1525-1497.2006.00358.x 543 41. Coles E, Anderson J, Maxwell M, et al. The influence of contextual factors on healthcare quality 544 improvement initiatives: a realist review. Systematic reviews 2020;9(1):94. doi: 10.1186/s13643-545 020-01344-3 [published Online First: 20200426] 546 42. The Ohio State University Wexner Medical Center. Quality Improvement Hub collaborative aims to 547 enhance health care across Ohio 2023 [Available from: 548 https://wexnermedical.osu.edu/departments/innovations/urology/quality-improvement-hub 549 accessed October 20, 2023. 550 43. Jesus TS, Papadimitriou C, Pinho CS, Hoenig H. Key Characteristics of Rehabilitation Quality 551 Improvement Publications: Scoping Review From 2010 to 2016. Arch Phys Med Rehabil 552 2018;99(6):1141-48.e4. doi: 10.1016/j.apmr.2017.08.491 [published Online First: 20170928] 44. Berkowitz RE, Fang Z, Helfand BK, et al. Project ReEngineered Discharge (RED) lowers hospital 553 554 readmissions of patients discharged from a skilled nursing facility. Journal of the American 555 Medical Directors Association 2013;14(10):736-40. doi: 10.1016/j.jamda.2013.03.004 [published 556 Online First: 2013/04/24] 557 45. Gardner RL, Pelland K, Youssef R, et al. Reducing Hospital Readmissions Through a Skilled Nursing 558 Facility Discharge Intervention: A Pragmatic Trial. Journal of the American Medical Directors

559	Association 2020;21(4):508-12. doi: 10.1016/j.jamda.2019.10.001 [published Online First:
560	2019/12/10]
561	46. Oh EG, Lee HJ, Yang YL, Kim YM. Effectiveness of Discharge Education With the Teach-Back Method
562	on 30-Day Readmission: A Systematic Review. <i>J Patient Saf</i> 2021;17(4):305-10. doi:
563	10.1097/pts.000000000000596
564	47. Meddings J, Manojlovich M, Ameling JM, et al. Quantitative Results of a National Intervention to
565	Prevent Hospital-Acquired Catheter-Associated Urinary Tract Infection: A Pre-Post Observational
566	Study. Ann Intern Med 2019;171(7_Suppl):S38-s44. doi: 10.7326/m18-3534
567	48. Swindle T, Selig JP, Rutledge JM, et al. Fidelity monitoring in complex interventions: a case study of
568	the WISE intervention. Arch Public Health 2018;76:53. doi: 10.1186/s13690-018-0292-2
569	[published Online First: 20180829]
570	49. Jesus TS, Papadimitriou C, Bright FA, et al. Person-Centered Rehabilitation Model: Framing the
571	Concept and Practice of Person-Centered Adult Physical Rehabilitation Based on a Scoping
572	Review and Thematic Analysis of the Literature. Arch Phys Med Rehabil 2022;103(1):106-20. doi:
573	10.1016/j.apmr.2021.05.005 [published Online First: 20210703]
574	50. Green T, Bonner A, Teleni L, et al. Use and reporting of experience-based codesign studies in the
575	healthcare setting: a systematic review. BMJ Qual Saf 2020;29(1):64-76. doi: 10.1136/bmjqs-
576	2019-009570 [published Online First: 20190923]
577	51. Jesus TS, Stern BZ, Struhar J, et al. The use of patient experience feedback in rehabilitation quality
578	improvement and codesign activities: Scoping review of the literature. Clin Rehabil
579	2023;37(2):261-76. doi: 10.1177/02692155221126690 [published Online First: 20220916]
580	52. Quigley DD, Qureshi N, AlMasarweh L, Hays RD. Using CAHPS patient experience data for patient-
581	centered medical home transformation. The American journal of managed care
582	2021;27(9):e322-e29. doi: 10.37765/ajmc.2021.88745 [published Online First: 2021/09/18]

583