

1 **Causal relationship between mental disorders and abdominal aortic aneurysm: insights**
2 **from the genetic perspective**

3

4 **Short title:** Mental disorders and abdominal aortic aneurysm

5

6 Ming-Gang Deng^{1,2}, Chen Chai³, Kai Wang⁴, Zhi-Hui Zhao⁵, Jia-Qi Nie⁶, Fang Liu⁷, Yuehui
7 Liang⁷, Jiewei Liu^{1,2}

8

9 ¹ Department of Psychiatry, Wuhan Mental Health Center, Wuhan 430012, Hubei, China.

10 ² Department of Psychiatry, Wuhan Hospital for Psychotherapy, Wuhan 430012, Hubei,
11 China.

12 ³ Emergency Center, Hubei Clinical Research Center for Emergency and Resuscitation,
13 Zhongnan Hospital of Wuhan University, Wuhan 430071, Hubei, China.

14 ⁴ Department of Public Health, Wuhan Fourth Hospital, Wuhan 430000, Hubei, China.

15 ⁵ School of Nursing, Shanghai Jiao Tong University, Shanghai 200025, China.

16 ⁶ Xiaogan Municipal Center for Disease Control and Prevention, Xiaogan 432000, Hubei,
17 China.

18 ⁷ School of Public Health, Wuhan University, Wuhan 430071, Hubei, China.

19

20 **Correspondence:**

21 **Ming-Gang Deng**

22 ORCID: 0000-0002-3154-6909

23 Email: deng.minggang@outlook.com

24 Address: Department of Psychiatry, Wuhan Mental Health Center, Wuhan 430012, Hubei,
25 China.

26

27 **Jiewei Liu**

28 ORCID: 0009-0009-8265-8984

29 Email: liujiewei@hust.edu.cn

30 Address: Department of Psychiatry, Wuhan Mental Health Center, Wuhan 430012, Hubei,
31 China.

32

33 Word count: 5950

34 **Abstract**

35 **Objective:** This study investigated the genetic link between mental disorders—depression,
36 schizophrenia (SCZ), and bipolar disorder (BIP)—and abdominal aortic aneurysm (AAA).

37 **Methods:** We used global/local genetic correlation analyses and identified shared genomic
38 loci. Bidirectional univariable Mendelian Randomization (UMR) assessed causal directions,
39 with multivariable MR (MVMR) refining the analysis. Mediation analyses examined if
40 known AAA risk factors influenced this relationship.

41 **Results:** Global correlations showed positive links between depression, SCZ, and AAA, but
42 not BIP. Local analyses identified specific genomic regions of correlation. We found 26, 141,
43 and 10 shared loci for AAA with depression, SCZ, and BIP, respectively. UMR indicated
44 significant associations between genetically predicted depression (OR 1.270; 95% CI
45 1.071-1.504; $p = 0.006$) and SCZ (OR 1.047; 95% CI 1.010-1.084; $p = 0.011$) with AAA, but
46 not BIP. These results were confirmed by MVMR analyses. Mediation analyses showed that
47 smoking, hypertension, hyperlipidemia, and coronary atherosclerosis mediated the impact of
48 depression on AAA while smoking mediated SCZ's impact.

49 **Conclusion:** This study provides evidence that genetically predicted depression and SCZ are
50 linked to an increased risk of AAA, mediated by traditional AAA risk factors.

51 **Keywords:** Mental disorders; Abdominal aortic aneurysm; Mendelian Randomization;
52 Genetic association.

53 **1. Introduction**

54 Abdominal aortic aneurysm (AAA) is a prevalent vascular disease marked by localized
55 dilation of the abdominal aorta and is linked to an approximately 90% mortality risk due to
56 rupture¹. While recent open surgical repair and endovascular stent graft therapies have notably
57 enhanced the treatment of AAA, there is currently no approved and effective
58 pharmacotherapy to restrain and prevent AAA progression and rupture^{2,3}. Consequently,
59 primary preventive measures targeting the risk factors of AAA are necessary to reduce its
60 incidence and adverse effects on health.

61 Mental disorders, such as depression, schizophrenia (SCZ), and bipolar disorder (BIP), have
62 been found to lead to various adverse outcomes. For example, our previous studies revealed
63 that individuals with depression or SCZ were more likely to be frail^{4,5}. In addition, current
64 evidence suggests that mental disorders are linked to higher risks of cardiovascular disease,
65 which include but are not limited to heart failure, coronary artery disease, and hypertension⁶⁻⁸.
66 Therefore, investigating the association between mental disorders and AAA is of vital
67 importance for public health.

68 To our knowledge, the associations of AAA with SCZ and BIP are currently unclear, as no
69 studies have reported them. In the case of depression and AAA, a prospective study based on
70 the Korean National Health Insurance System database revealed that patients with AAA were
71 more likely to be depressed⁹. Conversely, depressed individuals were found to have a higher
72 risk of incident AAA in the Norwegian Nord-Trøndelag Health Study¹⁰. Despite this, the
73 causative association between depression and AAA needs further investigation, as
74 observational studies were susceptible to confounding bias, reverse causality, etc.

75 Recent large-scale genome-wide association studies (GWASs) have identified numerous
76 genetic loci associated with depression¹¹, SCZ¹², BIP¹³, and AAA¹⁴. These findings enhance
77 the statistical power and offer the opportunity to explore their genetic associations. In addition,
78 Mendelian Randomization (MR), which uses genetic variants (single nucleotide
79 polymorphisms, SNPs) as instruments, is a powerful method for obtaining robust causal
80 inference¹⁵ and has been widely utilized in recent epidemiological research¹⁶⁻¹⁸. As a result,
81 this study aims to investigate the genetic correlation between AAA and mental disorders and
82 to utilize the MR design to evaluate their causality.

83 **2. Methods and materials**

84 **2.1. Study design**

85 An overview of this research is presented in **Fig 1**. Initially, we explored the genetic

86 association between AAA and mental disorders from three perspectives: global genetic
87 correlation, local genetic correlation, and identification of shared genomic loci. Following that,
88 we utilized a bidirectional MR design to evaluate the causal association between AAA and
89 mental disorders. Furthermore, mediation MR analyses were performed to ascertain whether
90 the causality was mediated by known risk factors.

91 **2.2. Data sources and ethical approval**

92 The data sets utilized in this research were all publicly available from the previous studies,
93 and the details were described in the **Supplementary Methods**. The ethical approvals were
94 all obtained by the original papers, and thus the ethical approval for our study could be
95 waived.

96 **2.3. Genetic association**

97 2.3.1. Global genetic correlation

98 We conducted a global genetic correlation analysis to estimate the correlation between the
99 GWAS summary statistics of AAA and three mental disorders. Two different software tools
100 were utilized to perform this analysis. The first one is Linkage disequilibrium score regression
101 (LDSC, <https://github.com/bulik/ldsc>)¹⁹. We used the cross-trait LDSC which only requires
102 the GWAS summary statistics that account for sample overlap. Cross-trait LDSC analysis
103 requires the GWAS summary statistics to be from the same ethnic group. Since the GWAS
104 summary statistics used in this study are largely from Europeans, we used the 1000 Genomes
105 Project European samples as the reference sample for LD score calculation.

106 Another complementary method for estimating global genetic correlation is the Genetic
107 Covariance Analyzer (GNOVA)²⁰. GNOVA (available at
108 <https://github.com/xtonyjiang/GNOVA>) also requires only the GWAS summary statistics as
109 input to perform global genetic correlation analysis. The major advantage of GNOVA is its
110 ability to include omic datasets (e.g., transcriptomic and epigenomic) in the GNOVA model.
111 The LD score was also calculated using the 1000 Genomes European samples.

112 2.3.2. Local genetic correlation

113 In addition to estimating the global genetic correlation between the GWAS summary statistics
114 of AAA and mental disorders, we also attempted to estimate the local genetic correlation
115 between them in specific genomic regions. To perform this analysis, we used SUPERGNOVA
116 (available at <https://github.com/qlu-lab/SUPERGNOVA>). SUPERGNOVA was developed
117 within the previous GNOVA statistical framework, which includes functional annotation when
118 estimating genetic correlation²¹. Similar to GNOVA, only GWAS summary statistics and an

119 LD calculation reference panel were needed as input to run SUPERGENOVA.

120 2.3.3. Shared genomic loci

121 To enhance the statistical power of detecting loci associated with both AAA and mental
122 disorders, we utilized the conjunctive false discovery rate (conjFDR) method for this
123 analysis. This method is based on standard FDR analysis and aims to identify SNPs that are
124 associated with both target traits (e.g., SNPs that are jointly associated with AAA and SCZ).
125 The conjFDR analysis is implemented in the pleioFDR software (available at
126 <https://github.com/precimed/pleiofdr>), and it runs in the MATLAB environment²². The
127 analysis followed the instructions of the pleioFDR tool with default parameters.

128 2.4. Causal inference

129 2.4.1. Univariate Mendelian Randomization

130 We initially performed the univariate Mendelian Randomization (UMR) to assess the causal
131 association between mental disorders and AAA. The statistical analyses were conducted in
132 two directions: 1) mental disorders as the exposures to check the effect of mental disorders on
133 AAA, and 2) AAA as exposures to evaluate whether people with AAA had a likelihood to
134 develop mental disorders.

135 Independent SNPs ($R^2 < 0.001$, window size = 10MB) were firstly screened at the
136 genome-wide significance threshold ($p < 5 \times 10^{-8}$) to ensure their robust association with
137 exposure and freedom from linkage disequilibrium. Subsequently, the F statistic for each SNP
138 was calculated using the formula β^2/se^2 , and those with an $F < 10$ were considered weak
139 instruments and were excluded. The remaining SNPs were harmonized with the outcome data,
140 and the palindromic genetic variants were eliminated at this stage. If any genetic variants were
141 unavailable in the outcome data, they will be replaced by the proxied SNPs if possible. After
142 the aforementioned procedures, the remaining SNPs were used for causal inference.

143 The inverse variance weighted (IVW) method was used as the primary approach to obtain the
144 causal estimate, assuming the absence of average pleiotropic effects, and in this scenario, it is
145 the most efficient method. Cochran's Q statistic was computed to assess the heterogeneity
146 among the SNPs. If significant heterogeneity was detected ($p < 0.05$), the IVW method would
147 be applied under a random-effects model; otherwise, it would be applied under a fixed-effects
148 model. Alongside the IVW method, several other approaches, such as MR-Egger, weighted
149 median, MR-cML (constrained maximum likelihood), and MR-RAPS (robust adjusted profile
150 score), were employed to evaluate the robustness of the findings. In the meantime, the
151 MR-Egger regression intercept term was utilized to evaluate the presence of horizontal
152 pleiotropy among the instruments.

153 2.4.2. Multivariate Mendelian Randomization

154 The multivariate Mendelian Randomization (MVMR), which employs SNPs associated with
155 various and potentially related exposures, is an extension of UMR that can detect the causal
156 effects of many risk factors simultaneously²³. Considering that the instrumental variables in
157 the UMR may meanwhile be linked to secondary phenotypes and even to the risk factor of
158 outcomes, which may violate the critical assumption of MR, we further carried out the
159 MVMR to ascertain if the UMR estimates were affected by any confounders. The
160 PhenoScanner V2, an expanded tool for searching human genotype-phenotype associations,
161 was employed in advance to screen the secondary phenotypes with which the instrumental
162 variables were associated²⁴. The significant confounding phenotype ($p < 1 \times 10^{-5}$) would then be
163 included as a covariate in the MVMR model. Similar to the UMR analyses, the
164 multivariable-adjusted effect sizes were primarily estimated by the IVW method, and
165 supplemented by the MR-Egger, weighted median, and MR-cML methods.

166 2.4.3. Mediation effect of known risk factors

167 After establishing the causal relationship between mental disorders and AAA, our objective is
168 to delve deeper into whether this causality is mediated by known risk factors. For example,
169 smoking, hypertension, hyperlipidemia, and coronary atherosclerosis (CA) have been
170 identified as risk factors for AAA³. To assess their mediation role, a two-step UMR analysis
171 was performed. The first step involved estimating the causal effects of mental disorders on
172 smoking, hypertension, hyperlipidemia, and coronary atherosclerosis, and the second step
173 involved estimating the causal effects of smoking, hypertension, hyperlipidemia, and CA on
174 AAA. The mediation proportion was calculated based on the causal effect estimated by the
175 IVW method, while other methods, including MR-Egger, weighted median, MR-cML, and
176 MR-RAPS, were employed as sensitivity analyses.

177 **3. Results**

178 **3.1. Genetic association**

179 3.1.1. Global genetic correlation

180 The estimates of the global genetic correlation between mental disorders and AAA are
181 presented in **Fig 2**. The LDSC method indicated that genetic variants related to depression
182 [$r_g = 0.157$, $se = 0.023$, $p = 3.52 \times 10^{-12}$] and SCZ ($r_g = 0.042$, $se = 0.020$, $p = 0.042$) were
183 positively correlated with AAA, while BIP was not ($r_g = 0.013$, $se = 0.023$, $p = 0.591$).
184 Additionally, the GNOVA method estimates further supported these findings, indicating a
185 positive association of AAA with depression ($r_g = 0.152$, $se = 0.019$, $p = 8.41 \times 10^{-16}$) and SCZ

186 ($r_g = 0.053$, $se = 0.017$, $p = 0.002$). Conversely, BIP did not demonstrate a significant
187 association with AAA ($r_g = 0.021$, $se = 0.019$, $p = 0.272$).

188 3.1.2. Local genetic correlation

189 The local genetic correlation analyses revealed that mental disorders exhibited local
190 correlations with AAA in various genomic regions, as detailed in **Table 1**. Following
191 correction for multiple tests, depression was identified to have a positive association with
192 AAA in four genomic regions (chr9: 126927204-128926989, chr10: 106139294-107874146,
193 chr11: 42244050-42854352, and chr18: 77151867-78017074). Additionally, SCZ
194 demonstrated a positive correlation with AAA in 9 out of 10 significant regions. While the
195 global correlation did not indicate an association between BIP and AAA, the local analyses
196 revealed correlations in 7 genomic regions, with 6 of them also showing a positive
197 association.

198 3.1.3. Shared genetic variants

199 In line with the preceding analyses, a multitude of shared genetic variants have been discerned
200 between mental disorders and AAA. The findings are meticulously outlined in **Fig. 3** and
201 **Table S1**. Specifically, there are 26 independent SNPs shared between depression and AAA,
202 with 21 of them demonstrating a consistent directional effect on both conditions. Similarly,
203 SCZ and AAA exhibit the sharing of 141 independent loci, among which 21 loci manifest an
204 opposing effect. Notably, only 10 genetic variants are concurrently associated with BIP and
205 AAA, with 4 of them displaying the same directional effect.

206 3.2. Causal inference

207 3.2.1. Univariate Mendelian Randomization

208 We initially examined the bidirectional causal relationship between mental disorders and AAA
209 using the UMR, and the results are illustrated in **Fig. 4**. In the forward analyses, the
210 MR-Egger regression intercept suggested the absence of horizontal pleiotropy among the
211 selected SNPs (p values all > 0.05). The Cochran's Q test revealed significant heterogeneity
212 among the instruments concerning depression (Cochran's Q = 163.448, $p < 0.001$), SCZ
213 (Cochran's Q = 333.578, $p < 0.001$), and BIP (Cochran's Q = 32.604, $p = 0.002$). Consequently,
214 the primary causal inference utilized the random-effect IVW method. The IVW method
215 revealed that people with depression (OR: 1.270, 95% CI: 1.071-1.504, $p = 0.006$) and SCZ
216 (OR: 1.047, 95% CI: 1.010-1.084, $p = 0.011$) have a heightened susceptibility to AAA, while
217 no causal effect of BIP on AAA was found (OR: 0.980, 95% CI: 0.896-1.121, $p = 0.653$).

218 The findings mentioned above are supported by the weighted median, MR-CML, and
219 MR-RAPS methods.

220 For the reverse analyses, the MR-Egger regression intercept also demonstrated there was no
221 horizontal pleiotropy among the AAA-associated instrumental variables, and obvious
222 heterogeneity was identified. The IVW method revealed that genetically predicted AAA
223 showed no association with the risks of depression (OR: 1.010, 95% CI: 0.995-1.025, $p =$
224 0.190), SCZ (OR: 1.007, 95% CI: 0.968-1.048, $p = 0.740$), as well as BIP (OR: 0.989, 95% CI:
225 0.944-1.037, $p = 0.646$), and these null associations were consistently supported by estimates
226 from the MR-Egger, weighted median, MR-cML, and MR-RAPS methods.

227 3.2.2. Multivariate Mendelian Randomization

228 After performing the UMR, we proceeded with the MVMR to evaluate potential bias in the
229 causal estimates due to confounding effects. The PhenoScanner revealed that instruments
230 associated with mental disorders were predominantly linked to body mass index (BMI) (**Table**
231 **S2**). Additionally, SNPs associated with AAA were found to be correlated with BMI, coronary
232 heart disease (CAD), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein
233 cholesterol (LDL-C), and total cholesterol (TC) (**Table S3**). Consequently, we conducted
234 bidirectional MVMR to evaluate the multivariate-adjusted causal estimates between mental
235 disorders and AAA. The forward analyses aimed to assess the effect of mental disorders on
236 AAA with adjustments for BMI, while the reverse analyses aimed to evaluate the effect of
237 AAA on mental disorders with adjustments for BMI, CAD, HDL-C, LDL-C, and TC.

238 The results of MVMR are presented in **Table S4**. The IVW method revealed that individuals
239 with depression (OR: 1.315, 95% CI: 1.095-1.578, $p = 0.003$) and SCZ (OR: 1.042, 95%
240 CI: 1.0004-1.084, $p = 0.048$) were correlated with a higher risk of AAA even after controlling
241 the effect of BMI, while people with BIP were not (OR: 1.014, 95% CI: 0.955- 1.076, $p =$
242 0.652). Consistent with the results of UMR, the IVW methods also presented that genetically
243 predicted AAA still showed no association with the risks of depression, SCZ, and BIP after
244 adjustments for BMI, CAD, HDL-C, LDL-C, and TC. These bidirectional relationships were
245 supported by the sensitivity analysis methods.

246 3.2.3. Mediation Mendelian Randomization

247 Since the UMR and MVMR only supported the causal effects of depression and SCZ on AAA,
248 we would only investigate whether these two causalities were mediated by the known risk
249 factors of AAA. As described in the methods section, we selected smoking, hypertension,
250 hyperlipidemia, and CA as the four candidate mediating factors.

251 The results of mediation MR analyses are presented in **Table S5**. The first step MR analyses
252 presented that genetically predicted depression was associated with increased odds ratio of
253 smoking (OR: 1.236, 95% CI: 1.140-1.340, $p < 0.001$), hypertension (OR: 1.206, 95% CI:

254 1.069-1.361, $p = 0.002$), hyperlipidemia (OR: 1.200, 95% CI: 1.037-1.388, $p = 0.015$), and
255 CA (OR: 1.210, 95% CI: 1.078-1.359, $p = 0.001$). In contrast, genetically predicted SCZ was
256 only associated with an increased likelihood of smoking (OR: 1.047, 95% CI: 1.023-1.071, p
257 <0.001). The second step MR analyses demonstrated that genetically predicted smoking (OR:
258 1.610, 95% CI: 1.449-1.788, $p <0.001$), hypertension (OR: 1.218, 95% CI: 1.142-1.300, p
259 <0.001), hyperlipidemia (OR: 1.547, 1.288-1.857, $p <0.001$), and CA (OR: 1.385, 95% CI:
260 1.204-1.594, $p <0.001$) were all associated with elevated risks of AAA. Based on the results
261 of mediation MR findings, the causal effect of depression on AAA was mediated by smoking,
262 hypertension, hyperlipidemia, and CA, while the causal effect of SCZ on AAA was mediated
263 by smoking only (**Fig 5**).

264 **4. Discussion**

265 We conducted an investigation into the relationship between mental disorders and AAA from
266 the genetic perspective. Our findings provided solid evidence of the genetic correlation
267 between AAA and depression as well as SCZ. Besides, the causal effects of depression and
268 SCZ on AAA were identified, with known risk factors for AAA playing a mediating role.

269 Few studies have investigated the associations between mental disorders and AAA, with the
270 evidence supporting the correlations between depression and AAA being limited. Furthermore,
271 a study involving 574 participants from four centers in Australia suggested an insignificant
272 association between depression and AAA growth²⁵, contradicting conclusions drawn in
273 previous studies from another perspective. Given the scarcity of research and the
274 inconsistency of results, we conducted comprehensive genetic association, and univariate,
275 multivariate as well as mediating MR to evaluate the correlation and causal links between
276 mental disorders and AAA.

277 The global genetic correlation analyses indicated positive associations of AAA with
278 depression and SCZ, and meanwhile, numerous locally correlated genomic regions and shared
279 loci were identified. The genetic association analyses presented that AAA had plenty of
280 genetic basis with mental disorders. The results of univariate and multivariate MR analyses
281 demonstrated robust and statistically significant associations of depression, SCZ, and AAA,
282 and we further extended our investigation through mediation analysis to delineate potential
283 pathways underlying these associations. By incorporating well-established risk factors of
284 AAA as potential mediators, significant mediation effects of smoking, hyperlipidemia, CA,
285 and hypertension were identified in the causal pathways between depression and AAA, and
286 the proportions of mediation effects were calculated as 42.22%, 32.20%, 26.05%, and 15.41%,

287 respectively. In addition, smoking exhibited a substantial mediation effect proportion of 47.60%
288 in the association between SCZ and AAA. These findings underscore the pivotal role of
289 smoking in the nexus between mental disorders and AAA, offering crucial insights for further
290 etiological explorations.

291 While the pathological mechanism of AAA remains elusive, existing literature has elucidated
292 a series of contributing processes, notably the infiltration and subsequent destruction of the
293 aortic wall mediated by immune and inflammatory mechanisms²⁶. Both innate and adaptive
294 immune cells, including macrophages and T cells, have undergone extensive scrutiny in the
295 pathogenesis of AAA, with several inflammasomes suggested to be involved in both the
296 inhibition and amelioration of AAA^{27,28}. NF- κ B, the primary regulator of inflammation and
297 immune response, is activated by multiple factors including angiotensin II, triggering
298 macrophage infiltration, inflammation, and subsequent aortic deterioration²⁹. Numerous
299 studies have explored the impact of pharmacological inhibition of nucleotide-binding
300 oligomerization domain-like receptor family protein 3 (NLRP3) inflammasome on
301 experimental AAAs, revealing preventive effects characterized by reduced AAA diameter and
302 incidence³⁰⁻³², etc. Furthermore, the remarkable parallels between the immune system and the
303 brain, both intricately integrated and complex systems associated with memory, imply
304 intimate connections between immune responses and mental disorders. For instance, some
305 researchers consider depression as an inflammatory disease^{33,34}, with inflammation likely
306 playing a critical role in heightening individual susceptibility to depression³⁵. Similarly,
307 extensive reviews have highlighted the central role of inflammation and immunity in SCZ,
308 findings supported by both human and animal studies^{36,37}. Several two-sample MR studies
309 indicate the causal associations of inflammatory biomarkers such as C-reactive protein and
310 soluble interleukin-6 receptor with SCZ, with anti-inflammation strategies emerging as
311 therapeutic considerations^{38,39}. Therefore, inflammation and immune responses may function
312 as a conduit linking mental disorders and AAA.

313 Similar to other cardiovascular diseases (CVD), the relationship between mental disorders and
314 AAA may also be explained by the shared etiological factors, encompassing genetic,
315 biological, and behavioral mechanisms. Early in 2002, Bondy et al. found that the same allelic
316 combination of two genes could increase the vulnerability for depressive disorder while also
317 being associated with a higher risk of myocardial infarction⁴⁰, suggesting the genetic
318 associations between depression and CVD. Molecular genetic studies have further
319 corroborated modest genetic correlations between severe mental illness and CVD⁴¹.

320 Furthermore, biological constituents such as neutrophil gelatinase associated lipocalin⁴² and
321 physiological processes like platelet receptors⁴³ and function have been implicated as
322 mediating factors linking CVD and depression. Moreover, the overlap of behavioral
323 determinants including smoking, physical inactivity, obesity, and poor diet was observed
324 among participants affected by mental disorders and CVD. Similarly, GWAS studies have
325 unveiled extensive genetic convergence between SCZ and CVD risk factors, with notable
326 associations observed with smoking initiation and body mass index⁴⁴.

327 Our study had several strengths worth pointing out. Firstly, we utilized the genetic data from
328 the large-scale GWASs, which could significantly enhance the statistical power and avoid the
329 confounding bias and reverse causality of observational studies. Additionally, we conducted
330 two parts of analyses, including genetic correlation and causality inference, and the results
331 provided consistent evidence on the causal effects of depression and SCZ on AAA.
332 Furthermore, we identified the mediation effect, which highlighted the interactive impacts of
333 mental disorders and other risk factors on the risk of AAA.

334 Despite these advantages, our findings should be interpreted in the context of limitations.
335 Firstly, the most recent GWAS of BIP have identified various significant SNPs⁴⁵, but the
336 samples overlapped with the AAA. To avoid the causal inference bias due to sample overlap,
337 we selected the genetic variants associated with BIP from the GWAS with relatively lower
338 sample sizes¹³. Further studies were warranted to check the relationship between BIP and
339 AAA. In addition, only the summary-level data could be obtained thus making us unable to
340 perform the stratified analyses. At the same time, current knowledge indicates that the
341 prevalence of AAA in males was higher than in females. Further research was needed to
342 confirm our findings in specific sub-populations. Lastly, the samples were all of European
343 ancestry, making our findings may not apply to other populations.

344 **5. Conclusion**

345 Genetically predicted depression and schizophrenia were associated with elevated risk of
346 abdominal aortic aneurysm, and the known risk factors of abdominal aortic aneurysm played
347 mediation roles.

348 **6. Declarations**

349 **6.1. Acknowledgement**

350 None.

351 **6.2. Ethical approval**

352 The datasets used in this study were all publicly available from previous studies, and the

353 ethical approvals were obtained by the original researchers. Therefore, the ethical approval for
354 this study could be waived.

355 **6.3. Consent for publication**

356 Not applicable

357 **6.4. Data availability**

358 The datasets utilized in this study are publicly available; the details can be found in the
359 supplementary method.

360 **6.5. Code availability**

361 ldscr (<https://github.com/bulik/ldsc>), GNOVA (<https://github.com/xtonyjiang/GNOVA>), SU
362 PERGNOVA (<https://github.com/qiu-lab/SUPERGNOVA>), pleioFDR ([https://github.com/pr
364 ecimed/pleiofdr](https://github.com/pr
363 ecimed/pleiofdr)), TwoSampleMR (<https://mrcieu.github.io/TwoSampleMR/>), MendelianRa
364 ndomization (<https://github.com/cran/MendelianRandomization>), and mr.raps ([https://github
365 b.com/qingyuanzhao/mr.raps](https://github
365 b.com/qingyuanzhao/mr.raps)).

366 **6.6. Competing interests**

367 The authors declare no conflict of interest.

368 **6.7. Funding**

369 This work receives no external funding.

370 **6.8. Author contributions**

371 **Ming-Gang Deng:** Conceptualization, Methodology, Data Curation, Formal Analysis, and
372 Writing - Original Draft; **Chen Chai:** Writing - Original Draft; Kai Wang: Writing - Original
373 Draft; **Zhi-Hui Zhao:** Writing – review & editing; **Jia-Qi Nie:** Writing – review & editing;
374 **Fang Liu:** Writing – review & editing; **Yuehui Liang:** Writing – review & editing; **Jiewei
375 Liu:** Conceptualization, and Writing - Original Draft.

376 **7. Reference**

- 377 1. Summers KL, Kerut EK, Sheahan CM, Sheahan MG, 3rd. Evaluating the prevalence of
378 abdominal aortic aneurysms in the United States through a national screening database.
379 *Journal of vascular surgery*. 2021;73(1):61-68.
- 380 2. Davis FM, Rateri DL, Daugherty A. Abdominal aortic aneurysm: novel mechanisms and
381 therapies. *Current opinion in cardiology*. 2015;30(6):566-573.
- 382 3. Zhang F, Li K, Zhang W, et al. Ganglioside GM3 Protects Against Abdominal Aortic
383 Aneurysm by Suppressing Ferroptosis in Vascular Smooth Muscle Cells. *Circulation*.
384 2023.

- 385 4. Deng MG, Liu F, Liang Y, Wang K, Nie JQ, Liu J. Association between frailty and
386 depression: A bidirectional Mendelian randomization study. *Science advances*.
387 2023;9(38):eadi3902.
- 388 5. Deng M-G, Wang K, Liu F, Zhou X, Liu J. The Relationship Between Frailty and
389 Schizophrenia: A Genetic Association and Mendelian Randomization Study. 2023.
- 390 6. Li GH, Cheung CL, Chung AK, et al. Evaluation of bi-directional causal association
391 between depression and cardiovascular diseases: a Mendelian randomization study.
392 *Psychological medicine*. 2022;52(9):1765-1776.
- 393 7. Veeneman RR, Vermeulen JM, Abdellaoui A, et al. Exploring the Relationship Between
394 Schizophrenia and Cardiovascular Disease: A Genetic Correlation and Multivariable
395 Mendelian Randomization Study. *Schizophrenia bulletin*. 2022;48(2):463-473.
- 396 8. So HC, Chau KL, Ao FK, Mo CH, Sham PC. Exploring shared genetic bases and causal
397 relationships of schizophrenia and bipolar disorder with 28 cardiovascular and
398 metabolic traits. *Psychological medicine*. 2019;49(8):1286-1298.
- 399 9. Kim MH, Yoo JH, Cho HJ, et al. Increased depression risk in patients with abdominal
400 aortic aneurysm: a nationwide cohort study. *Annals of surgical treatment and research*.
401 2021;101(5):291-298.
- 402 10. Nyrønning L, Stenman M, Hultgren R, Albrektsen G, Videm V, Mattsson E. Symptoms
403 of Depression and Risk of Abdominal Aortic Aneurysm: A HUNT Study. *Journal of the*
404 *American Heart Association*. 2019;8(21):e012535.
- 405 11. Howard DM, Adams MJ, Clarke TK, et al. Genome-wide meta-analysis of depression
406 identifies 102 independent variants and highlights the importance of the prefrontal brain
407 regions. *Nature neuroscience*. 2019;22(3):343-352.
- 408 12. Trubetskoy V, Pardiñas AF, Qi T, et al. Mapping genomic loci implicates genes and
409 synaptic biology in schizophrenia. *Nature*. 2022;604(7906):502-508.
- 410 13. Stahl EA, Breen G, Forstner AJ, et al. Genome-wide association study identifies 30 loci
411 associated with bipolar disorder. *Nature genetics*. 2019;51(5):793-803.
- 412 14. Roychowdhury T, Klarin D, Levin MG, et al. Genome-wide association meta-analysis
413 identifies risk loci for abdominal aortic aneurysm and highlights PCSK9 as a therapeutic
414 target. *Nature genetics*. 2023;55(11):1831-1842.
- 415 15. Emdin CA, Khera AV, Kathiresan S. Mendelian Randomization. *Jama*.
416 2017;318(19):1925-1926.

- 417 16. Deng MG, Cui HT, Nie JQ, Liang Y, Chai C. Genetic association between circulating
418 selenium level and the risk of schizophrenia in the European population: A two-sample
419 Mendelian randomization study. *Frontiers in nutrition*. 2022;9:969887.
- 420 17. Deng MG, Liu F, Wang K, Liang Y, Nie JQ, Chai C. Genetic association between
421 coffee/caffeine consumption and the risk of obstructive sleep apnea in the European
422 population: a two-sample Mendelian randomization study. *European journal of nutrition*.
423 2023;62(8):3423-3431.
- 424 18. Yu X, Deng MG, Tang ZY, Zhang ZJ. Urticaria and increased risk of rheumatoid
425 arthritis: a two-sample Mendelian randomisation study in European population. *Modern
426 rheumatology*. 2022;32(4):736-740.
- 427 19. Bulik-Sullivan B, Finucane HK, Anttila V, et al. An atlas of genetic correlations across
428 human diseases and traits. *Nature genetics*. 2015;47(11):1236-1241.
- 429 20. Lu Q, Li B, Ou D, et al. A Powerful Approach to Estimating Annotation-Stratified
430 Genetic Covariance via GWAS Summary Statistics. *American journal of human
431 genetics*. 2017;101(6):939-964.
- 432 21. Zhang Y, Lu Q, Ye Y, et al. SUPERGNOVA: local genetic correlation analysis reveals
433 heterogeneous etiologic sharing of complex traits. *Genome biology*. 2021;22(1):262.
- 434 22. Andreassen OA, Thompson WK, Schork AJ, et al. Improved detection of common
435 variants associated with schizophrenia and bipolar disorder using pleiotropy-informed
436 conditional false discovery rate. *PLoS genetics*. 2013;9(4):e1003455.
- 437 23. Sanderson E. Multivariable Mendelian Randomization and Mediation. *Cold Spring
438 Harbor perspectives in medicine*. 2021;11(2).
- 439 24. Kamat MA, Blackshaw JA, Young R, et al. PhenoScanner V2: an expanded tool for
440 searching human genotype-phenotype associations. *Bioinformatics (Oxford, England)*.
441 2019;35(22):4851-4853.
- 442 25. Thanigaimani S, Phie J, Quigley F, et al. Association of Diagnosis of Depression and
443 Small Abdominal Aortic Aneurysm Growth. *Annals of vascular surgery*.
444 2022;79:256-263.
- 445 26. Márquez-Sánchez AC, Koltsova EK. Immune and inflammatory mechanisms of
446 abdominal aortic aneurysm. *Frontiers in immunology*. 2022;13:989933.
- 447 27. Yuan Z, Lu Y, Wei J, Wu J, Yang J, Cai Z. Abdominal Aortic Aneurysm: Roles of
448 Inflammatory Cells. *Frontiers in immunology*. 2020;11:609161.

- 449 28. Raffort J, Lareyre F, Clément M, Hassen-Khodja R, Chinetti G, Mallat Z. Monocytes
450 and macrophages in abdominal aortic aneurysm. *Nature reviews Cardiology*.
451 2017;14(8):457-471.
- 452 29. Li Z, Kong W. Cellular signaling in Abdominal Aortic Aneurysm. *Cellular signalling*.
453 2020;70:109575.
- 454 30. Tsuruda T, Kato J, Hatakeyama K, et al. Adventitial mast cells contribute to
455 pathogenesis in the progression of abdominal aortic aneurysm. *Circulation research*.
456 2008;102(11):1368-1377.
- 457 31. Yamanouchi D, Morgan S, Kato K, Lengfeld J, Zhang F, Liu B. Effects of caspase
458 inhibitor on angiotensin II-induced abdominal aortic aneurysm in apolipoprotein
459 E-deficient mice. *Arteriosclerosis, thrombosis, and vascular biology*.
460 2010;30(4):702-707.
- 461 32. Kaneko H, Anzai T, Morisawa M, et al. Resveratrol prevents the development of
462 abdominal aortic aneurysm through attenuation of inflammation, oxidative stress, and
463 neovascularization. *Atherosclerosis*. 2011;217(2):350-357.
- 464 33. Berk M, Williams LJ, Jacka FN, et al. So depression is an inflammatory disease, but
465 where does the inflammation come from? *BMC medicine*. 2013;11:200.
- 466 34. Halaris A. Inflammation and depression but where does the inflammation come from?
467 *Current opinion in psychiatry*. 2019;32(5):422-428.
- 468 35. Beurel E, Toups M, Nemeroff CB. The Bidirectional Relationship of Depression and
469 Inflammation: Double Trouble. *Neuron*. 2020;107(2):234-256.
- 470 36. Müller N. Inflammation in Schizophrenia: Pathogenetic Aspects and Therapeutic
471 Considerations. *Schizophrenia bulletin*. 2018;44(5):973-982.
- 472 37. Khandaker GM, Cousins L, Deakin J, Lennox BR, Yolken R, Jones PB. Inflammation
473 and immunity in schizophrenia: implications for pathophysiology and treatment. *The
474 lancet Psychiatry*. 2015;2(3):258-270.
- 475 38. Hartwig FP, Borges MC, Horta BL, Bowden J, Davey Smith G. Inflammatory
476 Biomarkers and Risk of Schizophrenia: A 2-Sample Mendelian Randomization Study.
477 *JAMA psychiatry*. 2017;74(12):1226-1233.
- 478 39. Williams JA, Burgess S, Suckling J, et al. Inflammation and Brain Structure in
479 Schizophrenia and Other Neuropsychiatric Disorders: A Mendelian Randomization
480 Study. *JAMA psychiatry*. 2022;79(5):498-507.

- 481 40. Bondy B, Baghai TC, Zill P, et al. Combined action of the ACE D- and the G-protein
482 beta3 T-allele in major depression: a possible link to cardiovascular disease? *Molecular*
483 *psychiatry*. 2002;7(10):1120-1126.
- 484 41. Amare AT, Schubert KO, Klingler-Hoffmann M, Cohen-Woods S, Baune BT. The
485 genetic overlap between mood disorders and cardiometabolic diseases: a systematic
486 review of genome wide and candidate gene studies. *Translational psychiatry*.
487 2017;7(1):e1007.
- 488 42. Gouweleeuw L, Naudé PJ, Rots M, DeJongste MJ, Eisel UL, Schoemaker RG. The role
489 of neutrophil gelatinase associated lipocalin (NGAL) as biological constituent linking
490 depression and cardiovascular disease. *Brain, behavior, and immunity*. 2015;46:23-32.
- 491 43. Amadio P, Zarà M, Sandrini L, Ieraci A, Barbieri SS. Depression and Cardiovascular
492 Disease: The Viewpoint of Platelets. *International journal of molecular sciences*.
493 2020;21(20).
- 494 44. Rødevand L, Rahman Z, Hindley GFL, et al. Characterizing the Shared Genetic
495 Underpinnings of Schizophrenia and Cardiovascular Disease Risk Factors. *The*
496 *American journal of psychiatry*. 2023;180(11):815-826.
- 497 45. Mullins N, Forstner AJ, O'Connell KS, et al. Genome-wide association study of more
498 than 40,000 bipolar disorder cases provides new insights into the underlying biology.
499 *Nature genetics*. 2021;53(6):817-829.
- 500

501 **Table 1. Locally correlated genomic regions between mental disorders and abdominal aortic aneurysm.**

Trait1&Trait2	chr	start	end	rho	corr	h ² _1	h ² _2	var	m	p-value	p.adjust
Depression & AAA	9	126927204	128926989	7.98E-05	0.945	7.26E-05	9.81E-05	2.47E-10	644	3.78E-07	8.57E-04
Depression & AAA	10	106139294	107874146	6.17E-05	1.112	1.40E-05	2.19E-04	1.72E-10	758	2.58E-06	2.93E-03
Depression & AAA	11	42244050	42854352	4.31E-05	1.130	2.05E-05	7.07E-05	9.28E-11	277	7.83E-06	5.92E-03
Depression & AAA	18	77151867	78017074	4.62E-05	1.159	2.74E-05	5.79E-05	1.12E-10	323	1.30E-05	7.37E-03
SCZ & AAA	2	73174848	74121000	7.41E-05	0.907	1.61E-05	4.15E-04	3.73E-10	277	1.24E-04	2.84E-02
SCZ & AAA	2	232791948	233964632	-1.38E-04	-1.021	2.56E-05	7.19E-04	6.93E-10	514	1.44E-07	1.64E-04
SCZ & AAA	5	73508509	75240469	1.21E-04	0.792	9.10E-05	2.57E-04	9.36E-10	766	7.55E-05	2.46E-02
SCZ & AAA	10	106139294	107874146	1.16E-04	1.355	1.40E-05	5.21E-04	7.46E-10	759	2.28E-05	1.30E-02
SCZ & AAA	10	130282817	131192968	1.08E-04	1.746	3.02E-05	1.27E-04	7.75E-10	576	1.02E-04	2.59E-02
SCZ & AAA	11	16790799	18788126	1.76E-04	0.722	6.85E-05	8.64E-04	1.90E-09	974	5.51E-05	2.19E-02
SCZ & AAA	12	2176265	2886299	9.31E-05	0.815	6.39E-06	2.04E-03	5.36E-10	367	5.76E-05	2.19E-02
SCZ & AAA	14	102341650	104759919	2.23E-04	0.671	8.11E-05	1.37E-03	2.12E-09	801	1.28E-06	9.75E-04
SCZ & AAA	15	78640760	79259009	2.80E-04	0.777	1.88E-04	6.91E-04	1.27E-09	202	3.75E-15	8.56E-12
SCZ & AAA	17	820511	1480393	1.05E-04	1.043	2.83E-05	3.56E-04	7.27E-10	264	1.01E-04	2.59E-02
BIP & AAA	2	73174848	74121000	8.47E-05	1.036	1.61E-05	4.15E-04	4.64E-10	277	8.37E-05	3.01E-02
BIP & AAA	8	24457122	25461315	-2.03E-04	-0.944	7.38E-05	6.29E-04	1.71E-09	466	9.08E-07	1.04E-03
BIP & AAA	12	47505424	48267360	1.90E-04	1.252	5.57E-05	4.15E-04	1.27E-09	379	9.37E-08	2.14E-04
BIP & AAA	15	66948951	68640917	2.27E-04	1.046	9.96E-05	4.72E-04	2.86E-09	789	2.18E-05	1.10E-02
BIP & AAA	15	78640760	79259009	1.71E-04	0.873	1.88E-04	2.03E-04	1.63E-09	202	2.41E-05	1.10E-02
BIP & AAA	19	10030690	11279257	2.63E-04	0.585	1.35E-04	1.49E-03	4.52E-09	482	9.23E-05	3.01E-02
BIP & AAA	22	43187900	43645335	1.21E-04	1.231	1.47E-05	6.53E-04	8.14E-10	257	2.31E-05	1.10E-02

502 **Abbreviations:** SCZ, schizophrenia; AAA, abdominal aortic aneurysm; BIP, bipolar disorder; chr, chromosome; start/end, the start or end
503 position of the genomic region from the input genome partition file; rho, the estimation of local genetic covariance; corr, the estimation of local
504 genetic correlation; h²_1/h²_2, The estimation of local heritability of trait 1/2; var, the variance of the estimation of local genetic covariance; m,
505 the number of SNPs involved in the estimation of local genetic covariance in the genomic region.

506 **Figure Legend**

507

508 **Fig 1. An overview of study design**

509

510 **Fig 2. Global genetic correlation between mental disorders and abdominal aortic aneurysm.**

511 LDSC, linkage disequilibrium score regression; GNOVA, GeNetic cOVariance Analyzer; SCZ, schizophrenia; BIP, bipolar disorder; *** indicates $p < 0.001$; ** indicates $p < 0.01$; * indicates $p < 0.05$; ^{ns} indicates no significance.

514

515 **Fig 3. ConjFDR Manhattan plots. The red dashed line represents the threshold of p -value for conjFDR < 0.05, and the black points are the independent SNPs; SCZ, schizophrenia; BIP,**

517 bipolar disorder; AAA, abdominal aortic aneurysm.

518

519 **Fig 4. The univariate bidirectional causal inference between mental disorders and abdominal aortic aneurysm. MDD, major depression disorder; SCZ, schizophrenia; BIP, bipolar disorder;**

521 AAA, abdominal aortic aneurysm.

522

523 **Fig 5. Mediation effects of risk factors in the causal associations of abdominal aortic aneurysm with depression and schizophrenia. SNPs, single nucleotide polymorphisms; AAA,**


525 abdominal aortic aneurysm; CA, coronary atherosclerosis.

Section 1: Genetic association


1. **Global correlation:** LDSC + GNOVA
2. **Local correlation:** SUPERGNOVA
3. **Shared loci:** ConjFDR

Section 2: Causal inference


1. Univariable Mendelian Randomization




2. Multivariable Mendelian Randomization



3. Mediation Mendelian Randomization





Global genetic correlation

Method
LDSC
GNOVA

Depression

SCZ

BIP


*

**


ns

ns


Depression & AAA



SCZ & AAA



BIP & AAA



Exposure	Outcome	Methods	OR (95% CI)	<i>p</i> -value		Cochran's Q	<i>p</i> value for heterogeneity* or pleiotropy†	
Forward analyses								
Depression	AAA	IVW	1.270 (1.071, 1.504)	0.006		163.448	<0.001	
		MR-Egger	0.865 (0.348, 2.149)	0.754	←→			0.404
		Weighted median	1.229 (1.063, 1.420)	0.005				
		MR-cML	1.184 (1.040, 1.349)	0.011				
		MR-RAPS	1.302 (1.190, 1.424)	<0.001				
SCZ	AAA	IVW	1.047 (1.010, 1.084)	0.011		333.578	<0.001	
		MR-Egger	1.018 (0.882, 1.174)	0.809	←→			0.692
		Weighted median	1.041 (1.002, 1.082)	0.041				
		MR-cML	1.041 (1.004, 1.079)	0.029				
		MR-RAPS	1.049 (1.025, 1.074)	<0.001				
BIP	AAA	IVW	0.980 (0.896, 1.071)	0.653	←→	32.604	0.002	
		MR-Egger	1.165 (0.656, 2.069)	0.601	←→			0.559
		Weighted median	1.024 (0.936, 1.121)	0.607				
		MR-cML	0.987 (0.915, 1.065)	0.737				
		MR-RAPS	0.978 (0.924, 1.035)	0.448				
Reverse analyses								
AAA	Depression	IVW	1.010 (0.995, 1.025)	0.190		183.629	<0.001	
		MR-Egger	1.006 (0.969, 1.046)	0.743				0.844
		Weighted median	1.001 (0.982, 1.020)	0.897				
		MR-cML	1.010 (0.995, 1.024)	0.187				
		MR-RAPS	1.010 (0.998, 1.022)	0.097				
AAA	SCZ	IVW	1.007 (0.968, 1.048)	0.740		328.312	<0.001	
		MR-Egger	0.989 (0.891, 1.096)	0.832	←→			0.711
		Weighted median	1.004 (0.962, 1.048)	0.843				
		MR-cML	0.985 (0.951, 1.020)	0.390				
		MR-RAPS	1.007 (0.983, 1.032)	0.562				
AAA	BIP	IVW	0.989 (0.944, 1.037)	0.646		183.276	<0.001	
		MR-Egger	1.005 (0.889, 1.137)	0.940	←→			0.786
		Weighted median	1.011 (0.950, 1.075)	0.735				
		MR-cML	0.991 (0.946, 1.039)	0.704				
		MR-RAPS	0.989 (0.952, 1.027)	0.560				

**p*-value for heterogeneity based on Cochran's Q statistic.

†*p*-value or pleiotropy based on MR-Egger regression intercept.

