Ageing Biomarkers Derived From Retinal Imaging: A Scoping Review

Michaela Grimbly¹, Sheri-Michelle Koopowitz¹, Alice Ruiye Chen², Zihan Sun³, Paul

Foster³, Mingguang He², Dan J. Stein¹, Jonathan Ipser¹, Lisa Zhuoting Zhu²

- 1. SAMRC Unit on Risk & Resilience for Mental Disorders, Department of Psychiatry and Neuroscience Institute, University of Cape Town, South Africa
- 2. Centre for Eye Research Australia; Ophthalmology, University of Melbourne, Australia
- 3. NIHR Biomedical Research Centre at Moorfields Eye Hospital and UCL Institute of Ophthalmology, London, England

Synopsis

This scoping review explores retinal age, a biological ageing marker derived from retinal images. Findings from 13 articles reveal links between advanced retinal ageing gaps and adverse outcomes, emphasizing its potential for improving health outcomes.

Abstract

Background/Aims: The emerging concept of retinal age, a biomarker derived from retinal images, holds promise in estimating biological age. The retinal age gap (RAG) represents the difference between retinal age and chronological age which serves as an indicator of deviations from normal ageing. This scoping review aims to collate studies on retinal age to determine its potential clinical utility and to identify knowledge gaps for future research.

Methods: Using the PRISMA checklist, eligible non-review, human studies were identified, selected, and appraised. Pubmed, Scopus, SciELO, PsycINFO, Google Scholar, Cochrane, CINAHL, Africa Wide EBSCO, MedRxiv, and BioRxiv databases were searched to identify literature pertaining to retinal age, the RAG, and their associations. No restrictions were imposed on publication date.

Results: Thirteen articles published between 2022 and 2023 were analysed, revealing four models capable of determining biological age from retinal images. Three models, 'Retinal Age', 'EyeAge' and a 'convolutional network-based model,' achieved comparable mean absolute errors (MAE): 3.55, 3.30 and 3.97 respectively. A fourth model, 'RetiAGE', predicting the NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

probability of being older than 65 years, also demonstrated strong predictive ability with respect to clinical outcomes. In the models identified, a higher predicted RAG demonstrated an association with negative occurrences, notably mortality and cardiovascular health outcomes.

Conclusion: This review highlights the potential clinical application of retinal age and RAG, emphasising the need for further research to establish their generalisability for clinical use, particularly in neuropsychiatry. The identified models showcase promising accuracy in estimating biological age, suggesting its viability for evaluating health status.

Key Message

What is already known on this topic: Retinal age has emerged as a promising ageing biomarker capable of determining biological age from retinal images.

What this study adds: This study presents a comprehensive scoping review of current literature concerning retinal age and the RAG, highlighting the reproducible association between advanced retinal age gap and increased mortality and cardiovascular disease risk.

How this study might affect research, practice, or policy: The findings underscore the paucity of knowledge on this topic, advocating for further research in this area to determine the potential clinical use of retinal age as a biomarker.

Introduction

Globally, the number of individuals aged over 60 is rising, leading to an increased burden to healthcare services and society. Ageing changes are heterogenous, with substantial variation in the health impacts of ageing across populations, individuals and tissues (Lowsky et al., 2014; Tian et al., 2023). Thus, biological ageing markers have emerged to better represent the ageing process and predict functional capability.

Retinal age, an imaging based biomarker, provides an estimate of biological age derived from retinal fundus photographs (Nusinovici et al., 2022; Zhu, Shi, et al., 2023). The rationale for relying on retinal age as a biomarker stems from the retina's shared embryological origin with the central nervous system (Ptito et al., 2021) and microvascular structure, which is closely related to that of the brain, heart and kidney (Elias et al., 2016; Farrah et al., 2020; London et al., 2013). Although retinal imaging has largely been used by ophthalmologists for improved understanding and treatment of ocular disease (Keane & Sadda, 2014), predictive retinal ageing extends this use by applying deep learning to retinal fundus photography. This greatly enhances the utility of retinal imaging beyond the realm of Ophthalmology.

The introduction of the retinal age gap (RAG), defined as the difference between calculated retinal age and chronological age, provides a valuable metric for assessing deviations from normal ageing. When compared to traditional biomarker approaches, often criticised for their cost, invasiveness, time-consuming nature, and suboptimal accuracy, the application of retinal age models provides a cost-effective, non-invasive, and readily accessible way of estimating biological age (Butler et al., 2004; Chen, Wang, et al., 2023). This makes it particularly suitable for large scale population studies.

To date, there is no specific review on the reliability of retinal age as a biomarker. Although several biomarker reviews have included retinal age as one of many biological age estimators, these reviews have not been able to provide a comprehensive summary of the accuracy, practical utilisation, or relevance of retinal ageing models in the healthcare domain. This scoping review seeks to consolidate what is known about retinal age, while identifying gaps for future research by collating studies published on retinal age.

Specifically, this review aims to answer the following questions:

- How extensive is the current literature pertaining to retinal age and RAG?
- How many models exist, and how accurate are they?
- What are the clinical associations of retinal age and RAG?
- Does this biomarker exhibit clinical utility?
- What are the most pressing future directions for research?

Methods

Protocol and registration

A scoping review is well suited to expand and consolidate what is known about the retinal age, as it allows synthesis of current literature. A protocol was developed for this purpose and registered on Open Science Forum (Available at: <u>https://osf.io/fse75/</u>DOI 10.17605/OSF.IO/FSE75). The format for this scoping review is based on the PRISMA-ScR checklist (Page et al., 2021) and has made use of the Janna Briggs Institute Manual for Evidence Synthesis (Peters et al., 2017).

Eligibility criteria

To ensure a broad search of current literature, all published literature, and preprints of primary studies of the retinal age in adults were included for analysis. There were no limitations imposed for publication language or publication date.

Search

A comprehensive literature search of the following electronic databases was conducted from 17 June 2023 – 19 June 2023: Pubmed, Scopus, Cochrane Library, CINAHL, SciELO, Google Scholar, PsycINFO, Africa Wide EBSCO Host, MedRxiv and BioRxiv. A librarian assisted with formulating the search strategy. Initial search terms for the Pubmed database included, "retinal age" AND "association", which were further refined to ((retinal age [Text Word]) OR (retinal age gap [Text Word])) AND (((association) OR (link)) OR (biomarker)) and adapted for each database searched. Please refer to appendix 1 for full search strategy used.

Selection

A two-stage selection process was employed to assess relevance of studies identified in the search. Articles identified through the above-mentioned search strategy were deduplicated,

after which two reviewers (MG, SK) independently screened titles and abstracts to ascertain eligibility and relevance, utilising the predetermined inclusion and exclusion criteria described above. For publications that met inclusion criteria, these criteria were re-applied to the fulltext articles. In cases where discrepancies in reviewers' ratings were observed, a coordinating investigator (JI) conducted a final review to determine inclusion eligibility. The citations within included articles were also scanned to identify any additional articles that may be suitable for inclusion. Email updates for Google Scholar were enabled to capture newly published literature between the initial search date and the writing phase. Subsequent articles found were subjected to the same review process.

Data charting process

Data was extracted from the articles deemed suitable for inclusion using a spreadsheet developed by the reviewers. Data extracted included title, publication date, authors, study design, aim of the study, type of retinal age model used, training and validation details of the model, outcome of interest, analytical approach used, and key findings of the papers.

Results

Selection of sources

The PRISMA flow diagram (Figure 1) outlines the article selection process. The search strategy yielded a total of 342 articles (Appendix 1). After initial application of inclusion criteria, a total of 41 full text articles were examined for eligibility. Additionally, four articles published after the literature search date were considered for inclusion. A total of 13 articles met criteria for inclusion in the scoping review.

Characteristics of the Articles

All articles included for review were published between 2022 and 2023. <u>Appendix 2</u> presents a summary of the studies.

Identification of studies via databases and registers



Figure 1. PRISMA 2020 flow diagram

Narrative Review of Study Findings

All included studies utilised deep learning algorithms for retinal age analysis based on retinal fundus photographs. Four distinct models capable of determining biological age from retinal fundus photographs are outlined, with their training, validating, and testing processes described below. This is followed by a description of the application of these models to clinical populations.

Model Development and Accuracy

Three models were designed to predict age from retinal images, namely: '**Retinal Age'** (Zhu, Shi, et al., 2023), '**EyeAge'** (Ahadi et al., 2023) and a '**convolutional network-based model'** (Abreu-Gonzalez et al., 2023). A fourth biological ageing model, '**RetiAGE'** (Nusinovici et al., 2022), was developed to estimate the probability of an individual being older than 65 years from a retinal image.

The 'Retinal Age' model was trained and validated on 19 200 fundus photographs from 11 052 healthy UK Biobank participants. The model underwent 5-fold cross validation and achieved a mean absolute error (MAE) of 3.55, and Pearson correlation coefficient (R) between estimated age and chronological age of 0.80 (Zhu, Shi, et al., 2023). The 'EyeAge' model was trained on 217 289 images from 100 692 individuals in the EvePACS dataset, validated on 54 292 images from 25 238 individuals the same dataset and tested in both the UK Biobank and EyePACS dataset. The model achieved a MAE of 3.30, and a Pearson R of 0.87 for the UK Biobank test dataset, with corresponding figures of 2.86 and 0.95 for the EyePACS dataset (Ahadi et al., 2023). The 'convolutional network-based model' was trained on 98 400 photos from patients diagnosed with diabetes who were enrolled in the Retisalud programme of the Canary Islands Health Service. One thousand images from the dataset were used to validate this model, achieving a MAE of 3.97. However, it is unclear whether these validation images were included in the initial training set. The 'RetiAGE' model was trained on 116 312 photographs from 36 432 participants of the Korean Health Screening Study and validated on 12 924 unseen photos from 4 048 participants from the same dataset. In an internal test on 32 318 photos of 10 171 participants, the model achieved an Area Under the Receiver Operating Characteristic (AUROC) curve of 0.968 (95% confidence interval (CI): 0.965-0.970) and an area under the precision-recall curve (AUPRC) of 0.83 (95% CI: 0.83-0.84). When

applied to the UK Biobank cohort, the model achieved an AUROC of 0.756 (95% CI: 0.753-0.759) and an AUPRC of 0.399 (95% CI: 0.388-0.410) with a correlation of 0.62 between 'RetiAGE' and chronological age (Nusinovici et al., 2022).

Clinical Utility and Model Associations

The RAG has conventionally been used as the metric for assessing the clinical utility of retinal age. Defined as the difference between the retinal age biomarker and chronological age, the RAG is used to assess the performance of the retinal age model in reflecting ageing. Eleven papers utilising two models, '**Retinal Age'** and **'convolutional network-based model'**, have been published with the RAG as the outcome measure of interest.

Ten association analyses were conducted using the 'Retinal Age' model to explore the relationship between RAG and age-related parameters, within the UK Biobank cohort. The introductory study, highlighting the development of the model, revealed a significant association of a 2% increase in mortality risk for each one-year increase in RAG [Hazards Risk (HR) = 1.02, 95% CI 1.00-1.03, p=0.020] (Zhu, Shi, et al., 2023). Beyond the risk stratification for mortality, several prospective studies have highlighted associations for each one-year increase in RAG with a 10% increase of Parkinson's disease [HR=1.10, 95% CI: 1.01-1.20, p=0.023] (Hu et al., 2022), a 4% increase of stroke [HR=1.04, 95% CI: 1.00-1.08, p=0.029] (Zhu, Hu, et al., 2022), a 3% increase of incident cardiovascular disease [HR = 1.03, 95% CI: 1.00-1.05, p=0.019] (Zhu, Chen, et al., 2022), a 10% increase in risk of incident kidney failure [HR = 1.10, 95% CI: 1.03-1.17, p=0.003] (Zhang et al., 2022), and a 7% increased risk of diabetic retinopathy in diabetic patients [HR = 1.07, 95% CI: 1.02-1.12, p=0.004] (Chen, Zhang, Hu, et al., 2023). Several cross-sectional studies utilising the 'Retinal Age' model explored the associations between certain lifestyle diseases and RAG. Central obesity [p<0.001] (Chen, Zhang, Shang, et al., 2023), higher glycemia levels [p<0.001] (Chen, Xu, Zhang, et al., 2023) and metabolic syndrome and inflammation [Odds Ratio = 1.01, 95% CI: 1.00-1.02, p= 0.0016] (Zhu, Liu, et al., 2023) were associated with higher RAGs, while a study aimed at correlating the RAG with cardiovascular health metrics, composed of 7 metrics totalling a score out of 14, determined that each 1-unit score increase in cardiovascular health was associated with a 13% decrease in calculated RAG [OR = 0.87, 95% CI: 0.85-0.90, p<0.001] (Chen, Xu, Shang, et al., 2023).

The **'EyeAge'** model evaluated its clinical performance by calculating EyeAge Acceleration, determined akin to RAG, as the difference between EyeAge (reflecting retinal age) and chronological age. The acceleration is derived through regressing EyeAge against chronological age for assessment of the model's clinical performance. In the UK Biobank cohort, adjusted EyeAge achieved an all-cause mortality hazard ratio of 1.03, while RAG (referred to as EyeAge Acceleration) was associated with a higher risk of chronic obstructive pulmonary disease [p=0.0048], myocardial infarction [p=0.049], elevated systolic blood pressure [p=1.025e-7] and fluid intelligence scores [p=5.34e-27]. Additionally, a genome wide association study (GWAS) was performed as part of the study and found that candidate genes identified for EyeAge acceleration are associated with eye function and age-related disorders (Ahadi et al., 2023).

The Abreu '**convolutional network-based model**' determined, from a case-control study, that in diabetic patients from the Retisalud programme of the Canary Islands, a higher RAG was associated with more severe, progressive diabetic retinopathy [p<0.001] (Abreu-Gonzalez et al., 2023), echoing the findings of the **'Retinal Age'** model in diabetic patients (Chen, Zhang, Hu, et al., 2023).

The **'RetiAGE**' model also used the UK Biobank cohort to assess its clinical performance. In contrast to **'Retinal Age'** and the **'convolutional network-based model**,' which utilised calculated RAG for outcome assessment, the developers of the 'RetiAGE' biological age marker directly investigated its association with different age-related outcomes. Individuals placed in higher quartiles of **'RetiAGE'** were considered to have accelerated ageing. Comparing those within the fourth quartile group to those in the first quartile, there was a 67% higher risk of 10 year all-cause mortality [HR = 1.67, 95% CI: 1.42-1.95], a 142% higher risk of cardiovascular related mortality [HR = 2.42, 95% CI: 1.69-3.48] and a 60% higher risk of cancer related mortality [HR=1.60, 95% CI: 1.31-1.96] after adjustment for chronological age and other established ageing biomarkers (Nusinovici et al., 2022).

Saliency Maps

Features that drive the retinal ageing estimates were identified for the '**Retinal Age**' and '**RetiAGE'** models. '**Retinal Age'** retrieved attention maps using guided Grad-CAM (Selvaraju

et al., 2020), to highlight pixels in the input retinal fundus image based on their contributions to the final evaluation. Areas around retinal vessels are highlighted, indicating that retinal microvasculature is used by the deep learning model for age prediction (Zhu, Shi, et al., 2023). '**RetiAGE**' generated saliency maps to localise anatomy contributing to retinal aging. They indicated that the model focuses on the macula, optic disc, and, echoing the findings from the **'Retinal Age'** model, retinal vessels for age determination (Nusinovici et al., 2022).

Discussion

This study aimed to qualitatively appraise existing research utilising retinal photography for the development of biological ageing markers. We sought to determine the accuracy of retinal age prediction models and evaluate their performance in reflecting age-related parameters, as well as their clinical associations. This scoping review identified several models which are able to estimate chronological age from retinal images with moderate to high accuracy, and identified several age-related associations. Here we present key findings from our review of the existing literature.

Four models are currently available to estimate biological age from retinal images, all based on deep learning algorithms: '**Retinal Age'** (Zhu, Shi, et al., 2023), '**EyeAge'** (Ahadi et al., 2023), the '**convolutional network-based model'** (Abreu-Gonzalez et al., 2023) and '**RetiAGE'** (Nusinovici et al., 2022). The '**Retinal Age'**, '**EyeAge'** and the '**convolutional network-based model**' were trained to predict numerical chronological age from retinal images, while '**RetiAGE'** was trained to predict the probability of an individual being over the age of 65 years.

All models were exclusively trained and validated using a single dataset, predominantly comprising Caucasian or Asian populations. To enhance robustness, both 'EyeAge' and 'RetiAGE' underwent additional internal testing on previously unseen images from the training and validation cohort. For model testing and outcome assessment, the UK Biobank cohort was utilised by three models: 'Retinal Age', 'EyeAge' and 'RetiAGE.' Although each of the four identified models demonstrated comparable model accuracy and performance, it is important to note that performance metrics were inconsistently reported, with some pertaining to validation performance, and others test performance. Consequently, the

generalisability of these models is uncertain, with further work warranted to assess the applicability of these models across diverse populations.

Nevertheless, the application of retinal age models in predicting mortality and morbidity carries significant clinical implications. A key finding from these thirteen selected papers emphasizes that accelerated ageing, whether calculated as a RAG, age acceleration, or other indices, consistently correlates with mortality risk across three models (Ahadi et al., 2023; Nusinovici et al., 2022; Zhu, Shi, et al., 2023). In addition, '**Retinal Age'** and '**EyeAge'** show consistent associations with cardiovascular disease, while '**Retinal Age'** and the '**convolutional network-based model'** show connections with the risk of diabetic retinopathy in diabetic patients. These findings highlight the potential of retinal age as an informative tool for quantifying the risk of mortality and cardiovascular morbidity. However, to date, no clinical trials have explored the utility or feasibility of the models, which is a crucial aspect for determining their clinical relevance. Furthermore, factors associated with higher RAG, including glycaemic status (Chen, Xu, Zhang, et al., 2023), central obesity (Chen, Zhang, Shang, et al., 2023) and metabolic syndrome (Zhu, Liu, et al., 2023), suggest that RAG may provide valuable insight into lifestyle habits and traits that accelerate ageing. Notably, these factors were only investigated using the '**Retinal Age'** model.

It is important to highlight the inadequate reporting of characteristics of the populations used for training the retinal age prediction models. Only the '**Retinal Age'** model mentions that it was trained on healthy populations. This distinction is important if one wishes to consider biological age as equal to chronological age in normal ageing individuals. The health status of the population used for training '**EyeAge'** and '**RetiAGE'** remains undisclosed, while the 'convolutional network-based model' used data from diabetic patients. This may lead to confounding the effects of diabetes on apparent aging with age itself. Such a discrepancy could lead to controversy over whether these three retinal age models are accurate predictors of biological age. Thus, there is a need for a standardized procedure of developing biological age which should be proposed for use in further studies.

Additionally, these models were trained on a limited set of retinal features with only two models, '**Retinal Age'** and '**RetiAge**', producing saliency or attention maps to identify the

potential features used by the deep learning models for age assessment. This links to concerns about regulatory compliance and interpretability of the use of artificial intelligence in health care (Albahri et al., 2023; Funer, 2022). However, both models alluded to retinal microvasculature being a key component of age ascertainment, indicating that retinal age may reflect ageing related to vascular status. This is supported by the finding that retinal age models are particularly associated with cardiovascular health (Chen, Xu, Shang, et al., 2023; Nusinovici et al., 2022). To improve understanding of retinal features that align with biological age, advanced or diverse visualization techniques are imperative.

The application of retinal age models in predicting neuropsychiatric diseases is relatively underexplored. Given that the retina is an extension of the central nervous system, it offers a unique and accessible 'window' to visualise cerebral neuronal health (Elias et al., 2016; Farrah et al., 2020; London et al., 2013). An extensive body of work has found that changes in the retina, most notably thinning in the retinal nerve fibre layer, may be associated with certain neuropsychiatric and neurodegenerative diseases (Gonzalez-Diaz et al., 2022; Kashani et al., 2021). In our review, only one paper using the '**Retinal Age'** model explored the RAG in the realm of neuropsychiatry, specifically in the context of Parkinson's disease, leaving this area underexplored (Hu et al., 2022). As neurodegeneration is an important aspect of ageing, future studies should concentrate on improving our understanding of the connections between retinal age and neuropsychiatric conditions.

Several limitations of this scoping review deserve emphasis. Publications in non-indexed journals and other 'grey literature' may have been missed. Due to insufficient data availability, it was not possible to quantitatively synthesis data using meta-analytic statistical techniques. As more literature becomes available on the topic, conducting a broader, more extensive review may unveil more diverse associations of retinal age, mechanisms for associations, and possibly link retinal age to other biomarkers. Strengths of our study included its development according to a predefined protocol, and application of the PRISMA extension for scoping reviews.

In conclusion, this scoping review identified four retinal ageing models derived from retinal images, with advancing retinal age gaps significantly associated with mortality and

cardiovascular disease. It highlights the scarcity of data in the realm of neuropsychiatry, emphasises the need for a standardised procedure in developing retinal ageing models, and shows that testing across different datasets is crucial to improve the generalisability and utility of the models. Improving our understanding of the biological underpinnings of how these models determine age may too improve their reliability in reflecting ageing processes. Nevertheless, the evidence highlights the potential of retinal age as a biomarker, suggesting its viability as a valuable, cost-effective tool for evaluating health status.

Funding

This research received no external funding.

Bibliography

- Abreu-Gonzalez, R., Rodríguez-Martín, J. N., Quezada-Peralta, G., Rodrigo-Bello, J. J., Gil-Hernández, M. A., Bermúdez-Pérez, C., & Donate-López, J. (2023). Retinal age as a predictive biomarker of the diabetic retinopathy grade. Archivos de La Sociedad Española de Oftalmología (English Edition), 98(5), 265–269. https://doi.org/10.1016/j.oftale.2023.04.008
- Ahadi, S., Wilson, K. A., Babenko, B., McLean, C. Y., Bryant, D., Pritchard, O., Kumar, A.,
 Carrera, E. M., Lamy, R., Stewart, J. M., Varadarajan, A., Berndl, M., Kapahi, P., &
 Bashir, A. (2023). Longitudinal fundus imaging and its genome-wide association
 analysis provide evidence for a human retinal aging clock. eLife, 12, e82364.
 https://doi.org/10.7554/eLife.82364
- Albahri, A. S., Duhaim, A. M., Fadhel, M. A., Alnoor, A., Baqer, N. S., Alzubaidi, L., Albahri, O.
 S., Alamoodi, A. H., Bai, J., Salhi, A., Santamaría, J., Ouyang, C., Gupta, A., Gu, Y., &
 Deveci, M. (2023). A systematic review of trustworthy and explainable artificial
 intelligence in healthcare: Assessment of quality, bias risk, and data fusion.
 Information Fusion, 96, 156–191. https://doi.org/10.1016/j.inffus.2023.03.008
- Butler, R. N., Sprott, R., Warner, H., Bland, J., Feuers, R., Forster, M., Fillit, H., Harman, S. M.,
 Hewitt, M., Hyman, M., Johnson, K., Kligman, E., McClearn, G., Nelson, J., Richardson,
 A., Sonntag, W., Weindruch, R., & Wolf, N. (2004). Biomarkers of aging: From
 primitive organisms to humans. The Journals of Gerontology. Series A, Biological
 Sciences and Medical Sciences, 59(6), B560-567.

https://doi.org/10.1093/gerona/59.6.b560

Chen, R., Wang, Y., Zhang, S., Bulloch, G., Zhang, J., Liao, H., Shang, X., Clark, M., Peng, Q., Ge, Z., Cheng, C.-Y., Gao, Y., He, M., & Zhu, Z. (2023). Biomarkers of ageing: Current

> state-of-art, challenges, and opportunities. MedComm – Future Medicine, 2(2), e50. https://doi.org/10.1002/mef2.50

Chen, R., Xu, J., Shang, X., Bulloch, G., He, M., Wang, W., & Zhu, Z. (2023). Association between cardiovascular health metrics and retinal ageing. GeroScience. https://doi.org/10.1007/s11357-023-00743-3

Chen, R., Xu, J., Zhang, X., Zhang, J., Shang, X., Ge, Z., He, M., Wang, W., & Zhu, Z. (2023). Glycemic status and its association with retinal age gap: Insights from the UK biobank study. Diabetes Research and Clinical Practice, 202, 110817. https://doi.org/10.1016/j.diabres.2023.110817

- Chen, R., Zhang, J., Hu, W., He, M., Zhu, Z., Chen, Y., & Wang, W. (2023). Retinal age gap as a predictive biomarker for future risk of clinically significant diabetic retinopathy. Acta Diabetologica, 1–8. https://doi.org/10.1007/s00592-023-02199-5
- Chen, R., Zhang, J., Shang, X., Wang, W., He, M., & Zhu, Z. (2023). Central obesity and its association with retinal age gap: Insights from the UK Biobank study. International Journal of Obesity (2005). https://doi.org/10.1038/s41366-023-01345-x
- Elias, M. F., Torres, R. V., & Davey, A. (2016). The Eye is the Window to the Kidney and Brain. eBioMedicine, 5, 24–25. https://doi.org/10.1016/j.ebiom.2016.02.010
- Farrah, T. E., Dhillon, B., Keane, P. A., Webb, D. J., & Dhaun, N. (2020). The eye, the kidney, and cardiovascular disease: Old concepts, better tools, and new horizons. Kidney International, 98(2), 323–342. https://doi.org/10.1016/j.kint.2020.01.039
- Funer, F. (2022). The Deception of Certainty: How Non-Interpretable Machine Learning
 Outcomes Challenge the Epistemic Authority of Physicians. A deliberative-relational
 Approach. Medicine, Health Care and Philosophy, 25(2), 167–178.
 https://doi.org/10.1007/s11019-022-10076-1

Gonzalez-Diaz, J. M., Radua, J., Sanchez-Dalmau, B., Camos-Carreras, A., Zamora, D. C., & Bernardo, M. (2022). Mapping Retinal Abnormalities in Psychosis: Meta-analytical Evidence for Focal Peripapillary and Macular Reductions. Schizophrenia Bulletin, 48(6), 1194–1205. https://doi.org/10.1093/schbul/sbac085

Hu, W., Wang, W., Wang, Y., Chen, Y., Shang, X., Liao, H., Huang, Y., Bulloch, G., Zhang, S.,
Kiburg, K., Zhang, X., Tang, S., Yu, H., Yang, X., He, M., & Zhu, Z. (2022). Retinal age
gap as a predictive biomarker of future risk of Parkinson's disease. Age and Ageing,
51(3), afac062. https://doi.org/10.1093/ageing/afac062

- Kashani, A. H., Asanad, S., Chan, J. W., Singer, M. B., Zhang, J., Sharifi, M., Khansari, M. M., Abdolahi, F., Shi, Y., Biffi, A., Chui, H., & Ringman, J. M. (2021). Past, present and future role of retinal imaging in neurodegenerative disease. Progress in Retinal and Eye Research, 83, 100938. https://doi.org/10.1016/j.preteyeres.2020.100938
- Keane, P. A., & Sadda, S. R. (2014). Retinal Imaging in the Twenty-First Century: State of the Art and Future Directions. Ophthalmology, 121(12), 2489–2500. https://doi.org/10.1016/j.ophtha.2014.07.054

London, A., Benhar, I., & Schwartz, M. (2013). The retina as a window to the brain—From eye research to CNS disorders. Nature Reviews Neurology, 9(1), Article 1. https://doi.org/10.1038/nrneurol.2012.227

Lowsky, D. J., Olshansky, S. J., Bhattacharya, J., & Goldman, D. P. (2014). Heterogeneity in Healthy Aging. The Journals of Gerontology: Series A, 69(6), 640–649. https://doi.org/10.1093/gerona/glt162

Nusinovici, S., Rim, T. H., Yu, M., Lee, G., Tham, Y.-C., Cheung, N., Chong, C. C. Y., Da Soh, Z., Thakur, S., Lee, C. J., Sabanayagam, C., Lee, B. K., Park, S., Kim, S. S., Kim, H. C., Wong, T.-Y., & Cheng, C.-Y. (2022). Retinal photograph-based deep learning predicts

> biological age, and stratifies morbidity and mortality risk. Age and Ageing, 51(4), afac065. https://doi.org/10.1093/ageing/afac065

- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D.,
 Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw,
 J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S.,
 ... Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting
 systematic reviews. BMJ, 372, n71. https://doi.org/10.1136/bmj.n71
- Peters, M., Godfrey, C., Khalil, H., Mcinerney, P., Soares, C., & Parker, D. (2017). 2017 Guidance for the Conduct of JBI Scoping Reviews.
- Ptito, M., Bleau, M., & Bouskila, J. (2021). The Retina: A Window into the Brain. Cells, 10(12), 3269. https://doi.org/10.3390/cells10123269
- Selvaraju, R. R., Cogswell, M., Das, A., Vedantam, R., Parikh, D., & Batra, D. (2020). Grad-CAM: Visual Explanations from Deep Networks via Gradient-based Localization.
 International Journal of Computer Vision, 128(2), 336–359.
 https://doi.org/10.1007/s11263-019-01228-7
- Tian, Y. E., Cropley, V., Maier, A. B., Lautenschlager, N. T., Breakspear, M., & Zalesky, A. (2023). Heterogeneous aging across multiple organ systems and prediction of chronic disease and mortality. Nature Medicine, 29(5), 1221–1231. https://doi.org/10.1038/s41591-023-02296-6
- Zhang, S., Chen, R., Wang, Y., Hu, W., Kiburg, K. V., Zhang, J., Yang, X., Yu, H., He, M., Wang, W., & Zhu, Z. (2022). Association of Retinal Age Gap and Risk of Kidney Failure: A UK Biobank Study. American Journal of Kidney Diseases.

https://doi.org/10.1053/j.ajkd.2022.09.018

Zhu, Z., Chen, Y., Wang, W., Wang, Y., Hu, W., Shang, X., Liao, H., Shi, D., Huang, Y., Ha, J., Tan,
 Z., Kiburg, K. V., Zhang, X., Tang, S., Yu, H., Yang, X., & He, M. (2022). Association of
 Retinal Age Gap With Arterial Stiffness and Incident Cardiovascular Disease. Stroke.
 https://doi.org/10.1161/STROKEAHA.122.038809

Zhu, Z., Hu, W., Chen, R., Xiong, R., Wang, W., Shang, X., Chen, Y., Kiburg, K., Shi, D., He, S.,
Huang, Y., Zhang, X., Tang, S., Zeng, J., Yu, H., Yang, X., & He, M. (2022). Retinal age
gap as a predictive biomarker of stroke risk. BMC Medicine, 20(1), 466.
https://doi.org/10.1186/s12916-022-02620-w

- Zhu, Z., Liu, D., Chen, R., Hu, W., Liao, H., Kiburg, K., Ha, J., He, S., Shang, X., Huang, Y., Wang,
 W., Yu, H., Yang, X., & He, M. (2023). The Association of Retinal age gap with
 metabolic syndrome and inflammation. Journal of Diabetes, 15(3), 237–245.
 https://doi.org/10.1111/1753-0407.13364
- Zhu, Z., Shi, D., Guankai, P., Tan, Z., Shang, X., Hu, W., Liao, H., Zhang, X., Huang, Y., Yu, H.,
 Meng, W., Wang, W., Ge, Z., Yang, X., & He, M. (2023). Retinal age gap as a predictive
 biomarker for mortality risk. British Journal of Ophthalmology, 107(4), 547–554.
 https://doi.org/10.1136/bjophthalmol-2021-319807

Supplementary Materials

Appendix 1:

Search of Published Works Databases

Preliminary search strategy formulated on Pubmed.

Initial search terms included: retinal age, retinal age gap, association, link, biomarker.

Database	Date of Search		Search Terms	Results	Articles Screened
Pubmed	17/06/2023	1	(retinal age) OR (retinal age gap)	49 442	14
		2	(retinal age[Text Word] OR (retinal age gap[Text Word])	23	
		3	(retinal age[Text Word] OR (retinal age gap[Text Word]) AND	14	
			(((association) OR (link) OR (biomarker)		
Scopus	17/06/2023	1	Retinal AND age OR retinal AND age AND gap	9 430	12
		2	TITLE-ABS-KEY ("retinal age" OR "retinal age gap")	29	
		3	TITLE-ABS-KEY (association OR biomarker OR link)) AND TITLE-ABS-	12	
			KEY ("retinal age" OR "retinal age gap")		
Cochrane	17/06/2023	1	(retinal age OR retinal age gap):ti,ab,kw (Word variations have been searched)	5362	1
		2	(retinal age OR retinal age gap):ti,ab,kw AND (association OR link OR biomarker):ti,ab,kw (Word variations have been searched)	1402	
		3	("retinal age" OR "retinal age gap" OR "retina age"):ti,ab,kw AND (association OR link OR biomarker):ti,ab,kw NOT (age related macular degeneration):ti,ab,kw (Word variations have been searched)	1	
CINAHL	17/06/2023	1	"retinal age" OR "retinal age gap"	3	11
		2	"retinal age" AND "association"	11	
SciELO	17/06/2023	1	Retinal age	223	16
		2	Retinal age gap	0	
		3	Retinal age OR retinal age gap	0	
		4	Retinal age AND association	16	
Google Scholar	18/06/2023	1	"retinal age" OR "retinal age gap" AND association OR link OR biomarker	48	48

PsychINFO	18/06/2023	1	(retinal age OR retinal age gap) AND (association OR link OR biomarker) Boolean/Phrase	32	32
Africa Wide EBSCO Host	18/06/2023	1	(retinal age OR retinal age gap) AND (association OR link OR biomarker) Boolean/Phrase	26	26
Total identified	from published data	oases			160

Search of Preprint Databases

Database	Date of Search		Search Terms	Results	Articles Screened
MedRxiv	19/06/2023	1	"retinal age gap"	161	161
NIH Reporter	19/06/2023	1	"retinal age gap" OR "retinal age" (advanced) Limit to: Project Title, Project Terms, Project Abstract	1 project 4 publications	0
BioArchives	19/06/2023	1	"retinal age"	21	21
Total identified	from preprint databa	ases			182

Appendix 2:

Table 1: Cross Sectional Studies – Data Extraction

Article ID	Year	Model Used	Training Dataset	Validating	Testing Dataset	Algorithm Performance	Outcome Measure	Disease of Interest	Associations	Summary
<u>Abreu 2023 Retinal</u>	2023	Based on convolutional networks	Diabetic patients; Retisalud programme of the Canary Islands Health Service; 40-90 years; 98 400 photos	1000 photos	Same as training: 40 - 90 years – 7694 without DR; 5850 with DR (mild - 4505; moderate - 1152; severe - 166; proliferative - 28) = 13544	MAE 3.97	Retinal Age Gap (RAG)	Diabetic Retinopathy (DR)	RAG without DR = 0.609 years; RAG with DR = 1.905 years (p<0.001) mild DR 1.541, mod 3.017, severe 3.117, proliferative 8.583; combined mild/no DR = 0.840, mod/severe = 3.131 (p<0.001)	More positive RAG: associated with more progressive DR (p<0.001)
Ahadi 2022 Longitudinal	2023	EyeAge: Deep learning - Inception v3 architecture	EyePACS Dataset; 217 289 images from 100 692 patients; mean age 54.21; 59% female	54 292 from 25 238 patients; mean age 54.2; 58% female	UK Biobank: mean age 56.85 and repeats from EyePACS UKB - 119 952 images from 64 019 patients Mean age 56.85; 55% female	EyePACS MAE 2.86; UKB MAE 3.30	EyeAge; EyeAgeAccel	Age prediction: genetic factors associated with EyeAgeAccel; mortality	EyeAge: All-cause mortality hazard ratio 1.026. EyeAge: morbidity + disability EyeAge: COPD (p=0.0048); MI (p=0.049). Increased EyeAgeAccel: increased systolic BP (p=1.025e-7); increased performance in fluid intelligence (p=5.34e- 27). GWAS: POC5; GJA3 - eye and age-related functions	EyeAge potential for studying aging; GWAS on 45 555 European individuals - BOLT-LMM v2.3.4 - association found with genetic loci

Chen 2023 Association 2023 Retinal Age: Deep learning UK Blobank; 19 200 images As training -5-fold UK Blobank; 26 354 MAE 3.55; Pearson R Retinal Age Gap (RAG) Cardiovascular Each 1 unit score increase in CVH: 13% decrease in RAG (D-7); intermediate/ideal CVH = lower RAG compared to por CVH (IR-08.7, 95% CI:0.55008). Bettra CVH associated with lower RAG View RAG recruitment; mean age 52.6 + 7.97; 53.7% female) 53.7% female -57.57 -57.57 -57.57 -57.57.57.57.57.57.57.57.57.57.57.57.57.5											
	Chen 2023 Association	2023	Retinal Age: Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40-69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank: 26 354 included mean age 56.6 +- 8.07; 53.7% female	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	Cardiovascular Health (CVH) - poor (0-7); intermediate (8-10); ideal (11-14)	Each 1 unit score increase in CVH: 13% decrease in RAG (OR=0.87, 95% CI: 0.85-0.90, p<0.001); intermediate/ideal CVH = lower RAG compared to poor CVH (OR=0.76, 95% CI:0.67-0.85, p<0.001; OR=0.58, 95% CI:0.50-0.67, p<0.001); intermediate and ideal CVH=reduced risk of accelerated retinal age compared to poor CVH (OR=0.83, 95%CI:0.77-0.90, p<0.001; OR=0.78, 95%CI:0.71-0.86,p<0.001) ; ideal status of smoking (OR=0.73, 95%CI:0.62-0.87), BMI (OR=0.80, 95%CI:0.71- 0.91), BP (OR=0.66, 95% CI: 0.66-0.89) and blood glucose (OR 0.66, 95% CI:0.55-0.80)	Better CVH associated with lower RAG

Chen 2023 Central 2023 Retinal Age: Deep learning UK Biobank; 19 200 images -Sception As training from 11 052 healthy architecture UK Biobank; 19 200 images healthy architecture As training UK Biobank; 35 500 Retinal Age Pearson B(0,001) Central Obesity-WC Gap (RAG) Overweight/high WC, mild obesity/high WC, severe obesity/high WC associated with Increased RAG Bigher RAG associated with eentral obesity 0.80 (p<0.001) mean age 52.6 + 7.97; 53.7% female) 55.6% female MAE 3.55; participants Retinal Age (p<0.001) Central Obesity-WC obesity/high WC, severe obesity/high WC associated with Increased RAG associated with eentral obesity 0.80 (p<0.001) mean age 52.6 + 7.97; 53.7% female) 55.6% female S5.6% female											
	hen 2023 Central	2023	Retinal Age: Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40-69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 35 550 participants; mean age 56.8 +- 8.04; 55.6% female	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	Central Obesity - WC + BMI: 7 groups	Overweight/high WC, mild obesity/high WC, severe obesity/high WC, severe obesity/high WC associated with increased RAG compared to normal weight/normal WC (B=0.333, 95%CI:0.191-0.474, p<0.001; B=0.383, 95% CI:0.257-0.509, p<0.001; B=0.440, 95% CI:0.278-0.602, p<0.001). Overweight/normal WC, mild obesity/normal WC, normal weight/high WC = no significant association with retinal age gaps. Overweight/high WC, severe obesity/high WC, severe obesity/high WC, severe obesity/high WC = higher risk of accelerated ageing compared to normal/normal (OR=1.18, 95% CI: 1.08-1.28, p<0.001; OR 1.20, 95% CI: 1.11-1.30, p<0.001, OR 1.27; 95% CI: 1.15-1.41, p<0.001)	Higher RAG associated with central obesity and higher WC

<u>Chen 2023 Glycemic</u>	2022	Retinal Age: Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40-69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 28 919 participants (mean 56.8, 55% female)	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	Normoglycaemia (HbA1c <5.7%), prediabetes (HbA1c >5.7 and <6.5%), DM2 (HbA1c >6.5% and/or insulin use) - glucose groups mmol/l (<5.5; 5.5- 6.4; 6.5-7.9; 8.0- 11.0; >11.1); HbA1c (<5.7%; 5.7-6.1%, 6.1-6.5%, 6.5-6.9%, >6.9%)	Prediabetes and DM2 = higher retinal age gap compared to normoglycemia (B=0.37, 95% CI: 0.24-0.49, p<0.001; B=1.16, 95%CI:0.96-1.35, p<0.001. Each 1 unit increase in HbA1C = higher RAG in all subjects and without DM2 (B=0.37, 95% CI:0.29-0.46, p<0.001; B=0.51, 95%CI: 0.36-0.66, p<0.001). Higher non-fasting plasma glucose associated with higher RAG. Each 1 SD increase of glucose level = significantly associated with increased RAG in all subjects and those without DM2 (B=0.17, 95% CI:0.12-0.22, p<0.001; B=0.23, 95%CI:0.13-0.33, p<0.001). Remained significant after excluding DR.	More positive RAG associated with dysglycemia

hu 2023 Association	2023	Deep learning - Xception	UK Biobank; 19 200 images	As training - 5-fold	UK Biobank 35 918	MAE 3.55; Pearson R	Retinal Age Gap (RAG)	Metabolic Syndrome (Met S) = >3/5 -	Each 1-year increase in RAG: 1% risk increase of MetS (OR	RAG associated with MetS and
		architecture	from 11 052	validation	participants;	0.80		abdominal obesity,	= 1.01, 95% CI: 1.00-1.02, p=	inflammation
			healthy		mean age	(p<0.001)		hypertension,	0.016). Each 1-year increase	
			participants		56.6 +- 8.04			elevated serum HDL,	In RAG: 1% risk increase in	
			(40-69 years at		years; 55.7%			elevated serum	Inflammation (OR=1.01, 95%	
			recruitment;		female			triglycerides,	CI:1.00-1.02, p=0.021) and	
			mean age 52.6					hyperglycaemia	1% risk increase of MetS and	
			+- 7.97; 53.7%						inflammation combined	
			female)						(OR=1.01, 95% CI: 1.00-1.02,	
									p=0.001). Per year increase	
									in RAG: associated with 2%	
									risk increase in abdominal	
									obesity (OR=1.02, 95%	
									Cl:1.01-1.02, p<0.001), 1%	
									risk increase in hypertension	
									(OR =1.01, 95% CI:1.00-1.02,	
									p=0.002) and 6% risk	

increase of hyperglycaemia (OR=1.06. 95% CI:1.04-1.07), p<0.001). When compared to participants with RAG in lower quartile, the risk of MetS was significantly increased by 10% in the 3rd quartile and 14% in the 4th quartile (OR=1.10, 95% CI:1.01-1.21, p=0.030, OR=1.14, 95% CI: 1.03-1.26, p=0.012)

Table 2: Prospective Studies – Data Extraction

Article ID	Year	Model Used	Training Dataset	Validating	Testing Dataset	Algorithm Performance	Outcome Measure	Disease of Interest	Associations	Summary
Chen 2023 Retinal	2023	Retinal Age: Deep learning – Xception Architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40- 69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 2311 diabetic patients without diabetic retinopathy (DR) at baseline. Mean age 58.5; 39.7% female	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	Incident diabetic retinopathy	Each 1-year increase in RAG: 7% adjusted increase in risk of incident DR [HR=1.07, 95% Cl: 1.02-1.12, p=0.004]. RAG in fourth quartile had higher DR risk [HR=2.88, 95% Cl: 1.61-5.15, p<0.001]. DR risk increased across RAG quartiles [HR=1.35, 95% Cl 1.11-1.64, p=0.002]	More positive RAG associated with increased risk of incident DR

<u>Hu 2022 Retinal</u>	2022	Retinal Age: Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40- 69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 35 834 participants free of Parkinson's disease (PD) at baseline; 56+- 8.04; 55.7% female	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	Incident Parkinson's disease - ICD 9/10 codes - history of PD; incident PD	Each 1-year increase in RAG: 10% increase in risk of PD (HR=1.10, 95% CI: 1.01-1.20, p=0.023). Compared with lowest RAG quartile, the risk of PD was increased in third and fourth quartiles (HR = 2.66, 95% CI: 1.13-6.22, p=0.024; HR = 4.86, 95% CI: 1.59-14.8, p=0.005). Predictive value of retinal age model (AUC = 0.708, 95% CI:0.638-0.778) and risk factor model (AUC=0.717, 95% CI:0.633-0.802) was similar (p=0.821).	More positive RAG associated with future risk of incident PD
<u>Jusinovici 2022 Retinal</u>	2022	RetiAge: Visual Geometry Group (VGG) - deep convolutional neural network architecture	Korean Health Screening Study; 129 236 photos from 40 480 participants; mean age 54 +- 9.01	12 924 photos from 4 048 participants	 56 301 from UK Biobank (mean age 57.1 +- 8.3 years; 46.5% female) used for all-cause mortality, CVD mortality and cancer mortality. 46 551 from Korean Health Screening Study (mean 53.6 +- 9.2 years; 45.4% female) used for all-cause mortality. 	AUROC 0.968 (95% CI: 0.965- 0.970); AUPRC 0.83)95% CI: 0.83-0.84) in internal test AUROC of 0.756 (95% CI: 0.753- 0.759) and an AUPRC of 0.399 (95% CI: 0.388- 0.410 in UK Biobank test	RetiAge	Predict old age from retina. All- cause mortality; CVD mortality and cancer mortality	Independent of chronological age and phenotypic ageing markers: when compared to RetiAge first quartile, those in the RetiAge fourth quartile had 67% higher risk of 10 year all-cause mortality (HR = 1.67, 95% CI: 1.42-1.95, p <0.001) 142% higher risk of CVD mortality (HR = 2.42, 95% CI: 1.69-3.48, p<0.001) and 60% higher risk of cancer mortality (HR=1.60, 95% CI: 1.31-1.96, p<0.001).	RetiAge can predict aging; higher RetiAge associated with mortality risk

Zhang 2023 Association	2023	Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40- 69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 35 864 participants with no kidney failure at baseline; mean age 56.75 +- 8.04; 55.7% female	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	Incident kidney failure - ICD 10 and OPCS 4 codes	Each 1-year increase in RAG: 9% increase in risk of incident kidney failure (HR = 1.09, 95% CI: 1.03-1.15, p<0.001). Retinal age gaps in the highest quartile had a significantly higher risk of incident kidney failure compared to those in the first quartile (HR = 2.77, 95% CI: 1.29-5.93, p=0.009). There was a graded association of incident kidney failure across retinal age gap quartiles (p=0.004). Participants in higher RAG quartile showed higher risk of kidney failure event.	More positive RAG associated with incident kidney failure
Zhu 2022 Association	2022	Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40- 69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 35 541 participants; mean age 56.8 +- 8.04; 55.6% female	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	ASI - arterial stiffness index; CVD events - cardiovascular disease	Each 1-year increase in RAG: increase in ASI (B=0.002, 95% Cl: 0.0001-0.003, p=0.001). Higher odd of having severe ASI with a larger RAG (OR=1.01, 95% Cl: 1.01-1.02, p<0.001). Each 1-year increase in RAG: predicts 3% increase in risk of incident CVD (HR = 1.03, 95% Cl: 1.00-1.05, p=0.019). Remains significant when ASI is incorporated (HR = 1.03, 95% Cl: 1.01-1.06, p=0.019)	More positive RAG significantly associated with ASI. RAG predictor of future risk of incident CVD.

Zhu 2022 Retinal	2022	Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40- 69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 35 304 without stroke history; mean age 56.7 +- 8.04; 55.9% female)	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	Stroke - ICD 9/10 codes	Each 1-year increase in RAG: associated 4% increase in risk of stroke when adjusting for confounding factors (HR=1.04, 95% Cl: 1.00-1.08, p=0.029). Retinal age gaps in the 5th quintile had higher risk of stroke compared to those whose RAG was in the first quintile (HR=2.37, 95% Cl:1.37-4.10, p=0.002). Predictive capability of retinal age alone in 10-year stroke risk was comparable to a well-established risk factor-based model (AUC=0.676 vs AUC 0.661, p=0.511)	More positive RAG associated with incident stroke
Zhu 2023 Retinal	2023	Deep learning - Xception architecture	UK Biobank; 19 200 images from 11 052 healthy participants (40- 69 years at recruitment; mean age 52.6 +- 7.97; 53.7% female)	As training - 5-fold validation	UK Biobank 35 913 photos from 35 917 participants; mean age 56.8 +- 8.04; 55.7% female	MAE 3.55; Pearson R 0.80 (p<0.001)	Retinal Age Gap (RAG)	All-cause mortality	Each 1-year increase in RAG: 2% increase in mortality risk (HR = 1.02, 95% CI 1.00- 1.03, p=0.020); Positive retinal age gap: substantially increased mortality; Higher retinal age gap (3rd and 4th quartile): higher non- CVD/cancer associated death (HR=1.49, 95% CI 1.13-1.96, p=0.005; HR=1.67, 95% CI 1.17-2.39, p=0.005)	More positive RAG associated with increased risk of mortality (>non- CVD/non- cancer)