

Anti-inflammatory effects of oral prednisolone at stable state in people treated with mepolizumab: a proteomic and bulk transcriptomics analysis

I. Howell¹, F. Yang^{2,3}, V. Brown⁴, J. Cane¹, E. Marchi¹, A. Azim⁵, J. Busby⁴, P.J. McDowell⁴, S.E. Diver⁶, C. Borg¹, L. G. Heaney⁴, I. D. Pavord¹, C. E. Brightling⁶, R. Chaudhuri², T.S.C. Hinks¹

1. Respiratory Medicine Unit, Experimental Medicine Division, Nuffield Department of Medicine, University of Oxford, John Radcliffe Hospital, Oxford, United Kingdom & National Institute for Health Research Oxford Biomedical Research Centre, John Radcliffe Hospital, Oxford, UK
2. University of Glasgow, Glasgow, UK.
3. The National Heart and Lung Institute, Imperial college London, London, UK
4. Queen's University Belfast, Belfast, UK.
5. Clinical and Experimental Sciences, University of Southampton Faculty of Medicine, Sir Henry Wellcome Laboratories and NIHR Southampton Respiratory Biomedical Research Unit, Southampton University, Southampton, UK
6. Institute for Lung Health, Leicester NIHR BRC, University of Leicester, Leicester, UK.

Abstract

Mepolizumab is an anti-interleukin-5 monoclonal antibody treatment for severe eosinophilic asthma (SEA) that reduces asthma exacerbations. Residual airway inflammation on mepolizumab may lead to persistent exacerbations. Oral corticosteroids have broad anti-inflammatory effects and remain the main treatment for these residual exacerbations. Our study aimed to explore the nature and corticosteroid-responsiveness of airway inflammation after mepolizumab treatment to find potentially treatable inflammatory mechanisms.

The MAPLE trial was a multi-centre, randomized, double-blind, placebo-controlled, crossover study of 2 weeks of high-dose oral prednisolone treatment at stable state in patients treated with mepolizumab for SEA. We analysed sputum and plasma samples from the MAPLE trial using high-throughput Olink® proteomics. We also analysed plasma microRNA, sputum proteins using ELISA, and nasal mucosal bulk RNA sequencing.

In patients receiving mepolizumab, prednisolone significantly downregulated sputum proteins related to type-2 inflammation and chemotaxis including IL-4, IL-5, IL-13, CCL24, CCL26, EDN, CCL17, CCL22, OX40 receptor, FCER2, and the ST2 receptor. Prednisolone also downregulated cell adhesion molecules, prostaglandin synthases, mast cell tryptases, MMP1, MMP12, and neuroimmune mediators. Tissue repair and neutrophilic pathways were upregulated. Type-2 proteins were also downregulated in plasma, combined with IL-12, IFN- γ , and IP-10. IL-10 and amphiregulin were upregulated. In the nasal transcriptome, prednisolone suppressed genes involved in leucocyte chemotaxis, mast cell tryptase, 15-lipoxygenase and *MMP12*. By contrast, mepolizumab differentially regulated only Galectin-10 in plasma and no sputum proteins, and in nasal tissue affected genes related to cilia, keratinisation, extracellular matrix formation, and IL-4/13 signalling.

At stable state, prednisolone has broad anti-inflammatory effects on top of mepolizumab.

Introduction

Oral corticosteroids (OCS) are a widely prescribed treatment for asthma exacerbations¹. Clinical evidence for OCS at exacerbation is based on trials demonstrating a reduction in relapse and accelerated symptom resolution². Biologically, OCS reduce eosinophilic inflammation, evidenced by a rapid decline in blood eosinophils through apoptosis and inhibition of interleukin-5 (IL-5)-induced production by the bone marrow^{3,4}. However, OCS have broader anti-inflammatory effects that may also be clinically important. The placebo-controlled BIOAIR study identified inflammatory proteins downregulated by prednisolone in both mild/moderate and severe asthma at stable state in plasma⁵. These included chemokines and proteins involved in recruiting eosinophils to the airways⁶. But anti-inflammatory effects of OCS also extend to the airway mucosa. A placebo-controlled human bronchoscopy study in stable, moderate asthma found that prednisolone reduced airway eosinophilia, tryptase single-positive mast cells, IL-4, IL-5, and CD3+ T lymphocytes⁷.

Systemic monoclonal antibody therapy (mAb) has revolutionised the treatment of severe eosinophilic asthma. Placebo-controlled trials of mepolizumab, an anti-IL5 mAb, demonstrated approximately a 50% reduction in the annualised exacerbation rate and maintenance OCS dose⁸. This is important because repeated OCS exposure is associated with serious acute and chronic adverse effects⁹. However, some patients continue to have asthma exacerbations despite mepolizumab therapy. The MEX study showed that roughly half of these events were sputum eosinophil high, which suggests that inflammatory pathways beyond IL-5 may remain upregulated and clinically relevant in some people¹⁰. The nature of these pathways is not understood.

The MAPLE trial was an exploratory, multi-centre, randomized, double-blind, placebo-controlled, crossover study of prednisolone in adults treated with mepolizumab at stable state. There was no significant clinical improvement with prednisolone¹¹. In this manuscript, we report the planned transcriptomic and proteomic analysis of sputum, plasma, and nasal samples from the MAPLE trial, to test the hypothesis that prednisolone exerts additional anti-inflammatory effects to mepolizumab at stable state.

Methods

Study design

The MAPLE study population and design were described previously¹¹. Briefly, participants were recruited before or after starting mepolizumab for usual clinical care. All participants received at least 12 weeks of mepolizumab, and were OCS-free for over 4 weeks, before they were randomised (1:1) to prednisolone (0.5 mg/kg/d, maximum dose of 40 mg/d, 14 ± 2 days) followed by a 4-week washout

period and then matched placebo or *vice versa* (figure 1). Prednisolone adherence was confirmed with tablet counts and serum prednisolone and cortisol levels.

Sputum, blood, and nasal sampling and processing

Figure 1 describes the trial sampling.

Sputum was induced using hypertonic saline (see Appendix). Samples were transferred to the laboratory within 2 hours for initial processing. Sputum plugs were selected from saliva, weighed, and centrifuged in phosphate-buffered saline (PBS) at 790 g for 10 minutes at 4°C. Samples were aliquoted and stored at -80°C in cryovials until further processing.

Nasal scrape samples were obtained using disposable curettes from the inferior turbinate under direct visualisation (see Appendix). Cells were stored in an RNAprotect cell reagent at -80 °C until processing.

Olink® proteomic analysis of sputum and plasma samples

Sputum supernatant and plasma from all visits were analysed by Olink® Explore 1536 proximity extension assays for 1536 proteins. Olink® hybridises DNA oligonucleotide-labelled antibody pairs to proteins, extends oligonucleotides by DNA polymerase, quantifies them with next generation sequencing, and log₂ transforms concentrations to a normalised protein expression (NPX) value.

Statistical analyses were performed with the Olink® Analyze package in R, according to manufacturer's guidance. We determined the effect of prednisolone *versus* placebo on paired sputum and plasma proteins using a change from baseline analysis in a linear mixed effects model¹². The type and sequence of treatment were fixed effects and participant ID was the random effect, with Benjamini-Hochberg (BH) correction for multiplicity of testing. An FDR <0.05 and absolute log₂fold change >0.5 was considered significant. The effect of mepolizumab on sputum and serum proteins was determined using paired t-tests with BH correction. Pathway analysis was performed using Cluster Profiler, heatmaps using Pheatmap, and correlation matrices using Corrplot.

ELISA of sputum samples

Sputum supernatants from all visits were assayed using the Ella® ELISA platform (Bio-Techne) for 15 analytes: IL-17E/IL-25, IL-1α, IL-1β, IL-2Ra, IL-33, IL-6, IL-6R, MPO, IL-3, YKL-40, IL-1RA, GM-CSF, TSLP, EDN, IL1R1. The effect of prednisolone *versus* placebo was analysed on paired samples using linear mixed effects models, and the effect of mepolizumab using paired t-tests, with BH correction (FDR<0.05).

Bulk transcriptomics of nasal tissue samples

RNA was extracted using Qiagen RNeasy mini kits. Samples with RNA integrity number <5 and/or undetectable RNA quantity were excluded from sequencing. RNA poly A sequencing was performed

by Novogene using Illumina Novaseq PE150. Gene alignment and quantification was performed by Novogene.

Differentially expressed genes were identified in DESeq2 using paired t-tests on gene count data between post-treatment placebo and prednisolone samples, and pre-mepolizumab to post-mepolizumab (post-placebo) samples. These underwent BH correction. An FDR <0.1 and absolute log2fold change >0.5 was considered significant.

MicroRNA sequencing of plasma samples

Plasma samples underwent Illumina TruSeq Small RNA analysis, read alignment, and quantification by Q² solutions[®]. Precursor and mature miRNA read counts were analysed for differential expression using a linear mixed effects model for the placebo *versus* prednisolone comparison, and paired t-tests comparing pre- to post-mepolizumab in DESeq2. These underwent BH correction. An FDR <0.1 and absolute log2fold change >0.5 was considered significant.

Sample size

As this was a secondary analysis on a trial powered on clinical endpoints, a power calculation was not performed. Findings are exploratory and descriptive.

Results

Comparison between prednisolone and placebo

Olink[®]

16 participants (supplementary table 1 and 2) had paired sputum samples before and after prednisolone and placebo in the crossover trial for Olink[®] analysis. 158 proteins were significantly downregulated by prednisolone and 51 proteins were significantly upregulated (figure 2, table 1, supplementary table 3 and 4).

Pathway analysis in sputum showed key downregulated biological processes including chemokine pathways crucial to subepithelial Th2/Tc2 cell, eosinophil and mast cell recruitment, and pathways related to IL-4 activity. Highly downregulated proteins related to type-2 inflammation and chemotaxis included IL-4, IL-13, IL-5, eotaxins 2 and 3 (CCL24 and CCL26), the chemokines CCL17 (TARC) and CCL22, OX40 receptor (TNFRSF4), TIMD4, the low-affinity IgE receptor FCER2, and the ST2 receptor for IL-33 (IL1RL1). Cell adhesion molecules involved in granulocyte trafficking, such as ICAM1, VCAM, TIE1, and e-selectin (SELE), were also downregulated. A reduction in the prostaglandin synthases PTGDS and HPGDS, SIGLEC6, and carboxypeptidase B1 (CPB1) that is closely related to mast cell tryptase (CPA3, not in the Olink[®] panel) suggests a downregulation of mast cell activity. Likewise, mast cell alpha-1/beta-1 tryptase (TPSAB1) was numerically reduced by over 50% though non-significant (adjusted p = 0.07). Proteins related to NF-kappaB (NF-κB) signalling were downregulated, including CD30 (TNFRSF8), CD69, and RANKL (TNFSF11), as were proteins related to tissue remodelling, including MMP12 and MMP1. NCAM1, NCAN, and GDNF, mediators of neuroimmune activity, were also downregulated.

Key upregulated pathways in sputum related to tissue repair and neutrophilic inflammation, such as IL-1β, IL-17C, and IL-22RA1. FKBP5, a classical hallmark of corticosteroid exposure, was upregulated.

Unsupervised clustering of the 40 proteins most downregulated in sputum after prednisolone revealed heterogeneity between individuals (n=16) in the anti-inflammatory effects of prednisolone (figure 2D). We analysed the correlation between clinical data and change in protein expression for biologically important sputum proteins after prednisolone (figure 3). This showed a reduction in sputum IL-5 and prostaglandin D synthase had an inverse correlation with FEV₁, change in RANKL was correlated with change in FeNO, and IL-33 correlated with the downregulation of other type-2 cytokines and chemokines, but TSLP did not.

25 participants (supplementary table 1 and 2) had paired plasma samples before and after prednisolone and placebo in the crossover trial analysed by Olink[®]. 74 proteins were significantly

downregulated by prednisolone and 53 proteins were significantly upregulated (figure 4, table 2, supplementary table 5 and 6).

The pathway analysis in plasma corresponded to that in sputum, showing downregulated biological processes included chemokines, type-2 inflammation pathways, and granulocyte trafficking. Relevant downregulated proteins included IL-4, eotaxin-3 (CCL26), CCL17 (TARC), CCL22, CCL19, CD83, OX40 receptor (TNFRSF4), ICAM1, VCAM. Proteins linked to remodelling such as MMP12, MMP1, and MMP10 were reduced. Proteins related to NF- κ B signalling were also downregulated, including CD30 (TNFRSF8) and B-cell activating factor (TNFSF13B). There was a decrease in plasma IL-6. Additionally, plasma IL-12 activity, and associated plasma IFN- γ and IP-10 (CXCL10) levels, were significantly reduced by prednisolone.

Key upregulated processes in plasma included IL-10 and amphiregulin (AREG), important in tissue repair. MMP3 and FKBP5 were raised by prednisolone; both are previously associated with corticosteroid exposure^{13,14}.

Unsupervised clustering of change in NPX after prednisolone and placebo in the 40 most downregulated plasma proteins suggested the anti-inflammatory effects of prednisolone in serum were more homogeneous compared to sputum (n=25) (figure 4D).

ELISA

16 participants provided sputum samples before and after prednisolone and placebo in the crossover trial that underwent ELISA. Eosinophil-derived neurotoxin (EDN) was significantly reduced after prednisolone (-1,180 ng/ml, adjusted p = 0.0001), while IL-1RA was significantly increased (16.9 ng/ml, adjusted p = 0.045). All other assays did not significantly change (supplementary figure 1, supplementary table 7).

Nasal tissue transcriptomics

6 participants provided good quality nasal tissue RNA post-prednisolone and post-placebo. 28 genes were significantly downregulated, and 3 genes significantly upregulated (Supplementary figure 2, supplementary table 8). Key genes suppressed by prednisolone included those involved in leucocyte chemotaxis (*CXCL2*, *CXCL3*, *CX3CR1*, *CXCL5*), mast cell tryptase (*TPSB2*), gasdermin D (*GSDMD*), 15-lipoxygenase (*ALOX15*), and matrix metalloproteinase 12 (*MMP12*).

Plasma microRNA

25 participants had paired plasma tested for miRNA. There were 19 significantly differentially expressed precursor miRNAs after prednisolone. However, there were no significantly differentially expressed mature miRNAs (supplementary table 9).

Comparison between pre-mepolizumab and post-mepolizumab

Olink®

Paired sputum samples from 10 participants were analysed by Olink® before and after starting mepolizumab. There were no significantly changed proteins. To understand the trend in pathways affected by mepolizumab, we performed pathway analysis using FDR <0.15 and abs log2fold >0.5. Upregulated pathways included MAPK signalling and regulation of leucocyte migration (supplementary figure 3).

Paired plasma samples from 10 participants were analysed by Olink® before and after starting mepolizumab. Galectin-10 (CLC) was the only significantly differing protein in plasma, reduced by 70%. Pathway analysis using FDR <0.15 and abs log2fold >0.2 showed mepolizumab initiation downregulated pathways related to lymphoid cell interactions and IL-3, IL-5, and GM-CSF signalling (supplementary figure 4).

ELISA

There were no significant differences associated with mepolizumab initiation in any of the sputum proteins tested.

Nasal tissue transcriptomics

5 participants provided good quality nasal tissue RNA before and after mepolizumab treatment. 272 genes were significantly downregulated and 105 significantly upregulated (supplementary table 10 and 11). In pathway analysis downregulated genes were related to ciliary structure and function. Upregulated genes were related to keratinisation, extracellular matrix formation, tight junctions, and IL-4 and IL-13 signalling pathways (figure 5).

Plasma microRNA

There were no significant differentially expressed precursor or mature miRNAs associated with mepolizumab initiation.

Discussion

Using a combination of high-throughput proteomics and bulk transcriptomics analysis of airway and blood samples at stable state, we found that oral prednisolone has extensive additional biological effects on top of mepolizumab.

The most striking effect of prednisolone on airway samples was the marked reduction in type-2 cytokines and chemokines (figure 6). Eotaxins, CCL17 and CCL22 are released by the airway epithelium and dendritic cells in response to IL-4 and TSLP. These chemokines have an important role in stimulating Th2/Tc2 cell and mast cell recruitment via CCR3 and CCR4 receptors. The resultant release of IL-4, IL-5, and IL-13 perpetuates Th2 cell differentiation, stimulates B cell maturation and production of IgE, and acts in concert with eotaxins, VCAM, ICAM1, E-selectin, and TIE1 to recruit eosinophils to the airway epithelium⁸. This self-perpetuating inflammation is analogous to an airway epithelial “magnet” recruiting effector granulocytes, analogous to “bombs”¹⁵. Our proteomic data demonstrate this process remains active and corticosteroid-responsive despite mepolizumab treatment, which selectively inhibits recruitment of eosinophils through IL-5. These findings are supported by the reduction of EDN in sputum, and CXCL2, CXCL3 and CXCL5 transcriptional suppression in nasal tissue.

Biological mechanisms underpinning these effects are complex. Corticosteroids impair dendritic cell maturation, downregulate IL-4 receptor expression, inhibit IL-4/IL-13 induced STAT6 signalling, and directly inhibit e-selectin, VCAM and ICAM1 expression¹⁶⁻¹⁹. Additionally, corticosteroids upregulate the production of IL-10, as observed in our plasma proteomics, which reduces Th2 cytokine production²⁰. Interestingly, the anti-inflammatory effects of prednisolone in the airway varied between individuals despite confirmed prednisolone adherence in all participants. This means that these inflammatory pathways were only active in some people at stable state, suggesting heterogeneity amongst the severe eosinophilic asthma phenotype.

Prednisolone also suppressed mast cell activity. Prostaglandin synthases (PTGDS, HPGDS) and Siglec-6 were reduced, and *15-lipoxygenase* was transcriptionally suppressed. Mast cell tryptase (CPA3) was not in the Olink[®] panel, but the related CPB1 which resides at the same gene locus as CPA3²¹, was significantly reduced. Additionally, the beta tryptase gene in mast cells (*TPSB2*) was transcriptionally suppressed in nasal tissue. Corticosteroids reduce mast cell signalling pathways and attenuate histamine release²². *Post-hoc* analysis of nasal transcriptome data in children suggested that mepolizumab may enhance mast cell and epithelial inflammation²³. If corticosteroid-responsive mast cell activation remains present in some individuals despite mepolizumab therapy, this may be clinically relevant due to mast cell mediated airway smooth muscle contraction, and may drive some residual exacerbations which occur despite effective eosinophil depletion.

Proteomics showed prednisolone reduced the matrix metalloproteinases MMP12 and MMP1 in sputum and plasma, which correlated with type-2 epithelial cytokines. Furthermore, MMP12 transcription was significantly downregulated in nasal tissue after prednisolone. MMP12 is produced by macrophages and bronchial epithelial cells²⁴. In murine models, it has been linked to airway remodelling via IL-13 dependent mechanisms^{25,26}. In humans, reduced *MMP12* expression due to a functional polymorphism was associated with improved pulmonary function²⁷. MMP1 expression is increased in the airways of people with asthma and is inversely associated with FEV₁²⁸. *In vitro*, MMP1 is induced via β 3-integrin-mediated MAPK signalling leading to reduced smooth muscle contraction²⁹. Additionally, MMP1 is activated by mast cell tryptase, which generates pro-proliferative extracellular matrix³⁰. Our data suggest that IL-13 and mast cell-dependent MMP12 and MMP1 activation remain in some patients with asthma despite anti-IL5 treatment. The consequent potential effect on long-term lung function decline needs further investigation.

Prednisolone also significantly reduced GDNF and NCAM, mediators of neuroimmune activity, and these reductions correlated strongly with other type-2 inflammatory proteins. GDNF signals through the NCAM receptor³¹. *In vitro*, GDNF expression is increased in human airway smooth muscle cells in asthma and influences intracellular calcium responses³². In rat asthma models, GDNF expression was significantly higher than in controls and central sensitisation to GDNF injected in the lateral ventricle led to a significant increase in bronchial IL-4 and Th2 cells, and significant decrease in pulmonary function³³. GDNF may contribute to smooth muscle hyperreactivity, and we found it was corticosteroid responsive.

Corticosteroids can inhibit signalling through the Toll-like receptor (TLR) cascade, including tethering and inhibiting transcription factors such as NF- κ B through glucocorticoid receptors³⁴. Our data suggest that components of NF- κ B signalling were downregulated by prednisolone and, interestingly, this was associated with reduction in FeNO. This association is plausible because inducible nitric oxide synthase (NOS2) production is dependent on NF- κ B signalling³⁵. The NOS2 protein was not in the Olink[®] panel so we cannot comment on its response to prednisolone. The mechanisms underpinning elevated FeNO in some people despite anti-IL5 treatment is of clinical interest. There was a suggestion in our proteomic and transcriptomic pathway analysis that MAPK – which is important in TLR signalling and transactivates STAT6^{36,37} – and IL-4/IL-13 signalling pathways were upregulated following mepolizumab initiation. We also found upregulation of ciliary and extracellular matrix genes in the nasal transcriptomics that may represent epithelial repair mechanisms. This would be consistent with previous nasal transcriptome data in mepolizumab-treated children where increased expression of epithelial signalling and extracellular matrix pathways were associated with more exacerbations, likely due to increased cycles of epithelial repair post-exacerbation³⁸. In adults, nasal epithelial cytokine and

alarmin activation (IL-4, TSLP, IL-33) after mepolizumab treatment was associated with non-responder status³⁹, suggesting a clinically relevant subgroup of severe eosinophilic asthma where inflammation is driven more by alarmins and the IL-4/13 pathway than by IL-5-mediated eosinophilia. As this epithelial signalling pathway was steroid-responsive, we speculate this inflammation could also be targeted using anti-IL4R α , anti-TSLP, anti-IL33, or JAK inhibitors.

Our study has several strengths. Firstly, the crossover study design allowed paired comparison of biological effects of prednisolone *versus* placebo within each participant, thus avoiding inter-participant variability. Secondly, high-throughput multiomic analyses of both airway and plasma samples enabled extensive exploration of prednisolone effects in both compartments. The only previous such placebo-controlled proteomic study of prednisolone in asthma analysed only plasma⁵. Thirdly, the 4-week washout and objective measurement of plasma prednisolone levels in participants provide confidence that effects are attributable to treatment.

The main limitation of our study was modest sample size, particularly for nasal transcriptomic analysis due to low quality nasal scrape samples. Whilst the paired crossover design improved power for prednisolone *versus* placebo assessment, the lack of placebo control limited assessment of mepolizumab effects. Our findings are exploratory and should be replicated in future cohorts. Furthermore, whilst sampling occurred within a clinical trial, allowing close comparison of biological and clinical effects, our study was undertaken at stable state without long-term follow-up during exacerbations, so we did not detect significant clinical changes.

The advent of anti-IL5 monoclonal antibody therapy has dramatically altered management of severe asthma. Patients rely less on oral corticosteroids, with their extensive associated toxicity, to achieve asthma control. However, severe asthma is a heterogeneous condition. Our data demonstrate that some individuals have IL-5-independent airway epithelial inflammatory pathways which remain active and are corticosteroid-responsive. The limited effect of prednisolone on symptoms and lung function in MAPLE suggests that these residual corticosteroid responsive pathways play a minor functional role in stable patients treated with mepolizumab. However, the effect of this persistent inflammation on key asthma outcomes such as long-term lung function decline and risk of exacerbations is unknown. Prospective studies are required to determine if there are clinically relevant anti-inflammatory effects of prednisolone at exacerbation in people treated with asthma monoclonal antibodies⁴⁰.

Contributors

RC, LGH, IDP, CEB, FY, JB contributed to conceptualisation and design of the protocol. FY, JPM, SED, CB contributed to acquisition of study data. Lab samples were processed by IH, VB and JC. Data were

analysed by IH and EM. IH and EM have accessed and verified the data. IH and TSCH drafted this submission which was approved by all authors.

Declaration of interests

IH has received a conference travel grant from GSK.

FY and SED have received speaker fees from AstraZeneca.

JC, VB, and EM report no declarations of interest.

AA is currently an employee of AZ.

PJM has received support to attend educational meetings from Chiesi.

JB has received personal fees from NuvoAir, and a research grant to his Institute from AstraZeneca, outside the submitted work.

CB has received speakers fees from AZ and GSK and has received advisory board fees from AZ.

LH has received grants from GSK, Astra Zeneca, Roche/Genentech, has given lectures supported by Astra Zeneca, Sanofi, Circassia, GlaxoSmithKline, has received travel grants from AstraZeneca and GSK, and has honoraria for Advisory Board Meetings from Novartis, Roche/Genentech, GSK, Teva and Celltrion.

IDP has received speaker's honoraria for speaking at sponsored meetings from Astra Zeneca, Boehringer Ingelheim, Aerocrine, Almirall, Novartis, Teva, Chiesi, Sanofi/Regeneron, Menarini and GSK and payments for organising educational events from AZ, GSK, Sanofi/Regeneron and Teva. He has received honoraria for attending advisory panels with Genentech, Sanofi/Regeneron, Astra Zeneca, Boehringer Ingelheim, GSK, Novartis, Teva, Merck, Circassia, Chiesi and Knopp and payments to support FDA approval meetings from GSK. He has received sponsorship to attend international scientific meetings from Boehringer Ingelheim, GSK, Astra Zeneca, Teva and Chiesi. He has received a grant from Chiesi to support a phase 2 clinical trial in Oxford. He is co-patent holder of the rights to the Leicester Cough Questionnaire and has received payments for its use in clinical trials from Merck, Bayer and Insmmed. In 2014-5 and 2019-20 he was an expert witness for a patent dispute involving Astra Zeneca and Teva.

CEB has received grants and consultancy fees from 4D Pharma, Areteia, AstraZeneca, Chiesi, Genentech, GlaxoSmithKline, Mologic, Novartis, Regeneron Pharmaceuticals, Roche and Sanofi.

RC has received lecture fees from GSK, AZ, Teva, Chiesi, Sanofi and Novartis; honoraria for Advisory Board Meetings from GSK, AZ and Celltrion; sponsorship to attend international scientific meetings from Chiesi, Sanofi and GSK and a research grant to her Institute from AZ for a UK multi-centre study.

TSCH has received grants from the Wellcome Trust, grants from The Guardians of the Beit Fellowship, and grants from the NIHR Oxford Biomedical Research Centre during the conduct of the study; and grants from Pfizer Inc., grants from University of Oxford, personal fees from Astra Zeneca, personal fees from TEVA, personal fees from Peer Voice outside the submitted work.

Data sharing

Anonymised patient level data analysed and presented in this study are available from the corresponding author on reasonable request, providing the request meets local ethical and research governance criteria after publication. Data will be available immediately after publication for 10 years.

Funding

This study was funded jointly by the Medical Research Council (MRC) UK (MR/M016579/1) and industrial partners within the MRC Refractory Asthma Stratification Programme consortium and by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). Proteomics and transcriptomics analysis was funded by GSK (ID: 215294). TSCH is supported by a Wellcome Trust Fellowship (211050/Z/18/z). All authors had full access to the full data in the study and accept responsibility to submit for publication.

Acknowledgements

The authors are grateful to all the participants who volunteered and to the clinical and research teams at all the participating centres. For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

References

1. Levy ML, Bacharier LB, Bateman E, et al. Key recommendations for primary care from the 2022 Global Initiative for Asthma (GINA) update. *NPJ Prim Care Respir Med* 2023;33:7.
2. Rowe BH, Spooner CH, Ducharme FM, Bretzlaff JA, Bota GW. Corticosteroids for preventing relapse following acute exacerbations of asthma. *Cochrane Database Syst Rev* 2007;Cd000195.
3. Sahid El-Radhi A, Hogg CL, Bungre JK, Bush A, Corrigan CJ. Effect of oral glucocorticoid treatment on serum inflammatory markers in acute asthma. *Arch Dis Child* 2000;83:158-62.
4. Moran AM, Ramakrishnan S, Borg CA, et al. Blood Eosinophil Depletion with Mepolizumab, Benralizumab, and Prednisolone in Eosinophilic Asthma. *Am J Respir Crit Care Med* 2020;202:1314-6.
5. Sparreman Mikus M, Kolmert J, Andersson LI, et al. Plasma proteins elevated in severe asthma despite oral steroid use and unrelated to Type-2 inflammation. *Eur Respir J* 2022;59.
6. Schleimer RP, Bochner BS. The effects of glucocorticoids on human eosinophils. *J Allergy Clin Immunol* 1994;94:1202-13.
7. Bentley AM, Hamid Q, Robinson DS, et al. Prednisolone treatment in asthma. Reduction in the numbers of eosinophils, T cells, tryptase-only positive mast cells, and modulation of IL-4, IL-5, and interferon-gamma cytokine gene expression within the bronchial mucosa. *Am J Respir Crit Care Med* 1996;153:551-6.
8. Howell I, Howell A, Pavord ID. Type 2 inflammation and biological therapies in asthma: Targeted medicine taking flight. *J Exp Med* 2023;220.
9. Skov IR, Madsen H, Henriksen DP, Andersen JH, Pottegård A, Davidsen JR. Low-dose oral corticosteroids in asthma associates with increased morbidity and mortality. *Eur Respir J* 2022;60.
10. McDowell PJ, Diver S, Yang F, et al. The inflammatory profile of exacerbations in patients with severe refractory eosinophilic asthma receiving mepolizumab (the MEX study): a prospective observational study. *Lancet Respir Med* 2021;9:1174-84.
11. Yang F, Busby J, Heaney LG, et al. Corticosteroid Responsiveness Following Mepolizumab in Severe Eosinophilic Asthma-A Randomized, Placebo-Controlled Crossover Trial (MAPLE). *J Allergy Clin Immunol Pract* 2022;10:2925-34.e12.
12. Mehrotra DV. A recommended analysis for 2 × 2 crossover trials with baseline measurements. *Pharm Stat* 2014;13:376-87.
13. James A, Kolmert J, Wheelock C, et al. Increased serum MMP-3 in severe asthma is associated with oral steroid use; data from U-BIOPRED and BIOAIR cohorts. *European Respiratory Journal* 2017;50:PA580.
14. Jääskeläinen T, Makkonen H, Palvimo JJ. Steroid up-regulation of FKBP51 and its role in hormone signaling. *Curr Opin Pharmacol* 2011;11:326-31.

15. Couillard S, Pavord ID, Heaney LG, Petousi N, Hinks TSC. Sub-stratification of type-2 high airway disease for therapeutic decision-making: A 'bomb' (blood eosinophils) meets 'magnet' (FeNO) framework. *Respirology* 2022;27:573-7.
16. Elftman MD, Norbury CC, Bonneau RH, Truckenmiller ME. Corticosterone impairs dendritic cell maturation and function. *Immunology* 2007;122:279-90.
17. So EY, Kim SH, Park HH, Cho BS, Lee CE. Corticosteroid inhibits IL-4 signaling through down-regulation of IL-4 receptor and STAT6 activity. *FEBS Lett* 2002;518:53-9.
18. Cronstein BN, Kimmel SC, Levin RI, Martiniuk F, Weissmann G. A mechanism for the antiinflammatory effects of corticosteroids: the glucocorticoid receptor regulates leukocyte adhesion to endothelial cells and expression of endothelial-leukocyte adhesion molecule 1 and intercellular adhesion molecule 1. *Proc Natl Acad Sci U S A* 1992;89:9991-5.
19. Atsuta J, Plitt J, Bochner BS, Schleimer RP. Inhibition of VCAM-1 expression in human bronchial epithelial cells by glucocorticoids. *Am J Respir Cell Mol Biol* 1999;20:643-50.
20. Ogawa Y, Duru EA, Ameredes BT. Role of IL-10 in the resolution of airway inflammation. *Curr Mol Med* 2008;8:437-45.
21. Atiakshin D, Kostin A, Trotsenko I, SamoiloVA V, Buchwalow I, Tiemann M. Carboxypeptidase A3-A Key Component of the Protease Phenotype of Mast Cells. *Cells* 2022;11.
22. Oppong E, Flink N, Cato AC. Molecular mechanisms of glucocorticoid action in mast cells. *Mol Cell Endocrinol* 2013;380:119-26.
23. Gaberino C, Segnitz RM, Cox M, et al. Mepolizumab Alters Regulation of Airway Type-2 Inflammation in Urban Children with Asthma by Disrupting Eosinophil Gene Expression but Enhancing Mast Cell and Epithelial Pathways. *Journal of Allergy and Clinical Immunology* 2023;151:AB125.
24. Lavigne MC, Thakker P, Gunn J, et al. Human bronchial epithelial cells express and secrete MMP-12. *Biochemical and biophysical research communications* 2004;324:534-46.
25. Pouladi MA, Robbins CS, Swirski FK, et al. Interleukin-13–dependent expression of matrix metalloproteinase-12 is required for the development of airway eosinophilia in mice. *American journal of respiratory cell and molecular biology* 2004;30:84-90.
26. Lanone S, Zheng T, Zhu Z, et al. Overlapping and enzyme-specific contributions of matrix metalloproteinases-9 and-12 in IL-13–induced inflammation and remodeling. *The Journal of clinical investigation* 2002;110:463-74.
27. Hunninghake GM, Cho MH, Tesfaigzi Y, et al. MMP12, lung function, and COPD in high-risk populations. *N Engl J Med* 2009;361:2599-608.

28. Cataldo DD, Gueders M, Munaut C, et al. Matrix metalloproteinases and tissue inhibitors of matrix metalloproteinases mRNA transcripts in the bronchial secretions of asthmatics. *Lab Invest* 2004;84:418-24.
29. Rogers NK, Clements D, Dongre A, Harrison TW, Shaw D, Johnson SR. Extra-cellular matrix proteins induce matrix metalloproteinase-1 (MMP-1) activity and increase airway smooth muscle contraction in asthma. *PLoS One* 2014;9:e90565.
30. Naveed S-u-n, Clements D, Jackson DJ, et al. Matrix Metalloproteinase-1 Activation Contributes to Airway Smooth Muscle Growth and Asthma Severity. *American Journal of Respiratory and Critical Care Medicine* 2017;195:1000-9.
31. Paratcha G, Ledda F, Ibáñez CF. The neural cell adhesion molecule NCAM is an alternative signaling receptor for GDNF family ligands. *Cell* 2003;113:867-79.
32. Bhallamudi S, Roos BB, Teske JJ, et al. Glial-derived neurotrophic factor in human airway smooth muscle. *J Cell Physiol* 2021;236:8184-96.
33. Wang X, Wang Y, Chen X, et al. The central glial cell line-derived neurotrophic factor (GDNF) regulates pulmonary function in asthmatic rats. *Ann Transl Med* 2023;11:113.
34. Cain DW, Cidlowski JA. Immune regulation by glucocorticoids. *Nat Rev Immunol* 2017;17:233-47.
35. Taylor BS, de Vera ME, Ganster RW, et al. Multiple NF-kappaB enhancer elements regulate cytokine induction of the human inducible nitric oxide synthase gene. *J Biol Chem* 1998;273:15148-56.
36. Peroval M, Boyd A, Young J, Smith A. A Critical Role for MAPK Signalling Pathways in the Transcriptional Regulation of Toll Like Receptors. *PloS one* 2013;8:e51243.
37. Athari SS. Targeting cell signaling in allergic asthma. *Signal Transduct Target Ther* 2019;4:45.
38. Jackson DJ, Bacharier LB, Gergen PJ, et al. Mepolizumab for urban children with exacerbation-prone eosinophilic asthma in the USA (MUPPITS-2): a randomised, double-blind, placebo-controlled, parallel-group trial. *Lancet* 2022;400:502-11.
39. Rakkar K, Pang YL, Rajasekar P, et al. Mepolizumab induced changes in nasal methylome and transcriptome to predict response in asthma. *medRxiv* 2023:2023.05.18.23290155.
40. Howell I, Mahdi M, Bafadhel M, et al. Recovery of Breakthrough Asthma Attacks Treated With Oral Steroids While on Monoclonal Antibody Therapy: Protocol for a Prospective Observational Study (BOOST). *JMIR Res Protoc* 2023;12:e46741.

Tables

Table 1 – Summary of 25 most downregulated proteins by prednisolone versus placebo in sputum (Olink®)


| Protein | Log2fold change | Adjusted p value | Function |
|----------|-----------------|------------------|---|
| IL4 | -3.37 | 4.81E-03 | Stimulates proliferation of Th2 cells, B cell class switching to IgE, and B cell differentiation to plasma cells. |
| SERPINA9 | -3.18 | 9.79E-06 | Inhibits trypsin-like serine proteases. |
| CCL24 | -2.6 | 2.47E-04 | Eotaxin-2. Induces eosinophil chemotaxis. |
| MMP1 | -2.6 | 1.44E-03 | Breaks down interstitial collagens in the ECM. |
| HLA-DRA | -2.46 | 1.88E-04 | Presents peptides from extracellular proteins. Expressed on B lymphocytes, dendritic cells and macrophages. |
| TIE1 | -2.44 | 3.70E-04 | Cell surface protein that upregulates cell adhesion molecules. |
| CCL26 | -2.43 | 5.02E-03 | Eotaxin-3. Induces eosinophil and basophil chemotaxis. |
| CCL22 | -2.43 | 2.79E-02 | Induces T cell, monocyte, and dendritic cell chemotaxis. |
| CCL17 | -2.37 | 6.68E-03 | Induces T cell and granulocyte chemotaxis. |
| GDNF | -2.29 | 3.96E-03 | Promotes neuronal growth and survival. |
| NCAM1 | -2.18 | 6.60E-04 | Mediates neurite outgrowth |
| TIMD4 | -2.16 | 8.56E-05 | Involved in recognition of apoptotic cells for macrophage phagocytosis. |
| CCL13 | -2.15 | 6.92E-03 | Induces T cell, monocyte, and granulocyte chemotaxis. |
| NCAN | -2.11 | 1.99E-04 | Component of the ECM and involved in neuronal growth |
| MCAM | -2.09 | 1.93E-04 | Vascular endothelial cell adhesion molecule |
| SELE | -2.07 | 3.74E-04 | Endothelial adhesion molecule involved in leukocyte recruitment. |
| DOK2 | -2.07 | 4.36E-03 | Scaffolding protein for multimolecular signalling complexes. |
| IL5 | -2.06 | 2.89E-04 | Mediates eosinophil production and activation |
| GP1BA | -2.04 | 2.65E-04 | Platelet surface glycoprotein |
| VCAM1 | -2.03 | 1.40E-03 | Mediates adhesion of lymphocytes, eosinophils, and basophils to the vascular endothelium. |
| CPA2 | -2 | 1.47E-02 | Pancreatic carboxypeptidase. |
| ARSB | -1.97 | 2.63E-03 | Enzyme that breaks down glycosaminoglycans. |
| MMP12 | -1.97 | 6.09E-03 | Breaks down elastin in the ECM. |
| TNFRSF8 | -1.95 | 5.57E-05 | Expressed by T and B cells. Leads to activation of NF-κB. |
| SPP1 | -1.94 | 8.87E-04 | Osteopontin. Anti-apoptotic factor in macrophages and T cells. Also involved in leukocyte chemotaxis. |

Table 2 – Summary of 25 most downregulated proteins by prednisolone versus placebo in plasma (Olink®)

| Protein | Log2fold change | Adjusted p value | Function |
|---------|-----------------|------------------|-------------|
| IL4 | -3.17 | 1.06E-05 | See table 1 |

| | | | |
|-------------|-------|----------|---|
| IL12A_IL12B | -2.24 | 1.55E-26 | Encodes a subunit of IL-12. IL-12 is produced by dendritic cells, macrophages, and neutrophils to promote Th1 differentiation. |
| IL12B | -2.09 | 3.88E-26 | Encodes a subunit of IL-12. |
| CRH | -1.46 | 3.80E-11 | Corticotrophin-releasing hormone. Stimulates ACTH production. |
| CCL22 | -1.42 | 4.18E-21 | See table 1 |
| MMP12 | -1.42 | 5.07E-14 | See table 1 |
| CCL19 | -1.24 | 2.94E-14 | Involved in T and B cell migration to secondary lymphoid organs. |
| TNFSF13B | -0.94 | 1.03E-20 | B cell activating factor. Important in B cell proliferation. |
| THBS4 | -0.93 | 7.59E-08 | Adhesive glycoprotein in cell-to-matrix interactions. |
| FGF19 | -0.93 | 1.63E-02 | Regulates bile acid synthesis. |
| GGH | -0.91 | 9.66E-18 | Key enzyme in intracellular folate homeostasis. |
| TNFRSF8 | -0.91 | 1.67E-16 | See table 1 |
| TNFRSF4 | -0.91 | 3.98E-19 | OX40 receptor. Combines with OX40 ligand to prolong T cell survival, develop memory T cells, and polarise to Th2 cells. |
| CCDC80 | -0.9 | 1.29E-14 | Promotes cell adhesion and ECM assembly. |
| SIGLEC6 | -0.86 | 2.87E-21 | Glycan-binding cell-surface protein highly expressed on mast cells. Inhibits mast cell activation after engagement with specific ligands. |
| TNFRSF9 | -0.85 | 9.89E-14 | Cell surface protein that stimulates T and B cells, and dendritic cells. |
| CD83 | -0.84 | 4.03E-13 | Stabilises MHC II. |
| IFNG | -0.84 | 6.50E-03 | Interferon gamma. Activates macrophages and MHC class II expression. |
| CLEC4C | -0.81 | 5.93E-06 | Cell surface receptor on dendritic cells involved in antigen capturing. |
| CCL18 | -0.81 | 2.03E-08 | Induces T cell, B cell, and dendritic cell chemotaxis. |
| CXCL10 | -0.79 | 3.74E-09 | Induces T cell, macrophage, and dendritic cell chemotaxis. |
| CCL17 | -0.77 | 4.81E-11 | See table 1 |
| PDCD1 | -0.77 | 2.35E-21 | Cell surface receptor on T and B cells that suppresses T cell activity. |
| CCL26 | -0.75 | 4.31E-06 | See table 1 |
| CCN3 | -0.74 | 5.80E-10 | Binds to integrin and involved ECM structure and function. |

Figures



Graphical abstract




Figure 1 – Schematic of the MAPLE study design and sampling

Participants entered the study at either visit 1 (pre-mepolizumab) or visit 2 (at least 12 weeks post-mepolizumab initiation). At visit 2 all participants were randomised to either prednisolone or placebo treatment for 2 weeks, followed by a 4-week washout, and at visit 4 crossed over to receive the opposite treatment for 2 weeks. At visits 1, 3 and 5 (red circle) participants had blood, induced sputum, and nasal scrape sampling. At visits 2 and 4 (blue circle) participants had blood and induced sputum sampling. Blood sampling consisted of EDTA and SST blood tubes.







Figure 2 – Summary of prednisolone effects on proteins measured by Olink® in sputum

A) volcano plot of differentially expressed proteins by prednisolone in sputum. An absolute log₂ fold >0.5 (vertical line) and FDR <0.05 (horizontal line) were considered significant. Blue, downregulated by prednisolone; red, upregulated by prednisolone; black, non-significant. B,C) Top 15 most over-represented Gene Ontology (GO) biological processes significantly downregulated (B) or (C) upregulated by prednisolone in sputum. Colour denotes adjusted p value (FDR<0.05). D) heatmap showing individual change in NPX (n=16 participants) after prednisolone and placebo in the 40 most significantly downregulated sputum proteins from the linear mixed effects model. Participants and proteins underwent unsupervised clustering. Top row denotes treatment received, either placebo (blue) or prednisolone (red). Other rows: red denotes increased protein expression, blue reduced expression.






Figure 3 - Correlation matrix of sputum proteins and clinical data

Matrix of Pearson correlation coefficients between clinical data and change in protein expression for biologically important sputum proteins after prednisolone. Only significant positive (blue) or negative (red) correlations are displayed (p<0.05). Key: feno_pre_treat, FeNO at baseline; delta_feno, change in FeNO after prednisolone; delta_fev1, change in FEV₁ after prednisolone; BMI, body mass index at baseline; sp_eos_pc_pre_treatment, sputum eosinophil count at baseline


A



B



C



D




Figure 4 - Summary of prednisolone effects on proteins measured by Olink® in plasma

A) volcano plot of differentially expressed proteins by prednisolone in plasma. An absolute log₂ fold >0.5 (vertical line) and FDR <0.05 (horizontal line) were considered significant. Blue, downregulated by prednisolone; red, upregulated by prednisolone; black, non-significant. B,C) Top 15 most over-represented Gene Ontology (GO) biological processes significantly downregulated (B) or (C) upregulated by prednisolone in plasma. Colour denotes adjusted p value (FDR<0.05). D) heatmap showing individual change in NPX (n=16 participants) after prednisolone and placebo in the 40 most significantly downregulated plasma proteins from the linear mixed effects model. Participants and proteins underwent unsupervised clustering. Top row denotes treatment received, either placebo (blue) or prednisolone (red). Other rows: red denotes increased protein expression, blue reduced expression.




Figure 5 – Differentially expressed nasal genes after mepolizumab

A) volcano plot of differentially expressed genes by mepolizumab in nasal tissue. An absolute log₂ fold >0.5 (vertical line) and FDR <0.1 (horizontal line) were considered significant. Blue, downregulated by prednisolone; red, upregulated by prednisolone; black, non-significant. B,C) Top 15 most over-represented Gene Ontology (GO) biological processes significantly downregulated (B) or (C) upregulated by mepolizumab in nasal tissue. Colour denotes adjusted p value (FDR<0.05).





Figure 6 - Epithelial effects of oral corticosteroids in stable asthma on mepolizumab treatment


Eotaxins (CCL24 and CCL26), CCL17 (TARC) and CCL22 are released by the airway epithelium and dendritic cells in response to IL-4 and TSLP. These chemokines have an important role in stimulating Th2/Tc2 cell and mast cell recruitment via CCR3 and CCR4 receptors. The resultant release of IL-4, IL-5, and IL-13 perpetuates Th2 cell differentiation, stimulates B cell maturation and production of IgE. IL-4 also acts in concert with eotaxins, VCAM, ICAM1, E-selectin, and TIE1 to recruit eosinophils, which are produced in the bone marrow after IL-5 stimulation, to the airway epithelium. Mepolizumab blocks IL-5, thereby reducing the production of eosinophils in the bone marrow. Oral corticosteroids have wider effects than mepolizumab, reducing IL-4, IL-13, eotaxins, CCL17, CCL22 and mast cell activity, and increasing IL-10 activity which inhibits Th2 cell production. CCL, C-C Motif Chemokine Ligand; C-C chemokine receptor, CCR; GM-CSF, Granulocyte-Macrophage Colony-Stimulating Factor; Interleukin, IL; ICAM, Intercellular Adhesion Molecule 1; Thymic Stromal Lymphopoietin, TSLP; VCAM, Vascular Cell Adhesion protein.

Supplementary material




Supplementary figure 1 – The effect of prednisolone on 15 selected proteins in sputum (ELISA)

Summary of the 15 proteins tested by ELISA in sputum samples. This box and whiskers plot shows concentration versus treatment (green, before mepolizumab; light blue, before placebo; dark blue, after placebo; light red, before prednisolone; dark red, after prednisolone). The p value from before to after mepolizumab was determined using paired t-tests, and between prednisolone *versus* placebo using linear mixed effects modelling. No proteins were significantly regulated by mepolizumab. Prednisolone significantly decreased EDN, and significantly increased IL1Ra.




Supplementary figure 2 – Heatmap showing effect of prednisolone and placebo on nasal transcriptome

Heatmap showing post treatment gene expression (n=6 participants) after prednisolone and placebo in the 31 significantly differentially expressed genes. Participants and genes underwent unsupervised clustering. Top row denotes treatment received, either placebo (blue) or prednisolone (red). Other rows: red denotes high gene expression, blue low expression.



Supplementary figure 3 – Pathways analysis of upregulated Olink® sputum proteins after mepolizumab

Top 10 most over-represented Gene Ontology (GO) biological processes for sputum proteins with a trend for upregulation by mepolizumab (FDR<0.15, abs log2fold >0.5). Colour denotes adjusted p value of pathway analysis (FDR<0.05).



Supplementary figure 4 – Pathways analysis of downregulated Olink® plasma proteins after mepolizumab

Top 10 most over-represented Reactome processes for plasma proteins with a trend for downregulation by mepolizumab (FDR<0.15, abs log2fold >0.2). Colour denotes adjusted p value of pathway analysis (FDR<0.05).

Supplementary table 1 – Baseline demographics of patients who had plasma and sputum at all visits in the crossover trial compared with all patients randomised

| | Plasma patients (N=25) | Sputum patients (N=16) | All randomised patients (N=27) |
|--|---------------------------|---------------------------|-----------------------------------|
| Gender | | | |
| Female | 8 (32.0%) | 6 (37.5%) | 9 (33.3%) |
| Male | 17 (68.0%) | 10 (62.5%) | 18 (66.7%) |
| Age Study Entry | | | |
| Mean (SD) | 56.4 (12.3) | 60.5 (8.46) | 56.9 (12.1) |
| Median [Min, Max] | 58.0 [22.0, 74.0] | 62.0 [47.0, 74.0] | 58.0 [22.0, 74.0] |
| Ethnicity | | | |
| African | 1 (4.0%) | 1 (6.3%) | 1 (3.7%) |
| Caucasian | 24 (96.0%) | 15 (93.8%) | 26 (96.3%) |
| Smoking Status | | | |
| Ex-smoker | 9 (36.0%) | 7 (43.8%) | 10 (37.0%) |
| Never smoker | 16 (64.0%) | 9 (56.3%) | 17 (63.0%) |
| Pack Years | | | |
| Mean (SD) | 4.26 (9.47) | 5.53 (11.4) | 4.13 (9.14) |
| Median [Min, Max] | 0 [0, 45.0] | 0 [0, 45.0] | 0 [0, 45.0] |
| Age Asthma Diagnosis | | | |
| Mean (SD) | 31.9 (22.2) | 39.3 (21.1) | 31.7 (22.1) |
| Median [Min, Max] | 34.0 [0, 66.0] | 43.5 [10.0, 66.0] | 34.0 [0, 66.0] |
| BMI_v1 | | | |
| Mean (SD) | 31.4 (5.38) | 32.3 (4.81) | 31.5 (5.19) |
| Median [Min, Max] | 31.9 [16.8, 40.5] | 32.2 [23.8, 40.5] | 31.9 [16.8, 40.5] |
| Missing | 1 (4.0%) | 1 (6.3%) | 1 (3.7%) |
| Exacerbations Pre Mepo | | | |
| Mean (SD) | 0.400 (1.26) | 0.188 (0.544) | 0.370 (1.21) |
| Median [Min, Max] | 0 [0, 6.00] | 0 [0, 2.00] | 0 [0, 6.00] |
| Exacerbations requiring ITU/HDU | | | |
| Mean (SD) | 0.160 (0.473) | 0.125 (0.342) | 0.148 (0.456) |
| Median [Min, Max] | 0 [0, 2.00] | 0 [0, 1.00] | 0 [0, 2.00] |
| Allergic Rhinitis | | | |
| No | 21 (84.0%) | 14 (87.5%) | 22 (81.5%) |
| Yes | 4 (16.0%) | 2 (12.5%) | 5 (18.5%) |
| Perennial Rhinitis | | | |
| No | 11 (44.0%) | 7 (43.8%) | 13 (48.1%) |
| Yes | 14 (56.0%) | 9 (56.3%) | 14 (51.9%) |
| Nasal Polyps | | | |
| No | 17 (68.0%) | 9 (56.3%) | 18 (66.7%) |
| Yes | 8 (32.0%) | 7 (43.8%) | 9 (33.3%) |

| | | | |
|-----------------------------|-------------------|-------------------|-------------------|
| Eczema | | | |
| No | 19 (76.0%) | 14 (87.5%) | 20 (74.1%) |
| Yes | 6 (24.0%) | 2 (12.5%) | 7 (25.9%) |
| Depression/Anxiety | | | |
| No | 20 (80.0%) | 12 (75.0%) | 22 (81.5%) |
| Yes | 5 (20.0%) | 4 (25.0%) | 5 (18.5%) |
| Bronchiectasis | | | |
| No | 23 (92.0%) | 14 (87.5%) | 24 (88.9%) |
| Yes | 2 (8.0%) | 2 (12.5%) | 3 (11.1%) |
| Reflux | | | |
| No | 18 (72.0%) | 12 (75.0%) | 20 (74.1%) |
| Yes | 7 (28.0%) | 4 (25.0%) | 7 (25.9%) |
| ICS Dose | | | |
| Mean (SD) | 2020 (300) | 2030 (379) | 2010 (288) |
| Median [Min, Max] | 2000 [1000, 3000] | 2000 [1000, 3000] | 2000 [1000, 3000] |
| LABA | | | |
| Yes | 25 (100%) | 16 (100%) | 27 (100%) |
| LTRA | | | |
| No | 9 (36.0%) | 5 (31.3%) | 9 (33.3%) |
| Yes | 16 (64.0%) | 11 (68.8%) | 18 (66.7%) |
| LAMA | | | |
| No | 5 (20.0%) | 3 (18.8%) | 6 (22.2%) |
| Yes | 20 (80.0%) | 13 (81.3%) | 21 (77.8%) |
| Nebuliser | | | |
| No | 22 (88.0%) | 15 (93.8%) | 24 (88.9%) |
| Yes | 3 (12.0%) | 1 (6.3%) | 3 (11.1%) |
| Long term antibiotic | | | |
| No | 22 (88.0%) | 13 (81.3%) | 23 (85.2%) |
| Yes | 3 (12.0%) | 3 (18.8%) | 4 (14.8%) |

Supplementary table 2 – Clinical measures at randomisation visit (V2) of patients who had plasma and sputum at all visits in the crossover trial compared with all patients randomised

| | Plasma patients (N=25) | Sputum patients (N=16) | All randomised patients (N=27) |
|-------------------|---------------------------|---------------------------|-----------------------------------|
| ACQ5 Score | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 1.09 [1.05] | 1.21 [1.14] | 1.16 [1.17] |
| Median [Q1, Q3] | 0.600 [0.200, 1.80] | 0.600 [0.500, 2.05] | 0.600 [0.200, 1.90] |
| Median [Min, Max] | 0.600 [0, 3.60] | 0.600 [0, 3.60] | 0.600 [0, 4.00] |

| mAQLQ Score | | | |
|----------------------------------|----------------------|----------------------|----------------------|
| N | 25 | 16 | 27 |
| Mean [SD] | 5.41 [1.38] | 5.41 [1.45] | 5.35 [1.48] |
| Median [Q1, Q3] | 5.73 [4.33, 6.53] | 5.67 [4.25, 6.63] | 5.73 [4.17, 6.57] |
| Median [Min, Max] | 5.73 [2.67, 7.00] | 5.67 [2.67, 7.00] | 5.73 [2.27, 7.00] |
| SGRQ Total | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 29.8 [20.3] | 30.8 [22.0] | 30.3 [21.9] |
| Median [Q1, Q3] | 30.7 [14.7, 46.2] | 33.1 [15.0, 46.3] | 30.7 [13.9, 46.3] |
| Median [Min, Max] | 30.7 [0.850, 69.6] | 33.1 [0.850, 69.6] | 30.7 [0, 70.6] |
| VAS Overall | | | |
| N | 23 | 16 | 25 |
| Mean [SD] | 2.47 [2.28] | 2.49 [2.34] | 2.47 [2.31] |
| Median [Q1, Q3] | 2.00 [0.450, 4.25] | 2.00 [0.475, 4.00] | 2.00 [0.300, 4.50] |
| Median [Min, Max] | 2.00 [0, 8.00] | 2.00 [0, 8.00] | 2.00 [0, 8.00] |
| Missing | 2 (8.0%) | | 2 (7.4%) |
| VAS SOB | | | |
| N | 23 | 16 | 25 |
| Mean [SD] | 2.58 [2.47] | 2.68 [2.58] | 2.65 [2.58] |
| Median [Q1, Q3] | 2.00 [0.400, 5.00] | 2.00 [0.500, 5.00] | 2.00 [0.200, 5.00] |
| Median [Min, Max] | 2.00 [0, 8.00] | 2.00 [0, 8.00] | 2.00 [0, 8.00] |
| Missing | 2 (8.0%) | | 2 (7.4%) |
| VAS Cough | | | |
| N | 23 | 16 | 25 |
| Mean [SD] | 2.29 [2.43] | 2.54 [2.68] | 2.39 [2.56] |
| Median [Q1, Q3] | 1.00 [0.100, 4.25] | 1.20 [0.150, 5.00] | 1.00 [0, 4.50] |
| Median [Min, Max] | 1.00 [0, 7.60] | 1.20 [0, 7.60] | 1.00 [0, 7.60] |
| Missing | 2 (8.0%) | | 2 (7.4%) |
| Airway oscillometry R5 | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 4.01 [1.72] | 3.78 [1.58] | 4.04 [1.73] |
| Median [Q1, Q3] | 3.53 [2.84, 4.85] | 3.47 [2.81, 4.71] | 3.53 [2.71, 4.94] |
| Median [Min, Max] | 3.53 [1.48, 7.78] | 3.47 [1.48, 7.64] | 3.53 [1.48, 7.78] |
| Airway oscillometry R5-20 | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 1.00 [1.09] | 0.963 [1.06] | 1.02 [1.11] |
| Median [Q1, Q3] | 0.670 [0.304, 1.36] | 0.720 [0.335, 1.34] | 0.670 [0.282, 1.62] |
| Median [Min, Max] | 0.670 [-0.460, 3.60] | 0.720 [-0.460, 3.60] | 0.670 [-0.460, 3.60] |
| AX | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 17.5 [18.0] | 19.2 [19.5] | 18.3 [18.8] |

| | | | |
|-------------------------------|------------------------|------------------------|-----------------------|
| Median [Q1, Q3] | 7.92 [4.88, 30.8] | 9.34 [4.70, 32.8] | 7.92 [4.89, 31.3] |
| Median [Min, Max] | 7.92 [0.730, 57.5] | 9.34 [0.730, 57.5] | 7.92 [0.730, 57.5] |
| Airway oscillometry X5 | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | -1.24 [2.48] | -1.08 [2.86] | -1.39 [2.50] |
| Median [Q1, Q3] | -0.870 [-2.57, -0.480] | -0.840 [-2.59, -0.475] | -1.11 [-2.61, -0.535] |
| Median [Min, Max] | -0.870 [-6.14, 7.71] | -0.840 [-6.14, 7.71] | -1.11 [-6.14, 7.71] |
| FEV1 (% predicted) | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 77.9 [18.9] | 81.2 [20.7] | 77.0 [20.2] |
| Median [Q1, Q3] | 79.0 [67.0, 88.0] | 85.0 [70.8, 99.3] | 79.0 [66.0, 91.5] |
| Median [Min, Max] | 79.0 [32.0, 107] | 85.0 [32.0, 107] | 79.0 [32.0, 107] |
| FVC (% predicted) | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 92.9 [15.6] | 96.5 [14.8] | 91.8 [16.6] |
| Median [Q1, Q3] | 95.0 [83.0, 104] | 101 [8.38, 108] | 95.0 [8.15, 104] |
| Median [Min, Max] | 95.0 [52.0, 118] | 101 [70.7, 118] | 95.0 [52.0, 118] |
| FEV1/FVC ratio | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 0.662 [0.122] | 0.661 [0.121] | 0.659 [0.122] |
| Median [Q1, Q3] | 0.670 [0.650, 0.720] | 0.670 [0.650, 0.703] | 0.670 [0.650, 0.730] |
| Median [Min, Max] | 0.670 [0.280, 0.830] | 0.670 [0.280, 0.830] | 0.670 [0.280, 0.830] |
| FEF (% predicted) | | | |
| N | 18 | 9 | 18 |
| Mean [SD] | 0.531 [0.245] | 0.607 [0.268] | 0.531 [0.245] |
| Median [Q1, Q3] | 0.510 [0.383, 0.603] | 0.520 [0.490, 0.610] | 0.510 [0.383, 0.603] |
| Median [Min, Max] | 0.510 [0.130, 1.17] | 0.520 [0.320, 1.17] | 0.510 [0.130, 1.17] |
| Missing | 7 (28.0%) | 7 (43.8%) | 9 (33.3%) |
| PEF (L/min) | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 457 [141] | 451 [143] | 446 [145] |
| Median [Q1, Q3] | 425 [343, 550] | 435 [337, 572] | 424 [341, 537] |
| Median [Min, Max] | 425 [224, 805] | 435 [224, 705] | 424 [202, 805] |
| FeNO (ppb) | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 47.2 [32.6] | 49.5 [34.0] | 44.9 [32.4] |
| Median [Q1, Q3] | 37.0 [26.0, 61.0] | 40.0 [26.3, 61.5] | 37.0 [23.0, 59.0] |
| Median [Min, Max] | 37.0 [3.00, 134] | 40.0 [3.00, 134] | 37.0 [3.00, 134] |

| Blood WCC | | | |
|-------------------------------------|-------------------------|-------------------------|-------------------------|
| N | 25 | 16 | 27 |
| Mean [SD] | 6.65 [1.48] | 6.96 [1.53] | 6.51 [1.52] |
| Median [Q1, Q3] | 6.20 [5.70, 7.82] | 6.20 [5.85, 8.04] | 6.00 [5.55, 7.46] |
| Median [Min, Max] | 6.20 [4.00, 10.2] | 6.20 [5.10, 10.2] | 6.00 [4.00, 10.2] |
| Blood Neutrophils | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 3.91 [1.10] | 4.01 [1.16] | 3.82 [1.11] |
| Median [Q1, Q3] | 3.70 [3.20, 4.40] | 3.75 [3.38, 4.18] | 3.70 [3.02, 4.25] |
| Median [Min, Max] | 3.70 [2.57, 6.51] | 3.75 [2.60, 6.51] | 3.70 [2.45, 6.51] |
| Blood Eosinophils | | | |
| N | 25 | 16 | 27 |
| Mean [SD] | 0.0784 [0.0657] | 0.0856 [0.0756] | 0.0763 [0.0637] |
| Median [Q1, Q3] | 0.0600 [0.0400, 0.0900] | 0.0600 [0.0400, 0.0950] | 0.0600 [0.0400, 0.0900] |
| Median [Min, Max] | 0.0600 [0.0200, 0.320] | 0.0600 [0.0200, 0.320] | 0.0600 [0.0200, 0.320] |
| Sputum Macrophages (% total) | | | |
| N | 20 | 15 | 22 |
| Mean [SD] | 24.2 [23.5] | 22.5 [21.1] | 28.8 [28.0] |
| Median [Q1, Q3] | 16.8 [8.16, 28.7] | 15.8 [9.45, 23.4] | 19.5 [9.08, 44.1] |
| Median [Min, Max] | 16.8 [1.10, 81.1] | 15.8 [1.10, 70.0] | 19.5 [1.10, 99.2] |
| Missing | 5 (20.0%) | 1 (6.3%) | 5 (18.5%) |
| Sputum Neutrophils (% total) | | | |
| N | 20 | 15 | 22 |
| Mean [SD] | 64.3 [28.6] | 68.3 [25.6] | 60.6 [30.6] |
| Median [Q1, Q3] | 72.4 [46.9, 84.4] | 75.8 [58.2, 86.1] | 69.8 [37.1, 82.6] |
| Median [Min, Max] | 72.4 [12.8, 98.8] | 75.8 [17.5, 97.0] | 69.8 [0.500, 98.8] |
| Missing | 5 (20.0%) | 1 (6.3%) | 5 (18.5%) |
| Sputum Eosinophils (% total) | | | |
| N | 20 | 15 | 22 |
| Mean [SD] | 6.90 [16.6] | 6.88 [18.6] | 6.29 [15.9] |
| Median [Q1, Q3] | 1.04 [0, 5.92] | 0.730 [0.100, 4.88] | 0.645 [0, 5.31] |
| Median [Min, Max] | 1.04 [0, 73.5] | 0.730 [0, 73.5] | 0.645 [0, 73.5] |
| Missing | 5 (20.0%) | 1 (6.3%) | 5 (18.5%) |
| Sputum Lymphocytes (% total) | | | |
| N | 20 | 15 | 22 |
| Mean [SD] | 0.370 [0.687] | 0.476 [0.767] | 0.390 [0.669] |
| Median [Q1, Q3] | 0 [0, 0.275] | 0 [0, 0.515] | 0 [0, 0.325] |
| Median [Min, Max] | 0 [0, 2.17] | 0 [0, 2.17] | 0 [0, 2.17] |
| Missing | 5 (20.0%) | 1 (6.3%) | 5 (18.5%) |

Supplementary table 3 – Proteins significantly downregulated by prednisolone versus placebo in sputum (Olink®)

| | Assay | UniProt | Log2fold change | Adjusted p value |
|----|----------|---------|-----------------|------------------|
| 1 | IL4 | P05112 | -3.37 | 4.81E-03 |
| 2 | SERPINA9 | Q86WD7 | -3.18 | 9.79E-06 |
| 3 | CCL24 | O00175 | -2.6 | 2.47E-04 |
| 4 | MMP1 | P03956 | -2.6 | 1.44E-03 |
| 5 | HLA-DRA | P01903 | -2.46 | 1.88E-04 |
| 6 | TIE1 | P35590 | -2.44 | 3.70E-04 |
| 7 | CCL26 | Q9Y258 | -2.43 | 5.02E-03 |
| 8 | CCL22 | O00626 | -2.43 | 2.79E-02 |
| 9 | CCL17 | Q92583 | -2.37 | 6.68E-03 |
| 10 | GDNF | P39905 | -2.29 | 3.96E-03 |
| 11 | NCAM1 | P13591 | -2.18 | 6.60E-04 |
| 12 | TIMD4 | Q96H15 | -2.16 | 8.56E-05 |
| 13 | CCL13 | Q99616 | -2.15 | 6.92E-03 |
| 14 | NCAN | O14594 | -2.11 | 1.99E-04 |
| 15 | MCAM | P43121 | -2.09 | 1.93E-04 |
| 16 | SELE | P16581 | -2.07 | 3.74E-04 |
| 17 | DOK2 | O60496 | -2.07 | 4.36E-03 |
| 18 | IL5 | P05113 | -2.06 | 2.89E-04 |
| 19 | GP1BA | P07359 | -2.04 | 2.65E-04 |
| 20 | VCAM1 | P19320 | -2.03 | 1.40E-03 |
| 21 | CPA2 | P48052 | -2 | 1.47E-02 |
| 22 | ARSB | P15848 | -1.97 | 2.63E-03 |
| 23 | MMP12 | P39900 | -1.97 | 6.09E-03 |
| 24 | TNFRSF8 | P28908 | -1.95 | 5.57E-05 |
| 25 | SPP1 | P10451 | -1.94 | 8.87E-04 |
| 26 | FCER2 | P06734 | -1.9 | 5.89E-04 |
| 27 | OGN | P20774 | -1.88 | 9.49E-05 |
| 28 | THBS4 | P35443 | -1.88 | 1.04E-03 |
| 29 | LEP | P41159 | -1.87 | 7.13E-04 |
| 30 | FABP1 | P07148 | -1.83 | 1.82E-02 |
| 31 | CPA1 | P15085 | -1.82 | 5.21E-03 |
| 32 | TNXB | P22105 | -1.78 | 6.98E-04 |
| 33 | LAP3 | P28838 | -1.74 | 4.13E-02 |
| 34 | TCL1A | P56279 | -1.73 | 4.80E-03 |
| 35 | KRT18 | P05783 | -1.72 | 5.54E-03 |
| 36 | TREM2 | Q9NZC2 | -1.7 | 4.96E-03 |
| 37 | UMOD | P07911 | -1.69 | 1.70E-03 |
| 38 | ANGPTL3 | Q9Y5C1 | -1.69 | 3.23E-03 |
| 39 | CD69 | Q07108 | -1.67 | 6.28E-03 |
| 40 | STC2 | O76061 | -1.65 | 3.08E-03 |

| | | | | |
|----|---------|--------|-------|----------|
| 41 | MB | P02144 | -1.63 | 2.74E-02 |
| 42 | HAVCR2 | Q8TDQ0 | -1.61 | 1.13E-02 |
| 43 | APEX1 | P27695 | -1.6 | 1.94E-02 |
| 44 | LILRB5 | O75023 | -1.6 | 2.03E-03 |
| 45 | DLK1 | P80370 | -1.6 | 2.58E-04 |
| 46 | FOLR3 | P41439 | -1.6 | 6.44E-03 |
| 47 | CR2 | P20023 | -1.6 | 7.73E-03 |
| 48 | CCL23 | P55773 | -1.59 | 1.33E-02 |
| 49 | FBP1 | P09467 | -1.57 | 2.12E-02 |
| 50 | VSTM2L | Q96N03 | -1.57 | 1.98E-02 |
| 51 | ENO1 | P06733 | -1.56 | 2.98E-04 |
| 52 | ICAM2 | P13598 | -1.56 | 5.49E-03 |
| 53 | ITIH3 | Q06033 | -1.56 | 3.19E-02 |
| 54 | AARSD1 | Q9BTE6 | -1.55 | 5.36E-03 |
| 55 | ICAM5 | Q9UMF0 | -1.52 | 2.12E-02 |
| 56 | FETUB | Q9UGM5 | -1.5 | 4.42E-02 |
| 57 | LAIR2 | Q6ISS4 | -1.49 | 4.46E-05 |
| 58 | TNFRSF4 | P43489 | -1.47 | 5.71E-04 |
| 59 | GP6 | Q9HCN6 | -1.47 | 2.40E-03 |
| 60 | VWF | P04275 | -1.45 | 4.92E-02 |
| 61 | PLPBP | O94903 | -1.45 | 4.60E-02 |
| 62 | CD209 | Q9NNX6 | -1.44 | 3.85E-02 |
| 63 | PROC | P04070 | -1.44 | 4.28E-02 |
| 64 | APOM | O95445 | -1.42 | 1.11E-02 |
| 65 | CASP3 | P42574 | -1.41 | 2.81E-02 |
| 66 | MEPE | Q9NQ76 | -1.4 | 8.37E-04 |
| 67 | AOC3 | Q16853 | -1.4 | 1.38E-02 |
| 68 | SIGLEC6 | O43699 | -1.37 | 1.18E-02 |
| 69 | SSC5D | A1L4H1 | -1.36 | 3.58E-04 |
| 70 | GPA33 | Q99795 | -1.36 | 4.58E-02 |
| 71 | COL6A3 | P12111 | -1.36 | 6.65E-03 |
| 72 | IL2RA | P01589 | -1.35 | 1.07E-02 |
| 73 | CDH5 | P33151 | -1.33 | 8.04E-03 |
| 74 | CA3 | P07451 | -1.33 | 3.76E-02 |
| 75 | IL1RL1 | Q01638 | -1.32 | 1.52E-02 |
| 76 | JAM2 | P57087 | -1.31 | 3.38E-03 |
| 77 | CXCL9 | Q07325 | -1.31 | 4.17E-02 |
| 78 | CRTAC1 | Q9NQ79 | -1.3 | 2.57E-02 |
| 79 | FOLR2 | P14207 | -1.3 | 6.18E-03 |
| 80 | IL20RA | Q9UHF4 | -1.29 | 1.13E-03 |
| 81 | PPP1R2 | P41236 | -1.29 | 1.23E-02 |
| 82 | PDGFRB | P09619 | -1.29 | 7.97E-04 |
| 83 | COMP | P49747 | -1.28 | 1.22E-02 |
| 84 | GUCA2A | Q02747 | -1.28 | 2.89E-02 |

| | | | | |
|-----|---------|--------|-------|----------|
| 85 | SCARF1 | Q14162 | -1.27 | 7.25E-03 |
| 86 | CTRC | Q99895 | -1.24 | 3.35E-02 |
| 87 | IL18BP | O95998 | -1.23 | 3.11E-02 |
| 88 | C2 | P06681 | -1.21 | 4.72E-02 |
| 89 | CPB1 | P15086 | -1.21 | 4.98E-03 |
| 90 | LILRB1 | Q8NHL6 | -1.18 | 2.30E-02 |
| 91 | MARCO | Q9UEW3 | -1.18 | 2.23E-02 |
| 92 | FST | P19883 | -1.18 | 2.13E-03 |
| 93 | CNDP1 | Q96KN2 | -1.17 | 4.20E-02 |
| 94 | CD70 | P32970 | -1.14 | 6.65E-03 |
| 95 | KLRB1 | Q12918 | -1.14 | 5.83E-03 |
| 96 | CDHR5 | Q9HBB8 | -1.14 | 1.55E-04 |
| 97 | SPON1 | Q9HCB6 | -1.13 | 3.13E-02 |
| 98 | REG3A | Q06141 | -1.13 | 3.10E-02 |
| 99 | CLEC14A | Q86T13 | -1.12 | 3.57E-02 |
| 100 | CLC | Q05315 | -1.1 | 1.65E-02 |
| 101 | FAM3C | Q92520 | -1.1 | 2.92E-03 |
| 102 | CD6 | P30203 | -1.09 | 2.84E-02 |
| 103 | WFIKKN2 | Q8TEU8 | -1.09 | 4.58E-02 |
| 104 | FCN2 | Q15485 | -1.09 | 2.97E-02 |
| 105 | IL13 | P35225 | -1.06 | 3.17E-02 |
| 106 | FABP2 | P12104 | -1.06 | 3.69E-02 |
| 107 | VASN | Q6EMK4 | -1.06 | 3.21E-02 |
| 108 | DPT | Q07507 | -1.05 | 1.01E-02 |
| 109 | IGFBP1 | P08833 | -1.04 | 2.67E-03 |
| 110 | TSHB | P01222 | -1.03 | 6.02E-03 |
| 111 | RUVBL1 | Q9Y265 | -1.02 | 1.90E-02 |
| 112 | CCL14 | Q16627 | -1.02 | 1.25E-02 |
| 113 | REG1A | P05451 | -1.01 | 1.05E-02 |
| 114 | NELL1 | Q92832 | -0.99 | 3.48E-02 |
| 115 | COL4A1 | P02462 | -0.98 | 1.74E-02 |
| 116 | PRDX5 | P30044 | -0.98 | 1.70E-03 |
| 117 | PTGDS | P41222 | -0.98 | 3.59E-03 |
| 118 | APOH | P02749 | -0.98 | 2.33E-03 |
| 119 | FABP4 | P15090 | -0.97 | 2.95E-04 |
| 120 | CD160 | O95971 | -0.97 | 4.59E-03 |
| 121 | STC1 | P52823 | -0.97 | 3.64E-02 |
| 122 | XPNPEP2 | O43895 | -0.95 | 1.37E-02 |
| 123 | ACP5 | P13686 | -0.94 | 1.69E-02 |
| 124 | OXT | P01178 | -0.93 | 2.76E-02 |
| 125 | AXL | P30530 | -0.92 | 1.72E-02 |
| 126 | CD22 | P20273 | -0.89 | 2.88E-02 |
| 127 | ENPP2 | Q13822 | -0.88 | 2.23E-02 |
| 128 | ADM | P35318 | -0.87 | 3.66E-02 |

| | | | | |
|-----|---------|--------|-------|----------|
| 129 | PVALB | P20472 | -0.86 | 2.38E-02 |
| 130 | REG1B | P48304 | -0.85 | 3.52E-02 |
| 131 | CNTN3 | Q9P232 | -0.84 | 3.41E-02 |
| 132 | PEAR1 | Q5VY43 | -0.84 | 1.21E-02 |
| 133 | SPINK1 | P00995 | -0.82 | 2.87E-02 |
| 134 | NRP1 | O14786 | -0.79 | 1.30E-02 |
| 135 | FASLG | P48023 | -0.79 | 4.23E-02 |
| 136 | CD300LF | Q8TDQ1 | -0.79 | 4.28E-02 |
| 137 | CFC1 | P0CG37 | -0.77 | 8.10E-03 |
| 138 | CD27 | P26842 | -0.76 | 2.68E-02 |
| 139 | VMO1 | Q7Z5L0 | -0.73 | 4.61E-02 |
| 140 | COL18A1 | P39060 | -0.72 | 4.52E-02 |
| 141 | CNST | Q6PJW8 | -0.71 | 1.46E-02 |
| 142 | IL12RB1 | P42701 | -0.7 | 9.54E-03 |
| 143 | CD79B | P40259 | -0.69 | 3.46E-02 |
| 144 | SCG2 | P13521 | -0.69 | 2.83E-02 |
| 145 | CDH17 | Q12864 | -0.66 | 6.90E-03 |
| 146 | PCDH1 | Q08174 | -0.65 | 6.29E-06 |
| 147 | ICAM1 | P05362 | -0.65 | 2.91E-02 |
| 148 | TNFSF11 | O14788 | -0.63 | 8.27E-03 |
| 149 | CD1C | P29017 | -0.62 | 1.21E-03 |
| 150 | CTSS | P25774 | -0.6 | 4.85E-02 |
| 151 | SFTPA2 | Q8IWL1 | -0.6 | 1.23E-02 |
| 152 | CBLN4 | Q9NTU7 | -0.59 | 5.34E-03 |
| 153 | VTCN1 | Q7Z7D3 | -0.58 | 3.64E-02 |
| 154 | LGALS3 | P17931 | -0.57 | 3.21E-02 |
| 155 | CD244 | Q9BZW8 | -0.56 | 2.00E-02 |
| 156 | MIF | P14174 | -0.54 | 3.59E-02 |
| 157 | FLT4 | P35916 | -0.52 | 1.30E-02 |
| 158 | CTSL | P07711 | -0.52 | 4.39E-02 |

Supplementary table 4 – Proteins significantly upregulated by prednisolone versus placebo in sputum (Olink®)

| | Assay | UniProt | Log2fold change | Adjusted p value |
|----|---------------|---------|-----------------|------------------|
| 1 | CES2 | O00748 | 2.35 | 2.65E-04 |
| 2 | EREG | O14944 | 1.98 | 4.81E-04 |
| 3 | DEFB4A_DEFB4B | O15263 | 1.73 | 2.56E-02 |
| 4 | DKK4 | Q9UBT3 | 1.65 | 1.41E-02 |
| 5 | GP2 | P55259 | 1.63 | 3.95E-06 |
| 6 | CALB1 | P05937 | 1.5 | 2.40E-02 |
| 7 | TIMP3 | P35625 | 1.44 | 1.51E-02 |
| 8 | FKBP5 | Q13451 | 1.43 | 3.29E-02 |
| 9 | XG | P55808 | 1.4 | 7.96E-03 |
| 10 | GFRA1 | P56159 | 1.4 | 2.40E-03 |

| | | | | |
|----|----------|--------|------|----------|
| 11 | NPPB | P16860 | 1.4 | 3.89E-02 |
| 12 | CRNN | Q9UBG3 | 1.39 | 4.73E-03 |
| 13 | PLAT | P00750 | 1.38 | 1.84E-02 |
| 14 | KLK8 | O60259 | 1.38 | 1.75E-04 |
| 15 | PLTP | P55058 | 1.35 | 2.48E-02 |
| 16 | SLC27A4 | Q6P1M0 | 1.33 | 1.46E-02 |
| 17 | IL1B | P01584 | 1.32 | 2.59E-02 |
| 18 | IL17C | Q9P0M4 | 1.28 | 3.11E-02 |
| 19 | EPHA2 | P29317 | 1.2 | 1.88E-02 |
| 20 | NDRG1 | Q92597 | 1.18 | 3.52E-03 |
| 21 | ARID4B | Q4LE39 | 1.18 | 4.88E-02 |
| 22 | DSG3 | P32926 | 1.16 | 4.00E-05 |
| 23 | PRSS27 | Q9BQR3 | 1.15 | 1.75E-02 |
| 24 | ANXA5 | P08758 | 1.13 | 2.24E-03 |
| 25 | CNTN5 | O94779 | 1.12 | 5.36E-03 |
| 26 | SNCG | O76070 | 1.11 | 2.89E-02 |
| 27 | ULBP2 | Q9BZM5 | 1.1 | 1.40E-02 |
| 28 | WFDC12 | Q8WWY7 | 1.08 | 5.30E-03 |
| 29 | CEACAM21 | Q3KPI0 | 1.05 | 3.74E-02 |
| 30 | FCAR | P24071 | 1.03 | 4.54E-02 |
| 31 | DSC2 | Q02487 | 1.02 | 1.88E-03 |
| 32 | SPINK5 | Q9NQ38 | 0.99 | 1.37E-03 |
| 33 | ADGRG2 | Q8IZP9 | 0.97 | 4.27E-03 |
| 34 | F2R | P25116 | 0.95 | 2.71E-04 |
| 35 | CA12 | O43570 | 0.92 | 4.93E-02 |
| 36 | KLK6 | Q92876 | 0.92 | 4.97E-04 |
| 37 | IVD | P26440 | 0.89 | 1.82E-04 |
| 38 | CDSN | Q15517 | 0.87 | 3.62E-03 |
| 39 | LDLR | P01130 | 0.83 | 3.69E-02 |
| 40 | DNER | Q8NFT8 | 0.81 | 7.91E-03 |
| 41 | PSMA1 | P25786 | 0.78 | 1.73E-02 |
| 42 | ESM1 | Q9NQ30 | 0.78 | 5.11E-03 |
| 43 | CHMP1A | Q9HD42 | 0.73 | 3.95E-03 |
| 44 | HS3ST3B1 | Q9Y662 | 0.72 | 6.73E-03 |
| 45 | IL22RA1 | Q8N6P7 | 0.67 | 8.90E-03 |
| 46 | DCN | P07585 | 0.67 | 2.24E-02 |
| 47 | PODXL2 | Q9NZ53 | 0.6 | 2.95E-02 |
| 48 | ACVRL1 | P37023 | 0.59 | 3.96E-02 |
| 49 | CTSV | O60911 | 0.58 | 1.05E-02 |
| 50 | KCNIP4 | Q6PIL6 | 0.54 | 4.69E-02 |
| 51 | PSG1 | P11464 | 0.51 | 1.44E-02 |

Supplementary table 5 – Proteins significantly downregulated by prednisolone versus placebo in plasma (Olink®)

| | Assay | UniProt | Log2fold change | Adjusted p value |
|----|--------------|----------------|------------------------|-------------------------|
| 1 | IL4 | P05112 | -3.17 | 1.06E-05 |
| 2 | IL12A_IL12B | P29459_P29460 | -2.24 | 1.55E-26 |
| 3 | IL12B | P29460 | -2.09 | 3.88E-26 |
| 4 | CRH | P06850 | -1.46 | 3.80E-11 |
| 5 | CCL22 | O00626 | -1.42 | 4.18E-21 |
| 6 | MMP12 | P39900 | -1.42 | 5.07E-14 |
| 7 | CCL19 | Q99731 | -1.24 | 2.94E-14 |
| 8 | TNFSF13B | Q9Y275 | -0.94 | 1.03E-20 |
| 9 | THBS4 | P35443 | -0.93 | 7.59E-08 |
| 10 | FGF19 | O95750 | -0.93 | 1.63E-02 |
| 11 | GGH | Q92820 | -0.91 | 9.66E-18 |
| 12 | TNFRSF8 | P28908 | -0.91 | 1.67E-16 |
| 13 | TNFRSF4 | P43489 | -0.91 | 3.98E-19 |
| 14 | CCDC80 | Q76M96 | -0.9 | 1.29E-14 |
| 15 | SIGLEC6 | O43699 | -0.86 | 2.87E-21 |
| 16 | TNFRSF9 | Q07011 | -0.85 | 9.89E-14 |
| 17 | CD83 | Q01151 | -0.84 | 4.03E-13 |
| 18 | IFNG | P01579 | -0.84 | 6.50E-03 |
| 19 | CLEC4C | Q8WTT0 | -0.81 | 5.93E-06 |
| 20 | CCL18 | P55774 | -0.81 | 2.03E-08 |
| 21 | CXCL10 | P02778 | -0.79 | 3.74E-09 |
| 22 | CCL17 | Q92583 | -0.77 | 4.81E-11 |
| 23 | PDCD1 | Q15116 | -0.77 | 2.35E-21 |
| 24 | CCL26 | Q9Y258 | -0.75 | 4.31E-06 |
| 25 | CCN3 | P48745 | -0.74 | 5.80E-10 |
| 26 | CD207 | Q9UJ71 | -0.71 | 9.97E-09 |
| 27 | OXT | P01178 | -0.69 | 1.11E-04 |
| 28 | ADAMTS8 | Q9UP79 | -0.68 | 4.01E-10 |
| 29 | CR2 | P20023 | -0.66 | 1.47E-13 |
| 30 | TPPP3 | Q9BW30 | -0.66 | 1.30E-04 |
| 31 | DLK1 | P80370 | -0.66 | 1.06E-12 |
| 32 | CD28 | P10747 | -0.66 | 1.73E-06 |
| 33 | OMG | P23515 | -0.65 | 8.41E-05 |
| 34 | OGN | P20774 | -0.65 | 4.84E-10 |
| 35 | RET | P07949 | -0.65 | 2.20E-13 |
| 36 | IL6 | P05231 | -0.65 | 2.51E-03 |
| 37 | CDON | Q4KMG0 | -0.64 | 7.29E-12 |
| 38 | PAEP | P09466 | -0.64 | 3.66E-06 |
| 39 | FOLR2 | P14207 | -0.64 | 3.48E-16 |
| 40 | SMAD5 | Q99717 | -0.64 | 5.05E-10 |
| 41 | CCL21 | O00585 | -0.64 | 9.54E-10 |

| | | | | |
|----|-----------|--------|-------|----------|
| 42 | CD70 | P32970 | -0.64 | 1.23E-10 |
| 43 | CD300E | Q496F6 | -0.63 | 6.27E-09 |
| 44 | ITIH3 | Q06033 | -0.63 | 2.09E-08 |
| 45 | CD1C | P29017 | -0.62 | 2.26E-18 |
| 46 | LAMP3 | Q9UQV4 | -0.62 | 6.56E-11 |
| 47 | MARCO | Q9UEW3 | -0.62 | 3.73E-18 |
| 48 | DRAXIN | Q8NBI3 | -0.62 | 5.19E-10 |
| 49 | CD200R1 | Q8TD46 | -0.62 | 5.60E-14 |
| 50 | MMP10 | P09238 | -0.61 | 8.08E-06 |
| 51 | FASLG | P48023 | -0.6 | 1.20E-16 |
| 52 | GSTA3 | Q16772 | -0.6 | 4.09E-03 |
| 53 | CXCL17 | Q6UXB2 | -0.59 | 1.16E-09 |
| 54 | CDCP1 | Q9H5V8 | -0.59 | 5.35E-09 |
| 55 | CD22 | P20273 | -0.59 | 3.33E-17 |
| 56 | GZMB | P10144 | -0.58 | 2.44E-04 |
| 57 | MSR1 | P21757 | -0.57 | 2.37E-11 |
| 58 | SERPINA9 | Q86WD7 | -0.56 | 1.39E-04 |
| 59 | CNTN2 | Q02246 | -0.55 | 3.55E-11 |
| 60 | LAG3 | P18627 | -0.54 | 1.43E-10 |
| 61 | IFNL1 | Q8IU54 | -0.54 | 8.64E-05 |
| 62 | KLK14 | Q9P0G3 | -0.54 | 1.79E-09 |
| 63 | LTA | P01374 | -0.54 | 2.19E-10 |
| 64 | FCRL1 | Q96LA6 | -0.53 | 2.29E-14 |
| 65 | CD79B | P40259 | -0.53 | 3.69E-15 |
| 66 | TNFRSF13B | O14836 | -0.52 | 1.20E-14 |
| 67 | ADAM23 | O75077 | -0.52 | 1.43E-08 |
| 68 | CXCL11 | O14625 | -0.51 | 2.10E-05 |
| 69 | DPT | Q07507 | -0.51 | 1.86E-10 |
| 70 | LRRC25 | Q8N386 | -0.51 | 8.36E-09 |
| 71 | COL1A1 | P02452 | -0.51 | 1.98E-07 |
| 72 | COL6A3 | P12111 | -0.51 | 1.39E-08 |
| 73 | VWC2 | Q2TAL6 | -0.5 | 1.71E-07 |
| 74 | MDGA1 | Q8NFP4 | -0.5 | 6.67E-13 |

Supplementary table 6 – Proteins significantly upregulated by prednisolone versus placebo in plasma (Olink®)

| | Assay | UniProt | Log2fold change | Adjusted p value |
|---|--------|---------|-----------------|------------------|
| 1 | MMP3 | P08254 | 2.29 | 1.21E-28 |
| 2 | ZBTB16 | Q05516 | 1.88 | 5.02E-14 |
| 3 | IL10 | P22301 | 1.63 | 1.98E-06 |
| 4 | GCG | P01275 | 1.41 | 3.93E-06 |
| 5 | PFKFB2 | O60825 | 1.41 | 7.54E-06 |
| 6 | AREG | P15514 | 1.37 | 8.78E-12 |
| 7 | FKBP5 | Q13451 | 1.3 | 1.72E-03 |

| | | | | |
|----|-----------|--------|------|----------|
| 8 | GP2 | P55259 | 1.16 | 3.41E-09 |
| 9 | MATN3 | O15232 | 1.03 | 9.59E-15 |
| 10 | TMPRSS15 | P98073 | 0.98 | 2.01E-04 |
| 11 | TNFAIP8 | O95379 | 0.9 | 4.60E-02 |
| 12 | SERPINA12 | Q8IW75 | 0.89 | 5.98E-09 |
| 13 | AGR3 | Q8TD06 | 0.87 | 3.16E-03 |
| 14 | OSM | P13725 | 0.86 | 8.28E-05 |
| 15 | TGFA | P01135 | 0.85 | 1.10E-05 |
| 16 | STAT5B | P51692 | 0.82 | 9.60E-04 |
| 17 | VSIG4 | Q9Y279 | 0.82 | 6.29E-13 |
| 18 | IL1RL1 | Q01638 | 0.81 | 7.26E-13 |
| 19 | CD177 | Q8N6Q3 | 0.79 | 9.72E-09 |
| 20 | CST7 | O76096 | 0.76 | 7.01E-08 |
| 21 | RBP2 | P50120 | 0.75 | 5.11E-04 |
| 22 | NAMPT | P43490 | 0.74 | 3.06E-03 |
| 23 | MFGE8 | Q08431 | 0.73 | 2.38E-10 |
| 24 | PADI2 | Q9Y2J8 | 0.68 | 8.22E-06 |
| 25 | IL17C | Q9P0M4 | 0.68 | 6.68E-03 |
| 26 | SLC39A5 | Q6ZMH5 | 0.67 | 7.27E-11 |
| 27 | CEACAM8 | P31997 | 0.65 | 4.45E-04 |
| 28 | CLPS | P04118 | 0.64 | 1.69E-06 |
| 29 | PIK3AP1 | Q6ZUJ8 | 0.63 | 6.57E-03 |
| 30 | IPCEF1 | Q8WWN9 | 0.63 | 2.71E-02 |
| 31 | MMP8 | P22894 | 0.63 | 8.89E-04 |
| 32 | SCP2 | P22307 | 0.61 | 1.17E-03 |
| 33 | PADI4 | Q9UM07 | 0.61 | 1.09E-02 |
| 34 | PROK1 | P58294 | 0.6 | 8.48E-05 |
| 35 | RGMA | Q96B86 | 0.6 | 8.78E-14 |
| 36 | CES1 | P23141 | 0.6 | 1.43E-04 |
| 37 | PM20D1 | Q6GTS8 | 0.59 | 4.24E-06 |
| 38 | HCLS1 | P14317 | 0.58 | 4.08E-03 |
| 39 | MME | P08473 | 0.58 | 1.34E-07 |
| 40 | MMP9 | P14780 | 0.57 | 1.44E-05 |
| 41 | ARG1 | P05089 | 0.55 | 3.25E-03 |
| 42 | CCL20 | P78556 | 0.55 | 4.47E-02 |
| 43 | CCL15 | Q16663 | 0.55 | 2.16E-07 |
| 44 | RASSF2 | P50749 | 0.55 | 3.90E-02 |
| 45 | LEP | P41159 | 0.54 | 4.29E-04 |
| 46 | CEP43 | O95684 | 0.54 | 3.13E-03 |
| 47 | REN | P00797 | 0.53 | 9.54E-07 |
| 48 | MVK | Q03426 | 0.53 | 2.45E-03 |
| 49 | PROC | P04070 | 0.53 | 3.80E-10 |
| 50 | YTHDF3 | Q7Z739 | 0.52 | 8.40E-03 |
| 51 | FEN1 | P39748 | 0.51 | 2.17E-02 |

| | | | | |
|----|--------|--------|------|----------|
| 52 | BAIAP2 | Q9UQB8 | 0.51 | 1.57E-03 |
| 53 | DNAJB1 | P25685 | 0.5 | 1.15E-02 |

Supplementary table 7 – The effect of prednisolone versus placebo on sputum proteins (ELISA)

| Assay | Change in protein concentration | Log2fold change | Adjusted p value |
|----------------------|---------------------------------|-----------------|------------------|
| IL17/25 (pg/ml) | -1.63 | -0.40 | 0.41 |
| IL1 α (pg/ml) | 0.658 | 0.014 | 0.98 |
| IL1 β (pg/ml) | -24.2 | -0.18 | 0.59 |
| IL2Ra (pg/ml) | -12.5 | -0.24 | 0.57 |
| IL3 (pg/ml) | -1.4 | -0.056 | 0.89 |
| IL33 (pg/ml) | -1.61 | -0.25 | 0.45 |
| IL6 (pg/ml) | 3.15 | 0.23 | 0.61 |
| IL6R (pg/ml) | 43.6 | 0.18 | 0.67 |
| MPO (ng/ml) | 833 | 0.89 | 0.27 |
| YKL40 (ng/ml) | -19.6 | -0.57 | 0.52 |
| IL1Ra (ng/ml) | 16.9 | 1.00 | 0.045 |
| GMCSF (pg/ml) | 0.022 | 0.46 | 0.45 |
| TSLP (pg/ml) | 1.25 | 1.30 | 0.37 |
| EDN (ng/ml) | -1181 | -1.04 | 0.0001 |
| IL1R1 (pg/ml) | 222 | 0.51 | 0.125 |

Supplementary table 8 - The effect of prednisolone versus placebo on nasal tissue gene transcription

| | Log2fold change | Adjusted p value | Gene |
|----|------------------------|-------------------------|-------------|
| 1 | -2.93 | 7.42E-02 | CXCL5 |
| 2 | -2.88 | 9.25E-02 | SRARP |
| 3 | -2.69 | 5.47E-02 | TNFAIP6 |
| 4 | -2.62 | 8.88E-02 | GNAO1 |
| 5 | -2.55 | 8.43E-02 | IGLC2 |
| 6 | -2.41 | 5.37E-02 | MMP12 |
| 7 | -2.38 | 2.84E-02 | ADCYAP1 |
| 8 | -2.03 | 2.84E-02 | CXCL3 |
| 9 | -1.96 | 2.84E-02 | HOXB2 |
| 10 | -1.80 | 2.84E-02 | TPSB2 |
| 11 | -1.75 | 5.47E-02 | CXCL2 |
| 12 | -1.73 | 2.84E-02 | LGI1 |
| 13 | -1.73 | 9.97E-02 | AC016868.1 |
| 14 | -1.71 | 2.84E-02 | FCGBP |
| 15 | -1.67 | 9.27E-02 | CD48 |
| 16 | -1.64 | 3.32E-02 | ADRA2A |
| 17 | -1.54 | 2.84E-02 | B3GNT6 |
| 18 | -1.45 | 2.84E-02 | PPP1R16B |
| 19 | -1.43 | 3.30E-02 | CX3CR1 |
| 20 | -1.38 | 5.04E-02 | DUSP2 |
| 21 | -1.37 | 2.84E-02 | GSDMD |
| 22 | -1.33 | 2.84E-02 | C3 |
| 23 | -1.21 | 9.37E-02 | BPIFA1 |
| 24 | -1.12 | 9.37E-02 | CYR61 |
| 25 | -0.91 | 7.29E-02 | FER1L6 |
| 26 | -0.91 | 5.47E-02 | HIPK2 |
| 27 | -0.90 | 2.84E-02 | ALOX15 |
| 28 | -0.83 | 5.37E-02 | HDAC9 |
| 29 | 0.71 | 2.84E-02 | RBP1 |
| 30 | 0.71 | 9.37E-02 | ALOX15B |
| 31 | 1.44 | 2.84E-02 | PANK4 |

Supplementary table 9 – Precursor miRNA significantly differentially expressed by prednisolone versus placebo

| | Pre-cursor miRNA | Difference in | Adjusted p value |
|--|-------------------------|----------------------|-------------------------|
|--|-------------------------|----------------------|-------------------------|

| | | normalised gene count | |
|----|-----------------|------------------------------|---------|
| 1 | hsa-mir-6511a-3 | -58.8 | 0.00252 |
| 2 | hsa-mir-6511a-2 | -58.8 | 0.00252 |
| 3 | hsa-mir-6511a-1 | -58.8 | 0.00252 |
| 4 | hsa-mir-6511a-4 | -58.8 | 0.00252 |
| 5 | hsa-mir-516a-1 | 111 | 0.00592 |
| 6 | hsa-mir-516a-2 | 111 | 0.00592 |
| 7 | hsa-mir-5701-2 | -77.5 | 0.0114 |
| 8 | hsa-mir-5701-3 | -77.5 | 0.0114 |
| 9 | hsa-mir-5701-1 | -77.5 | 0.0114 |
| 10 | hsa-mir-518a-2 | -84.4 | 0.0405 |
| 11 | hsa-mir-518a-1 | -84.4 | 0.0405 |
| 12 | hsa-mir-3670-2 | 24.0 | 0.0448 |
| 13 | hsa-mir-3670-4 | 24.0 | 0.0448 |
| 14 | hsa-mir-3670-3 | 24.0 | 0.0448 |
| 15 | hsa-mir-3670-1 | 24.0 | 0.0448 |
| 16 | hsa-mir-4520-1 | 44.9 | 0.0687 |
| 17 | hsa-mir-4520-2 | 39.9 | 0.0687 |
| 18 | hsa-mir-4650-2 | -30.9 | 0.0930 |
| 19 | hsa-mir-4650-1 | -30.9 | 0.0930 |

Supplementary table 10 – Nasal tissue genes significantly downregulated by mepolizumab

| | Log2fold change | Adjusted p value | Gene |
|----|------------------------|-------------------------|-------------|
| 1 | -1.38 | 3.05E-03 | CROCC2 |
| 2 | -1.05 | 3.05E-03 | KCNH3 |
| 3 | -1.17 | 3.05E-03 | HMCN2 |
| 4 | -1.17 | 7.35E-03 | BEST4 |
| 5 | -0.84 | 7.90E-03 | KNDC1 |
| 6 | -1.03 | 9.41E-03 | HAGHL |
| 7 | -0.85 | 1.32E-02 | GRIN3B |
| 8 | -0.87 | 1.36E-02 | BAIAP3 |
| 9 | -1.04 | 1.41E-02 | CFAP65 |
| 10 | -0.94 | 1.56E-02 | WDR97 |
| 11 | -0.82 | 1.56E-02 | RHPN1 |
| 12 | -0.65 | 1.58E-02 | MUC16 |
| 13 | -1.54 | 1.58E-02 | MAPK15 |
| 14 | -0.90 | 1.58E-02 | RSPH14 |
| 15 | -1.70 | 1.60E-02 | LKAAEAR1 |
| 16 | -1.28 | 1.60E-02 | TGFBR3L |
| 17 | -0.97 | 1.87E-02 | TTL10 |
| 18 | -0.83 | 1.87E-02 | CD164L2 |
| 19 | -1.11 | 1.87E-02 | CCDC114 |

| | | | |
|----|-------|----------|-----------|
| 20 | -0.99 | 1.87E-02 | SPEF1 |
| 21 | -1.00 | 1.87E-02 | CDHR4 |
| 22 | -0.98 | 1.88E-02 | DLEC1 |
| 23 | -0.97 | 1.88E-02 | DNAI1 |
| 24 | -0.97 | 1.90E-02 | DNAI2 |
| 25 | -0.64 | 1.96E-02 | SCGB2A1 |
| 26 | -1.55 | 1.97E-02 | RNF31 |
| 27 | -1.01 | 1.97E-02 | ZFH2 |
| 28 | -0.99 | 2.01E-02 | LRR46 |
| 29 | -1.11 | 2.04E-02 | PPP1R32 |
| 30 | -0.92 | 2.09E-02 | TEKT2 |
| 31 | -1.01 | 2.11E-02 | STK33 |
| 32 | -0.87 | 2.11E-02 | CCDC153 |
| 33 | -0.82 | 2.11E-02 | NAT14 |
| 34 | -1.42 | 2.11E-02 | FAM95C |
| 35 | -0.90 | 2.23E-02 | RFX2 |
| 36 | -1.54 | 2.35E-02 | FAM238C |
| 37 | -0.91 | 2.35E-02 | UBXN11 |
| 38 | -1.13 | 2.38E-02 | JPH3 |
| 39 | -0.73 | 2.54E-02 | WDR90 |
| 40 | -0.89 | 2.54E-02 | DENND6B |
| 41 | -1.01 | 2.54E-02 | CATIP |
| 42 | -0.83 | 2.55E-02 | ZNF473 |
| 43 | -1.15 | 2.55E-02 | LRR47B |
| 44 | -0.96 | 2.55E-02 | LRR23 |
| 45 | -0.91 | 2.60E-02 | MAP6 |
| 46 | -0.72 | 2.60E-02 | HSD11B1L |
| 47 | -0.63 | 2.60E-02 | WDR34 |
| 48 | -0.94 | 2.77E-02 | RNFT2 |
| 49 | -1.03 | 2.82E-02 | CCDC151 |
| 50 | -0.73 | 2.91E-02 | DNER |
| 51 | -0.81 | 3.01E-02 | ODF3B |
| 52 | -1.32 | 3.04E-02 | TNNI3 |
| 53 | -0.90 | 3.04E-02 | FAM92B |
| 54 | -1.27 | 3.11E-02 | SERPINI2 |
| 55 | -0.97 | 3.11E-02 | TCTEX1D4 |
| 56 | -0.95 | 3.11E-02 | LRR71 |
| 57 | -0.89 | 3.12E-02 | RSPH9 |
| 58 | -0.89 | 3.12E-02 | CCDC78 |
| 59 | -0.95 | 3.27E-02 | FNDC11 |
| 60 | -1.19 | 3.27E-02 | PCSK1N |
| 61 | -0.61 | 3.27E-02 | PPP1R16A |
| 62 | -0.86 | 3.28E-02 | MORN1 |
| 63 | -1.15 | 3.28E-02 | HHATL-AS1 |

| | | | |
|-----|-------|----------|------------|
| 64 | -1.30 | 3.28E-02 | TLL6 |
| 65 | -0.99 | 3.33E-02 | DNAAF1 |
| 66 | -0.74 | 3.33E-02 | PRUNE2 |
| 67 | -0.91 | 3.33E-02 | C21orf58 |
| 68 | -0.98 | 3.33E-02 | C2orf70 |
| 69 | -1.22 | 3.33E-02 | AC004836.1 |
| 70 | -1.03 | 3.33E-02 | TMC4 |
| 71 | -0.71 | 3.33E-02 | ANKMY1 |
| 72 | -0.94 | 3.33E-02 | PCSK4 |
| 73 | -1.22 | 3.33E-02 | SYT5 |
| 74 | -0.54 | 3.36E-02 | SPPL2B |
| 75 | -0.94 | 3.38E-02 | CCDC13 |
| 76 | -0.91 | 3.41E-02 | LINC02166 |
| 77 | -0.68 | 3.42E-02 | SSBP4 |
| 78 | -0.86 | 3.42E-02 | TTC21A |
| 79 | -1.65 | 3.42E-02 | DOC2A |
| 80 | -0.87 | 3.42E-02 | ZMYND10 |
| 81 | -0.97 | 3.45E-02 | CCDC162P |
| 82 | -1.38 | 3.48E-02 | MUC12 |
| 83 | -0.91 | 3.50E-02 | DNAAF3 |
| 84 | -1.11 | 3.50E-02 | SSUH2 |
| 85 | -1.02 | 3.50E-02 | DZIP1L |
| 86 | -2.24 | 3.50E-02 | U62317.1 |
| 87 | -0.75 | 3.50E-02 | ENKD1 |
| 88 | -1.02 | 3.53E-02 | TMEM52 |
| 89 | -1.20 | 3.56E-02 | SQSTM1 |
| 90 | -0.91 | 3.57E-02 | KIF19 |
| 91 | -0.89 | 3.57E-02 | FAM183A |
| 92 | -1.01 | 3.57E-02 | SPAG8 |
| 93 | -0.96 | 3.57E-02 | DRC1 |
| 94 | -0.84 | 3.70E-02 | GOLGA2P5 |
| 95 | -1.04 | 3.70E-02 | AC027281.1 |
| 96 | -1.01 | 3.72E-02 | CRIP1 |
| 97 | -0.82 | 3.76E-02 | CCDC17 |
| 98 | -0.84 | 3.76E-02 | LRRC10B |
| 99 | -0.69 | 3.76E-02 | GAS8 |
| 100 | -0.92 | 3.76E-02 | CFAP73 |
| 101 | -0.81 | 3.76E-02 | OSCP1 |
| 102 | -0.89 | 3.76E-02 | TBC1D9B |
| 103 | -0.61 | 3.76E-02 | ZNF688 |
| 104 | -1.01 | 3.76E-02 | AC010754.1 |
| 105 | -0.85 | 3.98E-02 | DRC3 |
| 106 | -0.89 | 4.15E-02 | GPR162 |
| 107 | -0.59 | 4.18E-02 | ZNHIT2 |

| | | | |
|-----|-------|----------|------------|
| 108 | -1.00 | 4.19E-02 | CFAP74 |
| 109 | -0.87 | 4.23E-02 | WDR54 |
| 110 | -0.88 | 4.31E-02 | TTC25 |
| 111 | -0.70 | 4.36E-02 | LRRRC61 |
| 112 | -0.91 | 4.40E-02 | C2orf40 |
| 113 | -0.79 | 4.48E-02 | GIPR |
| 114 | -0.88 | 4.56E-02 | C1orf189 |
| 115 | -1.11 | 4.69E-02 | FABP6 |
| 116 | -0.76 | 4.69E-02 | C1orf87 |
| 117 | -0.69 | 4.69E-02 | MAPK8IP1 |
| 118 | -0.91 | 4.69E-02 | CFAP99 |
| 119 | -0.85 | 4.69E-02 | CFAP100 |
| 120 | -0.73 | 4.72E-02 | PSENN |
| 121 | -0.80 | 4.72E-02 | PRR7 |
| 122 | -0.74 | 4.72E-02 | TPPP3 |
| 123 | -0.68 | 4.72E-02 | GALNS |
| 124 | -0.89 | 4.72E-02 | EFCAB12 |
| 125 | -0.64 | 4.72E-02 | WHRN |
| 126 | -0.69 | 4.72E-02 | UCN3 |
| 127 | -0.62 | 4.80E-02 | MZF1 |
| 128 | -0.95 | 4.86E-02 | HHLA2 |
| 129 | -0.98 | 4.86E-02 | NPR2 |
| 130 | -0.89 | 4.87E-02 | CFAP157 |
| 131 | -0.87 | 4.94E-02 | LRRRC73 |
| 132 | -1.13 | 4.95E-02 | CCDC180 |
| 133 | -0.94 | 4.95E-02 | TEKT4 |
| 134 | -0.97 | 4.95E-02 | LRRRC43 |
| 135 | -1.35 | 5.02E-02 | CLDN9 |
| 136 | -0.83 | 5.10E-02 | CTXN1 |
| 137 | -0.64 | 5.10E-02 | CLDN3 |
| 138 | -0.94 | 5.12E-02 | GAS2L2 |
| 139 | -0.81 | 5.13E-02 | MORN5 |
| 140 | -0.99 | 5.13E-02 | TTC16 |
| 141 | -2.13 | 5.13E-02 | KNCN |
| 142 | -0.81 | 5.13E-02 | EFCAB1 |
| 143 | -1.42 | 5.13E-02 | AL357093.1 |
| 144 | -1.39 | 5.19E-02 | AQP10 |
| 145 | -0.94 | 5.32E-02 | C4orf47 |
| 146 | -0.79 | 5.37E-02 | LDLRAD1 |
| 147 | -0.87 | 5.43E-02 | CFAP46 |
| 148 | -0.89 | 5.51E-02 | WDR38 |
| 149 | -0.69 | 5.60E-02 | MAPRE3 |
| 150 | -0.78 | 5.80E-02 | CYP2F1 |
| 151 | -0.79 | 5.80E-02 | DNAH2 |

| | | | |
|-----|-------|----------|------------|
| 152 | -0.97 | 5.86E-02 | Mar-10 |
| 153 | -1.04 | 5.86E-02 | SLC7A2 |
| 154 | -0.84 | 5.86E-02 | C1orf228 |
| 155 | -0.87 | 5.86E-02 | CDHR3 |
| 156 | -0.87 | 5.86E-02 | DNAH6 |
| 157 | -0.99 | 5.86E-02 | AL121899.1 |
| 158 | -1.47 | 5.89E-02 | VWA3A |
| 159 | -0.77 | 5.89E-02 | AK8 |
| 160 | -0.81 | 5.91E-02 | CROCC |
| 161 | -0.83 | 5.93E-02 | RSPH1 |
| 162 | -0.85 | 5.98E-02 | CFAP45 |
| 163 | -0.84 | 5.98E-02 | MAP1A |
| 164 | -0.63 | 5.98E-02 | CCDC24 |
| 165 | -0.85 | 5.98E-02 | C20orf85 |
| 166 | -0.83 | 6.04E-02 | AL031283.1 |
| 167 | -0.75 | 6.06E-02 | CCDC96 |
| 168 | -0.83 | 6.06E-02 | IQCD |
| 169 | -1.09 | 6.19E-02 | AC073439.1 |
| 170 | -0.86 | 6.22E-02 | RAB36 |
| 171 | -0.75 | 6.25E-02 | SRGAP3-AS2 |
| 172 | -0.77 | 6.28E-02 | C9orf24 |
| 173 | -1.15 | 6.40E-02 | AC104809.1 |
| 174 | -0.83 | 6.40E-02 | IQCA1 |
| 175 | -0.82 | 6.43E-02 | CCDC74A |
| 176 | -0.78 | 6.43E-02 | CCDC33 |
| 177 | -0.56 | 6.43E-02 | RPRM |
| 178 | -1.63 | 6.54E-02 | CRISP2 |
| 179 | -0.50 | 6.56E-02 | LRRC45 |
| 180 | -0.83 | 6.57E-02 | C1orf194 |
| 181 | -0.82 | 6.65E-02 | CFAP54 |
| 182 | -0.59 | 6.69E-02 | TMEM234 |
| 183 | -1.05 | 6.76E-02 | AC113349.1 |
| 184 | -0.86 | 6.76E-02 | TUBA4B |
| 185 | -0.83 | 6.78E-02 | CFAP52 |
| 186 | -1.12 | 6.81E-02 | PTCH2 |
| 187 | -0.82 | 6.85E-02 | RIIAD1 |
| 188 | -0.84 | 6.88E-02 | C11orf88 |
| 189 | -0.63 | 6.88E-02 | AC073335.2 |
| 190 | -0.72 | 7.06E-02 | CD38 |
| 191 | -1.28 | 7.06E-02 | MAT1A |
| 192 | -0.71 | 7.06E-02 | IQCN |
| 193 | -0.58 | 7.06E-02 | COPRS |
| 194 | -0.69 | 7.22E-02 | WDR66 |
| 195 | -0.80 | 7.27E-02 | SPEF2 |

| | | | |
|-----|-------|----------|------------|
| 196 | -1.61 | 7.27E-02 | ARSEP1 |
| 197 | -1.02 | 7.30E-02 | ALG1L8P |
| 198 | -0.86 | 7.36E-02 | SCARF2 |
| 199 | -0.84 | 7.36E-02 | NHLRC4 |
| 200 | -0.87 | 7.41E-02 | NME9 |
| 201 | -0.92 | 7.44E-02 | C11orf16 |
| 202 | -0.77 | 7.56E-02 | CABCOCO1 |
| 203 | -0.77 | 7.60E-02 | KLHDC9 |
| 204 | -0.82 | 7.62E-02 | MAATS1 |
| 205 | -0.90 | 7.67E-02 | RBM24 |
| 206 | -0.91 | 7.75E-02 | C1orf158 |
| 207 | -0.97 | 7.83E-02 | UMODL1-AS1 |
| 208 | -0.53 | 7.83E-02 | RAD9A |
| 209 | -0.52 | 7.83E-02 | ZDHHC1 |
| 210 | -1.40 | 7.83E-02 | Mar-04 |
| 211 | -0.54 | 7.93E-02 | ADPRHL2 |
| 212 | -0.68 | 7.93E-02 | TUBB4B |
| 213 | -0.84 | 8.16E-02 | PRR29 |
| 214 | -1.04 | 8.20E-02 | CCDC184 |
| 215 | -0.64 | 8.21E-02 | METTTL27 |
| 216 | -0.79 | 8.24E-02 | FANK1 |
| 217 | -1.61 | 8.30E-02 | FAM120B |
| 218 | -0.95 | 8.30E-02 | FAM227A |
| 219 | -0.51 | 8.42E-02 | MGAT3 |
| 220 | -0.83 | 8.47E-02 | CCDC30 |
| 221 | -0.75 | 8.47E-02 | LRRC27 |
| 222 | -0.77 | 8.47E-02 | C9orf116 |
| 223 | -1.21 | 8.47E-02 | DCAF11 |
| 224 | -0.63 | 8.47E-02 | C21orf59 |
| 225 | -0.74 | 8.56E-02 | NPHP1 |
| 226 | -0.80 | 8.56E-02 | DNAH12 |
| 227 | -0.57 | 8.57E-02 | P4HTM |
| 228 | -0.78 | 8.59E-02 | DNAH9 |
| 229 | -1.05 | 8.72E-02 | KLHDC7B |
| 230 | -0.79 | 8.86E-02 | TUBA1A |
| 231 | -0.54 | 8.87E-02 | CKB |
| 232 | -0.86 | 8.88E-02 | CFAP161 |
| 233 | -0.59 | 8.90E-02 | ANKK1 |
| 234 | -0.88 | 8.93E-02 | TSPAN19 |
| 235 | -0.79 | 8.93E-02 | CCDC157 |
| 236 | -1.17 | 8.94E-02 | AC116562.3 |
| 237 | -0.78 | 8.95E-02 | TSNAXIP1 |
| 238 | -0.70 | 9.02E-02 | PPOX |
| 239 | -0.73 | 9.02E-02 | PACRG |

| | | | |
|-----|-------|----------|------------|
| 240 | -1.07 | 9.02E-02 | ZNF295-AS1 |
| 241 | -0.51 | 9.02E-02 | MED25 |
| 242 | -0.63 | 9.10E-02 | ANKRD54 |
| 243 | -0.66 | 9.14E-02 | WDR19 |
| 244 | -0.82 | 9.24E-02 | TOGARAM2 |
| 245 | -1.05 | 9.24E-02 | LRRIQ1 |
| 246 | -0.83 | 9.24E-02 | CAPS |
| 247 | -0.62 | 9.24E-02 | ORAI2 |
| 248 | -0.79 | 9.26E-02 | MS4A8 |
| 249 | -0.89 | 9.27E-02 | PTPRT |
| 250 | -1.01 | 9.27E-02 | AL645924.1 |
| 251 | -0.78 | 9.30E-02 | CAPSL |
| 252 | -0.66 | 9.34E-02 | TCTN1 |
| 253 | -0.82 | 9.35E-02 | NEK5 |
| 254 | -0.83 | 9.54E-02 | FOXJ1 |
| 255 | -0.92 | 9.63E-02 | TTLL9 |
| 256 | -0.64 | 9.63E-02 | NPM2 |
| 257 | -0.64 | 9.64E-02 | CETN2 |
| 258 | -0.70 | 9.64E-02 | CFAP57 |
| 259 | -0.69 | 9.64E-02 | POMT2 |
| 260 | -0.81 | 9.64E-02 | CCDC189 |
| 261 | -0.80 | 9.64E-02 | EFHC1 |
| 262 | -0.75 | 9.64E-02 | RIBC1 |
| 263 | -0.67 | 9.64E-02 | CBY1 |
| 264 | -0.79 | 9.69E-02 | C5orf49 |
| 265 | -0.67 | 9.80E-02 | ICA1L |
| 266 | -0.80 | 9.80E-02 | VPS28 |
| 267 | -0.59 | 9.80E-02 | FBXW9 |
| 268 | -0.97 | 9.80E-02 | ENO4 |
| 269 | -1.14 | 9.80E-02 | AL353747.3 |
| 270 | -0.94 | 9.80E-02 | C22orf15 |
| 271 | -0.66 | 9.80E-02 | DPCD |
| 272 | -0.78 | 9.80E-02 | KIF24 |

Supplementary table 11 - Nasal tissue genes significantly upregulated by mepolizumab

| | Log2fold change | Adjusted p value | Gene |
|----|------------------------|-------------------------|-------------|
| 1 | 4.88 | 2.00E-02 | KRT14 |
| 2 | 4.14 | 3.00E-02 | LIPK |
| 3 | 3.85 | 1.00E-02 | KRT16P6 |
| 4 | 3.84 | 1.00E-02 | LIPG |
| 5 | 3.76 | 7.00E-02 | KRT16 |
| 6 | 3.7 | 4.00E-02 | SPRR1B |
| 7 | 3.54 | 0.00E+00 | LUM |
| 8 | 3.2 | 4.00E-02 | KRT6A |
| 9 | 2.97 | 3.00E-02 | CXCL5 |
| 10 | 2.96 | 2.00E-02 | NES |
| 11 | 2.87 | 3.00E-02 | LPL |
| 12 | 2.84 | 1.00E-02 | PDGFRB |
| 13 | 2.55 | 0.00E+00 | COL5A3 |
| 14 | 2.52 | 4.00E-02 | KLK14 |
| 15 | 2.51 | 2.00E-02 | NNMT |
| 16 | 2.46 | 0.00E+00 | EREG |
| 17 | 2.43 | 1.00E-02 | TGM1 |
| 18 | 2.4 | 1.00E-02 | COL6A3 |
| 19 | 2.39 | 3.00E-02 | SFRP2 |
| 20 | 2.33 | 2.00E-02 | HCAR3 |
| 21 | 2.11 | 4.00E-02 | COL1A2 |
| 22 | 2.06 | 0.00E+00 | SPARC |
| 23 | 2.01 | 7.00E-02 | SLC25A24 |
| 24 | 1.98 | 4.00E-02 | OLR1 |
| 25 | 1.9 | 9.00E-02 | MMP19 |
| 26 | 1.89 | 6.00E-02 | S100A12 |
| 27 | 1.83 | 1.00E-02 | FOSL1 |
| 28 | 1.81 | 3.00E-02 | DCN |
| 29 | 1.77 | 0.00E+00 | IL19 |
| 30 | 1.77 | 3.00E-02 | GJB6 |
| 31 | 1.76 | 8.00E-02 | DUSP9 |
| 32 | 1.75 | 4.00E-02 | TINF2 |
| 33 | 1.74 | 6.00E-02 | S100A2 |
| 34 | 1.74 | 4.00E-02 | MCM10 |
| 35 | 1.73 | 0.00E+00 | GJB2 |
| 36 | 1.73 | 7.00E-02 | DOC2B |
| 37 | 1.66 | 4.00E-02 | PTHLH |
| 38 | 1.64 | 7.00E-02 | ELOVL4 |
| 39 | 1.62 | 0.00E+00 | MYBL2 |
| 40 | 1.59 | 5.00E-02 | KRT16P2 |
| 41 | 1.56 | 0.00E+00 | RRM2 |


| | | | |
|----|------|----------|----------|
| 42 | 1.55 | 2.00E-02 | MRC1 |
| 43 | 1.54 | 0.00E+00 | SLC7A5 |
| 44 | 1.53 | 6.00E-02 | GOLGA7B |
| 45 | 1.53 | 1.00E-02 | FOSB |
| 46 | 1.5 | 0.00E+00 | KRT17 |
| 47 | 1.47 | 3.00E-02 | SERPINB7 |
| 48 | 1.47 | 0.00E+00 | EGR1 |
| 49 | 1.44 | 6.00E-02 | VSIG4 |
| 50 | 1.43 | 6.00E-02 | EXO1 |
| 51 | 1.39 | 4.00E-02 | COL6A2 |
| 52 | 1.38 | 2.00E-02 | HBEGF |
| 53 | 1.38 | 2.00E-02 | C5AR2 |
| 54 | 1.37 | 0.00E+00 | THBD |
| 55 | 1.28 | 4.00E-02 | ADAMTSL4 |
| 56 | 1.26 | 5.00E-02 | CEMIP |
| 57 | 1.22 | 5.00E-02 | CDC45 |
| 58 | 1.21 | 4.00E-02 | PBK |
| 59 | 1.2 | 5.00E-02 | IL1R2 |
| 60 | 1.2 | 5.00E-02 | LZTS1 |
| 61 | 1.18 | 9.00E-02 | ANGPTL4 |
| 62 | 1.18 | 2.00E-02 | CEP55 |
| 63 | 1.15 | 6.00E-02 | TIMP3 |
| 64 | 1.13 | 4.00E-02 | CDC6 |
| 65 | 1.1 | 1.00E-01 | FAM126A |
| 66 | 1.09 | 9.00E-02 | BIRC5 |
| 67 | 1.09 | 8.00E-02 | SOCS3 |
| 68 | 1.06 | 8.00E-02 | AMOTL1 |
| 69 | 1.05 | 4.00E-02 | ADM |
| 70 | 1.05 | 8.00E-02 | ABCA12 |
| 71 | 1.01 | 8.00E-02 | MMP2 |
| 72 | 0.98 | 5.00E-02 | FSCN1 |
| 73 | 0.95 | 9.00E-02 | TPX2 |
| 74 | 0.95 | 1.00E-02 | LGALS7 |
| 75 | 0.93 | 5.00E-02 | MXD1 |
| 76 | 0.92 | 9.00E-02 | C1S |
| 77 | 0.9 | 5.00E-02 | GINS2 |
| 78 | 0.89 | 1.00E-02 | PLAU |
| 79 | 0.88 | 1.00E-02 | FLNA |
| 80 | 0.88 | 3.00E-02 | DUSP7 |
| 81 | 0.88 | 6.00E-02 | AKR1B10 |
| 82 | 0.88 | 1.00E-02 | SERPINB4 |
| 83 | 0.86 | 8.00E-02 | CALD1 |
| 84 | 0.85 | 9.00E-02 | NDRG1 |
| 85 | 0.84 | 1.00E-01 | SAPCD2 |

| | | | |
|-----|------|----------|---------|
| 86 | 0.81 | 2.00E-02 | JUNB |
| 87 | 0.8 | 6.00E-02 | TK1 |
| 88 | 0.78 | 6.00E-02 | FERMT1 |
| 89 | 0.76 | 2.00E-02 | MCM6 |
| 90 | 0.76 | 4.00E-02 | MDFI |
| 91 | 0.76 | 6.00E-02 | TCN1 |
| 92 | 0.75 | 4.00E-02 | OSMR |
| 93 | 0.74 | 7.00E-02 | TINAGL1 |
| 94 | 0.71 | 5.00E-02 | LDHA |
| 95 | 0.7 | 6.00E-02 | CD44 |
| 96 | 0.69 | 3.00E-02 | ZFP36 |
| 97 | 0.69 | 4.00E-02 | DAB2 |
| 98 | 0.68 | 3.00E-02 | STMN1 |
| 99 | 0.68 | 5.00E-02 | PTGS2 |
| 100 | 0.67 | 3.00E-02 | NFKBIA |
| 101 | 0.66 | 7.00E-02 | TPM4 |
| 102 | 0.61 | 8.00E-02 | FAM83D |
| 103 | 0.58 | 5.00E-02 | SIRPA |
| 104 | 0.57 | 6.00E-02 | MSN |
| 105 | 0.53 | 7.00E-02 | AKR1C2 |

Appendix

Summaries of RASP-UK standard operating procedures for sputum induction, nasal scrape sampling, and sputum processing in the MAPLE trial are as follows:

Sputum induction



*Start rescue procedure in section 4.7

⁵Re-measure vital signs. Compare to pre-induction values. If stable, procedure is finished. If not, re-measure in 10 minutes. If still not stable, consult with PI/medic.

Nasal scrape sampling

1. Under direct visual inspection, insert cupped probe into the mid- inferior portion of the inferior turbinate.
2. Gently press the cupped tip on mucosal surface and move outwards 2-3mm (from only one nostril). Repeat motion 2-3 times. If the first scrape does not provide a visible sample, a second scrape should be done on the same nostril. If a second scrape is performed, the cells should be added to the same RNA protect as the first scrape. Make sure the second scrape is not traumatic.
3. The nasal cytology curette can be bent on an angle for ease of use.
4. Add 500µl of RNA protect cell reagent into an 2mL RNase/DNase-free cryovial.
5. Dip the curette into the cryovial containing RNA protect reagent. Lightly shake to free the sample from the curette. Discard the used curette.
6. Store samples at -80°C until shipment to central laboratory .

Sputum processing

