

1 **Dietary diversity and the risk of metabolic syndrome in a Japanese population: An**
2 **analysis of baseline cross-sectional data from the J-MICC study**

3

4 Zin Wai Htay^{1,2}, Nobuaki Michihata¹, Yohko Nakamura¹, Yoshitaka Hippo¹, Jun Otonari³,
5 Hiroaki Ikezaki^{4,5}, Yuichiro Nishida⁶, Chisato Shimanoe⁷, Takashi Tamura⁸, Mako
6 Nagayoshi⁸, Yasufumi Kato⁸, Yudai Tamada⁸, Asahi Hishida⁹, Shiroh Tanoue¹⁰, Daisaku
7 Nishimoto^{10,11}, Teruhide Koyama¹², Etsuko Ozaki¹², Kiyonori Kuriki¹³, Naoyuki
8 Takashima,^{14,15} Naoko Miyagawa^{16,17}, Sakurako Katsuura-Kamano¹⁸, Takeshi Watanabe¹⁸,
9 Kenji Wakai⁸, Keitaro Matsuo^{19,20}

10

11 ¹ Cancer Prevention Center, Chiba Cancer Center Research Institute, 666-2 Nitona-cho,
12 Chuo-ku, Chiba, 260-8717, Japan.

13 ² Department of Global Health Policy, Graduate School of Medicine, The University of
14 Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan

15 ³ Department of Psychosomatic Medicine Graduate School of Medical Sciences, Kyushu
16 University, 3-1-1 Maidashi, Higashi-ku, Fukuoka, 812-8582, Japan

17 ⁴ Department of Comprehensive General Internal Medicine, Kyushu University Faculty of
18 Medical Sciences, 3-1-1 Maidashi, Higashi-ku, Fukuoka, 812-8582, Japan

19 ⁵ Department of General Internal Medicine, Kyushu University Hospital, 3-1-1 Maidashi,
20 Higashi-ku, Fukuoka, 812-8582, Japan

21 ⁶ Department of Preventive Medicine, Faculty of Medicine, Saga University, 5-1-1
22 Nabeshima, Saga, 849-8501, Japan.

23 ⁷ Department of Pharmacy, Saga University Hospital, 5-1-1 Nabeshima, Saga, 849-8501,
24 Japan.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

- 25 ⁸ Department of Preventive Medicine, Nagoya University Graduate School of Medicine, 65
26 Tsurumai-cho, Showa-ku, Nagoya, Aichi, 466-8550, Japan.
- 27 ⁹ Department of Public Health, Aichi Medical University, 1-1 Yasago Karimata. Nagakute,
28 Aichi, 480-1103, Japan
- 29 ¹⁰ Department of Epidemiology and Preventive Medicine, Kagoshima University Graduate
30 School of Medical and Dental Sciences, 8-35-1, Sakuragaoka, Kagoshima, 890-8544, Japan.
- 31 ¹¹ School of Health Sciences, Faculty of Medicine, Kagoshima University, 8-35-1,
32 Sakuragaoka, Kagoshima, 890-8544, Japan.
- 33 ¹² Department of Epidemiology for Community Health and Medicine, Kyoto Prefectural
34 University of Medicine, 465 Kajii-cho, Kamigyō-ku, Kyoto, 602-8566, Japan.
- 35 ¹³ Laboratory of Public Health, Division of Nutritional Sciences, School of Food and
36 Nutritional Sciences, University of Shizuoka, 52-1 Yada, Suruga-ku, Shizuoka, 422-8526,
37 Japan.
- 38 ¹⁴ Department of Epidemiology for Community Health and Medicine, Kyoto Prefectural
39 University of Medicine, 465 Kajii-cho, Kamigyō-ku, Kyoto, 602-8566, Japan.
- 40 ¹⁵ NCD Epidemiology Research Center, Shiga University of Medical Science, Seta-
41 Tsukiwacho, Otsu, Shiga, 520-2192, Japan
- 42 ¹⁶ Department of Preventive Medicine and Public Health, Keio University School of
43 Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo, 160-8582, Japan
- 44 ¹⁷ Department of Public Health, Shiga University of Medical Science, Seta Tsukiwacho, Otsu,
45 Shiga, 520-2192, Japan
- 46 ¹⁸ Department of Preventive Medicine, Tokushima University Graduate School of Biomedical
47 Sciences, 3-18-15 Kuramoto-cho, Tokushima, 770-8503, Japan.
- 48 ¹⁹ Division of Cancer Epidemiology and Prevention, Aichi Cancer Center, 1-1 Kanokoden,
49 Chikusa-ku, Nagoya, Aichi, 464-8681, Japan.

50 ²⁰ Department of Cancer Epidemiology, Nagoya University Graduate School of Medicine, 65
51 Tsurumai-cho, Showa-ku, Nagoya, Aichi, 466-8550, Japan.

52

53 Short title: Diet diversity and metabolic syndrome in Japan

54

55 Address correspondence to:

56 Nobuaki Michihata, Cancer Prevention Center, Chiba Cancer Center Research Institute, 666-
57 2 Nitona-cho, Chuo-ku, Chiba 260-8717, Japan

58 E-mail: nmichihata@chiba-cc.jp

59 Phone: +81-43-264-5431 (PHS:8543), facsimile: +81-43-265-4459

60

61 Word count

62 Abstract: 162 Words

63 Main text without references: 2567 Words

64 Total word count, including abstract and references: 3650 Words

65

66 **Authorship and contributions**

67 Conception and design: ZWH, NM1

68 Data acquisition: JO, HI, YN2, CS, TT, MN, YK, YT, AH, ST, DN, TK, EO, KK, NT, NM2,

69 SK, TW, KW, KM, J-MICC members

70 Analysis and interpretation of data; ZWH, NM1

71 Drafting the manuscript: ZWH

72 Revising the manuscript: NM1, YN1, YH, JO, HI, YN2, CS, TT, MN, YK, YT, AH, ST, DN,

73 TK, EO, KK, NT, NM2, SK, TW, KW, KM, J-MICC members

74

75 **Abstract**

76 **Background:** With the increasing burden of metabolic syndrome, it is crucial to focus on
77 lifestyle interventions to reduce the risk. Diet is a modifiable factor that can reduce the risk of
78 metabolic syndrome.

79 **Methods:** We examined the association between the dietary diversity score (DDS) and risk
80 of metabolic syndrome using baseline data from the Japan Multi-Institutional Collaborative
81 Cohort (J-MICC) study. In total, 75,332 participants were included in this study. A multiple
82 logistic regression analysis was conducted to analyze the association between the DDS and
83 metabolic syndrome.

84 **Results:** Inverse associations were observed between a high DDS and metabolic syndrome
85 (adjusted odds ratio, 0.83 [95% confidential interval 0.76-0.92]). Likewise, a high DDS was
86 associated with reduced odds of a high body mass index and hypertension. No significant
87 associations were observed between the DDS and serum triglyceride, fasting blood glucose,
88 or high-density lipoprotein cholesterol values.

89 **Conclusion:** To reduce the risk of metabolic syndrome, public health interventions should
90 focus on promoting a diverse and balanced diet.

91

92 **Keywords:** dietary diversity score, food variety, nutritional adequacy

93

94 **List of Abbreviations**

95 aOR adjusted odds ratio

96 BMI body mass index

97 DDS dietary diversity score

98 J-MICC Japan Multi-institutional Collaborative Cohort Study

99 KOPS area Kyushu and Okinawa Population Study area

100 HDL-C high-density lipoprotein cholesterol

101 **Introduction**

102 Metabolic syndrome is a cluster of interconnected non-communicable diseases, including
103 insulin resistance, obesity, dyslipidemia, low high-density serum cholesterol, and
104 hypertension.^{1,2} Metabolic syndrome is highly prevalent in both low- and high-income
105 countries and is intricately linked with the incidence of cardiovascular diseases. According to
106 the Global Burden of Disease Study 2019, stroke, cardiovascular diseases, and diabetes were
107 the leading contributors of high disability-adjusted-life-years worldwide.³ Furthermore, in
108 Japan, the age-adjusted prevalence of metabolic syndrome was reported to be 19.3% in
109 2005.⁴

110 Lifestyle modification is crucial in the management of metabolic syndrome. The
111 dietary intake has a direct impact on metabolic risk factors. The intake of high-fiber, high-
112 carbohydrate, and low-fat diets was reported to have positive effects on metabolic diseases.⁵
113 A single-center study in Japan also reported that intake of vegetable-rich diets has an inverse
114 association with the risk of metabolic syndrome.⁶ A fast eating speed is reportedly associated
115 with an increased risk of metabolic syndrome.⁷⁻⁹ However, previous studies have mainly
116 explored the content and quality of the food intake, with little research into dietary diversity.

117 The dietary diversity score (DDS) is a composite measure of the diet quality by
118 counting the number of food groups consumed. Consumption of diverse food groups can also
119 result in balanced nutrient intake¹⁰ and thus can help reduce the risk of metabolic syndrome.
120 The literature mainly explains the associations between DDS and the components of
121 metabolic syndrome, such as obesity and diabetic conditions. However, inconsistent
122 associations have been reported in different studies. Previous studies conducted among
123 Iranian adolescents and adults reported a positive association between DDS and obesity.^{11,12}
124 However, another study in the UK reported inverse associations between DDS and diabetic
125 incidence¹³. Similarly, a study in Korea also stated that DDS was associated with a reduced

126 likelihood of abdominal obesity and high triglyceride levels.¹⁴ Such variations in findings
127 might be due to small sample populations and differing study designs among previously
128 conducted studies, resulting in inconclusive evidence.

129 The present study therefore explored the association between the DDS and risk of
130 metabolic syndrome in a Japanese population.

131

132 **Methods**

133 *Study setting and participants*

134 Our study analyzed baseline data from the Japan Multi-institutional Collaborative Cohort (J-
135 MICC) study (version.20210901 dataset). Baseline surveys were collected during 2005-2014
136 at 14 different sites in Japan, including the Aichi Cancer Center, Chiba, Kanagawa,
137 Takashima, Okazaki, Shizuoka-Daiko, Kyoto, Fukuoka, Saga, Kagoshima, Tokushima,
138 Kyushu and Okinawa Population Study (KOPS) area, Shizuoka-Sakuragaoka, and Iga. The
139 current study excluded the Aichi Cancer Center, Chiba, and Kanagawa data due to a lack of
140 surveys on height, weight, blood tests, and other data necessary for defining metabolic
141 syndrome.

142 Ethical approval was obtained from the ethics committee of the Aichi Cancer Center
143 and each study site. Written informed consent was obtained from all participants. The details
144 of the J-MICC study were reported in a previously published study profile.¹⁵ Participants 35-
145 69 years old were recruited, and information was collected via a structured questionnaire. In
146 addition, peripheral blood samples and anthropometric measurements were collected at the
147 time of enrollment.¹⁵ The response rates at the study sites ranged from 19.7% to 69.8%.

148

149 *Study measurements*

- 150 • Dietary diversity assessments

151 Participants were administered a short food frequency questionnaire containing 47 items as
152 part of the survey questions to obtain information on dietary habits.^{16,17} The short food
153 frequency questionnaire assessed the food intake frequency on an 8-point scale from "not at
154 all" to "1 to 3 or more times per day." In this study, the food intake frequency was treated as a
155 dichotomous variable of whether or not the participants consumed food "at least once a
156 week". We classified foods into five major groups (bread and grains, vegetables, meat, dairy,
157 and fruit) based on Kant et al.¹⁸ To date, no universal definition of dietary diversity has been
158 proposed. The DDS was calculated for each participant according to a counting approach that
159 codes the intake frequency as 1 if the food group was consumed at least once a week and 0 if
160 not.^{13,19} The DDS categorizes diets into three groups (group 3 or less, group 4, or group 5)
161 based on the number of the five food groups (bread and grains, vegetables, meat, dairy, and
162 fruit) consumed. To understand the intake quantity in addition to the variety, we calculated
163 the cumulative score of five food groups based on the numbers of items consumed in each
164 food group as the DDS-2.²⁰ Each food group was assigned a maximum number of two, and
165 the score was calculated by dividing the number of times a given item was consumed by the
166 total number of items in each group. The obtained number was then multiplied by 2. A total
167 score ranging from 0-10 was obtained by combining the five food groups, which were
168 classified into four quantiles.²⁰

169

- 170 • Metabolic syndrome assessments

171 We followed the criteria of the National Cholesterol Education Program Adult Treatment
172 Panel III (NCEP-ATP III)²¹ for the diagnosis of metabolic syndrome. Participants with
173 biochemical data were diagnosed with metabolic syndrome if they met three of the following
174 five conditions: 1) body mass index (BMI) ≥ 25 kg/m²; 2) systolic blood pressure ≥ 130
175 mmHg and/or diastolic blood pressure ≥ 85 mmHg or the use of antihypertensive medication;

176 3) serum triglyceride level ≥ 150 mg/dL; 4) serum high-density lipoprotein cholesterol (HDL-
177 C) level < 40 mg/dL in men or < 50 mg/dL in women; and 5) blood fasting glucose level ≥ 100
178 mg/dL. Based on a previous study, the BMI was used instead of the waist circumference
179 because waist circumference data were unavailable in the current study.²²

180

- 181 • Other variables

182 The survey questions assessed the demographic characteristics (age, sex, years of education),
183 lifestyle factors (physical activity, history of smoking, and drinking alcohol), and family
184 history of hypertension and diabetes mellitus. Based on previous studies on metabolic
185 syndrome, these variables were considered possible confounders.^{14,20}

186

187 *Statistical analyses*

188 Descriptive analyses were conducted for each variable in order to understand their
189 distribution. We also performed cross-tabulations and chi-square tests to understand the
190 distribution of each variable with metabolic syndrome occurrence. To avoid underestimation,
191 a multiple imputation method was applied when the number of missing observations
192 exceeded 5% of the total observations.²³ This method utilizes a Markov chain Monte Carlo
193 algorithm to generate 20 complete datasets by replacing missing values with a series of
194 predicted values. Multiple imputation assumes that the data are randomly missing. The
195 covariates included in the imputation were the age, sex, BMI, systolic blood pressure,
196 diastolic blood pressure, triglyceride level, HDL-C level, fasting glucose level, smoking
197 status, alcohol intake status, family history of hypertension, family history of diabetes, DDS,
198 and DDS-2. The input generated 20 complete datasets. The results from each input dataset
199 were combined into a single overall estimate and standard error using Rubin's law to reflect

200 the uncertainty associated with missing values. This method allows for a comprehensive
201 analysis that accounts for potential variation due to missing data.²³

202 We conducted multivariable logistic regression to obtain the risk estimates of
203 metabolic syndrome in relation to the two types of DDS. The following variables were
204 controlled for in both models as possible confounders: age, sex, years of education, history of
205 smoking, alcohol consumption, and physical activity. The site variable was also included in
206 the model to account for variations in the data collection at different centers. Analyses were
207 repeated for stratification according to metabolic syndrome risk factors.

208 All analyses were conducted using the Stata statistical software program (17.1;
209 StataCorp LLC, College Station, TX, USA). The threshold level was set to 5%.

210

211 **Results**

212 In total, 75,332 participants were included in the analysis. Table 1 shows the distribution of
213 the participants. The majority (58.1%) of the population was ≥ 55 years old. More than half of
214 the participants were women (55.3%). The average age of the study population was 55 years
215 old. On average, 76.5% of the population engaged in physical exercise. The percentage of
216 participants with a DDS of ≤ 3 was significantly higher in men (76.4%) than in women.
217 Similar results were obtained for DDS-2, with the lowest quintile of DDS-2 predominantly
218 comprising men (64.3%) (Supplementary Table S1).

219 Table 2 shows the distribution of the DDS and metabolic syndrome. Overall,
220 metabolic syndrome was reported to be the highest among subjects 55-69 years old (66.3%)
221 and men (60.7%). The distribution of metabolic syndrome among the study sites also varied,
222 with the highest percentage reported in Saga (17.2%) and the lowest in Tokushima (2.0%).
223 The distribution of the DDS among the five major food groups is reported in Supplementary
224 Table S2. In the group with a DDS ≤ 3 , 97% and 37.6% of participants reported not eating

225 fruits and vegetables, respectively. Similarly, participants in the lowest quantile (quantile 1)
226 of DDS-2 reported a reduced intake of fruits (66.7%) and dairy products (15.4%).

227 Table 3 presents the results of the multivariate logistic regression analysis. The total
228 number of observations in this logistic regression model was imputed because more than
229 28.3% of the observations were missing for education and fasting blood glucose variables.
230 Participants with a high DDS were significantly less likely to have metabolic syndrome than
231 those with lower values, as evidenced by an adjusted odds ratio (aOR) of 0.83 (95%
232 confidence interval [CI] 0.76-0.92). A similar estimate was observed for the DDS-2, with the
233 rate of metabolic syndrome being lowest among those in the highest DDS-2 quantile, as
234 indicated by an aOR of 0.90 (95% CI 0.82-0.99). Similarly, we found that a high DDS was
235 associated with a reduced occurrence of a high BMI (aOR 0.91 [95% CI 0.84-0.99]),
236 hypertension (aOR 0.84 [95% CI 0.77-0.92]), and high serum triglyceride levels (aOR 0.90
237 [95% CI 0.82-0.99]) (Supplementary Table S3). A similar pattern was observed for the DDS-
238 2 and metabolic syndrome risk factors, except for high serum triglyceride levels
239 (Supplementary Table S4). However, we found no association between the DDS and low
240 HDL-C levels or high fasting glucose levels (Supplementary Tables S5 and S6). We also
241 conducted stratification analyses by sex group; however, no effect modification was observed
242 (Supplementary Tables S5 and S6).

243

244 **Discussion**

245 In this study, we collected data from various regions of Japan to examine the association
246 between the DDS and metabolic syndrome. This is one of the few studies to focus on dietary
247 diversity and metabolic syndrome risk factors in Japan. We found that the overall DDS had
248 an inverse relationship with the risk of metabolic syndrome. We also found that a high DDS

249 was associated with a reduced likelihood of reporting a high BMI, hypertension, high serum
250 triglyceride levels, and high fasting blood glucose levels.

251 Both a high DDS and high DDS-2 were associated with a reduced likelihood of
252 reporting metabolic syndrome. Our findings were consistent with those of a previous study in
253 a Korean population, in which a lower risk of metabolic syndrome and its risk factors were
254 reported in the higher DDS group than in the lower DDS group.¹⁴ This supports our
255 hypothesis that a diverse diet is a protective factor against the risk of metabolic syndrome.
256 The underlying assumption for the current finding is that dietary diversity is highly associated
257 with the consumption of a balanced and adequate diet.²⁴ Our study also showed that those in
258 the diverse diet groups were more likely to consume fruits and vegetables than others. Our
259 descriptive analyses revealed that men had a higher percentage of lower DDS scores than did
260 women. This observation aligns with previous studies conducted in China and Australia,^{25,26}
261 which also reported higher meat consumption among men than among women. These
262 findings emphasize the need to prioritize the male population when promoting dietary
263 diversity, warranting increased attention, and targeted interventions. In our study, we
264 compared the risk estimates of the two DDS indicators, with the second indicator considering
265 the cumulative index of dietary diversity. However, since the risk estimates were similar for
266 both DDS indicators, we were unable to determine whether or not consuming a wide range of
267 food groups within a single food category was particularly advantageous for health.

268 Analyses of metabolic syndrome risk factors also showed that a high DDS was
269 associated with a reduced likelihood of reporting hypertension and high serum triglyceride
270 levels. The finding that DDS is associated with a reduced likelihood of hypertension was
271 supported by a previous study,²⁷ which reported that DDS was inversely associated with
272 serum triglyceride levels and systolic blood pressure. In addition to the carbohydrate intake,
273 the meat intake was also reported to be associated with the risk of peripheral

274 atherosclerosis.²⁸ Furthermore, low-carbohydrate diets and calorie restriction have been
275 reported to be effective interventions in reducing serum triglyceride levels.²⁹ Similar to our
276 findings, a previous Japanese study reported that greater dietary diversity was associated with
277 the more frequent consumption of fiber in the diet,³⁰ which can be a possible explanation to
278 reduced serum triglyceride concentrations, thereby leading to a preventive effect.

279 In addition, we found that a higher DDS was associated with a reduced risk of
280 reporting a high BMI. One underlying assumption is that a higher DDS is associated with a
281 lower consumption of a carbohydrate-rich diet, which is the main factor leading to obesity.³¹
282 However, contrary to our findings, a previous systematic review reported no associations
283 between the DDS and BMI.³² Similarly, another systematic review reported no conclusive
284 evidence concerning the DDS and obesity associations.¹⁹ These diverse associations might be
285 related to different cultural aspects of the involved participants. In our study, we investigated
286 the associations by comparing the two DDS scores, and the results from both scores were
287 consistent, indicating that consuming a diverse diet is advantageous for BMI management.

288 Our study had several strengths and limitations. This study offers a comprehensive
289 evaluation of dietary diversity in the Japanese population by employing two different
290 approaches to define the DDS. The findings from both approaches are comparable, revealing
291 a consistent association between increased dietary diversity and a decreased risk of metabolic
292 syndrome. In addition, our study incorporated a substantial sample size, which enhanced the
293 reliability and statistical power of our findings. However, this study was an analysis of
294 baseline cross-sectional data, and because of the nature of the cross-sectional study, we were
295 unable to establish causal relationships. Although we controlled for educational status and
296 basic demographic characteristics in our model, we were unable to control for the wealth
297 index of the study population, which might have resulted in residual confounding. In addition,
298 a self-reported questionnaire was used to collect data on dietary factors, the history of

299 smoking, drinking, physical exercise, and the family history, so there was a possibility of
300 inevitable misclassification or measurement error. This might have led to the underestimation
301 of true associations, with the estimates biased towards the null hypothesis. Despite the
302 potential for misclassification or measurement errors due to the use of self-reported
303 questionnaires, our findings were relatively uninfluenced, as we identified significant
304 associations.

305 Our findings revealed variations in site-specific risks for the association between DDS
306 and metabolic syndrome. Further exploration of the prefectural-level economic situation and
307 other possible influencing factors is recommended to explain such variations. This study did
308 not explore the determinants of dietary diversity in the Japanese population, so further studies
309 are recommended to determine policy interventions to implement and promote dietary
310 diversity.

311

312 **Conclusion**

313 In conclusion, our study found that consumption of a diverse range of foods was associated
314 with a reduced risk of metabolic syndrome and its associated risk factors. To mitigate the risk
315 of metabolic syndrome in the Japanese population, it is advisable to avoid restricting food
316 intake to specific groups. Instead, promoting a varied and balanced diet may be
317 recommended as a preventive measure against metabolic syndrome risk factors.

318

319 **Sources of funding**

320 This study was supported by Grants-in-Aid for Scientific Research for Priority Areas of
321 Cancer (No.17015018), Innovative Areas (No.221S0001), and Platform of Supporting Cohort
322 Study and Biospecimen Analysis (Japan Society for the Promotion of Science [JSPS])

323 KAKENHI Grants No.16H06277 and 22H04923 [CoBiA]) from the Japanese Ministry of
324 Education, Culture, Sports, Science, and Technology.

325

326 **Acknowledgements**

327 We would like to thank the participants of the J-MICC study for providing the necessary
328 information to conduct the current research.

329

330 **Disclosures**

331 The authors declare no conflict of interests.

332

333 **Supplementary Materials**

334 Tables S1-S6

335 **References**

- 336 1. Saklayen MG. The global epidemic of the metabolic syndrome. *Curr Hypertens Rep.*
337 2018;20. doi:10.1007/S11906-018-0812-Z
- 338 2. Kassi E, Pervanidou P, Kaltsas G, Chrousos G. Metabolic syndrome: definitions and
339 controversies. *BMC Med.* 2011;9:1-13. doi:10.1186/1741-7015-9-48/TABLES/2
- 340 3. Abbafati C, Abbas KM, Abbasi-Kangevari M, et al. Global burden of 369 diseases and
341 injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global
342 Burden of Disease Study 2019. *The Lancet.* 2020;396(10258):1204-1222.
343 doi:10.1016/S0140-6736(20)30925-9
- 344 4. Liu L, Miura K, Fujiyoshi A, et al. Impact of metabolic syndrome on the risk of
345 cardiovascular disease mortality in the United States and in Japan. *American Journal of*
346 *Cardiology.* 2014;113:84-89. doi:10.1016/j.amjcard.2013.08.042
- 347 5. Calton EK, James AP, Pannu PK, Soares MJ. Certain dietary patterns are beneficial for
348 the metabolic syndrome: reviewing the evidence. *Nutrition Research.* 2014;34:559-568.
349 doi:10.1016/J.NUTRES.2014.06.012
- 350 6. Bahari T, Uemura H, Katsuura-Kamano S, et al. Nutrient-derived dietary patterns and
351 their association with metabolic syndrome in a Japanese population. *J Epidemiol.*
352 2018;28:194. doi:10.2188/JEA.JE20170010
- 353 7. Garcidueñas-Fimbres TE, Paz-Graniel I, Nishi SK, Salas-Salvadó J, Babio N. Eating
354 speed, eating frequency, and their relationships with diet quality, adiposity, and metabolic
355 syndrome, or its components. *Nutrients.* 2021;13. doi:10.3390/NU13051687
- 356 8. Nanri A, Miyaji N, Kochi T, Eguchi M, Kabe I, Mizoue T. Eating speed and risk of
357 metabolic syndrome among Japanese workers: the Furukawa Nutrition and Health Study.
358 *Nutrition.* 2020;78:110962. doi:10.1016/J.NUT.2020.110962

- 359 9. Zhu B, Haruyama Y, Muto T, Yamazaki T. Association between eating speed and
360 metabolic syndrome in a three-year population-based cohort study. *J Epidemiol.*
361 2015;25:332. doi:10.2188/JEA.JE20140131
- 362 10. Kennedy GL, Pedro MR, Seghieri C, Nantel G, Brouwer I. Dietary diversity score is a
363 useful indicator of micronutrient intake in non-breast-feeding Filipino children. *J Nutr.*
364 2007;137:472-477. doi:10.1093/JN/137.2.472
- 365 11. Karimbeiki R, Pourmasoumi M, Feizi A, et al. Higher dietary diversity score is
366 associated with obesity: a case-control study. *Public Health.* 2018;157:127-134.
367 doi:10.1016/j.puhe.2018.01.028
- 368 12. Golpour-Hamedani S, Rafie N, Pourmasoumi M, Saneei P, Safavi SM. The association
369 between dietary diversity score and general and abdominal obesity in Iranian children
370 and adolescents. *BMC Endocr Disord.* 2020;20:1-8. doi:10.1186/S12902-020-00662-
371 W/TABLES/4
- 372 13. Conklin AI, Monsivais P, Khaw KT, Wareham NJ, Forouhi NG. Dietary diversity, diet
373 cost, and incidence of type 2 diabetes in the United Kingdom: a prospective cohort study.
374 *PLoS Med.* 2016;13. doi:10.1371/JOURNAL.PMED.1002085
- 375 14. Kim J, Kim M, Shin Y, Cho JH, Lee D, Kim Y. Association between dietary diversity
376 score and metabolic syndrome in Korean adults: a community-based prospective cohort
377 study. *Nutrients.* 2022;14. doi:10.3390/NU14245298/S1
- 378 15. Takeuchi K, Naito M, Kawai S, et al. Study profile of the Japan Multi-institutional
379 Collaborative Cohort (J-MICC) Study. *J Epidemiol.* 2021;31:660-668.
380 doi:10.2188/JEA.JE20200147
- 381 16. Tokudome S, Ikeda M, Tokudome Y, Imaeda N, Kitagawa I, Fujiwara N. Development of
382 data-based semi-quantitative food frequency questionnaire for dietary studies in middle-
383 aged Japanese. *Jpn J Clin Oncol.* 1998;28:679-687. doi:10.1093/jjco/28.11.679

- 384 17. Tokudome Y, Goto C, Imaeda N, et al. Relative validity of a short food frequency
385 questionnaire for assessing nutrient intake versus three-day weighed diet records in
386 middle-aged Japanese. *J Epidemiol.* 2005;15:135. doi:10.2188/JEA.15.135
- 387 18. Kant AK, Schatzkin A, Harris TB, Ziegler RG, Block G. Dietary diversity and
388 subsequent mortality in the First National Health and Nutrition Examination Survey
389 Epidemiologic Follow-up Study. *Am J Clin Nutr.* 1993;57:434-440.
390 doi:10.1093/AJCN/57.3.434
- 391 19. de Oliveira Otto MC, Anderson CAM, Dearborn JL, et al. Dietary diversity: implications
392 for obesity prevention in adult populations: a science advisory from the American Heart
393 Association. *Circulation.* 2018;138:e160. doi:10.1161/CIR.0000000000000595
- 394 20. Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi F. Dietary diversity score and
395 cardiovascular risk factors in Tehranian adults. *Public Health Nutr.* 2006;9:728-736.
396 doi:10.1079/PHN2005887
- 397 21. Grundy SM, Cleeman JI, Daniels SR, et al. Diagnosis and management of the metabolic
398 syndrome. *Circulation.* 2005;112:2735-2752.
399 doi:10.1161/CIRCULATIONAHA.105.169404
- 400 22. Van Nguyen T, Arisawa K, Katsuura-Kamano S, et al. Associations of metabolic
401 syndrome and metabolically unhealthy obesity with cancer mortality: The Japan Multi-
402 Institutional Collaborative Cohort (J-MICC) study. *PLoS One.* 2022;17:e0269550.
403 doi:10.1371/JOURNAL.PONE.0269550
- 404 23. Li P, Stuart EA, Allison DB. Multiple imputation: a flexible tool for handling missing
405 data. *JAMA.* 2015;314:1966. doi:10.1001/JAMA.2015.15281
- 406 24. Oldewage-Theron WH, Kruger R. Food variety and dietary diversity as indicators of the
407 dietary adequacy and health status of an elderly population in Sharpeville, South Africa. *J*
408 *Nutr Elder.* 2008;27:101-133. doi:10.1080/01639360802060140

- 409 25. Tian X, Wu M, Zang J, Zhu Y, Wang H. Dietary diversity and adiposity in Chinese men
410 and women: an analysis of four waves of cross-sectional survey data. *European Journal*
411 *of Clinical Nutrition* 2017 71:4. 2016;71(4):506-511. doi:10.1038/ejcn.2016.212
- 412 26. Xu X, Inglis SC, Parker D. Sex differences in dietary consumption and its association
413 with frailty among middle-aged and older Australians: a 10-year longitudinal survey.
414 *BMC Geriatr.* 2021;21(1):1-12. doi:10.1186/S12877-021-02165-2/TABLES/4
- 415 27. Farhangi MA, Jahangiry L. Dietary diversity score is associated with cardiovascular risk
416 factors and serum adiponectin concentrations in patients with metabolic syndrome. *BMC*
417 *Cardiovasc Disord.* 2018;18:1-6. doi:10.1186/S12872-018-0807-3/FIGURES/1
- 418 28. Ogilvie RP, Lutsey PL, Heiss G, Folsom AR, Steffen LM. Dietary intake and peripheral
419 arterial disease incidence in middle-aged adults: the Atherosclerosis Risk in
420 Communities (ARIC) Study. *Am J Clin Nutr.* 2017;105:651.
421 doi:10.3945/AJCN.116.137497
- 422 29. Luna-Castillo KP, Olivares-Ochoa XC, Hernández-Ruiz RG, et al. The effect of dietary
423 interventions on hypertriglyceridemia: from public health to molecular nutrition
424 evidence. *Nutrients.* 2022;14. doi:10.3390/NU14051104
- 425 30. Otsuka R, Tange C, Nishita Y, et al. Dietary diversity and all-Cause and cause-specific
426 mortality in Japanese community-dwelling older adults. *Nutrients.* 2020;12.
427 doi:10.3390/NU12041052
- 428 31. Miyamoto K, Kawase F, Imai T, Sezaki A, Shimokata H. Dietary diversity and healthy
429 life expectancy—an international comparative study. *European Journal of Clinical*
430 *Nutrition* 2018 73:3. 2018;73:395-400. doi:10.1038/s41430-018-0270-3
- 431 32. Salehi-Abargouei A, Akbari F, Bellissimo N, Azadbakht L. Dietary diversity score and
432 obesity: a systematic review and meta-analysis of observational studies. *European*
433 *Journal of Clinical Nutrition* 2016 70:1. 2015;70:1-9. doi:10.1038/ejcn.2015.118

434 Table 1. Basic characteristics of participants and distribution of the dietary diversity score

Variables	Total N=75,332	Dietary diversity score			p-value
		≤3 N=2,781	4 N=18,394	5 N=54,147	
Age, n (%)					<0.001
35-44 years	12,205 (16.2)	556 (20.0)	4,404 (23.9)	7,245 (13.4)	
45-54 years	19,374 (25.7)	820 (29.5)	5,616 (30.5)	12,938 (23.9)	
55-69 years	43,753 (58.1)	1,405 (50.5)	8,374 (45.5)	33,974 (62.7)	
Gender, n (%)					<0.001
Female	41,694 (55.3)	655 (23.6)	7,832 (42.6)	33,207 (61.3)	
Male	33,638 (44.7)	2,126 (76.4)	10,562 (57.4)	20,950 (38.7)	
Body mass index, mean (SD), kg/m ²	23.1 (3.3)	23.50 (3.5)	23.2 (3.4)	23.0 (3.3)	<0.001
Systolic blood pressure, mean (SD), mmHg	127.8 (20.0)	129.9 (19.8)	127.7 (20.0)	127.8 (20.0)	<0.001
Diastolic blood pressure, mean (SD), mmHg	78.0 (12.1)	80.24 (12.1)	78.5 (12.4)	77.8 (12.0)	<0.001
Triglycerides, mean (SD), mg/dL	125.1 (98.0)	141.10 (120.3)	130.6 (108.3)	122.4 (92.8)	<0.001
HDL-C, mean (SD), mg/dL	63.4 (16.6)	59.84 (16.6)	62.5 (16.8)	63.9 (16.6)	<0.001
Fasting glucose, mean (SD), mg/dL	98.0 (21.0)	100.62 (22.8)	98.1 (21.3)	97.8 (20.8)	<0.001
Smoking status, n (%)					<0.001
Current	12,883 (17.2)	1,024 (37.0)	5,014 (27.4)	6,845 (12.7)	
Past	16,455 (21.9)	846 (30.6)	4,749 (26.0)	10,860 (20.1)	
Never	45,631 (60.9)	897 (32.4)	8,531 (46.6)	36,203 (67.2)	
Drinking status, n (%)					<0.001
Current	40,635 (54.0)	1,942 (69.9)	11,590 (63.1)	27,103 (50.1)	
Past	1,925 (2.6)	91 (3.3)	468 (2.5)	1,366 (2.5)	
Never	32,686 (43.4)	746 (26.8)	6,316 (34.4)	25,624 (47.4)	
Regular physical exercise, n (%)	57,647 (76.5)	1,758 (63.3)	12,729 (69.4)	42,692 (79.0)	<0.001
Years of education, n (%)					<0.001
≤9	5,613 (10.4)	373 (15.0)	1,457 (10.7)	3,783 (10.0)	
10-15	23,152 (42.9)	1,151 (46.2)	6,026 (44.1)	15,975 (42.2)	
≥16	25,030 (46.4)	956 (38.4)	6,131 (44.9)	17,943 (47.4)	
Unknown	201 (0.4)	11 (0.4)	52 (0.4)	138 (0.4)	
History of hypertension, n (%)	34,146 (46.9)	1,342 (51.2)	8,201 (46.5)	24,603 (46.8)	<0.001
Family history, n (%)					
Hypertension	33,300 (47.7)	1,132 (45.6)	8,031 (47.7)	24,137 (47.8)	<0.001
Diabetes	12,789 (18.3)	479 (19.1)	3,302 (19.6)	9,008 (17.8)	<0.001

435 SD: standard deviation; HDL-C: high-density lipoprotein cholesterol

436 Table 2. Distributions of the dietary diversity score and metabolic syndrome

Variables	Metabolic syndrome		p-value
	Yes, n (%)	No, n (%)	
Dietary diversity score			< 0.001
≤3	549 (4.8)	2,232 (3.5)	
4	2,980 (26.2)	15,414 (24.1)	
5	7,847 (69.0)	46,310 (72.4)	
Age			< 0.001
35-44 years	1,086 (9.6)	11,119 (17.4)	
45-54 years	2,743 (24.1)	16,631 (26.0)	
55-69 years	7,547 (66.3)	36,206 (56.6)	
Gender			< 0.001
Female	4,466 (39.3)	37,228 (58.2)	
Male	6,910 (60.7)	26,728 (41.8)	
Years of education			< 0.001
≤9	1,128 (14.6)	4,485 (9.7)	
10-15	3,428 (44.3)	19,724 (42.6)	
≥16	3,150 (40.7)	21,880 (47.3)	
Unknown	34 (0.4)	167 (0.4)	
Smoking status			< 0.001
Never	5,734 (50.6)	39,897 (62.7)	
Past	3,253 (28.7)	13,202 (20.8)	
Current	2,348 (20.7)	10,535 (16.6)	
Drinking status			< 0.001
Never	4,602 (40.5)	28,084 (44.0)	
Past	343 (3.0)	1,582 (2.5)	
Current	6,413 (56.5)	34,222 (53.6)	
Regular physical exercise			< 0.001
No	2,931 (25.8)	15,046 (23.6)	
Yes	8,410 (74.2)	48,769 (76.4)	
Site			< 0.001
Takashima	460 (4.0)	3,065 (4.8)	
Okazaki	982 (8.6)	5,568 (8.7)	
Shizuoka-Daiko	968 (8.5)	9,189 (14.4)	
Kyoto	917 (8.1)	5,276 (8.3)	
Fukuoka	1,444 (12.7)	8,649 (13.5)	
Saga	1,953 (17.2)	10,115 (15.8)	
Kagoshima	1,572 (13.8)	6,066 (9.5)	
Tokushima	222 (2.0)	2,218 (3.5)	
KOPS area	1,948 (17.1)	7,782 (12.2)	
Shizuoka-Sakuragaoka	680 (6.0)	4,837 (7.6)	
Iga	230 (2.0)	1,191 (1.9)	

437

438 Table 3. Associations between dietary diversity scores and metabolic syndrome

Variables	Multivariable logistic regression	
	aOR (95% CI)*	<i>p</i> -value
DDS		
≤ 3	Reference	
4	0.88 (0.80-0.97)	0.013
5	0.83 (0.76-0.92)	< 0.001
DDS-2		
Quantile 1	Reference	
Quantile 2	0.98 (0.92-1.03)	0.359
Quantile 3	0.92 (0.87-0.98)	0.005
Quantile 4	0.91 (0.84-0.98)	0.011

439 aOR: adjusted odds ratio, CI: confidence interval, DDS: dietary diversity score

440 *Adjusted for the age, sex, years of education, smoking status, drinking status, regular

441 physical exercise, and site