

1 **Associations between Diabetes Self-Management and Microvascular** 2 **Complications Among Patients Living in Rural Areas, in Kenya**

3 Authors: Rachael Ileri^{1,2}, Gideon Kikvi¹, Susan Mambo¹, Besty Cheriro¹

4 ¹School of Public Health; Jomo Kenyatta University of Agriculture and
5 Technology

6 ²Department of Clinical Medicine; Kenya Medical Training College

7 **Introduction:** Diabetes is a major public health concern worldwide. Low and middle-
8 income countries are the most affected. Diabetes self-management can significantly
9 reduce the burden of diabetes complications and mortality.

10 **Methods:** This cross-sectional study was conducted at the outpatient department of a
11 county referral hospital in Kenya, from 1st August 2022 to 30th October 2022. Patients
12 with known type II diabetes of age ≥ 20 years visiting the hospital for routine follow-up
13 visits were included. A 7-item Summary of Diabetes Self-care Activities (SDSAC)
14 Questionnaire was used to assess Diabetes self-care activities. For data entry and
15 statistical analysis, SPSS for Windows version 27.0 was used.

16 **Results:** There were 96 (39.2%) males and 149 (60.8%) females. Most of the
17 participants were more than 61 years 148 (60.4%). Prevalence of Diabetes microvascular
18 complications among the study participants was 56.78%, 25.3%, and 14.7% for
19 neuropathy, retinopathy, and nephropathy respectively. significant association was
20 found between the sum scale scores of dietary activities, blood glucose testing, physical
21 activity, foot care, and neuropathy at 95% CI and ($P < 0.001$). Scores on the Blood
22 glucose testing and foot care activities subscale were significantly associated with

23 Retinopathy ($p < 0.001$). Additionally, significant associations were found between the
24 presence of nephropathy and dietary activities, foot care, and physical activity subscale
25 ($p = < 0.001$).

26 **Conclusions:** Diabetes self-management activities have an impact on microvascular
27 complications in patients with diabetes.

28 **Keywords:** self-management activities, diabetes, Microvascular complications,
29 Summary of diabetes self-care activities.

30 **Introduction**

31 Diabetes is a major public health concern that is approaching epidemic proportions
32 worldwide(1). Levels of morbidity and mortality attributable to diabetes remain high.
33 Though diabetes and its related complications is a global problem low and middle-
34 income countries (LMICs) are the most affected(2). Diabetes treatment imposes a huge
35 economic burden in developing countries and the costs are projected to rise further(3).
36 Among the chronic complications of diabetes, microvascular complications of kidney,
37 retina, and peripheral nerves are a significant cause of mortality and disability in
38 patients with diabetes. This greatly affects their quality of life and care burden. the
39 Diabetes self-management can reduce complications and mortality in patients living
40 with diabetes(4).

41 Unlike previously believed that it is more of a problem of cities and towns, the rural
42 communities have experienced an increase as well(3). The responsibility of providing
43 preventive and therapeutic healthcare services falls on the healthcare workers in the
44 primary health centers and dispensaries as there are not many private or tertiary-level

45 hospitals to care for the health demands of diabetic patients(5). The situation is worse
46 because these communities are faced with multiple challenges, including a lack of access
47 to diabetes education, clinical services, limited smart cellphone coverage, and internet
48 access, limited transportation and long travel distances, as well as higher rates of
49 poverty(6).

50 The needs of diabetic patients are not only limited to adequate glycemic control but also
51 prevent complications and disability(7). One of the biggest challenges for healthcare
52 providers is addressing the continued needs and demands of individuals with
53 diabetes(8). The regular follow-up of diabetic patients by a multidisciplinary team is
54 vital in averting long-term complications(9,10). This is met with difficulties as more
55 than 80% of consultants in Kenya are found in Urban areas and tertiary hospitals, this
56 in turn contributes to suboptimal diabetes management and higher rates of diabetes-
57 related complications(11).

58 **Diabetes Self-Management Activities**

59 Self-care in diabetes is a process of development of knowledge or awareness by learning
60 to survive with the complex nature of diabetes in a social context(12). The majority of
61 daily diabetes care is handled by patients and/or families. Seven essential self-care
62 behaviors in people with diabetes predict outcomes. They include; healthy eating, being
63 physically active, monitoring of blood sugar, compliance with medications, good
64 problem-solving skills, healthy coping skills, and risk-reduction behaviors(13–16). These
65 measures can be useful for both clinicians and educators treating individual patients
66 and for researchers evaluating new approaches to care(17). Self-report is an important
67 practical and cost-effective approach for self-management assessment(18).

68 Diabetes self-management activities are behaviors undertaken by people with diabetes
69 to successfully manage the disease on their own. All seven behaviors are positively
70 correlated with good glycemic control, reduction of complications, and improvement in
71 quality of life(9,19,20). In addition, it was observed that self-care encompasses not only
72 performing these activities but also the interrelationships between them. Diabetes self-
73 care requires the patient to make major lifestyle modifications enhanced with the
74 supportive role of healthcare staff for maintaining a higher level of self-confidence
75 leading to a successful behavior change(21–24).

76 Diabetes education is important but it must be transferred to action or self-care
77 activities to fully benefit the patient. The American Association of Clinical
78 Endocrinologists and WHO emphasize the importance of patients becoming active and
79 knowledgeable participants in their care(10,25). Emphasis should be placed on;
80 understanding the disease and the role of drugs, diet plans, foot, and eye care, role of
81 continuous and sustained lifestyle modification, physical activities, and self-glucose
82 monitoring(26).

83 Given the prevalence of DM and its complications in Nyeri county, there had been
84 limited studies conducted to determine diabetes self-management among its
85 population. Therefore, study aimed to assess the diabetes self-management practices
86 among diabetes patients living in rural areas in Nyeri county, Kenya.

87 **Materials and Methods**

88 A facility-based, cross-sectional study was conducted at the outpatient department of
89 Nyeri County referral hospital in Kenya. The study duration was from 1st August 2022

90 till 30th October 2022. In this study, patients of both genders, of age 20 years and
91 above, known cases of type II DM visiting the hospital for routine follow-up visit were
92 included. Patients being seen in the emergency or in-patient department, patients who
93 did not have at least one available record of HbA1c from the last six months were
94 excluded. All patients were included after attaining informed consent. A semi-structured
95 questionnaire was used to record patient information. It included biodata information,
96 Duration of DM and treatment and HbA1c record from the last six months.

97 In order to assess their status of self-care, Summary of Diabetes self-care activities
98 (SDSCA) Questionnaire was used. The seven items of SDSCA are used to assess self-care
99 activities. They are divided into 7 subscales – blood glucose testing subscale (item
100 7,8,3A), Diet sub scale (item 1,2,3,4,1A,5A), Physical exercise subscale (item 5,6, 2A),
101 smoking subscale (item 11,12A,13A, 14A), foot care subscale (item 9,10,9A,10A,11A),
102 medications subscale (item 6A,7A,8A) and self-care recommendations subscale (item
103 1A,2A,3A,4A). The questionnaire ask patient about their diabetes self-care activities
104 during the past 7 days. If they were sick during the past 7 days, we went back to the last
105 7 days that they were not sick.

106 Scoring instructions were. Step 1: For items 1–10, use the number of days per week on a
107 scale of 0–7. Step 2: Scoring Scales; General Diet = Mean number of days for items 1
108 and 2 and Specific Diet = Mean number of days for items 3, and 4 where the scale was
109 (0=7, 1=6, 2=5, 3=4, 4=3, 5=2, 6=1,7=0); Exercise = Mean number of days for items 5
110 and 6; Blood-Glucose Testing = Mean number of days for items 7 and 8. Foot-Care =
111 Mean number of days for items 9 and 10; Smoking Status = Item 11 (0 = nonsmoker,1 =
112 smoker), and number of cigarettes smoked per day.

113 Scoring for Additional Items such as; Recommended regimen = Items 1A - 4A, and
114 items 12A - 14A, no scoring required; Diet = Use total number of days for item 5A;
115 Medications = Use item 6A - OR - 7A AND 8A, use total number of days for item 6A, use
116 mean number of days if both 7A and 8A are applicable. Lastly; Foot-Care = Mean
117 number of days for items 9A - 11A, after reversing 10A and including items 9 and 10.

118 For data entry and statistical analysis SPSS for Windows version 27 was utilized. Patient
119 characteristics were presented as frequency and percentages and compared using Chi
120 square test. For SDSCA overall and its subscales mean and standard deviation (SD) were
121 calculated and for comparison independent sample t-test was applied. The p value
122 ≤ 0.05 was taken as significant.

123 **Ethics statement**

124 Both written and verbal informed consent for participation was also obtained from all
125 the participants. For written consent all participants signed the consent form in the
126 questionnaire. Verbal consent was witnessed by the nurse or clinical officer in-charge of
127 the clinic. The study was approved by the Ethical Review Committee of Jomo Kenyatta
128 University of Agriculture and Technology, and the National Commission for Science,
129 Technology and Innovation (NACOSTI/P/22/18990)

130 **Results**

131 **Participants characteristics, both overall and according to the presence of**
132 **the microvascular complications of diabetes**

133 Two hundred and forty-five people with Diabetes (PWD) participated in the study.
 134 Among them, 96 (39.2%) were males and 149 (60.8%) were females. Most of the
 135 participants were more than 61 years 148 (60.4%). Prevalence of Diabetes microvascular
 136 complications among the study participants were 56.78%, 25.3% and 14.7% for
 137 neuropathy, retinopathy and nephropathy respectively. Significant association was
 138 found between age and level of education with all the three microvascular
 139 complications. Furthermore, marital status was associated nephropathy and neuropathy
 140 whereas Hb1Ac was associated with nephropathy (Table 1).

Table 1: Participants socio-demographic characteristics association with the presence of the microvascular complications of diabetes							
Characteristic variable	Total (n=245)	Neuropathy (n= 139; 56.7%)	P value	Retinopathy (n= 62; 25.3%)	P value	Nephropathy (n= 36 ;14.7%)	P value
Age							
20-35	12 (4.9%)	1(0.7)	0.001	0 (0%)	0.009	2(5.6%)	0.02
36-45	21 (8.6%)	7(5.0%)		3(4.8%)		1(2.8%)	
46-60	64 (26.1%)	27(19.4)		11(17.7%)		3(8.3%)	
61 and above	148 (60.4%)	104(74.8)		48(77.4%)		30(83.3%)	
Sex							
Male	96 (39.2%)	51 (36.7%)	0.36	19 (30.6%)	0.11	14 (38.9%)	0.97
Female	149 (60.8%)	88 (63.3%)		43 (69.4%)		22 (61.1%)	
Marital status							
Single	23 (9.4%)	7 (5.0%)	0.003	3 (4.8%)	0.08	2 (5.6%)	0.007
Married	168 (68.6%)	93 (66.9%)		39 (62.9%)		20 (55.6%)	
Divorced/Separated	8 (3.8%)	4 (2.9%)		2 (3.2%)		0 (0%)	
Widowed	46 (18.8%)	35 (25.2%)		18 (29.0%)		14 (38.9%)	
Level of Education							
Primary	140 (57.1%)	83 (59.7%)	0.003	39 (62.9%)	0.003	19 (52.8%)	0.008
Secondary	64 (26.1%)	28 (20.1%)		7 (11.3%)		6 (16.7%)	
Tertiary	13 (5.3%)	5 (3.6%)		3 (4.8%)		1 (2.8%)	

No schooling	28 (11.4%)	23 (16.1%)		13 (21.0%)		10 (27.8%)	
Occupation							
Office job	9 (3.7%)	3 (2.2%)	0.17	3 (4.8%)	0.13	0 (0%)	0.06
Outdoor job	30 (12.2%)	13 (19.4%)		4 (6.5%)		5 (13.9%)	
Farmer	175 (71.4%)	104 (74.8%)		43 (69.4%)		22 (61.1%)	
Stay at home	31 (12.7%)	19 (13.7%)		12 (19.4%)		9 (25.0%)	
Duration of diabetes							
Below 5 years	76 (31.0%)	35 (25.2%)	0.05	19 (30.6%)	0.26	8 (22.2%)	0.44
5-10 years	89 (36.3%)	52 (37.4%)		18 (29.1%)		14 (38.9%)	
More than 10 years	80 (32.7%)	52 (37.4%)		25 (40.3%)		14 (38.9%)	
Diabetes treatment regimen							
Oral Hypoglycemics	165 (67.3%)	96 (69.1%)	0.73	41 (66.1%)	0.90	21 (58.3%)	0.27
Insulin	24 (9.8%)	12 (8.6%)		7 (11.3%)		6 (16.7%)	
Insulin plus oral hypoglycemics	56 (22.9%)	31 (22.3%)		14 (22.6%)		9 (25.0%)	
Hb1aC level							
Good control	101 (41.2%)	56 (40.3%)	0.73	21 (33.9%)	0.17	8 (22.2%)	0.01
Poor Control	144 (58.8%)	83 (59.7%)		41 (66.1%)		28 (77.8%)	
Values are presented as number (%) and P value for associations.							

141

142 **Associations between Summary of diabetes self-care activities and the**
 143 **microvascular complications of diabetes**

144 As shown in Table 2, a significant association was found between the sum scale scores of
 145 dietary activities, blood glucose testing, physical activity, foot care and neuropathy at
 146 95% CI and ($P < 0.001$). Scores on the Blood glucose testing and foot care activities
 147 subscale were significantly associated with Retinopathy ($p < 0.001$). Additionally,
 148 significant associations were found between the presence of nephropathy and dietary
 149 activities, foot care and physical activity subscale ($p < 0.001$) (Table 2).

Table: 2 Associations between Summary of diabetes self-care activities and the microvascular complications of diabetes

SDSAC Variable	Total (n=245) mean and SD	Neuropathy OR (95% CI)	P value	Retinopathy OR (95% CI)	P value	Nephropathy OR (95% CI)	P value
Dietary Practice	4.44 ±1.51	0.07 (0.03,0.11)	<0.001*	0.04 (-0.001,0.07)	0.06	0.06 (0.03,0.08)	<0.001*
Blood glucose monitoring	1.52 ±2.15	0.06 (0.03,0.09)	<0.001*	0.03 (0.02,0.07)	<0.001*	-0.003 (-0.02,0.02)	0.76
Physical Activity	2.39 ±1.86	0.06 (0.03,0.09)	<0.001*	0.03 (0.003,0.06)	0.08	0.05 (0.03,0.08)	<0.001*
Foot care	4.99 ±1.63	0.08 (0.02,0.09)	<0.001*	0.06 (0.02,0.09)	<0.001*	0.06(0.03,0.08)	<0.001*
DM Medication Adherence	7.28± 1.55	-0.02(-0.07,0.02)	0.29	0.01 (-0.03,0.05)	0.55	0.007 (0.02,0.04)	0.62

SD, standard deviation; OR, odds ratio; CI, confidence interval.
*p<0.05.

150

151

Discussion

152 In this study, we sought to assess associations between diabetes self-care activities and
 153 the microvascular complications of diabetes in patients with diabetes living in rural
 154 area. In our study, Prevalence of Diabetes microvascular complications among the
 155 study participants were 56.78%, 25.3% and 14.7% for neuropathy, retinopathy and
 156 nephropathy respectively. The proportion is higher as compared to other
 157 studies(27,28). This study mainly focused on participants living in the rural area unlike
 158 other studies. This difference could be because rural populations with diabetes
 159 experience unique challenges and substantial inequities in the achievement of diabetes
 160 care(29).

161 In the present study, participants were further categorized into strata according to their
 162 various demographic characteristics. It was observed that patients who were older than

163 sixty-one years were more likely to have Neuropathy, Retinopathy compared to those
164 who were younger ($p < 0.05$). This finding is consistent with other studies that have
165 shown age is an important factor that affects the increased risk of prediabetes and
166 diabetes (30). Additionally, microvascular events occur more commonly in populations
167 with diabetes and the risk of these events are predominantly related to duration of
168 diabetes(31).

169 Level of education was associated with all the MVCs. This study agrees with other
170 studies that have revealed educational status influences the awareness about diabetes
171 self-care, compliance to drugs and the health seeking behavior of an individual (32).
172 Studies have observed that lower the education higher is the risk of developing diabetes
173 complications(33).

174 In the present study, it was observed that dietary practices and physical activity was
175 significantly associated neuropathy and nephropathy complications. This can be
176 explained by a consistent healthy dietary pattern is associated with a reduced risk of
177 microvascular complications in patients with diabetes. In this study we also assessed
178 carbohydrate spacing and fats intake; studies have shown that high carbohydrate and
179 high monounsaturated fat diets improve insulin sensitivity (34–36). These findings
180 provide a reasonable basis for dietary recommendations aimed at preventing diabetic
181 microvascular complications.

182 Physical activity such as walking, gardening, leisure activity, exercise, or movement for
183 transportation or an occupation among other aerobic and resistance exercise are known
184 to improve insulin sensitivity. Additionally, it improves the lipid profile, BMI and blood

185 pressure. This study is consistent with other study findings that have reported that;
186 physical activity improves glycemic control and better disease control(37–39).

187 Foot care practices was significantly associated with all the three microvascular
188 complications. Others studies have demonstrated that patients with peripheral
189 neuropathy presented with significantly higher rates of development of DR,
190 macroalbuminuria and the existence of macrovascular complications (40–42). The
191 results suggest that neuropathy, nephropathy and retinopathy are signs of a generalized
192 diabetic microangiopathic process whose progression may be influenced by factors other
193 than diabetes duration and metabolic and therefore when a patient takes measures to
194 prevent one micro vascular complication it has direct impact to prevent of the similar
195 complications (43,44).

196 The findings of association between blood glucose testing and neuropathy and
197 retinopathy in this study are in agreement with previous literature. Blood glucose
198 monitoring is an essential part of management in patients with diabetes mellitus (45).
199 It helps to identify patterns in the fluctuation of blood glucose levels that occur in
200 response to diet, exercise, medications, and pathological processes. It enhances early
201 intervention in case of hypoglycemia or hyperglycemia which impacts on various body
202 system such as the retina and neurons (46,47). It supports improvement of patient
203 outcomes because critical decision can be made in time.

204 **Study Limitations**

205 This study is a cross-sectional study hence cannot establish causal inferences but rather
206 associations. Nonetheless, this study provides the extent of association between diabetes
207 self-management and MVCs to inform development of interventions.

208 **Conclusion**

209 The present study indicates that the diabetes-related self-care activities was directly
210 linked with diabetes microvascular complications. Foot care, dietary practices, physical
211 activity, blood glucose testing, Age, marital status, Hb1Ac and level of education were
212 significantly associated with MVCs among patients. It is essential to promote Diabetes
213 self-care education related to DM as a strategy to improve self-management and
214 improve the overall patient outcome.

215 **Acknowledgments**

216 Gratitude goes to workers at Diabetic Clinic and authorities of Nyeri county Referral
217 Hospital for their great support during the study.

218 **Financial obligation**

219 No funding was obtained for this research.

220 **Conflict of Interest**

221 None to Declare

222 **Authors' contributions**

223 Rachael Ileri contributed to the conception of the research idea, design data analysis
224 and interpretation, paper drafting and revision. Gideon Kikvi contributed to the

225 conception of the research idea, design and revision of the final draft. Susan Mambo
226 contributed to the conception of the research idea, design, and revision of final draft.
227 Besty Cheriro contributed to the conception of the research idea, design, and revision of
228 final draft. All authors approved the final manuscript before publication and agree to be
229 accountable for all aspects of the work.

230 REFERENCES

- 231 1. Mathur P, Leburu S, Kulothungan V. Prevalence, Awareness, Treatment and
232 Control of Diabetes in India From the Countrywide National NCD Monitoring
233 Survey. *Front Public Heal.* 2022 Mar 14;10.
- 234 2. Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, et al. IDF
235 Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates
236 for 2021 and projections for 2045. *Diabetes Res Clin Pract.* 2022 Jan 1;183.
- 237 3. Marsh Z, Teegala Y, Cotter V. Improving diabetes care of community-dwelling
238 underserved older adults. *J Am Assoc Nurse Pract.* 2022 Oct 7;34(10):1156–66.
- 239 4. Tang J, Wu T, Hu X, Gao L. Self-care activities among patients with type 2
240 diabetes mellitus: A cross-sectional study. *Int J Nurs Pract.* 2021 Dec 1;27(6).
- 241 5. Alanazi M. Determinants of successful diabetes self-management behaviors
242 among women of Arab descent with Type 2 Diabetes. *Prim Care Diabetes.* 2021
243 Apr 1;15(2):306–13.
- 244 6. Ary D V., Toobert D, Wilson W, Glasgow RE. Patient perspective on factors
245 contributing to nonadherence to diabetes regimen. *Diabetes Care.* 1986;9(2):168–

- 246 72.
- 247 7. Shrivastava SRBL, Shrivastava PS, Ramasamy J. Role of self-care in management
248 of diabetes mellitus. *J Diabetes Metab Disord*. 2013 Mar 5;12(1).
- 249 8. Lin K, Yang X, Yin G, Lin S. Diabetes Self-Care Activities and Health-Related
250 Quality-of-Life of individuals with Type 1 Diabetes Mellitus in Shantou, China. *J*
251 *Int Med Res*. 2016 Feb 1;44(1):147–56.
- 252 9. Aquino JA, Baldoni NR, Flôr CR, Sanches C, Di Lorenzo Oliveira C, Alves GCS, et
253 al. Effectiveness of individual strategies for the empowerment of patients with
254 diabetes mellitus: A systematic review with meta-analysis. *Prim Care Diabetes*.
255 2018 Apr 1;12(2):97–110.
- 256 10. Aronson R, Brown RE, Jiandani D, Walker A, Orzech N, Mbuagbaw L. Assessment
257 of self-management in patients with diabetes using the novel LMC Skills,
258 Confidence and Preparedness Index (SCPI). *Diabetes Res Clin Pract*. 2018 Mar
259 1;137:128–36.
- 260 11. Afaya RA, Bam V, Lomotey AY, Afaya A. Clinical factors influencing knowledge
261 and self-care practice among adults with type 2 diabetes mellitus. *Nurs Open*.
262 2023 Apr 1;10(4):2492–500.
- 263 12. Khosravizadeh O, Ahadinezhad B, Maleki A, Yousefy S, Momeni Z. Diabetes self-
264 care activities among patients with type 2 diabetes: A systematic review and meta-
265 analysis. *Int J Diabetes Dev Ctries*. 2023;
- 266 13. Eva JJ, Kassab YW, Neoh CF, Ming LC, Wong YY, Hameed MA, et al. Self-care

- 267 and self-management among adolescent T2DM patients: A review. *Front*
268 *Endocrinol (Lausanne)*. 2018 Oct 18;9(OCT).
- 269 14. Maina PM, Pienaar M, Reid M. Self-management practices for preventing
270 complications of type II diabetes mellitus in low and middle-income countries: A
271 scoping review. *Int J Nurs Stud Adv*. 2023 Dec 1;5.
- 272 15. Young-Hyman D, De Groot M, Hill-Briggs F, Gonzalez JS, Hood K, Peyrot M.
273 Psychosocial care for people with diabetes: A position statement of the American
274 diabetes association. *Diabetes Care*. 2016;39(12):2126–40.
- 275 16. Toobert DJ, Hampson SE, Glasgow RE. The summary of diabetes self-care
276 activities measure: Results from 7 studies and a revised scale. *Diabetes Care*.
277 2000;23(7):943–50.
- 278 17. Llera-Fábregas A, Pérez-Ríos N, Camacho-Monclova DM, Ramirez-Vick M,
279 Andriankaja OM. Diabetes self-care activities and perception and glycemic control
280 in adult Puerto Rican residents with Type 2 Diabetes: The LLIPDS Study. *J Public*
281 *health Res*. 2022 Oct 1;11(4).
- 282 18. Lee J, Lee EH, Chae D, Kim CJ. Patient-reported outcome measures for diabetes
283 self-care: A systematic review of measurement properties. *Int J Nurs Stud*. 2020
284 May 1;105.
- 285 19. Chew BH, Vos RC, Pouwer F, Rutten GEHM. The associations between diabetes
286 distress and self-efficacy, medication adherence, self-care activities and disease
287 control depend on the way diabetes distress is measured: Comparing the DDS-17,
288 DDS-2 and the PAID-5. *Diabetes Res Clin Pract*. 2018 Aug 1;142:74–84.

- 289 20. Pintaudi B, Lucisano G, Gentile S, Bulotta A, Skovlund SE, Vespasiani G, et al.
290 Correlates of diabetes-related distress in type 2 diabetes: Findings from the
291 benchmarking network for clinical and humanistic outcomes in diabetes
292 (BENCH-D) study. *J Psychosom Res.* 2015;79(5):348–54.
- 293 21. Kim S, Love F, Quistberg DA, Shea JA. Association of health literacy with self-
294 management behavior in patients with diabetes. *Diabetes Care.* 2004
295 Dec;27(12):2980–2.
- 296 22. Bloomgarden Z, Gouller A. Prevention of Microvascular Complications of
297 Diabetes. *Princ Diabetes Mellit.* 2004;619–38.
- 298 23. Salmerón J, Hu FB, Manson JE, Stampfer MJ, Colditz GA, Rimm EB, et al.
299 Dietary fat intake and risk of type 2 diabetes in women. *Am J Clin Nutr.*
300 2001;73(6):1019–26.
- 301 24. Marshall JA, Bessesen DH. Dietary fat and the development of type 2 diabetes.
302 *Diabetes Care.* 2002;25(3):620–2.
- 303 25. Glasgow RE, Osteen VL. Evaluating diabetes education: Are we measuring the
304 most important outcomes? *Diabetes Care.* 1992;15(10):1423–32.
- 305 26. Glasgow RE, Fisher EB, Anderson BJ, LaGreca A, Marrero D, Johnson SB, et al.
306 Behavioral science in diabetes: Contributions and opportunities. *Diabetes Care.*
307 1999 May;22(5):832–43.
- 308 27. Govindarajan Venguidesvarane A, Jasmine A, Varadarajan S, Shriraam V,
309 Muthuthandavan AR, Durai V, et al. Prevalence of Vascular Complications Among

- 310 Type 2 Diabetic Patients in a Rural Health Center in South India. *J Prim Care*
311 *Community Heal* [Internet]. 2020 [cited 2024 Jan 29];11. Available from:
312 <https://doi.org/10.1177/2150132720959962>
- 313 28. Pelluri R, Srikanth K, Chimakurthy J, Nagasubramanian VR. Microvascular
314 Complications and Their Associated Risk Factors Among Rural Type 2 Diabetic
315 Population: A Cross-Sectional Study. *SN Compr Clin Med* 2021 32 [Internet].
316 2021 Feb 5 [cited 2024 Jan 29];3(2):625–31. Available from:
317 <https://link.springer.com/article/10.1007/s42399-021-00786-7>
- 318 29. Flood D, Geldsetzer P, Agoudavi K, Aryal KK, Campos L, Brant C, et al. Rural-
319 Urban Differences in Diabetes Care and Control in 42 Low-and Middle-Income
320 Countries: A Cross-sectional Study of Nationally Representative Individual-Level
321 Data. *Diabetes Care* [Internet]. 2022 [cited 2024 Jan 29];45. Available from:
322 <https://doi.org/10.2337/dc21-2342>
- 323 30. Yan Z, Cai M, Han X, Chen Q, Lu H. The Interaction Between Age and Risk
324 Factors for Diabetes and Prediabetes: A Community-Based Cross-Sectional Study.
325 *Diabetes, Metab Syndr Obes* [Internet]. 2023 [cited 2024 Jan 30];16:85–93.
326 Available from:
327 <https://www.tandfonline.com/action/journalInformation?journalCode=dms020>
- 328 31. Singh Jaggi A, Parkash Singh V, Bali A, Singh N. Advanced Glycation End
329 Products and Diabetic Complications. *Korean J Physiol Pharmacol* [Internet].
330 2014 [cited 2024 Jan 30];18:1–14. Available from:
331 <http://dx.doi.org/10.4196/kjpp.2014.18.1.1>

- 332 32. Chawla SPS, Kaur S, Bharti A, Garg R, Kaur M, Soin D, et al. Impact of health
333 education on knowledge, attitude, practices and glycemc control in type 2
334 diabetes mellitus. *J Fam Med Prim Care* [Internet]. 2019 [cited 2024 Jan
335 30];8(1):261. Available from: [/pmc/articles/PMC6396605/](#)
- 336 33. Hwang Y, Lee D, Kim YS. Educational Needs Associated with the Level of
337 Complication and Comparative Risk Perceptions in People with Type 2 Diabetes.
338 *Osong Public Heal Res Perspect* [Internet]. 2020 Aug 1 [cited 2024 Jan
339 30];11(4):170. Available from: [/pmc/articles/PMC7442446/](#)
- 340 34. Ghaemi F, Firouzabadi FD, Moosaie F, Shadnoush M, Poopak A, Kermanchi J, et
341 al. Effects of a Mediterranean diet on the development of diabetic complications:
342 A longitudinal study from the nationwide diabetes report of the National Program
343 for Prevention and Control of Diabetes (NPPCD 2016-2020). *Maturitas*. 2021 Nov
344 1;153:61–7.
- 345 35. Rydall AC, Rodin GM, Olmsted MP, Devenyi RG, Daneman D. Disordered Eating
346 Behavior and Microvascular Complications in Young Women with Insulin-
347 Dependent Diabetes Mellitus. *N Engl J Med*. 1997 Jun 26;336(26):1849–54.
- 348 36. Chudasama Y V., Khunti K. Healthy lifestyle choices and microvascular
349 complications: New insights into diabetes management. *PLoS Med*. 2023 Jan
350 1;20(1).
- 351 37. Zhu X, Zhang X, Zhou C, Li B, Huang Y, Li C, et al. Walking pace and
352 microvascular complications among individuals with type 2 diabetes: A cohort
353 study from the UK Biobank. *Scand J Med Sci Sport*. 2023 Jan 1;

- 354 38. Makura CBT, Nirantharakumar K, Girling AJ, Saravanan P, Narendran P. Effects
355 of physical activity on the development and progression of microvascular
356 complications in type 1 diabetes: Retrospective analysis of the DCCT study. *BMC*
357 *Endocr Disord.* 2013 Oct 2;13.
- 358 39. Blomster JI, Chow CK, Zoungas S, Woodward M, Patel A, Poulter NR, et al. The
359 influence of physical activity on vascular complications and mortality in patients
360 with type 2 diabetes mellitus. *Diabetes, Obes Metab.* 2013;15(11):1008–12.
- 361 40. Vithian K, Hurel S. Microvascular complications: pathophysiology and
362 management. *Clin Med (Northfield Il)* [Internet]. 2010 [cited 2024 Jan
363 30];10(5):505. Available from: [/pmc/articles/PMC4952418/](https://pubmed.ncbi.nlm.nih.gov/2418495/)
- 364 41. Mansour A, Mousa M, Abdelmannan D, Tay G, Hassoun A, Alsafar H.
365 Microvascular and macrovascular complications of type 2 diabetes mellitus:
366 Exome wide association analyses. *Front Endocrinol (Lausanne).* 2023;14.
- 367 42. Tuomilehto J, Schwarz P, Lindström J. Long-term benefits from lifestyle
368 interventions for type 2 diabetes prevention: Time to expand the efforts. *Diabetes*
369 *Care.* 2011 May;34(SUPPL. 2).
- 370 43. Geng T, Zhu K, Lu Q, Wan Z, Chen X, Liu L, et al. Healthy lifestyle behaviors,
371 mediating biomarkers, and risk of microvascular complications among individuals
372 with type 2 diabetes: A cohort study. *PLoS Med.* 2023 Jan 1;20(1).
- 373 44. Zhang Y, Pan XF, Chen J, Xia L, Cao A, Zhang Y, et al. Combined lifestyle factors
374 and risk of incident type 2 diabetes and prognosis among individuals with type 2
375 diabetes: a systematic review and meta-analysis of prospective cohort studies.

- 376 Diabetologia. 2020 Jan 1;63(1):21–33.
- 377 45. Turner R. Intensive blood-glucose control with sulphonylureas or insulin
378 compared with conventional treatment and risk of complications in patients with
379 type 2 diabetes (UKPDS 33). Lancet. 1998 Sep 12;352(9131):837–53.
- 380 46. Dahl-Jørgensen K, Brinchmann-Hansen O, Hanssen KF, Sandvik L, Aagenæs.
381 Rapid tightening of blood glucose control leads to transient deterioration of
382 retinopathy in insulin dependent diabetes mellitus: The Oslo study. Br Med J
383 (Clin Res Ed). 1985 Mar 16;290(6471):811–5.
- 384 47. Stratton IM, Adler AI, Neil HAW, Matthews DR, Manley SE, Cull CA, et al.
385 Association of glycaemia with macrovascular and microvascular complications of
386 type 2 diabetes (UKPDS 35): prospective observational study. BMJ Br Med J
387 [Internet]. 2000 Aug 8 [cited 2024 Jan 30];321(7258):405. Available from:
388 /pmc/articles/PMC27454/
389