1	Title: Non-Consensual Sex among Japanese Women in the COVID-19 Pandemic: A Large-
2	Scale Nationwide Survey-Based Study
3	
4	Authors:
5	Tomoya Suzuki, ¹ Yudai Kaneda, ² Yasuhiro Kotera, ³ Akihiko Ozaki, ^{1,4} Tetsuya Tanimoto, ^{1,5}
6	Divya Bhandari, ¹ Sayaka Horiuchi, ⁶ Takahiro Tabuchi ⁷
7	
8	Affiliations:
9	1. Medical Governance Research Institute, Minato, Tokyo, Japan
10	2. School of Medicine, Hokkaido University, Sapporo, Hokkaido, Japan
11	3. Faculty of Medicine and Health Sciences, University of Nottingham, Nottingham, UK
12	4. Department of Breast and Thyroid Surgery, Jyoban Hospital of Tokiwa Foundation, Iwaki,
13	Fukushima, Japan
14	5. Department of Internal Medicine, Navitas Clinic, Tachikawa, Tokyo, Japan
15	6. Graduate Faculty of Interdisciplinary Research Faculty of Medicine Basic Medicine
16	(Epidemiology/Environmental Medicine), University of Yamanashi
17	7. Cancer Control Center, Osaka International Cancer Institute, Osaka, Japan

19	Corresponding author:
20	Tomoya Suzuki
21	Medical Governance Research Institute, Minato, Tokyo, Japan
22	Tel: +81-80-9873-8361
23	E-mail: tomo1502@gmail.com
24	
25	Acknowledgments:
26	None
27	
28	Authors' Contributions
29	Conceptualization: TS, AO, and TakahiroT
30	Data analysis: TS and AO
31	Writing original draft: TS and AO
32	Funding acquisition: AO and TakahiroT
33	Project Administration: AO and TakahiroT
34	Review and editing: YudaiK, TetsuyaT, YasuhiroK, BD, SH, and TakahiroT
35	All authors read and approved the final manuscript for publication.

37 **Funding Information** 38 This study was funded by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grants (grant number 17H03589; 19K10671; 19K10446; 18H03107; 18H03062; 21H04856], 39 40 the JSPS Grant-in-Aid for Young Scientists (grant number 19K19439), Research Support Program to Apply the Wisdom of the University to tackle COVID-19-Related Emergency 41 42 Problems, University of Tsukuba and Health Labor Sciences Research Grant (grant number 19FA1005; 19FG2001]. 43 44 45 **Conflict of interest statement:** TT reports personal fees from Medical Network Systems, MNES Inc., and Bionics co. ltd., 46 outside the submitted work. AO reports personal fees from Medical Network Systems, MNES 47 48 Inc., outside the submitted work. All remaining authors have nothing to disclose. 49 50 **Keywords:** COVID-19, Non-consensual sex, sex consent, sexual abuse, sexual violence, rape, sexual education 51

53 **ABSTRACT Background:** Non-consensual sex including rape and sexual assault has been a global concern 54 and may have been influenced by the COVID-19 pandemic, however the information on this 55 56 topic is limited. Therefore, our objective was to survey the incidence rate of non-consensual sex among Japanese women aged 15-79 years between April to September 2020, following the 57 COVID-19 pandemic in Japan. 58 59 Materials and Methods: We utilized the data obtained from a nationwide, cross-sectional 60 61 internet survey conducted in Japan between August and September 2020. Sampling weights 62 were applied to calculate national estimates, and multivariable logistic regression was 63 performed to identify factors associated with non-consensual sex. Data was extracted from a cross-sectional, web-based, self-administered survey of approximately 2.2 million individuals 64 from the general public, including in men and women. 65 66 **Results:** Excluding men and responses with inconsistencies, the final analysis included 12,809 67 women participants, with 138 (1.1%) reporting experiencing non-consensual sex within a five-68 69 month period. Being aged 15–29 years and having a worsened mental or economic status were

70

associated with experiencing non-consensual sex.

Conclusions: Early intervention to prevent individuals from becoming victims of sexual harm should be extended to economically vulnerable and young women, especially during times of societal upheaval such as the COVID-19 pandemic. Additionally, Japan should prioritize the implementation of comprehensive education on the concept of sexual consent.

INTRODUCTION

Non-consensual sex refers to engaging in sexual behavior without obtaining consent to touch, observe, or perform sexual acts involving private body parts that exceed the boundaries of another person's body or mind (1,2). Non-consensual sex can take many forms, including rape, transactional sex, cross-generational sex, unwanted touch, and molestation. In the European Union, between 45% to 55% of women have experienced sexual harassment since the age of 15 (3). Previous research has identified common factors associated with victims, including ages between 10 and 30 years, mental vulnerability, and financial hardship (4). Women with these characteristics are at a higher risk of experiencing non-consensual sex compared to those without these traits (4). Furthermore, women with these characteristics often find it challenging to disclose incidents of non-consensual sex to others, making it difficult to address these issues (5).

Of note, non-consensual sex is reported to increase due to disruptions in social conditions, such as natural disasters and worsened economic conditions. For instance, incidents of non-consensual sex against women dramatically increased during times of economic and psychological turmoil, such as the 2005 Hurricane Katrina in the United States and the 2011 Great East Japan Earthquake (6–8). Therefore, it is crucial to investigate the prevalence of non-

consensual sex after the COVID-19 pandemic.

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

aged 15-49 years had been subjected to non-consensual sex worldwide (9). This was attributed to the increase in the mandatory stay-at-home measure due to lockdowns, and resulting in

At the onset of the COVID-19 pandemic, it was reported that 243 million girls and women

reduced access to social support from teachers, friends and caregivers (10). Awareness of the

status of non-consensual sex during the COVID-19 pandemic has grown in many countries

(11,12). However, the frequency and characteristics of non-consensual sex remain to be

elucidated, not only after but also before the COVID-19 pandemic in Japan to date, partly due

to traditional social norms that discourage open discussions about sexual issues (13).

The purpose of our study was to report the incidence of non-consensual during the early stages

of the COVID-19 pandemic in Japan. Additionally, we explored potential risk factors such as

mental health conditions, economic changes, and concerns about the lockdown, and anxiety

about the disease.

METHODS

Context

This cross-sectional study was conducted among a sample of approximately 2.2 million

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

perpetuity.

It is made available under a CC-BY 4.0 International license .

8

individuals from the general public who participated in a web-based self-reported questionnaire survey for the Japan COVID-19 and Society Internet Survey (JACSIS) project (14). The survey was administered by the well-established internet research agency, Rakuten Insight, Inc., which has been utilized in prior research studies (14,15). Data were collected from April 2020 to September 30, 2020. The prefectures where the selected individuals lived represented all prefectures in Japan, as indicated in previous studies (16). Patient and public involvement No patients were involved in this study. **Participants** The survey invitation was extended to 28,000 individuals out of 2.2 million registered with the survey agency, Rakuten Insight, Inc. Using a computer algorithm, study participants were selected through random sampling. **Analysis Subjects** Respondents who self-identified as women in the questionnaire were included in our analysis. Among all the survey respondents totaling 28,000, the number of valid women participants for

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

Outcome Variable: Experience of Non-Consensual Sex

The focus of this study was to examine instances of non-consensual sex occurring between from April 2020 to 30 September 2020. Participants were not questioned about specific details but were instead asked to respond with a simple "yes" or "no" to indicate whether they had experienced such incidents.

Exposure Variables

Sociodemographic characteristics

As independent variables, demographics were included, encompassing gender, age groups (15-19, 20-29, 30-39, 40-49, 50-59, 60-69, 70-79 years), marital status (unmarried, married and widowed/separated), having children (none or more than one), educational attainment (high school educated, college educated or higher), employment status (any type of employment, unemployed), household income level, which was calculated as dividing the household income by the square root of household size (categorized by the tertials of household equivalent income (low, <JPY2.5 million/US\$25 000/£16 667; intermediate, JPY2.5-JPY4.3 million/US\$25 000-US\$43 000/£16 667-£28 667; high, >JPY4.3

million/<US\$43 000/<£28 667; unknown/declined to answer)), and smoking status (not smoker or current smoker).(14)

Living Area Classification

Regarding living regions, all 47 prefectures were divided into the following three areas based on the date of the Declaration of the State of Emergency (DSE area) (17,18). "Designated as DSE on April 7, 2020", "Designated as DSE on April 16, 2020", and "Others", "Specific alert designated on April 7, 2020" included Tokyo, Kanagawa, Chiba, Saitama, Osaka, Hyogo, and Fukuoka. "Specific alert designated on April 16, 2020" included Hokkaido, Ibaraki, Ishikawa, Gifu, Aichi, and Kyoto. "Others" comprised the remaining 34 prefectures. These categorization were determined by the Japanese Government using three indicators: the cumulative number of infected people, epidemiological trend, medical capacity and surveillance system (17). During the target period, people living in the relevant prefectures were advised to avoid unnecessarily outings, limit the use of entertainment facilities, and adhere to restrictions on economic activities (19).

Fear of COVID-19 Scale

The Japanese version Fear of COVID-19 Scale (FCV-19S) was used to assess participants'

fear of the COVID-19 infection (20). This instrument comprises seven items rated on a 5-point scale (1 = "strongly disagree"; 5 = "strongly agree"). Consequently, the total score ranged from 7 to 35, with a higher score indicating a greater fear of COVID-19. Satisfactory internal consistency and validation of the scale were confirmed in the original seven-item scale (α = .82) (21).

Personal Health and Economic Statuses

Based on previous research, an examination was conducted to determine whether health and economic statuses were associated with non-consensual sex (4). The health-related questions included the following: Self-rated health (good or other than good), "Any change in mental state in the last month compared to before January 2020?" (Worse, No change, Getting better, I do not know)", Suicidal thoughts (Happened since before COVID-19 pandemic, First experienced during the COVID-19 pandemic or Never), Desire not to talk to anyone due to worries (Yes or No), Feeling isolated (Yes or No), Any cancellation of a family gathering due to the COVID-19 pandemic (Yes or No), Non-payment of salary (Happened since before COVID-19 pandemic, First happened during the COVID-19 pandemic, Never), and Lack of money to buy necessities of life (Happened since before COVID-19 pandemic, First Happened during the COVID-19 pandemic,

187 188 **Statistical Procedure** To achieve the purpose of identifying the incidence of non-consensual sexual activity and its 189 associated factors during the COVID-19 outbreak, two types of analysis methods were 190 191 employed. 192 193 First, to determine the incidence rate, we divided the sample into two groups: those who responded "Yes" to the question of "Any non-consensual sex" and who did not. The number 194 195 and percentage of individuals in these two categories, along with the χ -square test, were 196 calculated for each independent variable. The number, percentage, and p-value of each item are described in Table 1. 197 198 199 Second, a multivariate logistic regression analysis was conducted to identify potential 200 predictive factors in the occurrence of non-consensual sex. This analysis encompassed all the aforementioned exposure variables, and odds ratios with 95% confidence intervals (CIs) were 201 202 estimated (Table 2). As a sub-analysis, the incidence of suicidal ideation by age was also 203 calculated.

206

207

208

209

210

211

212

213

214

215

216

217

218

219

13

It should be noted that because the characteristics of the participants in this Internet survey may differ from the general population (i.e., individuals with low digital skills might be excluded), a weighted analysis was performed to minimize differences in demographics, economic status, and health-related characteristics. This approach aimed to approximate national estimates from a nationally representative survey. Statistical significance was defined as p<0.05. The data were analyzed using STATA V.16.1 (Stata Corp, College Station, Texas, USA). **Ethics** This study adhered to ethical standards in accordance with the 1975 Declaration of Helsinki, as revised in 2013. Prior to participating in the online survey, all participants provided informed consent through a web-based consent form. The Research Ethics Committee of the Osaka International Cancer Institute reviewed and approved the study (No. 20084).

Results

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

The total number of respondents, including both men and women, was 28,000. Out of these, 2,518 participants who provided invalid responses (e.g., the same number is used for all survey items; incomplete responses) and 12,673 men were excluded. Consequently, the final analysis sample consisted of 12,809 women, whose ages ranged from 15 and 79 years old.

Table 1 shows the characteristics of the weighted proportion of non-consensual sex among all participants. The overall weighted incidence rate for non-consensual sex during the five-month period (April to September 2020) was 1.1 %. The incidence rate of non-consensual sex was higher among individuals aged 15-19 years (1.1%) and 20-29 years (2.4%), which exceeded the rates in other age groups. Women with jobs (1.5%) had a higher incidence rate than those without jobs (0.7%).

Table 1: Demographics and Descriptive Statistics of Weighted Women Participants during the Five-Month from April to September 2020

Non-Consensual Sex April-Sept 2020

	Total (n	=12,809)
X7 : 11	Yes	No
Variable	N (%)	N (%)
Total	138 (1.1)	12,671 (98.9)
Age		, ,
15-19	8.0 (1.1)	707 (98.9)
20-29	37 (2.4)	1543 (97.7)
30-39	38 (2.0)	1845 (98.0)
40-49	25 (1.0)	2395 (98.9)
50-59	15 (0.7)	2097 (99.3)
60-69	15 (0.7)	2127 (99.3)
70-79	1.0 (0.1)	1956 (99.9)
Marital status	` '	. ,
Married	88 (1.1)	7929 (98.9)
Never married	29 (0.9)	3276 (99.1)
Widowed or separated	20 (1.4)	1467 (98.7)

Having children		
No children	56 (0.9)	6006 (99.1)
At least one or more	99 (1.5)	6648 (98.5)
Educational attainment	99 (1.3)	0048 (38.3)
High school or lower	54 (1.0)	4261 (99.0)
College/university or graduate school	83 (1.2)	7136 (98.9)
Employment status	83 (1.2)	/130 (36.3)
Unemployed	43 (0.7)	6259 (99.3)
In employment	95 (1.5)	6412 (98.5)
Smoking status	93 (1.3)	0412 (96.3)
Not smoker	117 (1.0)	11404(99.0)
Current smoker	` '	, ,
	20 (1.6)	1268 (98.4)
Household Income (million JPY)	52 (1.7)	2144 (00.2)
Low	53 (1.7)	3144 (98.3)
Moderate	43 (1.3)	3226 (98.7)
High	26 (1.0)	2669 (99.0)
Unknown	14 (0.4)	3632 (99.6)
Region	06 (1.2)	((20,(00.7)
Other	86 (1.3)	6628 (98.7)
DSE (April 16-)	16 (0.8)	2117 (99.2)
DSE (April 7-)	31 (0.9)	3568 (99.1)
Fear of COVID-19 [†]	41 (1.2)	2402 (00.0)
7–15	41 (1.2)	3402 (98.8)
16–20	26 (0.6)	4173 (99.4)
21–25	34 (1.0)	3405 (99.0)
26–35	37 (2.1)	1691 (97.9)
Self-rated health		
Other than good	94 (1.3)	7349 (98.7)
Good	44 (0.8)	5322 (99.2)
Mental state change in the last month compared to before January 2020		
Worse	59 (2.2)	2662 (97.8)
None	56 (0.6)	8998 (99.4)
Better	19 (3.3)	547 (96.7)
Unknown	4 (0.7)	500 (99.3)
Suicidal thoughts		
Yes, since before COVID-19	43 (4.0)	1039 (95.1)
First happened	20 (3.5)	539 (96.5)
Never	75 (0.7)	11093(99.3)
Desire not to talk to anyone due to worries		
Yes	76 (4.0)	2487 (97.0)
No	62 (0.6)	10184 (99.4)
Feeling isolated		
Yes	74 (3.7)	1925 (96.3)
No	64 (0.6)	10747 (99.4)
Cancellation of a family gathering due to the pandemic		
Yes	10 (0.6)	1730 (99.4)
No	127 (1.2)	10940(98.9)
Non-payment of salary		
Yes, since before COVID-19	15 (10.3)	133 (89.7)
First experience	8 (10.4)	67 (89.6)
Never	115 (0.9)	12471 (99.1)
Lack of money to buy necessities		
Yes, since before COVID-19	35 (4.8)	692 (95.2)
First experience	22 (6.1)	340 (94.0)
Never	81 (0.7)	11639 (99.3)

[†] Classification of the Fear of COVID-19 scale scores refers to previous studies (22).

All estimates account for survey weights.

235

236 237

238

239

240

241

There was no significant difference in the incidence rate of non-consensual sex among those living in areas designated as DSE (0.9 % on April 7, 2020, 0.8 % on April 16, 2020, 1.3%

others). A higher FCV-19S (2.1%, [26-35]) was associated with a higher incidence rate

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

compared to other scores (1.2% [score from 7 to 15], 0.6% [score from 16 to 20], 1.0% [score from 21 to 25]). The incidence rate of participants who reported feeling "worse" (2.2%) or "getting better" (4.4%) before the COVID-19 pandemic was higher than those who "No change" (0.6%) and "I do not know" (0.7%). Participants who had suicidal thoughts (4.0% since before COVID-19, 3.5% first happened) had a higher incidence rate than those who had never experienced them (0.7%). Participants who reported feeling isolated (3.7%) had a higher incidence rate than those who had never felt isolated (0.6%). Additionally, participants who had experienced nonpayment of salary (10.3% since before COVID-19, 10.4% first experience) had a higher incidence rate than those who had never experienced it (0.9%). Similarly, participants who had experienced a lack of money to buy necessities (4.8% since before COVID-19, 6.1% first experience) had a higher incidence rate than those who had never experienced it (0.7%). Table 2 shows the factors associated with non-consensual sex among participants. Participants aged 15 to 19 years and those aged 20 to 29 years had higher odds of experiencing nonconsensual sex compared to participants in other aging groups. (Aged 15- 19 years old: OR 4.74, 95%CI 1.37 – 16.4, Aged 20-29 years old: OR 3.20, 95%CI 1.09 – 9.38). Employed women participants were associated with higher odds of experiencing non-consensual sex than

unemployed participants (OR 1.68, 95% CI 1.36-2.07).

261

263 Table 2: Factors Associated with Non-Consensual Sex Among Weighted Participants During the-Five-Month from April to September 2020 264

	Weighted	Weighted	Crude Odds	Adjusted rate	Adjusted odds ratio	P-value
	sample	incidence	(95% CI)	% (95% CI)	(95% CI)	P-value
Total	12,809	138 (1.1)				
Age	-	, ,				
15- 19	715	8.0 (1.1)	2.26(1.29, 3.94)	3.4 (1.3, 5.5)	4.74(1.37, 16.4)	0.014
20-29	1,580	37.1(2.4)	1.74(1.07, 2.81)	2.4 (1.3, 3.4)	3.2 (1.09, 9.38)	0.034
30-39	1,883	37.7 (2.0)	1.37(0.84, 2.23)	1.5 (0.9, 2.1)	1.88 (0.63, 5.55)	0.256
40-49	2,420	24.5 (1.0)	Reference	0.8 (0.3, 1.3)	Reference	-
50-59	2,112	14.5 (0.7)	0.43(0.22, 0.83)	0.8 (0.5, 1.0)	0.90 (0.37, 2.19)	0.822
60-69	2,142	14.8 (0.7)	0.49(0.26, 0.92)	1.1 (0.6, 1.7)	1.35 (0.45, 4.06)	0.585
70-79	1,957	1.0 (0.1)	0.23(0.10, 0.55)	0.05 (0.0, 0.1)	0.05 (0.02, 0.19)	< 0.001
Marital status	-,,	-10 (01-)	**=*(****, *****)	**** (***, ***)	**** (*****)	
Married	8,017	88.3 (1.2)	Reference	1.6 (1.2, 2.0)	Reference	
Never married	3,304	29.0 (0.8)	1.35(0.96, 1.90)	0.5 (0.1, 0.8)	0.26 (0.10, 0.70)	0.007
Widowed or separated	1,487	20.3 (1.2)	0.83(0.48, 1.43)	1.5 (0.8, 2.1)	0.92 (0.69, 1.24)	0.599
	1,407	20.3 (1.2)	0.05(0.40, 1.45)	1.5 (0.6, 2.1)	0.72 (0.07, 1.24)	0.377
Having children	6.202	42.9 (0.7)	Dafamanaa	0.9 (0.7, 0.0)	D ofonom a c	
No children	6,302	42.8 (0.7)	Reference	0.8 (0.7, 0.9)	Reference	-0.001
One or more	6,507	94.9 (1.4)	1.60(1.15, 2.22)	1.2 (1.1, 1.4)	1.58 (1.23, 2.02)	< 0.001
Educational Attainment						
College/university or graduate school	7,219	83.3 (1.0)	Reference	0.9 (0.8, 0.9)	Reference	
High school or less	5,590	54.4 (1.3)	0.76(0.53, 1.08)	1.3 (1.2, 1.3)	0.66 (0.57, 0.75)	< 0.001
Employment Status						
Unemployed	6,237	43.2 (0.7)	Reference	0.8(0.7, 0.9)	Reference	
Employed	6,572	94.5 (1.5)	1.701(1.23, 2.37)	1.3 (1.2, 1.4)	1.68 (1.36, 2.07)	< 0.001
Smoking status		` ′	` ' '	` ′ ′		
Not smoker	11,521	117.4 (1.0)	Reference	1.1 (1.0,1.2)	Reference	
Current smoker	1,288	20.3 (1.6)	1.63 (1.05, 2.53)	1.0 (0.3,1.6)	0.89 (0.39, 2.04)	0.779
Household Income (million JPY)	1,200	20.5 (1.0)	1.03 (1.03, 2.33)	1.0 (0.5,1.0)	0.05 (0.55, 2.01)	0.777
Low	3,579	57.2 (1.5)	Reference	1.4 (0.9, 1.8)	Reference	
Moderate	3,379			1.4 (0.9, 1.8)		0.715
		49.3 (1.6)	1.28(0.68, 1.56)	. , ,	1.06 (0.77, 1.46)	
High	2,313	16.6 (0.6)	1.01(0.66, 1.53)	0.8 (0.3, 1.3)	0.57 (0.21, 1.56)	0.275
Unknown	3,647	14.5 (0.5)	0.467(0.28, 0.79)	0.4 (0.2, 0.7)	0.29 (0.11, 0.75)	0.011
Region						
Other	6,715	86.4 (2.0)	Reference	1.1 (1.0, 1.1)	Reference	
DES (April 13-)	2,133	16.3 (0.8)	1.12(0.70, 1.78)	1.1 (1.1, 1.2)	1.07 (0.99, 1.15)	0.100
DES (April 7-)	3,599	31.2 (0.5)	1.03(0.72, 1.48)	1.0 (1.0, 1.1)	0.94 (0.68, 1.24)	0.261
Fear of COVID-19						
7–15	3,443	40.9 (1.2)	Reference	1.4 (1.1, 1.7)	Reference	
16–20	4,199	25.9 (0.6)	0.781(0.50, 1.21)	0.7 (0.4, 1.0)	0.45 (0.23, 0.88)	0.019
21–25	3,439	34.3 (0.9)	0.832(0.53, 1.30)	0.9 (0.9, 1.0)	0.62 (0.47, 0.81)	0.001
26–35	1,728	36.5 (2.3)	2.00(1.28, 3.11)	1.4 (1.3, 1.5)	1.01 (0.84, 1.21)	0.927
Self-rated health	1,720	30.3 (2.3)	2.00(1.20, 3.11)	1.4 (1.5, 1.5)	1.01 (0.04, 1.21)	0.727
Good	5,366	43.8 (0.7)	Reference	0.9 (0.8, 1.1)	Reference	
		. ,				0.069
Other than good	7,443	93.9 (1.5)	1.39(1.01, 1.91)	1.2 (1.1, 1.2)	1.25 (0.98, 1.48)	0.069
Mental state change in the last month compared						
to before January 2020						
Worse	9,054	56.0 (0.6)	Reference	0.9 (0.7, 1.2)	Reference	
None	2,686	59.3 (2.2)	3.14(2.19, 4.49)	1.1 (0.8, 1.5)	1.25 (0.67, 2.33)	0.480
		18.8 (3.0)	6.53(4.15, 10.30)			
Better	566			1.8 (1.4, 2.3)	2.11 (1.39, 3.18)	< 0.001
Unknown	504	3.5 (0.8)	1.55(0.622, 3.88)	0.8 (0.7, 0.8)	0.80 (0.53, 1.20)	0.286
Suicidal thoughts	11.460	55 A =	D 6	0.0 (0.0 1.0)	D 6	
Yes, since before COVID-19	11,169	75.2 (0.7)	Reference	0.9 (0.8, 1.0)	Reference	
First happened	1,082	42.8 (3.9)	5.85(4.06, 8.43)	1.6 (1.4, 1.8)	1.85 (1.50, 2.27)	< 0.001
Never	558	19.7 (4.3)	7.66(4.85, 12.01)	1.1 (0.7, 1.5)	1.25 (0.79, 1.97)	0.337
Desire not to talk to anyone due to worries						
Yes	10,246	61.5 (0.6)	Reference	0.8 (0.7, 0.9)	Reference	
No	2,563	76.2 (3.2)	7.27 (5.24, 10.08)	1.5 (1.3, 1.7)	1.98 (1.56, 2.51)	< 0.001
Feeling isolated	2,505	10.2 (3.2)	7.27 (3.24, 10.00)	1.5 (1.5, 1.7)	1.70 (1.30, 2.31)	-0.001

No Cancellation of a family gathering due to the	1,988	73.6 (3.4)	8.35 (6.02, 11.60)	1.8 (0.6, 2.9)	2.52 (0.70, 9.04)	0.156
pandemic						
Yes	1,741	83.3 (1.0)	Reference	0.5 (0.1, 0.9)	Reference	
No	11,068	54.4 (1.3)	1.035(0.64, 1.68)	1.2 (1.1, 1.3)	2.58 (1.07, 6.25)	0.035
Non-payment of salary						
Yes, since before COVID-19	12,586	7.8 (7.3)	Reference	1.0 (0.9, 1.0)	Reference	
First experience	148	114.6 (0.9)	12.15(7.02, 21.01)	3.9 (2.9, 4.9)	4.88 (3.42, 6.97)	< 0.001
Never	75	15.3 (10.4)	15.13(8.40, 27.28)	1.9 (1.1, 2.8)	2.17 (1.36, 3.48)	0.001
Lack of money to buy necessities						
Yes, since before COVID-19	11,720	80.5 (0.7)	Reference	0.8(0.7, 0.9)	Reference	
First experience	727	34.9 (5.7)	6.97(4.68, 10.38)	2.2 (1.9, 2.6)	2.94 (2.27, 3.81)	< 0.001
Never	362	22.2 (6.1)	7.62(4.74, 12.25)	1.9 (1.1, 2.8)	2.50 (1.76, 3.55)	< 0.001

266

267

268

269

270

271

272

273

274

275

276

277

278

279

Participants who did not receive salaries or compensations were more likely to experience nonconsensual sex than those who did: (experienced since before COVID-19) OR 4.88, 95% CI 3.42-6.97; (first experience during COVID-19) OR 2.17, 95% CI 1.36-3.48. Similarly, participants who could not afford to buy life necessities were more likely to experiencing nonconsensual sex than the others (OR 2.5, 95% CI 1.76-3.55).

Lastly, participants who had suicidal thoughts were more likely to experience non-consensual sex than those who did not: (since before COVID-19: OR 1.85, 95% CI 1.50-2.27). Participants with reported feeling isolated were more likely to experience non-consensual sex than those who did not (OR 1.98, 95% CI 1.56-2.51). The result of the sub-analysis showed suicidal ideation was present in 20% of all women aged between 15 and 19 years.

Figure 1: Forest plot illustrating incidence rate of non-consensual sex for each factor (reference) among the participants.

280 [Insert Figure 1] 281 282 283 Note: Significant positive factors associated with non-consensual sex included the age group 284 "Aged 15–29 years old", as well as worsened mental or economic status were associated with 285 an increased likelihood of experiencing non-consensual sex. 286 287 **Discussion** 288 This study assessed the incidence of non-consensual sex in Japan during COVID-19 and 289 explored associated factors while applying weighing to mitigate biases related to online 290 participation. The overall rate of participants reporting non-consensual sex was 1.2% over a 291 five-month period from April to September 2020. Factors associated with experiencing non-292 consensual sex among participants included ages between 15 and 29, employment status, 293 deteriorating mental health condition, suicidal thoughts, a desire not to talk to anyone, feeling isolated, and challenging economic circumstances. 294 295 296 In Japan, research related to sexual matters is limited. We believe this limitation stems from 297 the fact that sexual topics are often considered taboo with Japanese education institutions, and 298 sexual education in Japan lags behind that in Western countries (22–24). Additionally, the 299 understanding of obtaining sexual consent, which is essential in all sexual activities, lags 300 behind compared to countries that are known to be advanced in sex education in some of

Europe countries (13,22,25). For instance, the Netherlands had sex education compulsory from

elementary school, and Denmark was one of the European countries that criminalized nonconsensual sexual acts in 2020 at the time of our survey. Despite the efforts in Europe, conducting such a comprehensive study on a large scale was of significant importance as it ventured into uncharted territory in Japan.

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

Globally, concern arose about an increase in sexual violence against women around April 2020, during the early stages of the COVID-19 outbreak. However, specific incidence data were limited (12,26). While some countries reported an uptick in sexual violence cases (e.g., South Africa, Bangladesh (12,27)), others reported a significant decrease after implementing lockdown measures, as seen in Australia and Canada (28,29). In Japan, the available data included two key pieces of information from the police office: the reported prevalence of sexual violence among women, which stood at 6.9% (30), and a notable increase in the number of sexual assault victims seeking consultations, approximately 1.2 times higher in 2020 compared to 2019 (31,32). However, the incidence rate had not been thoroughly investigated during the COVID-19 pandemic, resulting in a scarcity of information regarding the incidence of sexual violence against women (33). It is important to note that our study did not have access to a comparative dataset from a different time point to evaluate percentage change in the incidence of sexual violence. Nevertheless, we were able to fundamental data concerning non-consensual

sex for the first time in Japan.

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

Approximately 80% of the women who reported experiencing non-consensual sex fell within the age range of 15 and 49 years old. Notably, the incidence of reported non-consensual sex dramatically decreased among women aged 50 to 79 years. There was a significant increase in the risk of non-consensual sex within the age group 15 to 29 years. This pattern is consistent with reports from the United States and Japan, where individuals in their late teens to early thirties are identified as the age group at the highest risk of experiencing rape (30,34). It is crucial to recognize that young women who have been victims of sexual assault are also more likely to experience lifetime suicide attempts and post-traumatic stress symptom. Specifically, research indicates that teenage sexual trauma is strongly correlate with suicide attempts (35–38). Additionally, our sub-analysis results revealed that suicidal ideation was present in 20% of all women aged between 15 and 19 years. Based on the aforementioned findings and considerations, we firmly believe that measures to protect the younger generation are imperative (39). As mentioned previously, Japan falls behind comprehensive sexual education for the younger generation. We contend that it is necessary to incorporate detailed

explanations and the factors and circumstances that lead to unwanted sexual experiences

involving the non-consensual sex into Japanese school curriculum.

Concretely, employed participants were more likely to be in contact with others than those who are unemployed. This time, we have not specifically investigated the locations where non-consensual sex has occurred. Therefore, it can be said that employed women may become victims of non-consensual sex, potentially by someone within their workplace or close social circle. Indeed, reports from both Japan and the United States have indicated that non-consensual sex is most frequently perpetrated by individuals known to the victims (30,40). Employed people have an extra circle of known individuals from their workplace compared to unemployed people.

Intriguingly, women who did not receive a salary and lacked the financial means to purchase necessities were at an elevated risk of experiencing non-consensual sex. While this appears to be the opposite of the above finding suggesting that being employed could be associated with a higher odds of experiencing non-consensual sex, this aligns with previous studies showing a connection between deteriorating economic conditions and an increased incidence of sexual assault that have established a connection between deteriorating economic conditions and an increased incidence of sexual assault (41,42), which is consistent with our findings. There are

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

23

two possible explanations for this association: First, worsening personal economic circumstances may lead women to engage in occupations with a higher risk of non-consensual sex, such as working in the sex industry. Second, women in precarious financial situation often experience psychological pressure to secure and maintain employment (42). Under such pressure, they may become psychologically vulnerable to sexual coercion from their employers in exchange for job security (42). Future research should explore people in these circumstances in depth to better under-explored areas in Japan.

While 66% of the participants lived in areas under DSE, our study did not find a significant association between non-consensual sex and living in the DSE area. However, we did identify a significant correlation between experiencing a moderate level of fear related to the COVID-19 pandemic (scoring 16-25 points) and instances of non-consensual sex. This suggests that participants who stayed at home more frequently might have become more vulnerable to nonconsensual sexual experiences by their partner. In fact, non-consensual sex was more frequently reported as being perpetrated by partner during the COVID-19 pandemic, and there was an observed increase in the number of domestic problems during this period (31).

Based on the considerations outlined above, there is a heightened risk of non-consensual sexual

activity occurred during the COVID-19 pandemic, particularly among young, impoverished

women. Consequently, it is of utmost importance to identify and provide economic and

psychosocial support to these vulnerable individuals during times of social disruption.

Furthermore, a notable deficiency in the current landscape of Japan is the absence of

comprehensive education regarding the concept of sexual consent. Specifically, there is an

urgent need for thorough instruction emphasizing the importance of men obtaining consent

from women before engaging in any sexual activities (43). This education should be

disseminated at both the local and national levels, as it may necessitate a societal shift in

awareness at all levels.

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

In a noteworthy development, on June 16, 2023, Japan amended the legal term "Forced Sexual

Intercourse Crime" to "Non-consensual Sexual Intercourse Crime", explicitly stipulating that

any sexual activity without consent is now considered a criminal offense (44). We anticipate

that this alteration will provide a means of support to victims who have endured their suffering

in silence and were unable to report their experiences.

Limitations

This study has several limitations. First, as it is a web-based survey, participants in an internet-

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

25

but we did not specifically ask about non-consensual sexual activity in the workplace. However,

it is made available under a CC-BY 4.0 International license.

we observed a significant correlation between being employed and experiencing non-consensual sexual activity. Also, we did not inquire the place of experienced non-consensual sex such as workplace. We observed a significant correlation between being employed and experiencing non-consensual sexual activity. However, it was not possible to ascertain whether

non-consensual sex had occurred in the workplace.

The Forth, as mentioned in the section on "Analysis Subjects", our questionnaire only asked whether respondents self-identified as male or female. Therefore, among those who self-identified as female, there is a possibility that transgender women or individuals who identify as bigender may also be included. This follows the convention in Japan, where many national surveys still only inquire about male or female gender identities.

Conclusion

Early intervention to prevent individuals from becoming victims of sexual harm should be extended to economically vulnerable and young women, especially during times of societal upheaval such as the COVID-19 pandemic. Additionally, Japan should prioritize the implementation of comprehensive education on the concept of sexual consent.

References

- 429 Jozkowski KN, Sanders S, Peterson ZD, Dennis B, Reece M. Consenting to sexual 1. activity: The development and psychometric assessment of dual measures of consent. 430 431 Arch Sex Behav [Internet]. 2014 Jan 23 [cited 2022 Nov 27];43(3):437–50. Available 432 from: https://link.springer.com/article/10.1007/s10508-013-0225-7 2. Jozkowski KN, Peterson ZD. Assessing the Validity and Reliability of the Perceptions 433 of the Consent to Sex Scale. http://dx.doi.org/101080/002244992012757282 [Internet]. 434 2014 Aug 1 [cited 2022 Nov 27];51(6):632–45. Available from: 435 https://www.tandfonline.com/doi/abs/10.1080/00224499.2012.757282 436 437 European Union Agency for Fundamental Rights. Violence against women: an 3. EU-wide survey. Main results report. Hum Rights Doc online [Internet]. 2014 [cited 438 2022 Dec 15]; Available from: 439 https://www.academia.edu/31782914/Violence against women an EU wide survey 440 441 Main results report 442 Tharp AT, DeGue S, Valle LA, Brookmeyer KA, Massetti GM, Matjasko JL. A 4. 443 Systematic Qualitative Review of Risk and Protective Factors for Sexual Violence 444 Perpetration. Trauma, Violence, Abus. 2022 Apr;14(2):133-67.
- 5. Delker BC, Salton R, McLean KC, Syed M. Who has to tell their trauma story and

how hard will it be? Influence of cultural stigma and narrative redemption on the 446 storying of sexual violence. PLoS One [Internet]. 2020 Jun 5 [cited 2022 Nov 447 2];15(6):e0234201. Available from: 448 449 https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0234201 450 Anastario M, Lawry L, Shehab N. Increased Gender-based Violence Among Women 6. 451 Internally Displaced in Mississippi 2 Years Post-Hurricane Katrina. Disaster Med Public Health Prep [Internet]. 2009 Mar [cited 2022 Jun 23];3(1):18–26. Available 452 from: https://www.cambridge.org/core/journals/disaster-medicine-and-public-health-453 preparedness/article/abs/increased-genderbased-violence-among-women-internally-454 displaced-in-mississippi-2-years-posthurricane-455 katrina/D576CBEB73639E69823A1391B5BF160B 456 457 7. Thornton W, Voigt L. Disaster rape: Vulnerability of women to sexual assaults during Hurricane Katrina. J Public Manag Soc ... [Internet]. 2007 [cited 2022 Jun 23];13:23-458 459 49. Available from: http://sites.google.com/a/jpmsp.com/www/Vol13Iss2-DisasterRape-ThorntonandVo.pdf 460 461 8. Picardo CW, Burton S, Naponick J, Katrina Reproductive Assessment Team. 462 Physically and sexually violent experiences of reproductive-aged women displaced by Hurricane Katrina. J La State Med Soc [Internet]. 2010 Sep 1 [cited 2022 Jun 463

464		23];162(5):282, 284–8, 290. Available from:
465		https://europepmc.org/article/med/21141260
466	9.	Son C, Hegde S, Smith A, Wang X, Sasangohar F. Effects of COVID-19 on College
467		Students' Mental Health in the United States: Interview Survey Study. J Med Internet
468		Res 2020;22(9)e21279 https://www.jmir.org/2020/9/e21279 [Internet]. 2020 Sep 3
469		[cited 2022 Feb 14];22(9):e21279. Available from:
470		https://www.jmir.org/2020/9/e21279
471	10.	Wakata S, Nishioka D, Takaki Y. Changes in health-related quality of life scores
472		among low-income patients on social welfare programs in Japan during the COVID-19
473		pandemic: a single-center repeated cross-sectional study. BMC Public Health
474		[Internet]. 2022 Dec 1 [cited 2022 Dec 7];22(1):1–8. Available from:
475		https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-022-14597-5
476	11.	European Union Agency for Fundamental Rights. Violence against women: An EU-
477		wide survey main results, European Union Agency for Fundamental Rights [Internet].
478		2014 [cited 2022 Jun 23]. Available from:
479		https://fra.europa.eu/en/publication/2014/violence-against-women-eu-wide-survey-
480		main-results-report
481	12.	al Mamun F, Hosen I, Mamun MA. Sexual violence and rapes' increment during the

482 COVID-19 pandemic in Bangladesh [Internet]. Vol. 34, EClinical Medicine. Lancet Publishing Group; 2021 [cited 2022 Nov 2]. Available from: 483 484 http://www.thelancet.com/article/S2589537021000973/fulltext 485 13. Nishioka E. Historical Transition of Sexuality Education in Japan and Outline of Reproductive Health/Rights. Nihon Eiseigaku Zasshi [Internet]. 2018 [cited 2022 Nov 486 27];73(2):178–84. Available from: 487 https://www.researchgate.net/publication/325459300 Historical Transition of Sexual 488 ity Education in Japan and Outline of Reproductive HealthRights 489 490 14. Miyawaki A, Tabuchi T, Tomata Y, Tsugawa Y. Association between participation in 491 the government subsidy programme for domestic travel and symptoms indicative of COVID-19 infection in Japan: Cross-sectional study. BMJ Open [Internet]. 2021 Apr 492 493 13 [cited 2021 Jun 24];11(4):49069. Available from: http://dx.doi.org/10.1136/bmjopen-2021-049069 494 495 15. Rakuten Insight. Rakuten Insight - Asia's Leader in Online Market Research [Internet]. 2021 [cited 2022 Nov 2]. p. 1. Available from: https://insight.rakuten.com/ 496 497 16. Hosokawa Y, Okawa S, Hori A, Morisaki N, Takahashi Y, Fujiwara T, et al. The 498 Prevalence of COVID-19 Vaccination and Vaccine Hesitancy in Pregnant Women: An Internet-based Cross-sectional Study in Japan. J Epidemiol [Internet]. 2022 [cited 2023] 499

500		Nov 12];32(4):188–94. Available from: https://doi.org/10.2188/jea.JE20210458
501	17.	Prime Minister of Japan and His Cabinet. Guidelines for Lifting the State of
502		Emergency [Internet]. 2020 [cited 2021 Jul 7]. Available from:
503		https://japan.kantei.go.jp/ongoingtopics/_00025.html
504	18.	Kyodo News+. Abe declares coronavirus emergency over in Japan [Internet]. [cited
505		2021 Jul 7]. Available from:
506		https://english.kyodonews.net/news/2020/05/a1f00cf165ae-japan-poised-to-end-state-
507		of-emergency-over-coronavirus-crisis.html
508	19.	Sieg L. Explainer: As COVID-19 cases surge, Japan sticks to "lockdown-lite"
509		Reuters [Internet]. 2021 [cited 2022 Dec 15]. Available from:
510		https://www.reuters.com/world/asia-pacific/covid-19-cases-surge-japan-sticks-
511		lockdown-lite-2021-08-06/
512	20.	Wakashima K, Asai K, Kobayashi D, Koiwa K, Kamoshida S, Sakuraba M. The
513		Japanese version of the Fear of COVID-19 scale: Reliability, validity, and relation to
514		coping behavior. PLoS One [Internet]. 2020 Nov 5 [cited 2022 Dec
515		15];15(11):e0241958. Available from:
516		https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0241958
517	21.	Ahorsu DK, Lin C-Y, Imani V, Saffari M, Griffiths MD, Pakpour AH. The Fear of

518		COVID-19 Scale: Development and Initial Validation. Int J Ment Heal Addict 2020
519		[Internet]. 2020 Mar 27 [cited 2021 Jul 9];1–9. Available from:
520		https://link.springer.com/article/10.1007/s11469-020-00270-8
521	22.	Hashimoto N, Ushitora K, Morioka M, Motegi T, Tanaka K, Tashiro M, et al. Sex
522		Education Sexuality, Society and Learning School education and development of
523		gender perspectives and sexuality in Japan. 2017 [cited 2022 Nov 27]; Available from:
524		http://www.tandfonline.com/action/journalInformation?journalCode=csed20
525	23.	MARCH 2021 Global Gender Gap Report The analysis presented in the Global
526		Gender Gap Report 2021. 2021 [cited 2022 Nov 27]; Available from:
527		http://reports.weforum.org/global-
528	24.	Fu H. The bumpy road to socialise nature: sex education in Japan. Cult Health Sex
529		[Internet]. 2011 Sep [cited 2022 Dec 15];13(8):903–15. Available from:
530		https://pubmed.ncbi.nlm.nih.gov/21787254/
531	25.	Ponzetti JJ. Evidence-based approaches to sexuality education: A global perspective.
532		Evidence-based Approaches to Sex Educ A Glob Perspect. 2015 Jan 1;359–84.
533	26.	UN Women. COVID-19 and Ending Violence Against Women and Girls [Internet].
000		
534		United Nations Entity for Gender Equality and the Empowerment of Women (UN

536		/media/head quarters/attachments/sections/library/publications/2020/issue-brief-covid-decorations/2020/issue-brief-covi
537		19-and-ending-violence-against-women-and-girls-en.pdf?la=en&vs=5006
538	27.	Ditekemena JD, Luhata C, Mavoko HM, Siewe Fodjo JN, Nkamba DM, Van Damme
539		W, et al. Intimate Partners Violence against Women during a COVID-19 Lockdown
540		Period: Results of an Online Survey in 7 Provinces of the Democratic Republic of
541		Congo. Int J Environ Res Public Heal 2021, Vol 18, Page 5108 [Internet]. 2021 May
542		12 [cited 2022 Nov 2];18(10):5108. Available from: https://www.mdpi.com/1660-
543		4601/18/10/5108/htm
544	28.	Payne JL, Morgan A, Piquero AR. COVID-19 and social distancing measures in
545		Queensland, Australia, are associated with short-term decreases in recorded violent
546		crime. J Exp Criminol [Internet]. 2022 Mar 1 [cited 2022 Nov 2];18(1):89–113.
547		Available from: https://link.springer.com/article/10.1007/s11292-020-09441-y
548	29.	Muldoon KA, Denize KM, Talarico R, Fell DB, Sobiesiak A, Heimerl M, et al.
549		COVID-19 pandemic and violence: rising risks and decreasing urgent care-seeking for
550		sexual assault and domestic violence survivors. BMC Med [Internet]. 2021 Dec 1
551		[cited 2021 Jun 24];19(1):1–9. Available from: https://doi.org/10.1186/s12916-020-
552		01897-z
553	30.	Gender Equality Bureau, Cabinet Office J. Survey on Violence between Men and

554		Women [Internet]. 2018 [cited 2022 Dec 15]. Available from:
555		http://www.gender.go.jp/policy/no_violence/e-vaw/chousa/h11_top.html
556	31.	Hashimoto S. Sexual violence consultations in Japan up 15 % during pandemic. 2020
557		[cited 2022 Dec 4];1–2. Available from:
558		https://english.kyodonews.net/news/2020/11/946cdc3cfe27-sexual-violence-
559		consultations-in-japan-up-15-during-pandemic.html?phrase=sex
560		crimes&words=sex,crime,crimes,Sex
561	32.	The White Paper on Gender Equality 2021 [Internet]. 2021 Jun [cited 2022 Dec 15].
562		Available from:
563		https://www.gender.go.jp/english_contents/about_danjo/whitepaper/pdf/ewp2021.pdf
564	33.	Mittal S, Singh T. Gender-Based Violence During COVID-19 Pandemic: A Mini-
565		Review. Front Glob Women's Heal. 2020 Sep 8;0:4.
566	34.	Morgan RE, Statistician B, Mason BJ, Intern B. Crimes Against the Elderly, 2003-
567		2013. 2014 Nov.
568	35.	Olshen E, McVeigh KH, Wunsch-Hitzig RA, Rickert VI. Dating violence, sexual
569		assault, and suicide attempts among urban teenagers. Arch Pediatr Adolesc Med
570		[Internet]. 2007 Jun 1 [cited 2021 Jun 24];161(6):539–45. Available from:
571		https://jamanetwork.com/journals/jamapediatrics/fullarticle/570505

572	36.	Davidson JRT, Hughes DC, George LK, Blazer DG. The Association of Sexual
573		Assault and Attempted Suicide Within the Community. Arch Gen Psychiatry
574		[Internet]. 1996 Jun 1 [cited 2022 Aug 22];53(6):550–5. Available from:
575		https://jamanetwork.com/journals/jamapsychiatry/fullarticle/497597
576	37.	Rosenberg HJ, Jankowski MK, Sengupta A, Wolfe RS, Wolford II GL, Rosenberg SD,
577		et al. Single and Multiple Suicide Attempts and Associated Health Risk Factors in
578		New Hampshire Adolescents. Suicide Life-Threatening Behav [Internet]. 2005 Oct 1
579		[cited 2022 Aug 22];35(5):547–57. Available from:
580		https://onlinelibrary.wiley.com/doi/full/10.1521/suli.2005.35.5.547
581	38.	McDougall EE, Langille DB, Steenbeek AA, Asbridge M, Andreou P. The
582		Relationship Between Non-Consensual Sex and Risk of Depression in Female
583		Undergraduates at Universities in Maritime Canada. J Interpers Violence [Internet].
584		2019 Nov 1 [cited 2022 Aug 23];34(21–22):4597–619. Available from:
585		https://journals.sagepub.com/doi/10.1177/0886260516675468?url_ver=Z39.88-
586		2003𝔯_id=ori%3Arid%3Acrossref.org𝔯_dat=cr_pub++0pubmed
587	39.	BRASOR P. In Japan no one wants to talk about sex education The Japan Times
588		[Internet]. 2018 [cited 2022 Dec 22]. Available from:
589		https://www.japantimes.co.jp/news/2018/04/07/national/media-national/japan-no-one-

590		wants-talk-sex-education/
591	40.	Breiding MJ, Smith SG, Basile KC, Walters ML, Chen J, Merrick MT. Prevalence and
592		Characteristics of Sexual Violence, Stalking, and Intimate Partner Violence
593		Victimization—National Intimate Partner and Sexual Violence Survey, United States,
594		2011. MMWR Surveill Summ [Internet]. 2014 Sep 9 [cited 2022 Aug 23];63(8):1.
595		Available from: /pmc/articles/PMC4692457/
596	41.	Markowitz FE. Socioeconomic disadvantage and violence: Recent research on culture
597		and neighborhood control as explanatory mechanisms. Aggress Violent Behav. 2003
598		Mar 1;8(2):145–54.
599	42.	Department of Reproductive Health and Research World Health Organization.
600		Violence against women [Internet]. 2014 [cited 2021 Jun 24]. Available from:
601		https://apps.who.int/iris/bitstream/handle/10665/112325/WHO_RHR_14.11_eng.pdf
602	43.	Willis M, Hunt M, Wodika A, Rhodes DL, Goodman J, Jozkowski KN. Explicit
603		Verbal Sexual Consent Communication: Effects of Gender, Relationship Status, and
604		Type of Sexual Behavior. https://doi.org/101080/1931761120191565793 [Internet].
605		2019 Jan 2 [cited 2022 Nov 27];31(1):60–70. Available from:
606		https://www.tandfonline.com/doi/abs/10.1080/19317611.2019.1565793
607	44.	Ministry of Justice. Ministry of Justice: FAQs on Amendments to Laws Related to

608 Sexual Crimes [Internet]. [cited 2023 Nov 12]. Available from: 609 https://www.moj.go.jp/keiji1/keiji12 00200.html Tabuchi T, Shinozaki T, Kunugita N, Nakamura M, Tsuji I. Study Profile: The Japan 610 45. 611 "Society and New Tobacco" Internet Survey (JASTIS): A Longitudinal Internet 612 Cohort Study of Heat-Not-Burn Tobacco Products, Electronic Cigarettes, and Conventional Tobacco Products in Japan. J Epidemiol [Internet]. 2019 Nov 5 [cited 613 2022 Dec 21];29(11):444–50. Available from: https://doi.org/10.2188/jea.JE20180116 614 Kotera Y, Van Laethem M, Ohshima R. Cross-cultural comparison of mental health 615 46. 616 between Japanese and Dutch workers: relationships with mental health shame, self-617 compassion, work engagement and motivation. Cross Cult Strateg Manag. 2020 Sep 3;27(3):511-30. 618

Figure 1. Forest plot of the incidence rate of non-consensual sex of each factor (reference) among the participants.

