1	Factors associated with commuting stress among tertiary education employees in
2	Georgetown, Guyana
3	
4	Davon Van-Veen <sup>1*</sup> , Hue-Tam Jamme <sup>2</sup> , Heather Ross <sup>3</sup> , Netra Chhetri <sup>3</sup>
5	
6	<sup>1</sup> School of Allied Health, College of Medical Sciences, University of Guyana
7	<sup>2</sup> School of Geographical Sciences and Urban Planning, Arizona State University
8	<sup>3</sup> School for the Future of Innovation in Society, Arizona State University
9	
10	*Corresponding author
11	Email: <u>davon.vanveen@uog.edu.gy</u>
12	

#### Abstract 13

14 The aim of this study was to identify factors associated with commuting stress using 15 symptomatology among tertiary education employees in Georgetown Guyana. A cross 16 sectional survey was conducted among 427 (100 male, 317 female) participants, with a mean 17 age of 29.6(sd=8.7) years. Data about their demographic characteristics, psychosocial measures (perceived stress, coping and resilience), characteristics about their commute 18 19 (length, mode, and satisfaction with commute infrastructure), and how often they experienced 20 selected symptoms associated with stress on the daily commute was collected using a self 21 administered questionnaire. A commuting stress score was calculated for each participant and 22 this was followed by regression analysis. The results showed that after controlling for 23 resiliency, coping, perception of job, total life stress, income levels and education all of 24 which has contribute to be write son a perceive iterstands less in their lives, the respective sign and clactice.

25 demonstrated that females, persons who used public transportation, persons who have longer 26 commute times and persons who are less satisfied with the commute infrastructure are more 27 likely to report that they experience symptoms associated with stress while engaged in the 28 daily commute compared to males, persons who use private transportation, persons who have 29 shorter commute times and persons who are more satisfied with the commute infrastructure respectively. Once all controlled variables were accounted for, it appears that for this study 30 31 population, age, where they live and whether or not they actively or passively commuted did 32 not have any meaningful impact on whether they would report if they experienced stress 33 associated symptoms while engaged in the daily commute. he results suggest the need for a 34 multi-pronged approach to address commuting stress, involving individual healthcare, mental-health-centric urban transport design, improved commute infrastructure, and 35 36 increased employer engagement. Future studies employing qualitative and experimental 37 methods are recommended. 38

Keywords: Stress, Commute, Urbanization, Daily hassles, Mental Health

# 41 Introduction

42 The majority of contemporary urbanisation, a complex social and economic process that 43 occurs through various forms of urban growth and expansion drastically altering the social 44 and physical environment in which people live and operate, is occurring in low and middle 45 income countries(LMICS).[1,2] Not surprisingly these changes can have a significant impact 46 on both physical and mental health on urban inhabitants.[3-5] Increased access to modern 47 technology in the urban environment has induced a fast-paced life which is also characterised by a heavy workload, fast paced lifestyle, communication, quick decisions, multitasking, 48 pressure to be active on social media, and high mobility, all of which have a direct impact on 49

people's health [6,7] however for the purposes of this paper, there is going to be an exclusivefocus on high mobility.

52

53 The majority of mobility in the urban environment consists of the daily commute which appears 54 to be intensifying in both the developed and developing world. This intensification is a result of an increase in both the frequency and length/duration of commutes to work.[8,9] There is a 55 56 significant body of literature that examines how the increased intensity of commuting to and from work can have detrimental effects on health. Impacts of lengthy commutes include an 57 58 increased incidence of respiratory illnesses, muscular injuries, and reduced physical activity 59 along with the associated undesirable health outcomes.[10-14] Additionally, an emerging body of scholarship reveals that a lengthy commute to workplaces is also associated with poor mental 60 61 health which may be expressed as stress.[15-19] The commute is more often than not a daily 62 activity, thus if it is stressful, a person may be placed under stress on an ongoing basis.

63

64 Stress is the physiological and psychological response to a situation that a person has perceived to be negative and has made a determination that they do not have the resources to address the 65 problem, accounting for all other factors. [20-22] In addition to being an indicator of poor 66 mental health, chronic exposure to stress is also associated with many chronic non-67 68 communicable diseases (CNCDs) including but not limited to diabetes, cardiovascular and 69 cerebrovascular pathologies, chronic obstructive pulmonary disease, and malignant 70 neoplasms.<sup>[23]</sup> CNCDs continue to be a significant public health concern globally, accounting for most of the reported global mortality and morbidity with the global burden rapidly shifting 71 72 towards the LMICs. [24-26] Furthermore, long-term exposure to stress is more likely to be in a 73 state of constant anger and irritability, leading to strained personal relationships which can

further lead to to potential isolation, anxiety, and depression. If it is severe enough, it may even
lead to suicide.[27-29]

76

77 The question therefore becomes who is more likely to experience commuter stress and there have been a few studies that attempted to examine these factors. For example, women are more 78 likely to experience commuter stress than men.[30-34] Similarly, active commuters are less 79 80 likely to experience stress than passive commuters.[35-37] The issue however with the majority of these studies either examined only the intrinsic factors (factors related to the commuter) or 81 82 the extrinsic factors (factors related to the commute process). Also, almost all of the previous studies were conducted in economies that were considered fully urbanised (that is, more than 83 50% of their population live in urban agglomerations) and there has been a dearth of studies 84 85 that examined this issue in economies that are in transition to urbanisation. Also, the majority 86 of previous studies used only Likert or binary scales to examine stress experienced during the commute. They did not focus on the physical and psychological symptoms of stress. In order 87 88 to address the preceding gaps identified, this paper aims to to identify what factors (both 89 extrinsic and intrinsic) are associated with experiencing stress while engaged in the daily commute using in an emerging economy that has less than 50% of its population living in an 90 urban environment using symptomatology to measure commuting stress. 91

92

The next section of this paper detailed the methods This is followed by the presentation of the results which consists of the presentation of the sample characteristics, the individual risk factor assessment and regression models that include all of the factors that were tested. These results and the implications with respect to the literature are then discussed in the next section after which some final conclusions along with some recommendations are made.

# 99 Method

# 100 Study design and setting

101 This observational cross-sectional study was conducted in Georgetown, the capital city of 102 Guyana, between March 1st to April 30th, 2023. As the country's political and economic 103 centre, it serves as a critical hub for administrative, commercial, and cultural activities. While 104 the reliable data on the population of Georgetown is limited, the National Census of 2012 puts 105 its population to be around 120,000 people.

106

# 107 Study Participants

Participants were drawn from among employees at the University and the University's affiliate partners. The University of Guyana serves as the National University of Guyana and has just about 8,000 students and employs about 1,000 teaching and non-teaching staff. Universities are normally microcosms of the wider society in which they operate, therefore it is to be expected that the demographic makeup of the university would be similar to that of the society in which they function.[38-39]

114

Using an online sample size calculator, (Raofost, 2004), assuming an unknown population a sample size of 377 was recommended. Assuming that all questionnaires would not be returned or only partially completed, it was determined that 33% more questionnaires should be distributed therefore 505 questionnaires were distributed. To be included in the study, participants must work in Georgetown for at least 30 hours per week at an in person job. The participant must also be at least 18 years of age and have worked on the job for at least 6 months.

123 Convenience sampling was used until the desired sample size was achieved. Employees would 124 be approached and given a short presentation about the rationale, objectives, and significance of the study along with guarantees that no identifiable information will be collected and that 125 126 all information collected would remain confidential. After the employee gave their verbal 127 consent to participate in the study, each employee was informed that there were three different 128 ways to participate in the study: a shared link, a tablet with the questionnaire on it, or a physical 129 paper questionnaire and it was for them to choose the method that they were most comfortable 130 with. Once a choice was made, the 83-item questionnaire was made available using the medium 131 of their choice, and the employee was thanked for their time.

132

#### 133 Variables

The dependent variable commuting stress was evaluated using symptomatology. This was done 134 by asking participants to report the frequency in which they experienced a variety of symptoms 135 136 associated with stress while engaged in the daily commute. These included physical symptoms 137 such as muscle pain, headaches, nausea, difficulty breathing, fatigue, chest pain, indigestion, 138 dizziness, sweating and constipation/diarrhoea and psychological symptoms such as feeling 139 irritable, overwhelmed, anxious, having uncontrollable thoughts, depressed, a sense of dread, 140 lonely, pressured, and distressed. For each symptom, they were asked whether it was 141 experienced very often, fairly often, sometimes, almost never, or never. Based on the response 142 a score was assigned to each symptom with very often=4 and never=1. A total commuting 143 stress score was then generated.

144

145 The independent variables aimed to assess personal (intrinsic) factors, commute specific 146 factors, and built environment factors. The demographic factors which were tested include age, 147 sex, place of residence (whether they lived in the city or the suburbs), and hours of work per

148 day. Other intrinsic factor included was the perceived quality of sleep which was measured on 149 a five-point Likert scale, the general life stress which was measured using the Cohen's Perceived Stress Scale (PSS-10), and how hassled a person perceived their job to be, which 150 151 was measured using a sub-scale of the Daily Hassles and Mood Ouestionnaire (DHMO). The 152 DHMQ is a self-report measure developed by Dr. Thomas W. Kamarck and his colleagues in 153 1991, which assesses the daily hassles, mood, and health status of individuals. The job sub-154 scale consists of seven questions that aimed to elicit how people felt about their fellow workers. 155 their supervisors, and their workload. It was scored from 0-21. Higher scores were associated 156 with negative feelings about the job. The DHMQ has been found to be a valid and reliable 157 measure, with high levels of internal consistency and test-retest reliability. It has been used in 158 various populations, including healthy adults, medical patients, and college students. The scale 159 has been found to be positively associated with a variety of health outcomes, such as 160 cardiovascular disease, hypertension, and depression. [40-42] The Perceived Life Stress 161 measured by Cohen's Perceived Stress Scale (PSS-10) is another independent variable. The 162 PSS-10 is a 10-item version of the scale, which was developed by Sheldon Cohen and his colleagues in 1983. The scale assesses the extent to which respondents find their lives 163 164 unpredictable, uncontrollable, and overloaded in the last month. Respondents are asked to rate the degree to which they agree with each statement on a 5-point Likert scale, where 0 = never 165 166 and 4 = very often. The PSS-10 has been found to be a valid and reliable measure of perceived 167 stress, with high levels of internal consistency and test-retest reliability. In addition, the scale 168 has been validated in various populations including, but not limited to, healthy adults, medical 169 patients, and college students. It has been found to be positively associated with a variety of 170 health outcomes, including cardiovascular disease, hypertension, and depression.[43-45]

The participants' coping capabilities were the third mediating variable that was tested using
the Brief Resilient Coping Scale (BRCS). This is a 4-item unidimensional scale that is
designed to capture tendencies to cope with stress in a highly adaptive manner. The scale
ranges from 0-20 and the higher the score, the greater the coping capabilities of the person
This instrument has also been successfully subjected to multiple reliability and validity
evaluations.[46-48]

178

179 Resiliency refers to the ability to adapt despite encountering less-than-ideal circumstances. 180 This was measured by using the Brief Resilience Scale (BRS) which is a 6-point 181 unidimensional scale that does not take into account external factors. [49] A person who is 182 resilient tends to demonstrate high levels of emotional stability, self-efficacy, and sociability 183 and thus better able to handle life stresses. The BRS score ranges from 1-5 with the higher 184 values indicating higher levels of resilience. Like the BCRS, the BRS was also subject to 185 multiple reliability and validity testing. [50-52]

186

187 The first commute-specific factors which were tested were the method of commuting whether 188 it was active or passive, or where commuting was done by public or private means. The second 189 commute-specific factor that was tested was the length of the commute. The study used 5-point 190 Likert Scale questions to assess commuters' satisfaction with various built environment factors. 191 These factors included greenery along the route, frequency of traffic congestion, and road-use 192 compliance by others. Additionally, satisfaction with transportation infrastructure was gauged 193 through multiple aspects like road conditions, street lighting, and public transport. A composite 194 score ranging from 9-45 was generated, with higher scores indicating greater satisfaction.

### 196 Data analysis

197 All data were analysed using Stata Version 17.0. Descriptive analysis was performed on all variables. Non numerical data was converted to binary. For gender, males were assigned a 198 value of 0 and females a value of 1. Those who lived in the city were assigned a value of 1 199 200 while those who lived in the suburbs were assigned a value of 0. Persons who used active and private commute modes were assigned a value of 0 while persons who used active and public 201 202 forms of transportation were assigned a value of 1. Regression analysis was conducted. All 203 values are reported with a 95% confidence interval. A *p*-value of less than 0.05 was considered to be statistically significant. 204

205

# 206 Ethical Issues

For each participant, the purpose of the study was explained. They were informed that 207 208 participation in this study was completely voluntary and that there were no consequences whether good or bad if they chose to participate in the study. They were also informed that they 209 210 do not have to answer any question they feel uncomfortable about asking and they can chose 211 to terminate their participation in the study at any point. Participants were also informed that no personally identifying data will be collected and therefore there was no way to trace a 212 213 particular submission to any one participant. This information was repeated on the cover of the 214 questionnaire that was provided. The first question of the questionnaire asked whether or not 215 they agreed to participate in the study.

This study was approved by the Institutional Review Board of Arizona State University.(Protocol number: STUDY00017507)

# 219 **Results**

Out of 505 questionnaires initially distributed, several were excluded from the final analysis 220 221 for various reasons: 22 had over 10% of questions unanswered; 24 were completed by 222 individuals working fewer than 30 hours per week; 2 were filled out by minors under the age of 18; 29 were from respondents who neither worked or lived in the study area; and 1 came 223 from a remote worker. This left 427 valid questionnaires for final analysis There participants 224 225 were relatively diverse with respect to gender, age, relationship status, ethnicity, religious affiliation, educational levels, length of commute, total life stress levels and where they resided. 226 227 Table 1 shows the nominal characteristics of the participants included in the final analysis while 228 table 2 provides a summary of the numerical characteristics.

229

#### 230 Table 1: Summary of nominal characteristics of study participants (n=427)

- 231
- 232

Parameter	Number (%)
Gender	
Male	100 (25.6)
Female	317 (74.4)
Relationship status	
Committed Relationship	205 (48.2)
Single	187 (44.1)
No response	35 (8.2)

Income level	
Low income	280 (65.5)
	280 (65.5)
High and middle income	147 (34.5)
Ethnicity	
Indigenous	3 (0.7)
African Guyanese	153 (36.0)
East Indian Guyanese	121 (28.5)
Mixed descent	148 (34.8)
No Response	2 (0.5)
Religious affiliation	
Christian	311 (73.3)
Hindu	64 (15.1)
Muslim	29 (6.8)
Other/ Non-affiliated	23 (5.4)
Educational Level Completed	
Tertiary	363 (85.2)
Non-Tertiary	64 (14.8)
Residence	
In Georgetown (main city)	149 (38.8)
Outside of Georgetown	278 (61.2)
Commute Mode (Private/Public)	
Private	240 (56.2)

187 (43.8)
23 (5.4)
404 (94.6)

#### Table 2: Summary of numerical characteristics of study participants (n=427)

Parameter	Range	Mean	Standard
			Deviation
Age	0-65	29.62	8.66
Perceived Stress Score (0-40)	0-37	19.03	6.97
Length of Commute (min)	5-165	39.20	25.67
Basic Coping Score (4-20)	4-20	15.08	3.04
Resiliency (1-5)	1-5	3.39	0.75
Job Perception (0-21)	0-21	7.47	7.45
Infrastructure Satisfaction (9-45)	9-45	21.37	7.53
Commute Stress Score (0-88)	1-88	36.77	16.01

The three most frequently reported symptoms associated with stress while engaged in the

daily commute are complaints about feeling fatigued, excessive sweating and feeling hassled. 

240 The four least commonly reported symptoms are feelings of indigestion, chest pains,

241 dizziness and bowel dysfunction. Figure 1 shows the distribution of all the symptoms

examined in this study.

243

#### 244 Table 3: Effect of intrinsic and extrinsic factors on commuting stress score

Commuting	Coef.	St.Err.	t-value	p-	[95% Conf	Interval]	Sig
Stress				value			
Female	4.606	1.614	2.85	.005	1.432	7.781	***
Age	.053	.092	0.58	.564	127	.233	
Lives outside of Georgetown	957	1.505	-0.64	.525	-3.917	2.003	
Storgets with							
Non-Active commute	3.265	3.243	1.01	.315	-3.113	9.644	

medRxiv preprint doi: https://doi.org/10.1101/2024.02.23.24303247; this version posted February 27, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license . Public 3.803 1.48 2.57 .011 .892 6.713 \*\* commute \*\* Time for .059 .03 1.97 .049 0 .117 commute (min) Satisfaction -.277 .095 -2.90 .004 -.465 -.089 \*\*\* with commute infrastructure Resilience \*\*\* -3.261 1.085 -3.01 .003 -5.395 -1.127 \*\* Coping .056 .576 2.18 .03 1.096 .264 \*\*\* Satisfaction 4.557 1.677 2.72 .007 1.257 7.856 with job \*\*\* Total life .917 .114 8.05 0 .693 1.14 stress

Low income	1.68	1.392	1.21	.228	-1.057	4.418	
Non tertiary educated	-2.026	1.911	-1.06	.29	-5.784	1.733	
Constant	11.309	7.62	1.48	.139	-3.679	26.297	
Mean dependent var	34.441	SD depend	dent var		15.997		
R-squared		0.409	Number o	f obs		363	
F-test		18.551	Prob > F			0.000	
Akaike crit. (AIC)		2879.231	Bayesian	crit. (BIC)		2933.753	

\*\*\*p<.01, \*\*p<.05, \*p<.1

The results in Table 3 shows that after controlling for resilience, coping, perceived life stress,
satisfaction with job, income levels and educational attainment, commuting stress levels
measured using symptomatology were significantly higher in females, persons who used
public means of transportation, persons with longer commute times and persons who were

not satisfied with commute infrastructure. Age and location of residence appear to have noeffect on the level of commuting stress as measured using symptomatology.

252

# 253 **Discussion**

254 The main aim of this paper was to determine what factors were associated with experiencing commuter stress during the daily commute using symptomatology. After controlling for 255 resiliency, coping, perception of job, total life stress, income levels and education all of which 256 257 can contribute to how persons perceive hassles in their lives, the regression model demonstrated that females, persons who used public transportation, persons who have longer 258 commute times and persons who are less satisfied with the commute infrastructure are more 259 260 likely to report that they experience symptoms associated with stress while engaged in the daily commute compared to males, persons who use private transportation, persons who have shorter 261 262 commute times and persons who are more satisfied with the commute infrastructure 263 respectively. Once all controlled variables were accounted for, it appears that for this study population, age, where they live and whether or not they actively or passively commuted did 264 265 not have any meaningful impact on whether they would report if they experienced stress associated symptoms while engaged in the daily commute. These findings are for the most part 266 consistent with what is already published in the literature that used alternative methods of 267 268 measuring commuting stress. In the present study, the non-association that was found between 269 the active/non-active commuting and commuting stress was discordant with previous studies, however this could be explained due to sampling inefficiencies which resulted in the majority 270 271 of the sample being users of non-active commuting modes.

273 The findings of this study are limited to persons who are employed with the University of 274 Guvana or one of its affiliate organisations/institutions for at least 6 months and have to commute to work on a daily basis. Despite these limitations however, the findings of this study 275 276 add to the existing knowledge in a few ways. Firstly, this appears to be the first time an attempt 277 was made to examine all of the factors (both the intrinsic and extrinsic) that could potentially 278 contribute to commuting stress in a single study. Previous research tended to only focus on a 279 few of the factors and there was very little consistency across studies as to how commuting 280 stress should be measured. Secondly, for the most part, commuting stress has been measured 281 either by using electrodermal activity or a single Likert-point scale, however, this study used 282 symptomatology. The only other study that used symptomology to assess commuting stress 283 was conducted by Rezapour et al., (2021), however, the focus was narrow with respect to both 284 the factors examined and the symptoms used when compared to the present study. Further, 285 Rezaour treated each symptom as an individual outcome whereas in the present study, all of 286 the symptoms were reduced to a single value that factored in any positive emotions that may 287 occur on the daily commute to give a more holistic view of the stress experienced on the daily 288 commute. The third way in which this paper contributes to the literature is the fact that 289 symptomology could now potentially be used as an alternative method for measuring 290 commuting stress as the findings of this study for the most part replicate that of what exists in 291 the available literature. Furthermore, in doing so, this study strengthens the findings in the 292 previously published studies as the findings are similar even when using a different method to measure the outcome variables. The fourth way in which this study contributes to the existing 293 294 literature is that it expands the list of factors that can be considered when thinking about 295 commuting stress. The two novel factors that emerged in this study are people's perception of 296 their jobs, their satisfaction with the commute infrastructure in place, and whether or not they 297 commute via public or private means. The fifth and final way in which this paper contributes

298 to the literature is that it adds to the geographical diversity in which this issue was examined. A cursory review of previous papers would reveal that the studies which have examined this 299 issue have been skewed towards North America, Europe and Asia, where there has been heavy 300 301 investment in public transportation, with little to no studies emanating from South and Central 302 America, the Caribbean and Africa. Furthermore, the majority of studies that are available were conducted in economies that have completed the urban transition, that is more than 50% of 303 304 their total population lives in urban areas. At the time of data collection, Guyana, located in South America is an economy in transition with less than 50% of its population living in urban 305 306 agglomerations.

307

# 308 Conclusions

309 We found that both the intrinsic and the extrinsic factors interact in complex ways to influence 310 the overall stress experienced during commuting. Understanding these can help in designing 311 interventions to make commuting a less stressful experience, whether it involves improving 312 public transit systems, encouraging flexible work arrangements, or promoting stress 313 management strategies for individuals. Interventions such as these can significantly reduce their 314 exposure to stress which stands as an independent risk factor for CNCDs that continues to 315 plague both the developed and developing world. As a policy response following three 316 measures can be implemented. First, at a personal level, health care providers could be provided with some tools that they can use on their own to reduce stressful situations. For example, one 317 could provide mindfulness - often used as a therapeutic technique to manage stress and promote 318 mental wellbeing. It is a practice that involves focusing one's awareness on the present moment, 319 320 while calmly acknowledging and accepting one's feelings, thoughts, and bodily sensations. [53-321 54]

322

323 The second, establishment of guidelines to ensure that the infrastructure that is being used by 324 commuters is mentally friendly, refers to physical infrastructure designed and constructed with 325 an understanding of the psychological and emotional impact it has on individuals who interact 326 with it. It aims to create an environment that promotes mental well-being and reduces stress 327 levels. City planners and transportation officials can create an infrastructure that supports mental 328 well-being and reduces the stress associated with commuting. In the long run, mentally friendly 329 infrastructure can contribute to happier, healthier, and more productive communities. This will 330 require high-level interventions as there may be a need to make the public transportation system 331 less chaotic, ensure that there is more greenery along major thoroughfares, and just simply ensure that the commuting infrastructure such as roads and footpaths are maintained. 332

333 The third and final way that is proposed to reduce the stress on the daily commute for organisations is to ensure that their employees are meaningfully engaged. An employee who is 334 335 meaningfully engaged at work is deeply connected and committed to their role, their team, and 336 their organisation. They find their work fulfilling and significant, feel that they are making a 337 valuable contribution, and are motivated to do their best. To foster meaningful engagement, 338 organisations need to provide supportive leadership, opportunities for growth and development, 339 a positive work environment, and recognition for contributions and achievements. While 340 everyone's circumstances are different, and a range of factors can influence commute-related 341 stress, a positive relationship with their work and their workplace can indeed help mitigate stress 342 during their commute. For example, when employees find meaning in their work and feel valued 343 in their roles, they may experience less anxiety about workplace challenges or pressures. This 344 positive mindset can extend to their commute, making them less stressed about potential delays or complications on their way to work. Similarly, employees who are empowered to manage their 345 346 workloads effectively may have more flexible schedules, allowing them to avoid peak commuting 347 times or opt for less stressful modes of transport. In an organisation that nurtures meaningful

engagement, the employee is likely to handle commuting stresses more effectively, viewing them
as minor inconveniences which can contribute to an overall reduction in stress, both within and
outside of work

351

# 352 **Reference**

353	1.	Zeng, C., Liu, Y., Liu, Y., & Qiu, L. Urban Sprawl and Related
354		Problems: Bibliometric Analysis and Refined Analysis from
355		<b>1991 to 2011.</b> <i>Chinese Geographical Science,</i> 2014, 24(2), 245–257
356	2.	United Nations, Department of Economic and Social Affairs, Population Division
357		2019. World Urbanization Prospects: The 2018 Revision (ST/ESA/SER.A/420). New
358		York: United Nations, 2019
359	3.	Aliyu, A. A., & Amadu, L. Urbanization, cities, and health: the challenges to
360		Nigeria-a review. Annals of African medicine, 2017, 16(4), 149.
361	4.	Li, X., Song, J., Lin, T., Dixon, J., Zhang, G., & Ye, H. Urbanization and health in
362		China, thinking at the national, local and individual levels. Environmental Health,
363		2016, 15, 113-123
364	5.	Eckert, S., & Kohler, S. Urbanization and health in developing countries: a systematic
365		review. World Health Popul, 2014, 15(1), 7-20.
366	6.	Lippke, S., Schalk, T. M., Kühnen, U., & Shang, B. Pace of life and perceived stress
367		in international students. PsyCh Journal, 2021, 10(3), 425-436.
368	7.	Qidwai, W., Khushk, I. A., Shamim, U., Altaf, S., Hadi, H., & Nanji, K. Fast pace of
369		life and its impact on health: results of a study from the largest city of Pakistan.
370		Pakistan journal of public health, 2016, 6(4), 10-16.

8.	Sultana, & Weber, J. The Nature of Urban Growth and the Commuting Transition:
	Endless Sprawl or a Growth Wave? Urban Studies, 2014, 51(3):544–576.
9.	Travisi, Camagni, R., & Nijkamp, P. Impacts of urban sprawl and commuting: A
	modelling study for Italy. Journal of Transport Geography, 2010, 18:383-392.
10	Ali, M., Ahsan, G. U., Uddin, Z., & Hossain, A. Road traffic delays in commuting
	workplace and musculoskeletal health among sedentary workers: a cross-sectional
	study in Dhaka city. Journal of occupational health, 2021, 63(1), e12289.
11	Kuenn. A. Does Commuting Affect Health? Health Economics, 2016, 25(8):984–
	1004
12	. Ramos, C. A., Wolterbeek, H. T., & Almeida, S. M. Air pollutant exposure and
	inhaled dose during urban commuting: a comparison between cycling and motorized
	modes. Air Quality, Atmosphere & Health, 2016, 9, 867-879.
13	. Raza, A., Pulakka, A., Hanson, L. L. M., Westerlund, H., & Halonen, J. I. Commuting
	distance and behavior-related health: a longitudinal study. Preventive Medicine, 2021,
	150, 106665.
14	. Wu, D. L., Lin, M., Chan, C. Y., Li, W. Z., Tao, J., Li, Y. P., & Bu, C. W.
	Influences of commuting mode, air conditioning mode and meteorological parameters
	on fine particle (PM2. 5) exposure levels in traffic microenvironments. Aerosol and
	Air Quality Research, 2013, 13(2), 709-720.
15	Karan, M., Rahal, D., Almeida, D. M., Bower, J. E., Irwin, M. R., McCreath, H.,
	Seeman, T., & Fuligni, A. J. School commute time, chronotype, and altered HPA axis
	functioning during adolescence. Psychoneuroendocrinology, 2021, 133, 105371-
	105371
	<ol> <li>9.</li> <li>10.</li> <li>11.</li> <li>12.</li> <li>13.</li> <li>14.</li> </ol>


394	16. Lucas, & Heady, R. B. Flextime Commuters and Their Driver Stress, Feelings of
395	Time Urgency, and Commute Satisfaction. Journal of Business and Psychology,
396	2002, 16(4), 565–571
397	17. Rosida, I., Sari, D. W., & Irjayanti, A. D. The Mode Choices And Commuting Stress:
398	Empirical Evidence From Jakarta And Denpasar. Jurnal Pengembangan Kota, 2019,
399	7(1), 68–76.
400	18. Rüger, H., Pfaff, S., Weishaar, H., & Wiernik, B. M. Does perceived stress mediate
401	the relationship between commuting and health-related quality of life? Transportation
402	Research Part F: Traffic Psychology and Behavior, 2017, 50, 100–108.
403	19. van Hooff. The Daily Commute from Work to Home: Examining Employees'
404	Experiences in Relation to Their Recovery Status. Stress and Health, 2015, 31(2),
405	124–137.
406	20. Hulbert, W. N. J., Morrison, V., Wilkinson, C., & Neal, R. D. Investigating the
407	cognitive precursors of emotional response to cancer stress: Re-testing Lazarus's
408	transactional model. British Journal of Health Psychology, 2013, 18(1), 97-121.
409	21. Lazarus R. S., Folkman S. Stress, appraisal, and coping. Springer Publishing
410	Company, 1984
411	22. Wang, F., Pan, F., Shapiro, L. A., & Huang, J. H. Stress Induced Neuroplasticity and
412	Mental Disorders 2018. Neural Plasticity, 2018, 2018:5382537.
413	23. Payne, J. K. State of the Science: Stress, Inflammation, and Cancer. Oncology
414	Nursing Forum, 2014, 41(5):533–540.
415	24. Shobeiri, Aali, A., Khan, I. A., Khan, M. A, Kim, M. S., et al. Global burden of
416	chronic respiratory diseases and risk factors, 1990–2019: an update from the Global
417	Burden of Disease Study 2019. EClinicalMedicine, 2023, 59, 101936–101936.

418	25. Simpson, & Camorlinga, S. G. A Framework to Study the Emergence of Non-
419	Communicable Diseases. Procedia Computer Science, 2017, 114, 116-125
420	26. Sommer, Griebler, U., Mahlknecht, P., Thaler, K., Bouskill, K., Gartlehner, G., &
421	Mendis, S. Socioeconomic inequalities in non-communicable diseases and their risk
422	factors: an overview of systematic reviews. BMC Public Health, 2015, 15(1), 914-
423	914
424	27. Asarnow, Goldston, D. B., Tunno, A. M., Inscoe, A. B., & Pynoos, R. Suicide, Self-
425	Harm, & Traumatic Stress Exposure: A Trauma-Informed Approach to the Evaluation
426	and Management of Suicide Risk. Evidence-Based Practice in Child and Adolescent
427	Mental Health, 2020, 5(4), 483–500.
428	28. Heidari, M., Ghodusi Borujeni, M., Kabirian Abyaneh, S., & Rezaei, P. The Effect of
429	Spiritual Care on Perceived Stress and Mental Health Among the Elderlies Living in
430	Nursing Home. Journal of Religion and Health, 2019, 58(4):1328–1339.
431	29. Kim, Jeon, SW., Cho, S. J., Shin, Y. C., & Park, JH. Comparison of the factors for
432	suicidal ideation and suicide attempt: a comprehensive examination of stress, view of
433	life, mental health, and alcohol use. Asian Journal of Psychiatry, 2021, 65, 102844
434	30. Jahangeer, S. M. A., Hasnain, N., Tariq, M. T., Jamil, A., Zia, S. Y., & Amir, W
435	Frequency and Association of Stress Levels with Modes of Commuting Among
436	Medical Students of a Developing Country. The Malaysian Journal of Medical
437	Sciences, 2021, 28(4), 113–122
438	31. Magaña, P, Roberto, G., Sara, P., & Laura, P. Beside and Behind the Wheel: Factors
439	that Influence Driving Stress and Driving Behavior. Sustainability (Basel,
440	Switzerland), 2021, 13(4775), 4775-


441	32. Plyushteva, A. Exploring the Role of the Workplace in Experiences of Commuter
442	Stress: A Mixed-Method Study from Sofia, Bulgaria. Social Sciences, 2021, 10(208),
443	208.
444	33. Rezapour, M., & Richard Ferraro, F. The impact of commuters' psychological
445	feelings due to delay on perceived quality of a rail transport. Humanities and Social
446	Sciences Communications, 2021, 8(1), 1–8.
447	34. Rosida, I., Sari, D. W., & Irjayanti, A. D. The Mode Choices And Commuting Stress:
448	Empirical Evidence From Jakarta And Denpasar. Jurnal Pengembangan Kota, 2019
449	7(1), 68–76.
450	35. Brutus, S., Javadian, R., & Panaccio, A. J. Cycling, car, or public transit: A study of
451	stress and mood upon arrival at work. International Journal of Workplace Health
452	Management, 2017, 10(1), 13-24
453	36. Legrain, A., Eluru, N., & El-Geneidy, A. M. Am stressed, must travel: The
454	relationship between mode choice and commuting stress. Transportation Research
455	Part F: Traffic Psychology and Behaviour, 2015, 34, 141–151.
456	37. Yang, X., McCoy, E., Anaya-Boig, E., Avila-Palencia, I., Brand, C., Carrasco-
457	Turigas, G., et al The effects of traveling in different transport modes on galvanic
458	skin response (GSR) as a measure of stress: An observational study. Environment
459	International, 2021, 156, 106764
460	38. Kaldis, B. The university as microcosm. Educational Philosophy and Theory, 2009,
461	41(5), 553-574.
462	39. Struppa, D. A university is a microcosm of the country: Daniele Struppa. Orange
463	County Register. (2021, Mar. 7). https://www.ocregister.com/2021/03/07/a-
464	university-is-a-microcosm-of-the-country-daniele-struppa/

465	40. Stone AA, Neale JM, Shiftman S. Daily assessments of stress and coping and their
466	association with mood. Annals of Behavioral Medicine. 1993 Mar 1;15(1):8-16
467	41. Maybery D, Maybery M, Bresnan R, Croft B, Graham R, Macaulay J, McQualter S,
468	Mitchell E, Sherwell K, Szakacs E. Responding to daily event questionnaires: The
469	influence of the order of hassle and uplift scales. Stress and health. 2002
470	Feb;18(1):19-26.
471	42. Udayar S, Urbanaviciute I, Morselli D, Bollmann G, Rossier J, Spini D. The LIVES
472	daily hassles scale and its relation to life satisfaction. Assessment. 2023
473	Mar;30(2):348-63.
474	43. Lee EH. Review of the psychometric evidence of the perceived stress scale. Asian
475	Nursing Research. 2012 Dec 1;6(4):121-7.
476	44. Reis RS, Hino AA, Añez CR. Perceived stress scale. J. health Psychol.
477	2010;15(1):107-14.
478	45. Smith KJ, Emerson DJ. An assessment of the psychometric properties of the
479	Perceived Stress Scale-10 (PSS10) with a US public accounting sample. Advances in
480	Accounting. 2014 Dec 1;30(2):309-14.
481	46. Fung. Validity of the brief resilience scale and brief resilient coping scale in a Chinese
482	sample. International Journal of Environmental Research and Public Health, 2020
483	17(4), 1265–.
484	47. Kocalevent, Zenger, M., Hinz, A., Klapp, B., & Brähler, E. Resilient coping in the
485	general population: Standardization of the brief resilient coping scale (BRCS). Health
486	and Quality of Life Outcomes, 2017 15(1), 251–251
487	48. Sinclair, & Wallston, K. A. The Development and Psychometric Evaluation of the
488	Brief Resilient Coping Scale. Assessment (Odessa, Fla.), 2004 11(1), 94-101.

489	49. Ye, Wu, CH., Huang, TY., & Yang, CT The difference between the Connor-
490	Davidson Resilience Scale and the Brief Resilience Scale when assessing resilience:
491	confirmatory factor analysis and predictive effects. Global Mental Health, 2022 9,
492	339–346.
493	50. Konaszewski, Niesiobędzka, M., & Surzykiewicz, J. Validation of the Polish version
494	of the Brief Resilience Scale (BRS). PloS One, 2020 15(8), e0237038-e0237038
495	51. Sánchez, Estrada-Hernández, N., Booth, J., & Pan, D. Factor structure, internal
496	reliability, and construct validity of the Brief Resilience Scale (BRS): A study on
497	persons with serious mental illness living in the community. Psychology and
498	Psychotherapy, 2021 94(3), 620-645
499	52. Surzykiewicz J, Konaszewski K, Skalski S, Dobrakowski PP, Muszyńska J.
500	Resilience and mental health in the polish population during the COVID-19
501	lockdown: A mediation analysis. Journal of clinical medicine. 2021 26;10(21):4974.
502	53. Baer, Carmody, J., & Hunsinger, M Weekly Change in Mindfulness and Perceived
503	Stress in a Mindfulness-Based Stress Reduction Program: Weekly Change in
504	Mindfulness and Stress in MBSR. Journal of Clinical Psychology, 2012, 68(7), 755-
505	765.
506	54. Uysal, & Çalışkan, B. B. The effects of mindfulness-based stress reduction on
507	mindfulness and stress levels of nursing students during first clinical experience.
508	Perspectives in Psychiatric Care, 2022 58(4), 2639–2645.



# Figure 1: Distribution of symptoms experienced by respondents



# Figure