

1 ***“If I am free from diabetes, that itself will be the happiest thing”:***

2 **A convergent mixed methods study of the lived experiences of young**

3 **adults with type 2 diabetes in Mysore district, India**

4

5 Nikhita R. Gopisetty^{1*}, Kiranmayee Muralidhar^{2,3}, Nagalambika Ningaiah², Rani

6 Chinnappa², Mia Buono¹, Poornima Jaykrishna², Purnima Madhivanan^{2,4}, Sumedha G.

7 Ariely^{1†}, Eve S. Puffer^{1,5†}

8

9 ¹ Duke Global Health Institute, Duke University, Durham, North Carolina, United States

10 of America

11 ² Public Health Research Institute of India, Mysore, Karnataka, India

12 ³ JSS Academy of Higher Education and Research, Mysore, Karnataka, India

13 ⁴ Mel & Enid Zuckerman College of Public Health, University of Arizona, Tucson,

14 Arizona, United States of America

15 ⁵ Department of Psychology and Neuroscience, Duke University, Durham, North

16 Carolina, United States of America

17

18 * Corresponding author: nikhita.gopisetty@duke.edu (NRG)

19 † These authors contributed equally to this work.

20 **Abstract**

21 Type 2 diabetes (T2D) has been occurring at younger ages of onset around the world.
22 India's population accounts for nearly 20% of the global disease burden. This study
23 investigated the occurrence of depressive symptoms and qualitatively explored the
24 lived experiences of 20 young adults living with T2D under the age of 35. We conducted
25 a convergent mixed-methods study with the Patient Health Questionnaire (PHQ-9) and
26 semi-structured interviews from June 2022 to July 2022 in Mysore district, India.
27 Guided by the World Health Organization's Commission on Social Determinants of
28 Health conceptual framework and biopsychosocial frameworks, areas of inquiry
29 included knowledge and perception about T2D, accessibility of healthcare resources,
30 T2D-related self-care activities, and the impact of the condition on their daily life.
31 Interviews were debriefed by the research team and analyzed thematically using NVivo
32 12. Participants were aged between 21 and 35 (mean: 30.8, SD: 4.2) and the majority
33 were female (75%). Overall, 55% reported mild depression symptoms, 15% reported
34 moderate to moderately severe depression symptoms; 5 participants (25%) reported
35 suicidality. Sex, living in rural Mysore district, socioeconomic status, T2D duration,
36 family history of T2D, T2D-induced complications, and T2D-related self-care behaviors
37 were associated with depressive symptoms. Thematic analysis revealed 1) low
38 knowledge about T2D, 2) substantial interpersonal and internalized stigma for having
39 T2D at a young age, 3) financial and time constraints to seek and receive care, 4) self-

40 perception as burdens to family members due to the cost and stress of living with T2D,
41 5) competing priorities with work and family, and 6) the power of social support in
42 managing T2D. These themes were consistent across the sample, regardless of severity
43 of depressive symptoms. Awareness campaigns and peer support programs may help
44 reduce depressive symptoms and increase self-efficacy in this population.

45

46 **Keywords:** Type 2 Diabetes; Young Adult; Mental Health; India; Stigma; Depressive
47 Symptoms

48 **Introduction**

49 Diabetes is a global epidemic – in 2019, over 450 million adults were estimated
50 to be living with type 2 diabetes (T2D) and this number is predicted to exceed 700
51 million by 2045 [1]. Almost 80% of people living with T2D live in low- and middle-
52 income countries (LMICs); over 77 million people with T2D live in India, of which over
53 50% are undiagnosed [2]. Diabetes is occurring at a younger age of onset and lower BMI
54 levels across India and around the world. In rural Pondicherry, India, one third of new
55 diabetes cases occur in people below 40 years old [3]. In younger populations living
56 with T2D, comorbidities also occur at a younger age, including fatty liver disease and
57 systolic hypertension [4]. Diabetes-related complications impact the physical, mental,
58 and social well-being of patients, placing a burden on the healthcare system [5]. Sub-
59 Saharan Africa, South America, and parts of Asia have yet to experience the
60 epidemiological transition from communicable diseases to non-communicable diseases
61 – India is an example to learn from before this occurs [6].

62 Prior research has established a bidirectional relationship between diabetes and
63 depression [7–9]. In urban-based clinics around India, researchers found that one-
64 quarter to one-third of patients with T2D were depressed. They predict that the
65 prevalence of depression in people living with T2D might be higher in LMICs than high
66 income countries [10]. Depressive disorders in people living with T2D are associated
67 with the female gender, young age of onset, lower education levels, worse glycemic

68 control, a lack of physician advice about lifestyle modifications, and low social support
69 [7,8,11–13]. Lower-income populations exhibit higher rates of depression, due to a lack
70 of access to healthcare, higher likelihood of developing complications, and a low social
71 awareness [14]. The comorbidity of depression and chronic disease is associated with
72 worse physical disorder outcomes and quality of life [7,13].

73 Previous research has shown younger age to be significantly associated with
74 diabetes distress in South India [15]. Young adults living with diabetes have been
75 shown to have lower self-esteem than those without, due to social perceptions and
76 preconceived notions about the condition [16]. Key emotions in young diabetics
77 discussing their disease included denial, anger, depression, and guilt. They feared
78 people would treat them differently and perceive them as sick, and experienced
79 unsolicited advice and instructions about household remedies from family members
80 and peers [17]. Diabetes occurring at a younger age is also associated with significant
81 social stigma, due to societal and cultural misconceptions about the disease [18].

82 As T2D becomes more prevalent in younger populations, it is crucial to
83 understand the various barriers to diabetes management and quality of life that they
84 face. A conceptual model, derived from the WHO's Commission on Social Determinants
85 of Health (CSDH) conceptual framework [19] and the biopsychosocial model [20], was
86 developed to guide this study (Fig 1). The CSDH framework presents the relationships
87 between structural and intermediary social determinants of health and equity in health

88 and well-being [19]. The biopsychosocial model of health and illness allows researchers
89 to model any disease caused and impacted by biological, psychological, and social
90 factors, revealing interactions between the mind and body [20]. The conceptual model
91 below maps out various pathways between sociodemographic characteristics, structural
92 and intermediary determinants, T2D-related self-care behaviors, physical health
93 outcomes, and quality of life. Intermediary determinants -- including health literacy,
94 access, social factors, and psychological factors -- are influenced by an individual's
95 environment and directly impact self-care behaviors and health outcomes. The purpose
96 of this study is to apply this conceptual model to explore the impact of intermediary
97 determinants of health on the prevalence of depressive symptoms and lived
98 experiences in young adults living with T2D in Mysore district, Karnataka, India.

99 **Fig 1. Conceptual model**

100

101 **Methods**

102 **Study design**

103 We used a convergent mixed-methods approach, with an emphasis on deeper
104 understanding and analysis of qualitative themes [21]. Qualitative and quantitative
105 data were collected concurrently and findings were interpreted with the intent of
106 validating and describing relationships among the data. Approval was obtained from

107 Duke Campus Institutional Review Board (2022-0358) and the Public Health Research
108 Institute of India's Institution Ethics Review Board (2022-01-29-64).

109 **Setting**

110 The study was conducted in Mysore, a district in southern Karnataka in India
111 between June and July 2022. The Indian National Census in 2011 reported a total
112 population of 3,001,127 in Mysore district, with approximately 58% living in rural
113 villages [22]. The National Family Health Survey-5 (NFHS-5) from 2019 to 2021
114 reported that India's population is young, which is typical of developing countries with
115 a low life expectancy; over 50% of India's population is below the age of 30 [23]. The
116 study was implemented in collaboration with the Public Health Research Institute of
117 India (PHRII), a research and service-based organization in Mysore, Karnataka. PHRII
118 has been conducting community-based research, providing health services, and
119 training public health professionals in Mysore district for the past 15 years.

120 **Author positionality**

121 It is important to recognize the authors' positionality and, therefore, their
122 interpretation of the data. The research team consisted of public health practitioners,
123 academic researchers, physicians, administrative staff, a graduate student, and an
124 undergraduate student. Seven members of the research team come from Indian
125 backgrounds, two born in the United States and five born in India. Two members of the
126 research team are United States-born White Americans with experience designing and

127 implementing research projects in global settings. All authors worked as a team to
128 ensure the study was guided by their collective cultural knowledge and expertise.
129 Positionality challenges related to sex, gender, race/ethnicity, power, socioeconomic
130 status, and privilege were intentionally addressed throughout the research design and
131 data collection processes through team-based reflection on how these issues might
132 impact the interview process and data analysis.

133 **Eligibility and participant recruitment**

134 Young adults with T2D were recruited using convenience sampling between June
135 6, 2022 and July 8, 2022. Participants were eligible to take part in the study if they were
136 between 18 to 35 years old, self-reported to have T2D, live in Mysore district, able to
137 speak English or Kannada, and able to provide informed consent. The research team
138 consulted with Primary Health Centres, Accredited Social Health Activists (ASHAs),
139 Urban Social Health Activists (USHAs), and the Karnataka Health Promotion Trust to
140 identify potential participants who met the inclusion criteria. Alongside the ASHAs and
141 USHAs, government-supported community health workers, the research team
142 physically approached potential participants in their communities to inform them of
143 the study. Written informed consent was obtained from the study participants after
144 explaining the purpose of the study and reading through the content of the study
145 information sheet and consent form. Of 21 eligible candidates that were approached,

146 one refused to participate in the study. This resulted in 20 eligible participants, 10 from
147 urban and 10 from rural Mysore district.

148 **Measures**

149 **Quantitative**

150 Participants completed a demographic questionnaire that included
151 sociodemographic characteristics, medical history information, and details regarding
152 their diagnosis with T2D and related management. Participants were screened for
153 depressive symptoms using the Patient Health Questionnaire-9 (PHQ-9) [24], a brief
154 self-report instrument of depression severity that has been translated and validated for
155 Indian populations in various languages [25]. The screening tool uses 9 questions
156 scored on a 4-point Likert-scale ranging from 0 (not at all) to 3 (nearly every day) to
157 examine symptoms over the past two-week period. Individual's PHQ-9 score was
158 categorized: 1-4 minimal depression, 5-9 mild depression, 10-14 moderate depression,
159 15-19 moderately severe depression, and 20-27 severe depression [24]. A single
160 interviewer (NN) and observing notetaker (RC) verbally administered the surveys in
161 *Kannada* and recorded responses through an offline Qualtrics survey.

162 **Qualitative**

163 A semi-structured interview guide (S1 File) was developed based on the
164 conceptual model. This guide covered the following domains and themes: general
165 knowledge and perception about T2D, accessibility to healthcare resources, self-care

166 behaviors related to T2D, impact of T2D on the individual, and impact of T2D on others
167 around them. The interview guide was developed in English by the authors (NRG, KM,
168 PJ, SA, ESP) and subsequently translated into *Kannada* by the interviewer. The guide
169 was back-translated into English to ensure language consistency and validity of the
170 questions. To assess the flow and colloquially adapt the language of the interview
171 guide, three mock interviews were conducted by the interviewer and staff members at
172 PHRII, who understood the local language and context where the study was conducted.
173 During each mock interview, staff members provided feedback on the language of the
174 questions and cultural appropriateness of the topics. Subsequent changes were
175 incorporated following each mock interview to make the wording of questions more
176 understandable and improve the flow of the conversation.

177 Interviews were completed either in a quiet, private room at PHRII or the
178 participant's home. Interviews were conducted by a single interviewer (NN) who was
179 trained in qualitative research methods and had experience conducting semi-
180 structured, in-depth interviews. Interviews lasted for approximately 60 minutes and
181 were audio recorded on a digital voice recorder. During the interviews, the interviewer
182 used probes to encourage participants to share anything relevant to the topics
183 discussed, beyond those laid out in the interview guide. The notetaker (RC) observed
184 and documented key observations from each interview.

185 **Analysis**

186 **Quantitative**

187 All quantitative analyses and data visualizations were conducted in Microsoft
188 Excel (2021) and RStudio (version 4.2.1; R Core Team 2021). Socioeconomic status was
189 classified by the modified BG Prasad scale, using annual household income and family
190 size to calculate per-capita income [26]. Individual's depressive symptoms were scored
191 by summing the total PHQ-9 values for all 9 questions, with higher scores
192 corresponding to increased symptomatology [24].

193 **Qualitative**

194 The research team debriefed each interview and discussed field notes within 48
195 hours of completion. Interviews were transcribed verbatim and translated to English.
196 The transcribed interviews were reviewed by the first author (NRG) to ensure
197 transcriptions were accurate and complete. Debrief notes were correlated to the
198 respective transcripts upon completion. Two researchers (NRG, KM) familiarized
199 themselves with the data and generated a codebook for thematic qualitative analysis,
200 based on the preconceived conceptual model (Fig 1). Two researchers (NRG, MB)
201 independently coded the transcripts and debrief notes with NVivo 12 software.
202 Interrater reliability was established with a kappa coefficient of 0.86, indicating almost
203 complete agreement [27]. Discrepancies in coding between the investigators were
204 resolved through consensus with a third researcher (KM). Emergent themes and

205 relevant sub-themes were then discussed among all authors to promote consistency
206 and minimize bias. Thematic saturation was reached after analyzing data from the
207 interviews, ensuring comprehensive coverage of the study's qualitative findings.

208

209 Results

210 Twenty participants with T2D participated in the study, of which 15 were female.
211 Twelve participants were diagnosed with T2D within the past two years. The age of
212 participants ranged from 21 to 35 years with a mean age of 30.8 (SD \pm 4.2) All spoke
213 Kannada as their first language. Seven were lower-middle class and six were middle
214 class, according to the modified BG Prasad scale. Table 1 presents key health and
215 demographic characteristics of the study participants, stratified by rural/urban
216 distinction.

217 **Table 1. Participant sociodemographic and health behavioral characteristics**

	Total (N=20) N (%)	Rural (N=10) N (%)	Urban (N=10) N (%)
Age (mean \pm SD)	30.8 \pm 4.19	29.2 \pm 4.73	32.4 \pm 2.99
T2D Duration (years)			
< 2	12 (60)	7 (70)	5 (50)
3+	8 (40)	3 (30)	5 (50)
Sex			
Female	15 (75)	8 (80)	7 (70)
Religion			
Hindu	20 (100)	10 (100)	10 (100)
Socioeconomic Status			
Lower-Middle Class	7 (35%)	3 (30)	4 (40)
Middle Class	6 (30%)	4 (40)	2 (20)
Upper-Middle Class	4 (20%)	2 (20)	2 (20)
Upper Class	3 (15%)	1 (10)	2 (20)

Education Level

High School	5 (25)	3 (30)	2 (20)
Secondary School	8 (75)	7 (70)	8 (80)

Marital Status

Married	16 (80)	9 (90)	7 (70)
Single	4 (20)	1 (0)	3 (30)

Children

0	6 (30)	2 (20)	4 (40)
1	3 (15)	2 (20)	1 (10)
2+	11 (55)	6 (60)	5 (50)

Family History of T2D

	15 (75)	6 (60)	9 (90)
--	---------	--------	--------

Medication^a

Oral Medication	17 (85)	8 (80)	9 (90)
Insulin Injections	2 (10)	0 (0)	2 (20)
No Medication	2 (10)	2 (20)	0 (0)
Ayurvedic Medication	2 (10)	0 (0)	2 (20)

Frequency of Doctor Visits

1-3 times a year	5 (25)	3 (30)	2 (20)
> 4 times a year	2 (10)	0 (0)	2 (20)
Monthly	13 (65)	7 (70)	6 (60)

Doctor Training

General Practitioner	13 (65)	7 (70)	6 (60)
Specialist	7 (35)	3 (30)	4 (40)

T2D-Related Complications^b

	5 (25)	4 (40)	1 (10)
--	--------	--------	--------

Frequency of Blood Sugar Testing

Only at doctor's appointment	7 (35)	2 (20)	5 (50)
Once a month	11 (55)	8 (80)	3 (30)
Once a week	2 (10)	0 (0)	2 (20)

Days of Physical Activity per Week

0	9 (45)	6 (60)	3 (30)
1-4	3 (15)	2 (20)	1 (10)
5-7	8 (40)	2 (20)	6 (60)

Follow a Diabetic Diet

	16 (80)	9 (90)	7 (70)
--	---------	--------	--------

218 ^a total percentage greater than 100 because of combination treatments

219 ^b eye-related complications (N = 3), reproductive health complications (N = 2)

220 **Depressive symptoms**

221 Table 2 presents the results of the PHQ-9 screening for the participants. While a
 222 small sample size, rates of symptoms were high. Of the 20 participants, over half (55%,
 223 N = 11) of the study participants had scores in the mild depression range, 15% (N = 3) in

224 the moderate to moderately severe depression range. Five participants reported having
225 suicidal thoughts in the past 2 weeks -- 3 with scores in the mild depression range and
226 2 in the moderately severe depression range. The highest scores were associated with
227 the following 3 prompts: 1) Little interest or pleasure in doing things; 2) Feeling down,
228 depressed, or hopeless; and 3) Feeling tired or having little energy.

229 **Table 2. Depression severity (PHQ-9)**

Depression Severity	N (%)
Minimal	6 (30)
Mild	11 (55)
Moderate / Moderately Severe	3 (15)
Suicidal Ideation (Past 2 Weeks)	5 (25)

230
231 Fig 2 visually presents the relationships between covariates and depressive
232 symptoms. Participants who identified as female, resided in rural Mysore district, and
233 belonged to a lower socioeconomic status were more likely to have depressive
234 symptoms. Participants who were diagnosed with T2D within 2 years, did not have a
235 family history of T2D, faced T2D-related complications, adhered to a diabetic diet, and
236 neglected physical activity in their routine were also more prone to depressive
237 symptoms.

238 **Fig 2. Association of covariates with depressive symptoms**

239 **Intermediary determinants of diabetes self-management**

240 **Health literacy**

241 When asked how many people in their community have diabetes, participants
242 predicted an average of 70% (SD 16.8%), indicating the seemingly high prevalence. In
243 2021, the prevalence of diabetes in India was reported at 9.6%, drastically different
244 from the participants' perception [2]. Participants across rural and urban Mysore
245 district showed varied levels of awareness about the prevalence of diabetes in younger
246 populations. In rural Mysore district, 90% of participants stated that they did not know
247 other people their age living with the condition:

248 *“People with very young age will not be affected by diabetes... I think I am*
249 *exceptional.”* (P10, Female, PHQ-9 = 6, Mild Depression)

250 However, in urban Mysore district, participants reported greater awareness about the
251 rising prevalence of diabetes at younger ages:

252 *“Now almost everyone is becoming diabetic at a younger age. If the parents have it,*
253 *the kids might acquire it too, and with that, the food we consume is also not good.*

254 *So, from that as well, people are getting sugar [diabetes] soon.”* (P15, Female, PHQ-
255 9 = 6, Mild Depression)

256 Overall, participants reported having a lack of trustworthy sources to learn about
257 diabetes. As a result, 7 (35%) felt scared to ask questions to better understand their
258 biological condition, due to false information or societal stigma:

259 *“Few of them will say that food may also cause diabetes, but few say that it is not*
260 *because of food. I don’t know whom I should listen and follow.”* (P10, Female, PHQ-
261 9 = 6, Mild Depression)

262 Participants reported that their healthcare providers recommended them to
263 maintain a diabetic diet, perform physical activity regularly, and to avoid stress to
264 manage their condition. However, due to the large quantity of patients in the clinic,
265 participants rarely experienced individualized care. One participant stated:

266 *“My doctor won’t specifically mention everything, they will just say to have proper*
267 *food intake and be calm without worrying about anything. Sometimes I feel bad that*
268 *they will not tell me properly about diabetes.”* (P9, Female, PHQ-9 = 15, Moderately
269 Severe Depression)

270 All of the participants knew that diabetes-related complications exist, but the majority
271 (65%) were not adequately informed about the complications and ways to prevent
272 them:

273 *“They [doctors] said that if there is a wound in the leg, they will amputate it and*
274 *there will be loss of vision. If there is a wound in the hand, they will amputate it.”*
275 (P11, Female, PHQ-9 = 9, Mild Depression)

276 Nearly all participants (90%) explicitly stated that they do not want to take medication
277 for the rest of their lives. These perceptions are influenced by participants’ community
278 members and even healthcare professionals.

279 *“They have said to control diabetes by taking proper food and stopping the tablets.*
280 *Even our family doctor suggested I should stop taking tablets and not develop a habit*
281 *of taking them forever.”* (P13, Male, PHQ-9 = 0, Minimal Depression)

282 Twelve participants (60%) used language about “addiction” when referring to
283 their diabetes medication, claiming that they do not want to become addicted to the
284 tablets:

285 *“If I get addicted to tablets, it will be difficult. I want to control it [diabetes] in food.*
286 *So, I am not taking any tablets.”* (P11, Female, PHQ-9 = 9, Mild Depression)

287 **Access**

288 In rural Mysore district, participants reported constant crowds at the
289 government hospitals, requiring patients to enter a queue just to meet with the doctor.
290 This comes alongside various challenges – women must consider who will accompany
291 them to their appointments, working professionals must request leave from their jobs
292 in advance, and all must find transportation to and from the clinic. One participant
293 described her challenges with seeing a healthcare provider:

294 *“Sometimes doctors will not be there. One day I waited for so many hours but the*
295 *doctor did not come to the hospital. So I came back home, if he is not there I will go*
296 *the next day.”* (P10, Female, PHQ-9 = 6, Mild Depression)

297 Seven participants (35%) stated that they prefer getting their healthcare at
298 private clinics because they believe that if they are paying for it, the service must be

299 better: *“I feel that whatever the money I spend, my health should be fine and good. So I will*
300 *be happy.”* (P2, Female, PHQ-9 = 15, Moderately Severe Depression).

301 With a greater presence of healthcare facilities, there are more opportunities to
302 meet with a doctor in urban Mysore district. Some doctors in the urban region even
303 adjust their hours to be available after the workday. However, participants reported
304 that crowds are still present at the clinics. Due to the crowds and long queues, some
305 patients reported feeling like a burden if they take too much time with the doctor,
306 which interferes with their own care. Some participants described:

307 *“There will be so many patients like me, if I take too much time to discuss my*
308 *problems, it will get very late for other patients. I will be troubling them. So I will*
309 *think that I can talk to him the next time when I go for a checkup.”* (P12, Female,
310 PHQ-9 = 7, Mild Depression)

311 Blame from healthcare providers emerged as a theme, illustrating some of the
312 complicated and diverse power dynamics between patients and their providers. Over
313 half of participants in urban Mysore district reported that they experienced blame from
314 their providers if their diabetes was not under control.

315 *“She [doctor] actually scolds me to take proper care of my health. She also*
316 *mentioned that due to my negligence I am unable to take proper care of my health. If*
317 *I don’t start taking care of it she has warned me to not visit the clinic.”* (P20,
318 Female, PHQ-9 = 7, Mild Depression)

319 Despite these challenges, some patients are willing to invest the time to take
320 care of their health. They navigate the difficulty with positive attitudes:

321 *“If we want to get the medicines we should wait, it’s not like they should only treat*
322 *me, there will be so many people like me. Even they will have some problem, we can’t*
323 *say that our time is getting wasted and blame doctors. If we are unable to contact*
324 *him today, we can visit the next day.”* (P6, Male, PHQ-9 = 0, Minimal Depression)

325 **Financial burden**

326 Due to competing financial priorities, most participants (65%) are unable to
327 afford their diabetes medication. All of the participants with children mentioned that
328 they did not want their diabetes to interfere with their children’s education:

329 *“I cannot spend my wages on my tablets. I have to take care of my children’s*
330 *education.”* (P18, Female, PHQ-9 = 6, Mild Depression).

331 Some participants reported skipping doses if they do not have the means to
332 purchase it, and others choose to not spend their limited expendable income on their
333 medications. One participant reported that she preferred government-subsidized
334 medications, but they are not available in her area. Three participants reported that
335 they try to maximize their limited supplies of medication by using it as a reactive
336 mechanism, rather than a preventative mechanism. They explained that they wait to
337 take the medication until they experience symptoms of high blood sugar levels:

338 *“When I get the burning sensation in my feet, I come to know that my sugar level is*
339 *high. Then, I take one tablet and gulp it. That is all.”* (P11, Female, PHQ-9 = 9, Mild
340 Depression)

341 Under financial constraints, people living with diabetes find themselves having
342 to choose between speaking to their doctor or purchasing their medication due to the
343 costs involved. In these scenarios, they are managing their condition blindly, until they
344 can visit their doctor again.

345 *“I will not go for a blood test every month because of financial problems. I will visit*
346 *the private hospital so the cost will be more, so I will just purchase tablets.”* (P9,
347 Female, PHQ-9 = 15, Moderately Severe Depression)

348 For some, their diabetes is still poorly controlled, so they questioned the purpose
349 of spending their money:

350 *“I purchase the tablets and I give the amount for appointments. Even though I spend*
351 *so much, my [blood] sugar will not get reduced. I am fed up with it.”* (P5, Female,
352 PHQ-9 = 13, Moderate Depression)

353 **Social factors**

354 During the interviews, all of the participants described encounters with people
355 who believed that diabetes was communicable, both in the rural and urban regions.
356 This misconception was so well-established in certain communities that 5 (25%)

357 participants reported never seeing their neighbors again after they shared their
358 diagnosis.

359 *“Once I was going out to the hospital and my neighbor asked what the problem was.*
360 *I said I have diabetes. They thought it was contagious because I have got diabetes at*
361 *an early age. They decided not to have any contact with us after that.”* (P8, Female,
362 PHQ-9 = 9, Mild Depression)

363 All of the participants reported experiences of teasing from community members
364 for having diabetes at a young age. Despite its ubiquity, it is still a taboo topic. The
365 young adults described feeling embarrassed to share their diagnosis because others talk
366 and laugh about them, even if they *“pretend to act like they care”* (P1, Female, PHQ-9 =
367 9, Mild Depression):

368 *“I do feel embarrassed to share. People who stay in the village will tease me, saying*
369 *that at an early age I have got diabetes. Some will understand, but not all will get to*
370 *know our problems.”* (P4, Female, PHQ-9 = 1, Minimal Depression)

371 Fear of this community gossip and stigma often deterred participants from going
372 to public health camps or visiting the community health workers, out of fear that
373 someone would *“watch and talk behind [her] back to the neighbors”* (P10, Female, PHQ-9
374 = 6, Mild Depression).

375 *“The ASHA (Accredited Social Health Activist) workers ask me to get the sugar*
376 *tested in camps, but I do not go and I do not like to go. When I have to go for a*

377 *checkup, I just tell others that I am going out. I will never disclose about sugar*
378 *[diabetes] checkup because I do not like to say it. The way they look at us is different.*
379 *So, I do not wish to disclose. That is sad.” (P18, Female, PHQ-9 = 6, Mild*
380 *Depression)*

381 Due to the prevalent social stigma, the 3 unmarried participants described their
382 fear of not finding a partner because of their diagnosis with diabetes. One married
383 participant shared that she did not inform her partner of her diagnosis until after the
384 wedding ceremony was complete, out of fear that his family would no longer approve of
385 the union.

386 **Barriers to diabetes self-management**

387 Sixteen participants (80%) self-reported following a diabetic diet, different from
388 the rest of their family members, increasing the burden on whoever is cooking the food.
389 Many participants spoke about how they would rather eat the same food as everyone
390 else than burden others. Some discussed the concept of portion control, while others
391 just eat the same to fit in.

392 *“When we are all together, I can’t ask for separate food. I should have what all*
393 *others are having.” (P3, Female, PHQ-9 = 0, Minimal Depression)*

394 Two participants (10%) described forgoing their recommended diabetic diet due
395 to fear of what others would think about them. If they were to change their lunches at

396 their workplace, they were afraid that their condition would be revealed to their
397 employer and coworkers:

398 *“I will have breakfast early in the morning. But for my workplace, I will carry rice.*

399 *They will think I am someone special if I don’t, so I have to carry rice for lunch.” (P3,*

400 Female, PHQ-9 = 0, Minimal Depression)

401 Some female participants do not feel comfortable going for a walk or performing
402 physical activity in public by themselves, especially at night. Then, they are reliant on
403 their husbands to accompany them on their walks.

404 *“In the evening it will be late and it becomes dark, so I will not go alone.” (P1,*

405 Female, PHQ-9 = 9, Mild Depression)

406 **Importance of positive social support**

407 The presence of familial support is known to be an important factor in diabetes
408 management and quality of life. Some female participants reported encouraging
409 support from their husbands to visit the hospital and reduce stress:

410 *“My husband will say that no matter how much money or time we spend, I should be*

411 *healthy without any problems. He is even ready to take me to the other hospital. He*

412 *is not worried about money.” (P10, Female, PHQ-9 = 6, Mild Depression)*

413 One participant described her close relationship with a friend who also has T2D.

414 Her friend was successfully managing her glycemic control and served as an inspiration

415 and motivation for her to take care of her own health.

416 *“One of my friends has got diagnosed with diabetes. She is having regular spouts in*
417 *her daily food routine and she is doing some exercise, going for a walk to maintain*
418 *her health. I feel if I do the same I may also have a proper control in sugar level and*
419 *can be healthy.”* (P10, Female, PHQ-9 = 6, Mild Depression)

420 **Psychological factors**

421 Feelings of sadness and depressive symptoms after diagnosis with diabetes was
422 common amongst study participants. Most participants (75%) described a sentiment of
423 “why me?” and “what else could happen to me?”

424 *“I felt sad and worried about the occurrence of diabetes at this young age, and also*
425 *felt sad that I cannot eat most of the things which I like.”* (P1, Female, PHQ-9 = 9,
426 Mild Depression)

427 Some participants relate their overall happiness to their diabetes diagnosis and
428 glycemic control, resulting in a lack of individual control over their general well-being.

429 *“If the sugar level is high, I will be very tensed. If it’s normal, I will be happy and*
430 *calm.”* (P10, Female, PHQ-9 = 6, Mild Depression)

431 A feeling of worry due to living with diabetes at a young age was common across
432 the study participants. This ranges from impeding routine activities, not being able to
433 manage their glycemic control, and troubling others in their family.

434 *“I got it from my mom, and I don’t want it to happen to my kids because of me. I have*
435 *two daughters. What if they have problems when they get married and have*

436 *complications during their pregnancies? Now, people are getting diabetic during*
437 *their pregnancy. I had two kids and after that I got diabetes, but people in the*
438 *present generation are getting diabetes during pregnancy. I am afraid about that.”*

439 (P15, Female, PHQ-9 = 6, Mild Depression)

440 All but one participant expressed fear about future complications and early
441 death due to poorly controlled diabetes:

442 *“I feel that I may die early, I will be worried that there will be no one to take care of*
443 *my daughter. They also say that there will be more consequences like kidney stone,*
444 *heart disease, and also some may lose their eyesight. So I am worried about all of*
445 *these.”* (P8, Female, PHQ-9 = 9, Mild Depression)

446 This fear of complications and death sometimes translates to a fear of going to
447 the doctor and addressing concerns about their own health.

448 *“I feel scared to discuss with them, I feel they might say something new. Now I am*
449 *already suffering from a lot, I don’t want to suffer even more.”* (P9, Female, PHQ-9 =
450 15, Moderately-Severe Depression)

451 Many participants expressed sentiments of feeling like a burden to their family
452 members:

453 *“I definitely feel that I am becoming a trouble to someone. I also feel I should not*
454 *depend on anyone, neither my husband nor son. I should take care of myself.”* (P12,
455 Female, PHQ-9 = 7, Mild Depression)

456 For some, this feeling had built over time to suicidal thoughts.

457 *“I feel that why should I live, all are facing problems because of me. No one will take*
458 *care of me except my husband. So I feel I shouldn’t be here, why should everyone*
459 *suffer because of me?”* (P8, Female, PHQ-9 = 9, Mild Depression)

460 Participants reported a significant impact to their energy levels and working
461 capacity since their diagnosis with diabetes:

462 *“I am not able to work as much as others, I get stressed quickly. I feel fatigued. I have*
463 *to take tablets on time, and after taking them, I feel really fatigued. If I happen to*
464 *take the tablets half an hour later because of some chores around the house, I will*
465 *feel really fatigued. I feel like eating immediately. I am not able to walk like others.*
466 *People who are my age and don’t have any sugar [diabetes], they can walk well. I do*
467 *not have as much strength as them.”* (P15, Female, PHQ-9 = 6, Mild Depression)

468

469 **Discussion**

470 With the rising prevalence of T2D in younger populations, it is critical to
471 understand their lived experiences and identify barriers to their effective diabetes
472 management and overall well-being. This study concurrently explores the occurrence
473 of depressive symptoms with the lived experiences, facilitators, and barriers to T2D-
474 related self-care activities in young adults under the age of 35 in Mysore district, South
475 India. Overall, participants exhibited high rates of depressive symptoms and suicidal

476 ideation. Key barriers and facilitators to diabetes self-management and overall quality
477 of life include knowledge and awareness about the condition, experiences with and in
478 the healthcare system, presence of depressive symptoms, and physical and external
479 manifestations and impact of the condition on other aspects of their lives.

480 All study participants described challenging circumstances in their daily lives
481 with the condition, regardless of their PHQ-9 score. Negative feelings towards the
482 condition, feeling like a burden, and worry about the future were sentiments expressed
483 by participants at all levels of depression symptoms. In this study, 14 participants
484 (70%) presented mild to moderately-severe depressive symptoms, with 5 participants
485 (25%) reporting recent suicidal thoughts. Compared to the national prevalence
486 estimate of 3.6% in India, young adults with T2D are at a greater risk of depressive
487 disorders [28]. Descriptively in this sample, depressive symptoms are associated with
488 sex, region, socioeconomic status, duration since T2D diagnosis, family history of T2D,
489 T2D-related complications, adherence to a diabetic diet, and physical activity. This
490 data is consistent with the established relationship between diabetes and depressive
491 symptoms recognized around the world as well as factors shown to be associated with
492 this relationship [7,8,10–14]. All people living with diabetes must manage the constant,
493 daily effects associated with the condition. Chronic illnesses are challenging for
494 everyone living with them, both physically and mentally, and it is important that both
495 are treated concurrently to improve overall health [12].

496 The interviews revealed several knowledge gaps and misconceptions related to
497 diabetes. Many participants lacked trustworthy sources of information about the
498 condition and were afraid to ask questions due to the stigma associated with it. The
499 belief that diabetes is communicable was particularly prevalent and concerning,
500 leading to experiences of social isolation and discrimination. Researchers in the United
501 States and Ghana have established that lower health literacy coupled with illness
502 perception is associated with lower self-management behaviors in adults living with
503 T2D [28–30]. Medication perception also played a significant role in participants’
504 experience with T2D. Most participants expressed reluctance to rely on medication for
505 the rest of their lives, influenced by both healthcare professionals and community
506 members. Similar results were found by Miglani et al. in India (2000), where young
507 adults living with diabetes reported fear that people would treat them differently and
508 experienced unsolicited advice and instructions about household remedies from family
509 members and peers [17]. Researchers in Saudi Arabia and Japan have found that
510 medication adherence is dependent on the beliefs toward the medications and
511 potential T2D complications [31,32]. This speaks to the urgency of implementing public
512 health campaigns to dispel myths circulating within local communities and provide
513 accurate information about T2D, including the importance of medication adherence for
514 long-term glycemic control and prevention of future complications. Adults with T2D
515 are more likely to form positive perceptions about T2D with increased health literacy,

516 encouraging self-care practices for glycemic control, eliminating cultural, social, and
517 personal convictions about the condition, and building patient compliance and self-
518 efficacy [33].

519 The healthcare system plays a pivotal role in T2D management. Participants
520 reported mixed interactions with healthcare providers; while they generally
521 recommended lifestyle modifications, including diet and exercise, individualized care
522 and evidence-based support was often lacking. In some cases, participants experienced
523 blame and scolding by their healthcare providers, which is shown to have detrimental
524 effects on patient motivation and self-efficacy [34,35]. Complex power dynamics
525 between patients and providers exist in both urban and rural Mysore district. A
526 qualitative study conducted in rural Australia found that power dynamics and trust
527 between healthcare providers and patients greatly influence patient experiences and
528 perceived care [36]. Furthermore, rural participants' preference for private clinics
529 underscores the need for improved accessibility and quality of public healthcare
530 services. Similar findings were established in Vietnam, where researchers found that
531 social factors including word-of-mouth, relationships with healthcare providers,
532 healthcare staff attitudes, and marketing influence the choice of private over public
533 services [37].

534 Self-care activities are essential components of diabetes management, but
535 participants expressed numerous barriers that they experience. Physical activity was

536 hindered by fatigue, concerns about social judgment, and a lack of time. Healthy diets
537 were limited by financial constraints and family priorities, as female participants
538 expressed difficulties in maintaining separate diets from their family members.
539 Medication adherence was also influenced by financial constraints, as some
540 participants had to choose between purchasing their medication or investing in
541 something else. Previous research amongst Asian young adults in China and Singapore
542 recognized that adherence to diabetes medication is poorer in younger adults than in
543 older adults [28]. Researchers found similar results in urban slums in South India,
544 where the overall prevalence of self-care activities was low, indicating a need for more
545 awareness and education about diabetes self-care management [38]. South Asian
546 populations with T2D have been shown to experience these barriers to self-care
547 activities, including communication discordance, lack of knowledge and
548 misperceptions about the condition and management, gender-specific roles, lack of
549 family support and cultural appropriateness [39]. Addressing these barriers requires a
550 multifaceted approach, including community support, increased awareness, and access
551 to safe physical activity opportunities, healthy foods, and affordable medications.

552 The presence or lack of social support played a critical role in the participants'
553 diabetes management. Supportive spouses and friends provided encouragement and
554 motivation to some participants, emphasizing the importance of involving peers in
555 diabetes care. Social and family support has been shown to improve motivation, self-

556 efficacy, and health outcomes in people living with T2D around the world [35,40,41].
557 However, competing priorities, particularly related to children’s education, often took
558 precedence over self-care. Previous research has shown that occupation, living
559 condition, comorbidity, current treatment, and history of hospitalization are also
560 predictors of perceived family burden of T2D [42]. It is important that young adults
561 with T2D feel empowered to prioritize their health, without feeling like a burden to
562 their families; increased social support amongst this rising population has the potential
563 to build community and foster solidarity to improve both physical and mental health
564 outcomes.

565 **Strengths and limitations**

566 A major strength of this study is the collaboration with local researchers in
567 Mysore district, who had existing trust and relationships within the community. All
568 interviews were conducted by one research assistant to maintain internal validity and
569 consistency across the interviews. Qualitative analysis was conducted by three
570 researchers; one participated in the interviews to account for contextual cues and the
571 other two were not present, minimizing bias and allowing for objectivity in the
572 outcomes. The local researchers were consulted closely to ensure adequate
573 representation of the local context in the presented qualitative and quantitative
574 findings.

575 Thematic saturation was reached with the sample of qualitative interviews, but
576 it was too small for robust quantitative analysis. There is inherent selection bias in who
577 chose to participate in the study, which may have positively skewed the results.
578 Additionally, there is gender asymmetry in the study participants; while we planned to
579 recruit an equal number of male and female T2D patients, more females volunteered
580 for the study. Some results may be biased by participants' perceptions of health
581 research, but we employed probing to minimize this bias as much as possible. Causal
582 inferences cannot be drawn, given the study's cross-sectional design. However, the
583 purpose of this study was not for generalizability, but rather an in-depth exploration of
584 lived experiences with T2D in Mysore district.

585 **Future directions**

586 Future research should consider larger and more diverse samples, from both
587 urban and rural Mysore district. Longitudinal studies could provide insights into how
588 young adults' experiences and perceptions of diabetes change over time. Additionally,
589 interventions tailored to address the identified barriers to self-care activities and
590 psychosocial well-being should be developed and evaluated for their effectiveness in
591 improving T2D management and overall quality of life in young adult populations.

592

593 **Conclusion**

594 The findings of this study highlight the multifaceted challenges faced by young
595 adults living with T2D in Mysore district, South India. Addressing these challenges
596 requires a holistic approach, including improved healthcare access, education and
597 awareness campaigns, mental health support, and interventions to overcome barriers
598 to self-care practices. As T2D becomes more prevalent across the world at younger
599 ages, it is imperative that we understand their lived experiences to best support them.
600 Young adulthood is an excellent opportunity to intervene and promote healthy
601 behaviors before life-changing complications occur. By understanding the unique
602 experiences and perceptions of young adults with T2D, healthcare systems and
603 policymakers can develop more effective strategies to support diabetes management
604 and improve the long-term well-being of those affected.

605

606 **Funding:** This project was funded by the Paul Farmer Global Health Fieldwork Grant (Duke
607 Global Health Institute), Janet B. Chiang Grant (Duke University Asian/Pacific Studies
608 Institute), Human Rights Summer Research Grant (Duke Human Rights Center at the Franklin
609 Humanities Institute), Undergraduate Summer Award (Duke University Center for International
610 & Global Studies), and Undergraduate Research Support Independent Study Grant (Duke
611 University Undergraduate Research Support Office). The funders had no role in study design,
612 data collection and analysis, decision to publish, or preparation of the manuscript.

613 **Competing Interests:** The authors declare no competing interests.

614 **References**

- 615 1. IDF Diabetes Atlas | Tenth Edition [Internet]. 2022 [cited 2022 Jan 17]. Available from:
616 <https://diabetesatlas.org/>
- 617 2. India diabetes report 2000 – 2045 [Internet]. [cited 2024 Jan 15]. Available from:
618 <https://www.diabetesatlas.org/data/>
- 619 3. Ghorpade A, Majgi S, Sarkar S, Kar S, Roy G, Ananthanarayanan P, et al. Diabetes in rural
620 Pondicherry, India: a population-based study of the incidence and risk factors. *WHO*
621 *South-East Asia J Public Health*. 2013;2(3):149.
- 622 4. Gurunathan U, Prasad HK, White S, Sangalalingam T, Krishnamoorthy N. Type 2 Diabetes
623 Mellitus in Adolescents From Southern India – A Single Center Experience. *Indian*
624 *Pediatr*. 2021;58(2):176–7.
- 625 5. Karthik RC, Jagadeesh S. Self-care practices among type II diabetics in rural area of
626 Kancheepuram district, Tamil Nadu. *J Fam Med Prim Care*. 2020;9(6):7.
- 627 6. Unnikrishnan R, Anjana RM, Mohan V. Diabetes mellitus and its complications in India.
628 *Nat Rev Endocrinol*. 2016 Jun;12(6):357–70.
- 629 7. Johnson LCM, Desloge A, Sathish T, Williams ED, Absetz P, Haregu T, et al. The
630 relationship between common mental disorders and incident diabetes among participants
631 in the Kerala Diabetes Prevention Program (K-DPP). *PLoS ONE*. 2021 Jul
632 23;16(7):e0255217.
- 633 8. Mukherjee N, Chaturvedi SK. Depressive symptoms and disorders in type 2 diabetes
634 mellitus. *Curr Opin Psychiatry*. 2019 Sep;32(5):416–21.
- 635 9. Weaver LJ, Mendenhall E. Applying Syndemics and Chronicity: Interpretations from
636 Studies of Poverty, Depression, and Diabetes. *Med Anthropol*. 2014 Mar 1;33(2):92–108.
- 637 10. Mendenhall E, Norris SA, Shidhaye R, Prabhakaran D. Depression and Type 2 Diabetes in
638 Low and Middle Income Countries: A Systematic Review. *Diabetes Res Clin Pract*. 2014
639 Feb;103(2):276–85.
- 640 11. Chaturvedi SK, Manche Gowda S, Ahmed HU, Alosaimi FD, Andreone N, Bobrov A, et al.
641 More anxious than depressed: prevalence and correlates in a 15-nation study of anxiety
642 disorders in people with type 2 diabetes mellitus. *Gen Psychiatry*. 2019 Aug
643 9;32(4):e100076.
- 644 12. Lloyd CE, Nouwen A, Sartorius N, Ahmed HU, Alvarez A, Bahendeka S, et al. Prevalence
645 and correlates of depressive disorders in people with Type 2 diabetes: results from the
646 International Prevalence and Treatment of Diabetes and Depression (INTERPRET-DD)
647 study, a collaborative study carried out in 14 countries. *Diabet Med*. 2018;35(6):760–9.
- 648 13. Mathew M, Abish A, Kuriakose A, Isaiah JR, A m K, K V. Predictors of depression among
649 patients with diabetes mellitus in Southern India. *Asian J Psychiatry*. 2013 Aug
650 1;6(4):313–7.
- 651 14. Mendenhall E, Shivashankar R, Tandon N, Ali MK, Narayan KMV, Prabhakaran D. Stress
652 and Diabetes in Socioeconomic Context: A Qualitative Study of Urban Indians. *Soc Sci*
653 *Med* 1982. 2012 Dec;75(12):2522–9.
- 654 15. Thomas L, Asha H, Amalraj R, Prakash R, Abraham P, Thomas N. The utility of Problem
655 Areas in Diabetes (PAID) scale amongst patients with Type 2 diabetes (T2DM): An
656 experience from a teaching hospital in southern India. *J Fam Med Prim Care*.

- 657 2021;10(4):1687.
- 658 16. Shobhana R, Rama Rao P, Lavanya A, Padma C, Vijay V, Ramachandran A. Quality of life
659 and diabetes integration among subjects with type 2 diabetes. *J Assoc Physicians India*.
660 2003 Apr;51:363–5.
- 661 17. Miglani S, Sood A, Shah P. Self reported attitude and behavior of young diabetics about
662 discussing their disease. *Diabetes Res Clin Pract*. 2000 Apr 1;48(1):9–13.
- 663 18. Kalra B, Kalra S, Kumar A. Social stigma and discrimination: A care crisis for young
664 women with diabetes in India. *Diabetes Voice*. 2009;54(special issue):37–9.
- 665 19. World Health Organization. A conceptual framework for action on the social determinants
666 of health [Internet]. World Health Organization; 2010 [cited 2024 Jan 15]. Available from:
667 <https://iris.who.int/handle/10665/44489>
- 668 20. Habtewold TD, Islam MdA, Radie YT, Tegegne BS. Comorbidity of depression and
669 diabetes: an application of biopsychosocial model. *Int J Ment Health Syst*. 2016 Dec
670 3;10(1):74.
- 671 21. Designing and Conducting Mixed Methods Research [Internet]. SAGE Publications Inc.
672 2011 [cited 2024 Jan 15]. Available from: [https://us.sagepub.com/en-us/nam/designing-](https://us.sagepub.com/en-us/nam/designing-and-conducting-mixed-methods-research/book241842)
673 [and-conducting-mixed-methods-research/book241842](https://us.sagepub.com/en-us/nam/designing-and-conducting-mixed-methods-research/book241842)
- 674 22. Census India [Internet]. [cited 2024 Jan 15]. Available from:
675 <https://censusindia.gov.in/census.website/>
- 676 23. National Family Health Survey [Internet]. [cited 2024 Jan 15]. Available from:
677 <https://rchiips.org/nfhs/about.shtml>
- 678 24. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity
679 measure. *J Gen Intern Med*. 2001 Sep;16(9):606–13.
- 680 25. De Man J, Absetz P, Sathish T, Desloge A, Haregu T, Oldenburg B, et al. Are the PHQ-9 and
681 GAD-7 Suitable for Use in India? A Psychometric Analysis. *Front Psychol*. 2021 May
682 13;12:676398.
- 683 26. Pentapati SSK, Debnath DJ. Updated BG Prasad’s classification for the year 2022. *J Fam
684 Med Prim Care*. 2023 Jan;12(1):189–90.
- 685 27. McHugh ML. Interrater reliability: the kappa statistic. *Biochem Medica*. 2012 Oct
686 15;22(3):276–82.
- 687 28. Alharbi S, Alhofaian A, Alaamri MM. Illness Perception and Medication Adherence among
688 Adult Patients with Type 2 Diabetes Mellitus: A Scoping Review. *Clin Pract*. 2023 Jan
689 3;13(1):71–83.
- 690 29. Kugbey N, Oppong Asante K, Adulai K. Illness perception, diabetes knowledge and self-
691 care practices among type-2 diabetes patients: a cross-sectional study. *BMC Res Notes*.
692 2017 Aug 10;10(1):381.
- 693 30. Ajuwon AM, Insel K. Health literacy, illness perception, depression, and self-management
694 among African Americans with type 2 diabetes. *J Am Assoc Nurse Pract*. 2022 Sep
695 1;34(9):1066–74.
- 696 31. Alatawi YM, Kavookjian J, Ekong G, Alrayees MM. The association between health beliefs
697 and medication adherence among patients with type 2 diabetes. *Res Soc Adm Pharm
698 RSAP*. 2016;12(6):914–25.
- 699 32. Hashimoto K, Urata K, Yoshida A, Horiuchi R, Yamaaki N, Yagi K, et al. The relationship
700 between patients’ perception of type 2 diabetes and medication adherence: a cross-

- 701 sectional study in Japan. *J Pharm Health Care Sci.* 2019;5:2.
- 702 33. Lee YJ, Shin SJ, Wang RH, Lin KD, Lee YL, Wang YH. Pathways of empowerment
703 perceptions, health literacy, self-efficacy, and self-care behaviors to glycemic control in
704 patients with type 2 diabetes mellitus. *Patient Educ Couns.* 2016 Feb;99(2):287–94.
- 705 34. Freeman-Hildreth Y, Aron D, Cola PA, Wang Y. Coping with diabetes: Provider attributes
706 that influence type 2 diabetes adherence. *PloS One.* 2019;14(4):e0214713.
- 707 35. Nam S, Chesla C, Stotts NA, Kroon L, Janson SL. Barriers to diabetes management: patient
708 and provider factors. *Diabetes Res Clin Pract.* 2011 Jul;93(1):1–9.
- 709 36. McDonald J, Jayasuriya R, Harris MF. The influence of power dynamics and trust on
710 multidisciplinary collaboration: a qualitative case study of type 2 diabetes mellitus. *BMC*
711 *Health Serv Res.* 2012 Mar 13;12:63.
- 712 37. Al-Balushi S, Khan FR. Factors Influencing the Preference of Private Hospitals to Public
713 Hospitals in Oman [Internet]. Rochester, NY; 2017 [cited 2024 Jan 15]. Available from:
714 <https://papers.ssrn.com/abstract=3065470>
- 715 38. Chandrika K, Das B, Syed S, Challa S. Diabetes self-care activities: A community-based
716 survey in an urban slum in Hyderabad, India. *Indian J Community Med.* 2020;45(3):307.
- 717 39. Sohal T, Sohal P, King-Shier KM, Khan NA. Barriers and Facilitators for Type-2 Diabetes
718 Management in South Asians: A Systematic Review. *PloS One.* 2015;10(9):e0136202.
- 719 40. Al-Dwaikat TN, Rababah JA, Al-Hammouri MM, Chlebowy DO. Social Support, Self-
720 Efficacy, and Psychological Wellbeing of Adults with Type 2 Diabetes. *West J Nurs Res.*
721 2021 Apr;43(4):288–97.
- 722 41. Pesantes MA, Del Valle A, Diez-Canseco F, Bernabé-Ortiz A, Portocarrero J, Trujillo A, et
723 al. Family Support and Diabetes: Patient’s Experiences From a Public Hospital in Peru.
724 *Qual Health Res.* 2018 Oct;28(12):1871–82.
- 725 42. Molla IB, Berhie MA, Debele KA, Germossa GN, Hailu FB. Persons with Diabetes’
726 Perceptions of Family Burden and Associated Factors. *J Diabetes Res* [Internet]. 2023
727 [cited 2024 Jan 15];2023. Available from:
728 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9836787/>




Fig 1

Association of Covariates with Depression Severity

Depression Severity ■ Moderate/Moderately Severe ■ Mild ■ Minimal




Fig 2