Prevalence of hepatitis A virus among migrant workers in Qatar: A national study Nadin Younes^{1,2}, Hiam Chemaitelly^{3,4,5}, Parveen Banu Nizamuddin¹, Tasneem Al-Hamad^{1,6}, Marah Abdallah^{1,7}, Hadi M Yassine^{1,2}, Laith J Abu-Raddad^{3,4,5,8,9}, Gheyath K. Nasrallah^{1,2,*} ¹Department of Biomedical Science, College of Health Sciences, QU Health, Qatar University, Doha, Qatar ²Biomedical Research Center, Oatar University, Doha, Oatar

- 9 ³Infectious Disease Epidemiology Group, Weill Cornell Medicine-Qatar, Cornell University,
- 10 Doha, Qatar
- ⁴World Health Organization Collaborating Centre for Disease Epidemiology Analytics on
- 12 HIV/AIDS, Sexually Transmitted Infections, and Viral Hepatitis, Weill Cornell Medicine–Qatar,
- 13 Cornell University, Qatar Foundation Education City, Doha, Qatar
- ⁵Department of Population Health Sciences, Weill Cornell Medicine, Cornell University, New
- 15 York, New York, USA
- 16 ⁶Qatar Biobank, Doha, Qatar
- 17 ⁷*Clinical Microbiology & Virology division, Hamad General Hospital, Doha, Qatar*
- 18 ⁸Department of Public Health, College of Health Sciences, QU Health, Qatar University, Doha,
- 19 Qatar
- ⁹College of Health and Life Sciences, Hamad bin Khalifa University, Doha, Qatar
- 21

1

2

3

4 5

6

7

- 22 Word count: Abstract: 276 words, Main Text: 2,274 words.
- **Number of figures:** 2.
- 24 Number of tables: 2.
- 25 Running head: HAV prevalence among workers in Qatar.
- 26 Disclose funding received for this work: Other
- 27
- 28 Correspondence
- 29 Dr. Gheyath K. Nasrallah, Department of Biomedical Science, College of Health Sciences, Qatar
- 30 University, Doha 2713, Qatar. Tel: +(974) 4403-4817, Email: <u>gheyath.nasrallah@qu.edu.qa</u>
- 31
- 32

33 ABSTRACT

Background: Hepatitis A virus (HAV) is the predominant cause of acute viral hepatitis worldwide; however, data on HAV antibody prevalence (seroprevalence) among migrant populations are limited. This study aimed to investigate HAV seroprevalence among Qatar's migrant craft and manual workers (CMWs), constituting approximately 60% of the country's population.

39 Methods: HAV antibody testing was conducted on stored serum specimens obtained from CMWs 40 during a nationwide severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) population-41 based cross-sectional survey between July 26 and September 9, 2020. Associations with HAV 42 infection were investigated through regression analyses.

Results: Of the 2,607 specimens with HAV antibody test results, 2,393 were positive, and 214 43 44 were negative. The estimated HAV seroprevalence among CMWs was 92.0% (95% CI: 90.9-93.1%). HAV seroprevalence was generally high but exhibited some variation, ranging from 45 70.9% (95% CI: 62.4-78.2%) among Sri Lankans to 99.8% (95% CI: 98.2-99.9%) among 46 Pakistanis. The multivariable regression analysis identified age, nationality, and educational 47 attainment as statistically significant factors associated with HAV infection. Relative to CMWs 48 aged <29 years, the adjusted relative risk (ARR) was 1.06 (95% CI: 1.03-1.10) in CMWs aged 30-49 39 years and reached 1.15 (95% CI: 1.10-1.19) in those aged >50 years. In comparison to Indians, 50 the ARR was lower among Sri Lankans, assessed at 0.81 (95% CI: 0.72-0.91), but higher among 51 Nepalese at 1.07 (95% CI: 1.04-1.11), Bangladeshis at 1.10 (95% CI: 1.07-1.13), Pakistanis at 1.12 52 (95% CI: 1.09-1.15), and Egyptians at 1.15 (95% CI: 1.08-1.23). No evidence for differences was 53 found by geographic location or occupation. 54

- 55 **Conclusions:** HAV seroprevalence among Qatar's CMW population is very high, with over nine
- 56 out of every ten individuals having been exposed to this infection, likely during childhood.
- 57 Keywords: Hepatitis; HAV; prevalence; infection; workers; cross-sectional; Qatar

59 INTRODUCTION

Hepatitis A virus (HAV) stands as the predominant cause of acute viral hepatitis globally (1).
While HAV often results in mild illness, there are rare instances where the infection progresses to
acute liver failure and, ultimately, death (2). Although vaccine preventable (3), the estimated
annual incidence of HAV infections exceeds 150 million, with an associated death toll surpassing
39,000 in 2019 (4).

HAV is mainly transmitted via the fecal-oral route through the ingestion of contaminated food and 65 water (5). Other modes of transmission include exposure to infected body fluids or close physical 66 contact such as oral or anal intercourse with an infectious person (6). The largest burden of 67 infection is concentrated in low- and middle-income countries where inadequate sanitary 68 69 conditions prevail, with most infections occurring asymptomatically during childhood (4). The endemic nature of HAV in these countries leads to frequent exposures in early life and a large 70 proportion of immune adults (7). In higher income countries, susceptible adults face the risk of 71 72 symptomatic infection, with outbreaks documented among international travelers and specific 73 populations at risk such as men who have sex with men, people who inject drugs, and homeless 74 populations (3, 6). Recent trends also indicate a rise in foodborne outbreaks due to the globalization 75 of trade (7, 8).

The World Health Organization (WHO) formulated the Global Health Sector Strategy 2022-2030, envisioning the elimination of viral hepatitis as a public health concern by 2030 (9). Currently, the WHO recommends universal vaccination against HAV but only in countries where individuals face an intermediate risk of infection, defined as an age-specific prevalence of infection of <90% by 10 years of age but \geq 50% by 15 years of age (3, 6, 10). The current approach includes targeted vaccination for high-risk populations in countries with lower infection rates, while routine

vaccination is discouraged in endemic countries to avoid paradoxical rise in disease incidence
among unvaccinated individuals (10) (3, 6, 10). As of May 2021, immunization against HAV for
children had been introduced or was planned to be introduced in only 34 countries (4).
Immunization against HAV has been successfully integrated into the national immunization
programs of Bahrain, Oman, Qatar, Saudi Arabia, and Tunisia (11). In Iraq, targeted vaccination
schemes for high-risk populations are also in place (11). Nevertheless, in most other countries in
the region, HAV vaccines are only available at a cost in the private sector (11).

In the Middle East and North Africa (MENA) region, a systematic review of evidence indicated a
decline in reported HAV incidence among children over the past two decades (11). However, HAV
remains endemic at high incidence in few countries, including Egypt, Morocco, and Pakistan, as
well as in geographic areas marked by conflict (11, 12).

Given the limited global data on viral hepatitis infections among migrant populations, this study 93 aimed to determine HAV antibody prevalence (seroprevalence), indicating a history of HAV 94 95 infection, among Qatar's craft and manual worker (CMW) population. Qatar, known for its diverse 96 demographics with 89% of the population comprising expatriates from over 150 countries (13), 97 has a large CMW population, constituting approximately 60% of the total population of the 98 country. This population group, primarily consisting of unmarried men aged 20-49 years, is recruited for employment in infrastructure and development projects, including those associated 99 with the World Cup 2022 (14-16). The majority of CMWs originate from countries where HAV 100 101 infection is endemic at high incidence, and the typical age for acquiring the infection is below 10 years (12, 17). The overarching objective of this study was to provide insights into HAV 102 epidemiology and contribute data to national initiatives aiming to meet global targets for viral 103 hepatitis elimination. 104

105 METHODS

106 Study design, sampling, and specimen collection and handling.

This study tested deidentified stored blood serum specimens collected during a nationwide serological survey conducted from July 26, 2020, to September 09, 2020 to determine the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) seroprevalence among CMWs (14, 18, 19). The survey's sampling strategy was formulated following an analysis of the registered users' database of the Qatar Red Crescent Society (QRCS), the primary healthcare provider for CMWs in the country (14).

QRCS manages four centers strategically spread across Qatar to address the healthcare needs of CMWs nationwide (14). These centers cover expansive catchment areas, operate extended hours, and provide services either free of charge or with substantial subsidies (14). To ensure sample representativeness, the probability distribution of CMWs by age and nationality from the QRCS database was cross-referenced with that of expatriate residents from the Ministry of Interior database (20).

With men constituting >99% of CMWs (21), the sampling strategy did not explicitly factor sex. 119 CMWs were recruited at QRCS centers using a systematic sampling approach, informed by the 120 average daily attendance at each center (14). At each center, recruitment proceeded by inviting 121 every 4th attendee to participate in the study until the required sample size was reached for all age 122 123 and nationality strata. To address challenges in recruiting participants in smaller age-nationality strata, especially among younger individuals of specific nationalities, the recruitment criteria were 124 adjusted towards the conclusion of the study. In these instances, all attendees in these strata were 125 invited to participate, rather than every 4th attendee. 126

Trained interviewers administered the written informed consent and the study instrument to participants, accommodating their language preferences among nine options: Arabic, Bengali, English, Hindi, Nepali, Sinhala, Tagalog, Tamil, and Urdu (14). The instrument, designed following WHO guidance for developing SARS-CoV-2 sero-epidemiological surveys (22), collected essential socio-demographic information. Certified nurse's drew a 10 mL blood specimen for serological testing, which was then stored in an icebox before transportation to the Qatar Biobank for long-term storage and subsequent testing.

134 Laboratory methods

Serum aliquots were extracted from the archived specimens at the Qatar Biobank (QBB) and subsequently transferred to the virology laboratory at Qatar University (QU) for serological testing. The sera at both the QBB and QU were maintained at -80°C until employed for serology testing.

139 HAV seroprevalence was determined by testing sera for HAV antibodies using Dia.Pro competitive enzyme linked Immunosorbent Assay (ELISA) From Diagnostic Bioprobes Srl, 140 141 Milano, Italy (23). This assay has a reported sensitivity of 100% and a specificity exceeding 98% (23). Interpretation of test results followed the manufacturer's guidelines. Specimens with a mean 142 cut-off/optical density (OD) value at 450nm (designated as cut-off index values) below 0.9 were 143 144 considered negative, those surpassing 1.1 were considered positive, and those within the range of 0.9 to 1.1 were considered equivocal (23). Equivocal samples were excluded from further analysis. 145 The mean cut-off value was calculated using the equation: mean cut-off value = (OD value for the 146 negative control well + OD value for the positive control well)/3 (23). 147

148 Oversight

This study received approval from the Institutional Review Boards of Hamad Medical Corporation, Qatar University, and Weill Cornell Medicine-Qatar. The reporting of the study adhered to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines, as detailed in Table S1 in the Online Supplementary Document.

153 Statistical analysis

Frequency distributions and measures of central tendency were used to describe study participants. HAV seroprevalence was estimated after applying probability weights to correct for participants' unequal selection and ensure the sample's representativeness of the broader CMW population. Weights were calculated using the population distribution of CMWs by age, nationality, and QRCS center, extracted from the QRCS registered-user database (14). HAV cut-off index values obtained from testing with the Dia.Pro competitive ELISA were illustrated using a histogram.

Associations with HAV infection were explored through Chi-square tests and univariable and 160 161 multivariable Poisson regression analyses with robust error variance, not logistic regression analyses, given the high observed HAV seroprevalence levels (24, 25). Variables with a p-value 162 163 ≤ 0.2 in the univariable analyses were included in the multivariable model. A p-value < 0.05 in the multivariable analysis was considered indicative of a statistically significant association. 164 Unadjusted and adjusted relative risks (RRs and ARRs, respectively), along with their respective 165 166 95% confidence intervals (CIs) and p-values, were reported. Interactions were not considered. All statistical analyses were conducted using Stata/SE version 18.0 (Stata Corporation, College 167 Station, TX, USA). 168

169 **RESULTS**

170 Study population.

171	Table 1 outlines the characteristics of the study participants. Among the 2,641 blood specimens
172	collected during the SARS-CoV-2 seroprevalence survey (14), only 2,616 specimens were
173	available and had sufficient sera for HAV testing. Of these, 2,393 specimens were positive, 214
174	were negative, and 9 were equivocal and therefore excluded from further analysis.

Table 1. Characteristics and seroprevalence of HAV infection among study participants. 175

Characteristics	Tested		HAV seroprevale	HAV seroprevalence	
Characteristics	N (%*)	Ν	%† (95% CI)†	Chi-square p-value	
Age (years)					
≤29	747 (27.7)	637	86.4 (83.7-88.7)	< 0.001	
30-39	966 (41.8)	893	92.5 (90.6-94.0)		
40-49	545 (21.4)	521	96.2 (94.2-97.5)		
50-59	258 (7.4)	251	96.8 (93.3-98.5)		
60+	91 (1.7)	91	100.0 (96.0-100.0)*		
Nationality					
All other nationalities [§]	231 (7.6)	210	90.6 (85.9-93.8)	< 0.001	
Bangladeshi	623 (26.0)	610	97.9 (96.4-98.8)		
Egyptian	90 (3.2)	86	95.7 (88.2-98.5)		
Filipino	102 (2.7)	78	78.3 (68.1-86.0)		
Indian	716 (29.2)	628	87.6 (84.9-89.9)		
Nepalese	564 (21.7)	539	95.6 (93.5-97.1)		
Pakistani	136 (4.9)	135	99.8 (98.2-99.9)		
Sri Lankan	145 (4.8)	107	70.9 (62.4-78.2)		
QRCS center (catchment area within Qatar)					
Fereej Abdel Aziz (Doha-East)	611 (23.5)	548	89.8 (87.1-92.0)	0.132	
Zekreet (North-West)	234 (2.3)	215	92.0 (87.7-94.9)		
Hemaila (South-West; "Industrial Area")	964 (41.9)	897	93.5 (91.8-94.9)		
Mesaimeer (Doha-South)	798 (32.4)	733	91.8 (89.6-93.5)		
Educational attainment					
Primary or lower	626 (24.6)	592	94.4 (92.2-96.1)	< 0.001	
Intermediate	429 (17.9)	417	97.4 (95.3-98.5)		
Secondary/High school/Vocational	1,085 (44.1)	1,003	92.7 (91.0-94.2)		
University	369 (13.4)	285	76.5 (71.6-80.8)		
Occupation					
Professional workers [¶]	136 (4.9)	107	76.9 (68.4-83.6)	< 0.001	
Food & beverage workers	91 (3.2)	79	89.0 (80.7-94.0)		
Administration workers	82 (3.0)	73	87.5 (77.2-93.5)		
Retail workers	169 (6.6)	149	90.1 (84.6-93.8)		
Transport workers	430 (16.4)	402	93.5 (90.6-95.6)		
Security workers	60 (2.3)	55	91.1 (80.2-96.3)		
Cleaning workers	105 (4.0)	96	91.7 (84.7-95.6)		
Technical and construction workers [£]	1,311 (52.8)	1,237	94.6 (93.1-95.7)		
Other workers**	174 (6.8)	151	86.5 (80.4-90.9)		
Total (%, 95% CI)	2,607 (100.0)	2,393	92.0 (90.9-93.1)		

CI, confidence interval; QRCS, Qatar Red Crescent Society.

176 177 178 179 180 181 182 183 184 185 *Percentage of the sample weighted by age, nationality, and QRCS center. Missing values for socio-demographic variables were excluded from the analysis.

[†]Percentage of positive out of the total sample weighted by age, nationality, and QRCS center.

*The 95% confidence intervals were calculated using the binomial distribution.

§Includes all other nationalities of craft and manual workers residing in Qatar.

Includes architects, designers, engineers, operation managers, and supervisors among other professions.

^fIncludes carpenters, construction workers, crane operators, electricians, foremen, maintenance/air conditioning/cable technicians, masons,

mechanics, painters, pipe-fitters, plumbers, and welders among other professions.

**Includes barbers, firefighters, gardeners, farmers, fishermen, and physical fitness trainers among other professions.

187	Around 70% of the study participants were younger than 40 years, with a median age of 35.0 years
188	and an interquartile range (IQR) spanning 29.0-43.0 years. Slightly more than 40% of participants
189	had educational attainment at intermediate or lower levels, 44.1% had high school education or
190	vocational training, and 13.4% had higher educational attainment. The distribution of nationalities
191	highlighted a prevalent presence of Indians (29.2%), Bangladeshis (26.0%), and Nepalese (21.7%),
192	aligning with the broader demographic composition of the CMW population in Qatar (20). Over
193	half of CMWs (52.8%) were employed in technical and construction jobs, which included
194	occupations such as carpenters, crane operators, electricians, masons, mechanics, painters,
195	plumbers, and welders.
196	HAV seroprevalence and associations with infection
197	HAV seroprevalence among CMWs was estimated at 92.0% (95% CI: 90.9-93.1%). HAV cut-off
198	index values in positive specimens ranged from 1.1 to 485.0 (Figure 1), with a median of 22.8

```
199 (IQR: 15.9-27.3).
```

Fig 1. Distribution of the mean cut-off/optical density (OD) values at 450nm among HAV
 antibody-positive individuals as measured using the Dia.Pro Competitive ELISA.

203	Table 1 shows HAV seroprevalence estimates across various socio-demographic characteristics
204	within the population. Figure 2 illustrates HAV seroprevalence by nationality group. Despite
205	variations, HAV seroprevalence was generally high ranging from 70.9% (95% CI: 62.4-78.2%)
206	among Sri Lankans to 99.8% (95% CI: 98.2-99.9%) among Pakistanis.
207 208	Fig 2. HAV seroprevalence by nationality group among the craft and manual worker population in Qatar.

210	The multivariable regression analysis identified age, nationality, and educational attainment as
211	statistically significant factors associated with HAV infection (Table 2). The ARR increased
212	gradually with age. Relative to CMWs \leq 29 years of age, the ARR was 1.06 (95% CI: 1.03-1.10)
213	in CMWs aged 30-39 years and reached 1.15 (95% CI: 1.10-1.19) in those aged \geq 50 years. There
214	were significant differences by nationality, where compared to Indians, the ARR was lower among
215	Sri Lankans assessed at 0.81 (95% CI: 0.72-0.91), but higher among Nepalese at 1.07 (95% CI:
216	1.04-1.11), Bangladeshis at 1.10 (95% CI: 1.07-1.13), Pakistanis at 1.12 (95% CI: 1.09-1.15), and

Egyptians at 1.15 (95% CI: 1.08-1.23). 217

Table 2. Associations with HAV infection among the craft and manual worker population in 218 Oatar 219

Characteristics	Univariable regression analysis			Multivariable regression analysis	
	RR* (95% CI*)	p-value	F test p-value [†]	ARR* (95% CI*)	p-value [‡]
Age (years)					
≤29	1.00		< 0.001	1.00	
30-39	1.07 (1.03-1.11)	< 0.001		1.06 (1.03-1.10)	< 0.001
40-49	1.11 (1.08-1.15)	< 0.001		1.12 (1.08-1.15)	< 0.001
50+	1.13 (1.09-1.17)	< 0.001		1.15 (1.10-1.19)	< 0.001
Nationality					
Indian [§]	1.00		< 0.001	1.00	
Bangladeshi	1.12 (1.08-1.15)	< 0.001		1.10 (1.07-1.13)	< 0.001
Egyptian	1.09 (1.03-1.15)	0.002		1.15 (1.08-1.23)	< 0.001
Nepalese	1.09 (1.06-1.13)	< 0.001		1.07 (1.04-1.11)	< 0.001
Pakistani	1.14 (1.11-1.17)	< 0.001		1.12 (1.09-1.15)	< 0.001
Sri Lankan	0.81 (0.72-0.91)	< 0.001		0.81 (0.72-0.91)	< 0.001
All other nationalities [¶]	1.00 (0.95-1.05)	0.920		1.05 (0.99-1.11)	0.122
QRCS center (catchment area within Qatar)	, í			l ì í	
Fereej Abdel Aziz (Doha-East)	1.00		0.084	1.00	
Zekreet (North-West)	1.03 (0.98-1.07)	0.302		1.01 (0.96-1.06)	0.691
Hemaila (South-West; "Industrial Area")	1.04 (1.01-1.08)	0.012		1.01 (0.98-1.05)	0.416
Mesaimeer (Doha-South)	1.02 (0.99-1.06)	0.208		1.01 (0.97-1.04)	0.690
Educational attainment	, í			l ì í	
Primary or lower	1.00		< 0.001	1.00	
Intermediate	1.03 (1.01-1.06)	0.019		1.04 (1.02-1.07)	0.002
Secondary/High school/Vocational	0.98 (0.96-1.01)	0.179		1.01 (0.98-1.04)	0.389
University	0.81 (0.76-0.86)	< 0.001		0.85 (0.79-0.91)	< 0.001
Occupation	, í			l ì í	
Professional workers**	1.00		< 0.001	1.00	
Food and beverage workers	1.16 (1.02-1.31)	0.019		1.05 (0.92-1.18)	0.487
Administration workers	1.14 (0.99-1.30)	0.059		1.08 (0.95-1.23)	0.231
Retail workers	1.17 (1.05-1.31)	0.005		1.07 (0.96-1.20)	0.225
Transport workers	1.22 (1.10-1.35)	< 0.001		1.07 (0.96-1.19)	0.200
Security workers	1.18 (1.04-1.35)	0.010		1.06 (0.93-1.22)	0.388
Cleaning workers	1.19 (1.06-1.34)	0.003		1.08 (0.96-1.21)	0.191
Technical and construction workers ^{††}	1.23 (1.11-1.36)	< 0.001		1.08 (0.98-1.20)	0.134
Other workers ^{‡‡}	1.12 (1.00-1.26)	0.047		1.03 (0.92-1.16)	0.634

ARR, adjusted relative risk; CI, confidence interval; RR, relative risk; QRCS, Qatar Red Crescent Society.

220 221 222 223 224 225 226 *Estimates weighted by age, nationality, and QRCS center.

*Covariates with p-value <0.05 in the multivariable analysis were considered to provide statistically significant evidence for an association.

[§]Indian was chosen as the reference group due to its comparatively low prevalence of HAV seroprevalence and a sufficiently large sample size,

making it a suitable reference group for the regression analysis.

Includes all other nationalities of craft and manual workers residing in Qatar.

[†]Covariates with p-value ≤ 0.2 in the univariable analysis were included in the multivariable analysis.

227 228 **Professional workers include architects, designers, engineers, operation managers, and supervisors among other professions.

- ^{††}Technical and construction workers include carpenters, construction workers, crane operators, electricians, foremen, maintenance/air
- 229 conditioning/cable technicians, masons, mechanics, painters, pipe-fitters, plumbers, and welders among other professions.
- 230 ^{‡‡}Other workers include barbers, firefighters, gardeners, farmers, fishermen, and physical fitness trainers among other professions.
- 231

Significant differences were observed based on educational attainment. CMWs with intermediate 232 education had an ARR of 1.04 (95% CI: 1.02-1.07) while those with university education had an 233 ARR of 0.85 (95% CI: 0.79-0.91), both compared to those with primary or lower education. No 234 evidence for differences was found by QRCS center (proxy of catchment area/geographic location) 235 or by occupation. 236

DISCUSSION 237

HAV seroprevalence is highly elevated among the CMW population in Oatar, with over nine out 238 of every ten individuals having experienced this infection. This observation suggests that nearly 239 240 the entire CMW population has acquired the infection, predominantly during childhood before relocating to Qatar, considering the very high HAV seroprevalence observed in the migrants' 241 countries of origin (12, 17). This underscores the significance of a comprehensive plan for HAV 242 prevention. Such a plan should incorporate measures to promote safe water and sanitation 243 practices, along with adherence to WHO recommendations for HAV vaccination in countries 244 where individuals face a heightened risk of symptomatic infection (3, 6, 10). 245

While HAV seroprevalence was generally very high, some variations were observed. These 246 variations included differences by nationality, with Sri Lankans exhibiting the lowest 247 seroprevalence and Nepalese, Bangladeshis, Pakistanis, and Egyptians having the highest 248 249 seroprevalence. Seroprevalence was at its lowest among individuals with a university education, suggesting an association between infection and lower socio-economic status, as observed 250 251 elsewhere (17). Seroprevalence increased with age, consistent with expectations for a measure of

cumulative exposure to the infection. However, the rate of increase with age was slow, indicatingthat only a minority of infections are acquired in adulthood.

Our study represents the first endeavor to estimate the seroprevalence of HAV in Qatar, filling a critical gap in the existing research. Its significance is underscored by the scarcity of studies on HAV seroprevalence in the Middle East, particularly among QRCS and the Gulf region, where CMWs constitute a significant demographic. Prominent strengths of our study include a substantial sample size and the utilization of a highly sensitive ELISA for detecting HAV antibodies, enhancing the reliability and accuracy of the seroprevalence results.

This study is subject to certain limitations. Originally designed to employ a probability-based 260 sampling strategy for CMW recruitment, logistical challenges led to the implementation of a 261 262 systematic sampling method, concentrating on QRCS attendees. In response to this modification, probability-based weights were introduced to produce a seroprevalence estimate that aligns with 263 264 the broader CMW population. To account for the representation of smaller age-nationality strata, 265 a deviation from the initial plan-selecting every fourth attendee-was introduced. Towards the 266 end of the study, all individuals in these strata were approached to participate, ensuring a more 267 comprehensive representation of the sampled population.

Operational challenges presented difficulties in monitoring and maintaining consistent records of the response rate among nurses in the QRCS centers. As a result, an exact estimate of the response rate could not be determined, but it was approximated to exceed 90% based on the interviewers' experience. While there is potential for the recruitment scheme to impact the generalizability of the study findings, this is considered less likely given the substantial daily influx of CMWs visiting these centers, surpassing 5,000 patients daily (14). Of note that these centers serve as the primary healthcare providers for CMWs in the country, offering a range of services beyond patient

treatment, including periodic health certifications, vaccinations, and pre-travel SARS-CoV-2
testing.

277 CONCLUSIONS

This study revealed a very high HAV seroprevalence among Qatar's CMW population, with more 278 than 90% having experienced the infection. This underscores that virtually the entire CMW 279 population contracted the infection, presumably during childhood before their migration to Qatar, 280 281 given the comparably elevated HAV seroprevalence in their countries of origin. Despite the overall elevated seroprevalence, notable variations were observed, particularly across nationalities and 282 socio-economic strata. These findings contribute considerations for healthcare service planning 283 and policy development. This aims to alleviate the burden of HAV-related illnesses and work 284 285 towards the 2030 targets for viral hepatitis elimination.

286 Acknowledgements

We thank all participants for their willingness to be part of this study. We thank all members of the Craft and Manual Workers Seroprevalence Study Group for their efforts in supporting the collection of the samples of this study. We also thank Dr. Nahla Afifi, Director of Qatar Biobank (QBB), Ms. Tasneem Al-Hamad, Ms. Eiman Al-Khayat, and the rest of the QBB team for their unwavering support in retrieving and analyzing samples. We also acknowledge the dedicated efforts of the Surveillance Team at the Ministry of Public Health for their support in sample collection.

294 Funding

295 The authors are also grateful for support from the Biomedical Research Program and the 296 Biostatistics, Epidemiology, and Biomathematics Research Core, both at Weill Cornell Medicine-Qatar, as well as for support provided by the Ministry of Public Health and Hamad Medical 297 Corporation. This work was supported by the National Priorities Research Program (NPRP) grant 298 numbers: 12S-0216-190094, 13S-0128-200185, GSRA8-L-1-0501-21022, and UREP30-041-3-299 300 014 from the Qatar National Research Fund (a member of Qatar Foundation). The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of 301 the article. Statements made herein are solely the responsibility of the authors. 302

303 Authorship contributions

GKN conceived the study and led the laboratory testing. NY, PBN, TAH, MA, and HMY conducted the lab testing of the specimens. GKN and NY validated and interpreted the laboratory results. HC developed the study design, managed the databases, performed the data analyses, and wrote the first draft of the article. LJA led the statistical analyses and contributed to the first draft

308 of the article. GKN and LJA critically reviewed the manuscript. All authors have read and 309 approved the final manuscript.

310 **Disclosure of interest**

- 311 I have read the journal's policy and the authors of this manuscript have the following competing
- 312 interests: Dr. Gheyath K. Nasrallah is currently an academic editor at PLOS ONE.

313 Data availability statement

All data are available in aggregate form within the manuscript.

315 **Ethics approval**

- Hamad Medical Corporation, Qatar University (QU-IRB 1558-EA/21), and Weill Cornell
- 317 Medicine-Qatar Institutional Review Boards approved this study.

318 ORCID iDs

- 319 Nadin Younes https://orcid.org/0000-0002-5241-944X
- Hiam Chemaitelly <u>https://orcid.org/0000-0002-8756-6968</u>
- 321 Laith Abu-Raddad <u>https://orcid.org/0000-0003-0790-0506</u>
- 322 Gheyath Nasrallah <u>https://orcid.org/0000-0001-9252-1038</u>

RFERENCES

1. Zeng DY, Li JM, Lin S, Dong X, You J, Xing QQ, et al. Global burden of acute viral hepatitis and its association with socioeconomic development status, 1990-2019. J Hepatol. 2021;75(3):547-56.

2. World Health Organization. Hepatitis A. Available from: <u>https://www.who.int/news-room/fact-sheets/detail/hepatitis-a</u>. Accessed on January 5, 2024. 2023.

3. World Health Organization. WHO Immunological Basis for Immunization Series, Module 18: Hepatitis A Update; WHO: Geneva, Switzerland. Available from:

https://www.who.int/publications/i/item/97892516327. Accessed on January 5, 2024. 2019.

4. Cao G, Jing W, Liu J, Liu M. The global trends and regional differences in incidence and mortality of hepatitis A from 1990 to 2019 and implications for its prevention. Hepatol Int. 2021;15(5):1068-82.

5. Wenzel JJ, Allerberger F. Hepatitis A as a foodborne infection. The Lancet Infectious Diseases. 2014;14(10):907-8.

6. Migueres M, Lhomme S, Izopet J. Hepatitis A: epidemiology, high-risk groups, prevention and research on antiviral treatment. Viruses. 2021;13(10):1900.

7. Jacobsen KH. Globalization and the Changing Epidemiology of Hepatitis A Virus. Cold Spring Harb Perspect Med. 2018;8(10).

8. Anita C, Tanisha B. Epidemiology of Hepatitis A: Past and Current Trends. In: Costin Teodor S, Cristin Constantin V, Ion R, Valeria T, Silvia L, editors. Hepatitis A and Other Associated Hepatobiliary Diseases. Rijeka: IntechOpen; 2019. p. Ch. 1.

9. World Health Organization. Global health sector strategies on, respectively, HIV, viral hepatitis and sexually transmitted infections for the period 2022-2030. Found at:

file:///C:/Users/gm15/Downloads/9789240053779-eng.pdf. Accessed June 7, 2023. 2022.

10. World Health Organization. Summary of WHO Position Paper on hepatitis A vaccines, October 2022. Available from:

efaidnbmnnnibpcajpcglclefindmkaj/<u>https://iris.who.int/bitstream/handle/10665/363396/WER9740-eng-fre.pdf?sequence=1</u>. Accessed on: January 10, 2024. 2022.

11. Badur S, Ozturk S, AbdelGhany M, Khalaf M, Lagoubi Y, Ozudogru O, et al. Hepatitis A in the Eastern Mediterranean Region: a comprehensive review. Hum Vaccin Immunother. 2022;18(5):2073146.

12. Nelson N., Weng M. Hepatitis A. CDC Yellow Book 2024. Available from: <u>https://wwwnc.cdc.gov/travel/yellowbook/2024/infections-diseases/hepatitis-a#map506</u>. Accessed on: January 8, 2024. 2024.

13. Abu-Raddad LJ, Chemaitelly H, Ayoub HH, Al Kanaani Z, Al Khal A, Al Kuwari E, et al. Characterizing the Qatar advanced-phase SARS-CoV-2 epidemic. Sci Rep. 2021;11(1):6233.

14. Al-Thani MH, Farag E, Bertollini R, Al Romaihi HE, Abdeen S, Abdelkarim A, et al. SARS-CoV-2 Infection Is at Herd Immunity in the Majority Segment of the Population of Qatar. Open Forum Infect Dis. 2021;8(8):ofab221.

15. AlNuaimi AA, Chemaitelly H, Semaan S, AlMukdad S, Al-Kanaani Z, Kaleeckal AH, et al. All-cause and COVID-19 mortality in Qatar during the COVID-19 pandemic. BMJ Glob Health. 2023;8(5).

16. Jeremijenko A, Chemaitelly H, Ayoub HH, Alishaq M, Abou-Samra AB, Al Ajmi J, et al. Herd Immunity against Severe Acute Respiratory Syndrome Coronavirus 2 Infection in 10 Communities, Qatar. Emerg Infect Dis. 2021;27(5):1343-52.

17. Mohd Hanafiah K, Jacobsen KH, Wiersma ST. Challenges to mapping the health risk of hepatitis A virus infection. Int J Health Geogr. 2011;10:57.

18. Nasrallah GK, Dargham SR, Al-Sadeq DW, Amanullah FH, Shurrab FM, Nizamuddin PB, et al. Seroprevalence of herpes simplex virus type 1 and type 2 among the migrant workers in Qatar. Virol J. 2023;20(1):188.

19. Younes N, Yassine HM, Nizamuddin PB, Kourentzi K, Tang P, Ayoub HH, et al. Seroprevalence of hepatitis E virus (HEV) among male craft and manual workers in Qatar (2020–2021). Heliyon. 2023;9(11):e21404.

20. Ministry of Interior-State of Qatar. Population distribution by sex, age, and nationality: results of Kashef database. 2020.

21. Planning and Statistics Authority-State of Qatar. Labor force sample survey. Available from: https://www.psa.gov.qa/en/statistics/Statistical%20Releases/Social/LaborForce/2017/statistical_analysi s_labor_force_2017_En.pdf. Accessed on May 23, 2022.; 2017.

22. World Health Organization. Population-based age-stratified seroepidemiological investigation protocol for COVID-19 virus infection. Available from: <u>https://apps.who.int/iris/handle/10665/331656</u>. Accessed on: April 15, 2020. 2020.

23. Diagnostic Bioprobes Srl. Dia.Pro HAV Ab Competitive Enzyme ImmunoAssay (ELISA). Available from: <u>http://npt.ir/uploads/AVAB.CE.pdf</u>. Accessed on: January 5, 2024. 2015.

24. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. BMC Med Res Methodol. 2003;3:21.

25. Zou G. A modified poisson regression approach to prospective studies with binary data. Am J Epidemiol. 2004;159(7):702-6.



Fig 1



Fig 2