- 1 Full Title: Evaluation of a Low-threshold Exercise And Protein
- 2 supplementation intervention for Women (LEAP-W) experiencing
- ³ homelessness and addiction: Protocol for a single site intervention
- 4 study.
- 5 **Short Title:** Protocol for a low threshold exercise intervention for
- 6 women experiencing homelessness and addiction.
- 7
- 8 Authors
- 9 Fiona Kennedy ^{1*}, Deirdre Murray, ², Clíona Ní Cheallaigh ^{3,4}, Roman Romero-Ortuno ^{5,6}, Julie
 10 Broderick ¹.
- ¹Discipline of Physiotherapy, School of Medicine, Trinity College Dublin, Dublin, D08W9RT,

12 Ireland

- 13 ²Academic Unit of Neurology, Trinity Biomedical Sciences Institute, Trinity College Dublin,
- 14 Dublin, D02 R590, Ireland.
- 15 ³Department of General Medicine and Infectious Diseases, St James's Hospital, Dublin,
- 16 D08W9RT, Ireland.
- 17 School of Medicine, Trinity College Dublin, Dublin, D08W9RT, Ireland.
- ⁵Discipline of Medical Gerontology, School of Medicine, Trinity College Dublin, Dublin, D08
- 19 NHY1, Ireland.
- ⁶Mercer's Institute for Successful Ageing, St James's Hospital, Dublin, D08 NHY1, Ireland.
- 21

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

- 22 *Corresponding author
- 23 E-mail: cromwelf@tcd.ie (FK)
- 24
- 25 FK and JB are Joint Senior Authors.
- ²⁶ [&]R.RO, CNC and DM contributed equally to this work.

27

- 28
- 29
- 30
- 31
- 32 33

34 Abstract

Background: Frailty is a complex multi-dimensional state of increased vulnerability to 35 adverse health outcomes and is usually associated with older age but there is growing 36 37 evidence of accelerated ageing and frailty in non-geriatric populations, including those 38 experiencing socio-economic deprivation and extreme social exclusion, such as people experiencing homelessness. Addiction, as a coping mechanism for prior trauma, is common 39 among people who are homeless and can have a gendered dimension. Women experiencing 40 homelessness and addiction have unique needs which require a gendered approach. The 41 aim of this study is to explore the effectiveness of an exercise intervention to target the 42 known physical functioning deficits and frailty which this population experiences. 43 **Methods:** This mixed-methods study will explore physical functioning deficits and frailty 44 in women experiencing homelessness and addiction, using a bespoke test battery and an 45 exercise intervention. Physical function (10m Walk Test, 2 Min Walk Test, Single Leg Stance, 46 47 Chair Stand Test, hand grip dynamometry), frailty (Clinical Frailty Scale and the SHARE-Frailty Instrument) and nutritional status (Mini-Nutritional Status), pain (Numerical Pain 48 49 Rating Scale) and quality of life (SF 12-V2) will be evaluated. The 10-week intervention will involve a 3-times weekly exercise programme with protein supplementation. Following this, 50 qualitative interviews, which will be thematically analysed using Braun & Clarke 51 52 methodology, will be conducted. This study will be conducted in Dublin from February to 53 July 2024.

54	Discussion: Little is known about frailty-focussed interventions in women experiencing
55	homelessness and addiction. This proposed study will help to increase the knowledge base
56	regarding the physical health burden and frailty experienced by this extremely vulnerable
57	population and will deliver a targeted intervention with a gendered dimension to mitigate
58	its affects. The findings of this research will help narrow this research gap and will guide
59	clinicians and policy makers to implement unique gender-based treatment strategies for this

60 population. (299 words)

61 Keywords

- 62 Physical function, frailty, homelessness, addiction, women.
- 63
- 64 Clinical Trials.gov number: *NCT05700305*

65 Introduction

66	The world's population is ageing rapidly (1). Frailty, a sometimes-preventable consequence
67	of a rapidly ageing population, is a complex state of cumulative decline across multiple
68	physiological systems which renders a person vulnerable to adverse health outcomes and
69	greatly challenges healthcare systems (2). Targeted interventions to manage frailty in older
70	adults are recommended and exercise with nutritional supplementation has proven
71	effectiveness in combatting and mitigating frailty (3, 4, 5).
72	Frailty as a construct is a geriatric concept which is strongly associated with increasing age
73	(6). Notwithstanding that biological age is closely linked to frailty, it is known that
74	chronological age alone will not identify frailty as the health and functional life status of
75	populations are so variable (7). Growing evidence indicates the prevalence of premature
76	ageing and frailty in non-geriatric populations (8, 9). 'Younger frailty' has been identified in
77	populations of lower socioeconomic status, who live in areas of greater deprivation and
78	experience an earlier onset of illness and disability (10, 11). In a study investigating people
79	experiencing homelessness (PEH) admitted for inpatient care, 83% of participants had
80	mobility problems and 70% were frail or pre-frail despite a low median age of 45 years (12,
81	13). In an analysis of UK Biobank participants aged 37–73 years, 42% of individuals living
82	with frailty were within the most socially deprived 20% of the cohort (10). Females are
83	known to experience higher levels of frailty compared to males (14, 15). It is therefore not

surprising that poor physical functioning and frailty is experienced by young women who are
homeless and use drugs (16, 17).

Health services for PEH frequently focus on crisis-based medical, nursing and addiction 86 services, often with a gender-blind lens (18, 19). High levels of stigma, shame and fear result 87 in women accessing services even less and seeking help at breaking point with complex 88 89 coinciding physical and mental health needs (20, 21). Homelessness and substance use in women is inter-sectoral and may overlap with trauma, abuse, domestic and sexual violence 90 and exploitation, and the needs of these women are complex and unique (22). Pathways 91 92 into homelessness are frequently different to that of men and require a different approach 93 (23). Internationally, gender-based services for women who are experiencing homelessness 94 and addiction are lacking and services are mostly delivered in mixed gender settings (18, 20). Women-specific services are recommended to increase safety, stability, wellbeing, and 95 psychosocial functioning to respond meaningfully to the serious health and social care 96 needs of women in this setting (24). Gender specific and gender sensitive research is 97 required to fill this knowledge gap (25). 98

99 Our prior research, the LEAP I study demonstrated high retention (83% female v 42% males) 100 with excellent engagement and feedback from its female participants, yet caution was 101 observed around engaging in a mixed exercised programme with men (26). Women have 102 traditionally been underserved in research and policy, yet multiple sources cite women as 103 the fastest growing homeless demographic (23). No study has examined how targeted

women-only exercise programmes with nutritional supplementation work in practice in thispopulation.

- 106 The overall aim of this study is to explore physical functioning and frailty outcomes of an
- 107 exercise intervention with protein supplementation in a group of (non-geriatric) women
- 108 experiencing homelessness and addiction challenges. The objectives are:
- 109 1. To evaluate recruitment, retention and adherence rates, and any adverse effects of the
- 110 intervention.
- 111 2. To evaluate pre- and post-intervention levels of physical function, nutritional and frailty
- 112 status, pain, and general health status.
- 113 3. To evaluate programme feedback and gain deeper insights into participants perspectives
- and experiences of the exercise intervention.

115

116 Materials and Methods

117 Study design, setting and time frame.

118 This mixed-methods study will involve a single-arm intervention which will take

119 place in Jane's Place, Merchants Quay Ireland (MQI), in Dublin city centre, a day-

- 120 service centre which provides services for women who are homeless and/or in
- addiction. Following the intervention qualitative research in the form of individual
- 122 exit interviews will be conducted with 20 participants or up to the point of data

123	saturation.	This study has	received ethical approval	from the Faculty of Health
-----	-------------	----------------	---------------------------	----------------------------

- 124 Sciences, Trinity College Dublin (Ethical Approval Reference Number: 211202).
- 125

126 Study population, eligibility criteria and recruitment strategy

127 Study population

- 128 This study will involve a non-geriatric population of women who attend Jane's
- 129 Place in Dublin city centre. These women have complex needs, many experiencing
- 130 homelessness and addiction challenges.

131 Eligibility criteria

- 132 Inclusion criteria:
- Adult women (18-65 years) accessing services in Jane's Place who consent to
- 134 participation.
- 135 Exclusion criteria:
- 136 Any person not meeting the age eligibility criteria.
- 137 > Any person identifying as a male.
- 138 Any person lacking capacity to give consent.
- 139 Any person with major physical/medical or cognitive challenges which would
- 140 preclude ability to safely complete the assessment or
- 141 Any person with insufficient English language ability to give fully informed
- 142 consent.

143 \succ Women with a confirmed pregnancy.

144 Recruitment Strategy

145	All potential	participants	will be provided	with a partici	pant information	leaflet
-----	---------------	--------------	------------------	----------------	------------------	---------

- 146 (PIL) detailing the purpose of the data collection, potential risks and benefits and
- 147 data protection rights. Where possible a seven-day gap, between receipt of the PIL
- 148 and obtaining consent, will be provided to allow people time to consider
- 149 participation. However, if it is their preference to be seen at an earlier and more
- 150 convenient time this will be facilitated due to the innocuous nature of the

151 evaluation.

- 152 Once the research physiotherapist is satisfied that the participant has read and
- 153 fully understands the PIL, they will proceed to obtain written informed consent.
- 154 Obtaining consent will take place at the first face to face interaction with the
- 155 participant prior to commencement of the assessment. The written consent
- informs participants that they are permitted to withdraw from the study at any
- 157 time. They will be provided with their own copy of this consent form and PIL with
- their own signature and that of the research physiotherapist.
- 159

160 Stakeholder engagement

161 Recommendations from the linked LEAP II trial (16) included health education and
162 awareness workshops preceding exercise interventions to enhance health literacy,

163 target lifestyle behavioural change and improve retention rates. Prior to the LEAP-

- 164 W programme, a number of exercise workshops will be delivered for staff and
- service users in Jane's Place. Feedback from this stakeholder engagement will
- 166 inform the design of the intervention.
- 167

168 Evaluation and test battery

- 169 Using a bespoke data collection form, the following demographic data will be collected:
- age, gender, ethnicity, living arrangement and environment, highest level of education,
- 171 marital status, employment status, history of incarceration, history of addiction, self-
- 172 reported health, and medical history. Physical function, frailty and other related
- variables will be evaluated using the test battery outlined in Table 1.
- 174 Table 1. Test Battery.

Construct Measured	Test
Sarcopenia and	Upper/Lower Limb circumference
upper limb strength	Hand grip dynamometry (27, 28)
Lower limb strength	30-sec Chair Stand Test (29)
and endurance	
Endurance	2-minute walk test (30)
Speed	10m walk test (31)
Balance	Single leg stance (32)

Frailty	SHARE-FI (33), Clinical Frailty Scale (6)
Pain	Numerical Pain Rating Scale (34)
Health	Short Form 12 (35)
status/Quality of Life	
Nutritional Status	Mini-Nutritional Assessment (36, 37)

175

176 **1. Strength and muscular mass:** Muscle strength will be measured, using

a Jamar Digital Hand Dynamometer, while sitting with the elbow flexed at 90°,

the forearm mid-prone, the wrist in neutral and the hand unsupported (27,

179 28). Results will be compared to normative reference values (38). Two

180 measurements will be recorded and also used for the SHARE-Frailty

181 Instrument (FI) assessment (33).

182 Mid-calf circumference girth will be evaluated as this measure correlates with

appendicular muscular mass (39). This will be measured using a flexible tape

184 measure at the level of the largest circumference of the calf. Measurements

185 will be compared to gendered cut-off values (40). Mid-arm muscle

186 circumference reflects both muscle mass and caloric and protein adequacy

and may be used to signify wasting or malnutrition (41). This test has been

recommended in situations where lower limb swelling is present (42). The

189 maximum upper arm muscular mass will be measured using a flexible tape

190 measure. Results will be compared to global reference values (43).

2. Physical performance and lower extremity physical function:

- 192 This will be measured using the following physical performance measures:
- (i) The 10m Walk Test (10mWT). This test measures gait speed and functional
- 194 mobility and is recorded in m/s. Gait speed is calculated as total distance/time
- 195 (31).
- 196 (ii) The 2 Minute Walk Test (2MWT). This test of self-paced walking ability and
- 197 functional capacity assesses a participants' ability to walk unaided over a 15m
- distance, as fast as possible, for two minutes. Rest breaks are permitted, and
- the distance covered is measured (30).
- (iii) The Chair Stand Test (CST). This tests lower limb strength and endurance
 and records the total number of sit to stand repetitions performed in 30
 seconds (29).
- (iv) The Single Leg Stance Test (SLST). This balance test is performed on each
 leg. The participant is timed standing unassisted on one leg, with eyes open
 and hands placed on the hips (32).
- 3. Pain: Each participant will be asked if they are experiencing pain and will
 be questioned about its location and duration. Pain severity will be assessed

using the Numerical Rating Scale (NRS) (34). The NRS is a unidimensional

209 measure of pain intensity from 0-10, with 0 being zero pain and 10 the worst

210 pain imaginable.

211 **4. Frailty:** Frailty will be assessed using the Clinical Frailty Scale (CFS) (6) and the SHARE-Frailty Instrument (FI) (33). The CFS is validated for people over 65 years. It is assessed by 212 the tester. Each point on the scale correlates with a level of frailty and a visual chart aids 213 214 classification from 1 (very fit) to 9 (terminally ill). Higher scores indicate higher levels of 215 frailty. The SHARE-FI is validated for people over 50 years (33). It consists of four brief questions related to the following variables: exhaustion, loss of appetite, walking difficulties 216 217 and low physical activity, and grip strength measurement. The five results are entered into a 218 freely available web calculator to generate a frailty score and a frailty category of non-frail, pre-frail and frail is also generated. A minor modification was made to the terminology of 219 220 the SHARE-FI to reflect the real-world experiences of this population. 221 5. Nutritional status will be assessed using the Mini-nutritional assessment (MNA) which assesses the risk of malnutrition (36). The short 222 form of the MNA (MNA-SF) (37) is an efficient screening tool consisting of six 223 224 questions on food intake, weight loss, mobility, psychological stress, or acute disease, the presence of dementia or depression, and body mass index (BMI). 225

The maximum score for this part is equal to 14. A score of 12 or higher

227 indicates a normal nutritional status thus excluding malnutrition and/or

228	malnutrition risk. A score of 11 of less indicates the requirement to proceed
229	with the complete version of the MNA (MNA-LF) (37). The terminology of two
230	of the questions of the MNA (regarding acuity of illness and psychological
231	stress) were slightly modified for the purposes of evaluation of this population
232	in this setting.
233	8. Short-Form 12 (SF-12) The SF-12 V2 is a self-report measure of health
234	used across age, disease, and treatment groups (35). It uses eight domains
235	including physical and social activities, pain, mental health, emotional health,
236	vitality and general health perceptions to measure health. The tester will read
237	and complete the 12-question survey with the participant. Results will be
238	entered into a software program provided by the licensing company
239	QualityMetric and two summary scores, mental health (MCS12) and physical
240	health (PCS12) will be generated.
241	The assessor will document any issues with terminology of the outcome
242	measures utilised, difficulties with completion or floor or ceiling effects
243	reached, as these instruments were designed with geriatric populations in

244 mind.

245 Intervention

246 This intervention study will be conducted for 10 weeks. It will involve a low threshold,

247 three-times weekly exercise intervention (two exercise classes with protein

248 supplementation and an outdoor 'Park Walk'). The intervention will be delivered by two 249 research physiotherapists who will adhere to all safety procedures. Flexibility will be 250 facilitated with the provision of a four-week window for pre-intervention evaluations 251 prior to commencement of the programme. Using a trauma-informed lens (20) flexibly arranged group or one-to one sessions will be delivered based on participant preference. 252 253 Exercise programming strategies and variables are based on our earlier work (LEAP I and LEAP II trials) (16, 17) and also informed by prior stakeholder involvement from people 254 255 with lived experience. The exercise classes will be multi-modal, with a primary focus on strength and based on core set of resistance exercises (Table 2). Aerobic, balance and 256 257 flexibility work will be integrated into the class and the exercises will be individualised based on initial assessment results and presentation of participants. Using a gender-258 based perspective, the following considerations and adaptions will be built into the class; 259 260 (i) core-stability exercises to target pelvic floor and/or abdominal muscle weakness, (ii) bone building exercises to target peri-and postmenopausal bone loss and (iii) age-261 associated muscle mass loss. Music, dance and fun orientated physical activity games, an 262 important feature of aforementioned linked studies, will be incorporated to optimize 263 264 enjoyment and self-esteem. Borg's Rate of Perceived Exertion Scale will be used to monitor effort and scale the intensity of the workout (44). To promote post-exercise 265 266 muscle protein synthesis, a nutritional supplement (200ml pre-prepared 'protein shake', Fresubin) which consists of 20g of protein will be offered immediately post exercise. The 267 'Park Walk' will focus on the aerobic component of the intervention, using open green 268

- space to maximise physical and mental health outcomes. It will be a flexibly arranged
- 270 20–30-minute self-paced group or one-to-one walk. To build sustainability beyond the
- 271 ten-week programme brief health promoting and physical activity educational
- 272 interventions will be included in the exercise class setting to empower people to engage
- in unsupervised exercise following the study.
- 274

275 Table 2. Exercise Circuit.

Core exercise	Initial Intensity	Progression/Adaptations*
Sit to stand/squats/lunges	2 sets 10-15 reps	3 sets of 15 reps
		use of weights/ball
Elbow Bends	2 sets 10-15 reps	3 sets of 15 reps
		use of weights/theraband
Step-ups	2 sets 10-15 reps	3 sets of 15 reps
		Vary height of step; use of weights
Arm elevations or boxing	2 sets 10-15 reps	3 sets of 15 reps
		use of weights or boxing gloves/pads
'Penguin waddle'-hip	2 sets 10-15 reps	3 sets of 15 reps
abduction		additional upper limb abduction and
		elevation; movement with 360° turns
Scapular retractions	2 sets 10-15 reps	3 sets of 15 reps
		Weights/theraband

Aerobic activity	5 mins	Exercise bike, ladders, hurdles,
		skipping ropes, end of session fun
		activity.
Core set	5 mins	Pelvic floor exercises; pelvic tilts,
		bridging, all 4's set

276 Adaptations: exercises individualised and progressed for each participant by research physiotherapist

277 Progression: 2-3 sets of 10-15 repetitions completed at each session; beyond that, low resistance weights

278 (1.1kg-4.5kg), theraband or increased step height introduced to achieve a strengthening effect.

279

Qualitative data 280

281 Following the intervention one-to-one in-person exit interviews will be conducted to

282 enable participants to share their feedback and express their views, in order to enhance

283 the understanding of behaviours, adherence and retention challenges and to provide

insights for future programmes in this area (45) (S3 Fig). Interviews will continue until 20 284

285 participants are seen or until data saturation is reached. The interviews will be audio-

recorded and transcribed verbatim and member checking will be utilised. 286

287

Sample size 288

As this is a novel field of research, sample size calculations are challenging. The sample 289

method used will be non-probability, convenience sampling, involving the recruitment of a 290

291 tailored sample that is in proportion to characteristics of the population of interest. This

approach may limit generalisability of results but will minimise selection bias (46). A sample
size of 30 or greater has been proposed for feasibility studies (47). Considering this and the
aim and objectives of this study a target of 40-50 participants has been selected, to allow for
drop-out rate of approximately 30%.

296 Data collection and management

297 Analytic plan

298 All data will be pseudonymised at point of entry into excel spreadsheets and then transferred into IBM SPSS V28 for analysis. For quantitative data, nominal or ordinal 299 variables will be reported as frequencies and percentages. Continuous variables will be 300 summarised as mean and standard deviation if normally distributed and median and inter-301 302 quartile range if non-normally distributed. Data will be tested for normality using the Kolmogorov–Smirnov/Shapiro Wilk test and will be compared across timepoints using the 303 general linear model procedure (normally distributed data) and the Friedman's test (non-304 305 normally distributed data). Chi-squared t-tests will be used where appropriate and some data may be categorised to investigate relationships between variables. Exploratory 306 regression models will be developed to explore correlates and predictors of frailty and poor 307 physical functioning. A p-value of <0.05 will be considered significant. For the qualitative 308 data, Braun and Clarke thematic analysis methodology will be employed to provide an in-309 depth analysis of the exit interview data (48). 310

311 Discussion

312	Poor physical functioning and frailty has been demonstrated in younger populations who
313	suffer from health disparities and an extreme form of social exclusion, such as PEH and
314	addiction challenges (12, 13, 49). Women who experience homelessness and addiction are
315	especially vulnerable and have unique and complex needs which require a targeted and
316	gender-sensitive approach. Little is known about frailty-focussed interventions in this
317	population, who are often excluded from research and services (18). However, this
318	population have engaged well in two previous linked studies (LEAP I and II). This study aims
319	to address the research gap by providing a targeted exercise intervention to women
320	experiencing homelessness and addiction challenges, thereby providing a gendered
321	dimension, with gendered sensitivity. Its findings will guide clinicians and policy makers in
322	implementing targeted interventions to assist in improving health outcomes in societies
323	most marginalised and vulnerable groups.

324

325 Study status

Recruitment and data collection will commence in February 2024 and will be completed byJuly 2024.

328 Funding

329 This has been funded by Trinity College Dublin and the Irish Research Council.

330 **Dissemination plans**

- 331 Conference presentations and publications in peer-reviewed journals will be one method of
- dissemination. These will be done following the data analysis.

333 Acknowledgments

- 334 The authors wish to extend their gratitude to the staff of Jane's Place, MQI who are helping
- to recruit participants.

336 Competing interests

337 The authors have no competing interests to declare in connection with this article.

338 Data availability

- 339 There are no underlying data associated with this protocol. Data from this study will be
- 340 available in open access form.

341 **References**

- 342 1. Nations U. Global Issues: Our growing population2023. Available from:
- 343 https://www.un.org/en/global-issues/population.
- Morley JE, Vellas B, van Kan GA, Anker SD, Bauer JM, Bernabei R et al Frailty consensus: a call to action. J Am Med Dir Assoc, 2013. (6): p. 392-7. doi: 10.1016/j.jamda.2013.03.022.
- Turner G and Clegg A. Best practice guidelines for the management of frailty: a British Geriatrics
 Society, Age UK and Royal College of General Practitioners report. Age Ageing. 2014;43(6):744-7. doi:
 10.1093/ageing/afu138.
- Travers J, Romero-Ortuno R, Bailey J & Cooney MT. Delaying and reversing frailty: a systematic
 review of primary care interventions. *Br J Gen Pract*, 2019. 69(678): p. e61-e69. doi:
 10.3399/bjgp18X700241.
- 3525.Travers J, Romero-Ortuno R, Langan J, MacNamara F, McCormack D, McDermott C, et al. Building353resilience and reversing frailty: a randomised controlled trial of a primary care intervention for older354adults. Age Ageing, 2023. 52(2). doi: 10.1093.

Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I et al. A global clinical measure
of fitness and frailty in elderly people. *Cmaj*, 2005. 173(5): p. 489-95. doi: 10.1503/cmaj.050051.

- Romero-Ortuno R and O'Shea D. Fitness and Frailty: opposite ends of a challenging continuum! Will
 the end of age discrimination make frailty assessments an imperative? Ae Ageing, 2013. 42(3) p. 279 10.1093/ageing/afs189.
- Loecker C, Schmaderer M, Zimmerman L. Frailty in Young and Middle-Aged Adults: An Integrative
 Review. J Frailty Aging. 2021;10(4):327-33. doi: 10.14283/jfa.2021.14.
- 3629.Spiers GF, Kunonga TP, Hall A, Beyer F, Boulton E, Parker S, et al. Measuring frailty in younger363populations: a rapid review of evidence. BMJ Open. 2021;11(3):e047051. doi: 10.1136/bmjopen-3642020-047051.
- Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS. Frailty and pre-frailty in middle-aged and
 older adults and its association with multimorbidity and mortality: a prospective analysis of 493 737
 UK Biobank participants. Lancet Public Health. 2018;3(7):e323-e32. doi: 10.1016/S24682667(18)30091-4
- Aldridge RW, Story A, Hwang SW, Nordentoft M, Luchenski SA, Hartwell G, et al. Morbidity and
 mortality in homeless individuals, prisoners, sex workers, and individuals with substance use disorders
 in high-income countries: a systematic review and meta-analysis. Lancet. 2018;391(10117):241-50.
 doi: 10.1016/S0140-6736(17)31869-X.
- Broderick J, Kiernan S, Murphy N, Dowds J, C NC. Feasibility of a Broad Test Battery to Assess Physical
 Functioning Limitations of People Experiencing Homelessness. Int J Environ Res Public Health.
 2021;18(3). doi: 10.3390/ijerph18031035.
- Kiernan S, C NC, Murphy N, Dowds J, Broderick J. Markedly poor physical functioning status of people
 experiencing homelessness admitted to an acute hospital setting. Sci Rep. 2021;11(1):9911. doi:
 10.1038/s41598-021-88590-0.
- Tembo MC, Holloway-Kew KL, Sui SX, Dunning T, Low ACH, Yong SJ, et al. Prevalence of Frailty in Older
 Men and Women: Cross-Sectional Data from the Geelong Osteoporosis Study. Calcif Tissue Int.
 2020;107(3):220-9. doi: 10.1007/s00223-020-00713-3.
- Zhang Q, Guo H, Gu H, Zhao X. Gender-associated factors for frailty and their impact on
 hospitalization and mortality among community-dwelling older adults: a cross-sectional population based study. PeerJ. 2018;6:e4326. doi: 10.7717/peerj.4326.
- Kennedy F, Romero-Ortuno R, Ní Cheallaigh C, Doyle S and Broderick, J. "It's like a free drug"-Low
 Threshold Exercise and Nutritional Programmes for People with Severe and Multiple Disadvantage
 Accessing Services for Addicition Chaleenges and Homelessness. Pathways from Homelessness.
 International Conference of the Faculty for Homeless and Inclusion Helath; 2023 March 16-17, 2023;
 London.
- 39017.Kennedy, F., Ní Cheallaigh, C., Romero-Ortuno, R., Doyle, S. and Broderick, J. Outcomes of the LEAP391feasibility trial-a low threshold exercise programme with protein supplementation to target frailty and392poor physical functioing in people experiencing homelessness and addiction issues. PLOS ONE (in393press)
- Addiction EMCfDaD. Health and Social responses to drug problems: A European Guide2022. Available
 from: https://www.emcdda.europa.eu/publications/mini-guides/homelessness-and-drugs-health and-social-responses_en.
- C NC, Cullivan S, Sears J, Lawlee AM, Browne J, Kieran J, et al. Usage of unscheduled hospital care by homeless individuals in Dublin, Ireland: a cross-sectional study. BMJ Open. 2017;7(11):e016420. doi: 10.1136/bmjopen-2017-016420.
- 400 20. FEANTSA EFONOWwtHFEWaH. 2019-2021. Available from:
- 401 https://www.feantsa.org/en/project/2019/01/03/women-and-homelessness.
- 402 21. The Department of Housing, Planning and Local Government (Feb 2018) Homelessness Report. 2018.
 403 Available from: https://www.gov.ie/en/publication/793bc-homeless-report-february-

404		2018/?referrer=http://www.housing.gov.ie/sites/default/files/publications/files/homelessness_repor
405	22	tfebruary_2018_0.pdf
406	22.	Mutatayi C MS, Robles Soto N, Palsdottor K & Vale Pires C. Implementing a Gender Approach in Drug
407		Policies: Prevention, Treatment and Criminal Justice. A handbook for practitioners and decision
408		makers. Council of Europe:Strasbourg.2022. Available from:
409		https://www.drugsandalcohol.ie/36974/
410	23.	Milani K, Ramage, K., Yang Fang, X. and Louis, M. Understanding Mothers experiencing homelessness:
411		A gendered approach to finding solutions for family homelessness. 2016. Available from:
412		https://www.homelesshub.ca/sites/default/files/attachments/Family_Homelessness_Report.pdf
413	24.	Morton SG, B. and McLoughlin, E "You can't fix this in six months": The intersectionality of women's
414		substance use in the Irish context In: Ireland UCDaMQ, editor. 2023. Available from:
415		https://www.drugsandalcohol.ie/38492/
416	25.	European Commission, Directorate-General for Research and Innovation, Gender in EU-funded
417		research – Toolkit, Publications Office, 2014, https://data.europa.eu/doi/10.2777/74150
418	26.	Kennedy F, Ní Cheallaigh, C., Romero-Ortuno et al. A study to explore the role of a low
419		threshold, fitness focussed physical rehabilitation intervention with protein supplementation
420		to target physical function and frailty in people with problematic substance use and
421		homelessness; protocol for a single-arm feasibility cohort study. HRB Open Research.
422		2023.6 doi org/10 12688/hrbonenres 13678 1
122	27	Leong DP. Teo KK. Rangarajan S. Lonez-Jaramillo P. Avezum A. Jr. Orlandini A. et al
425	27.	Drognostic value of grin strength: findings from the Prospective Urban Bural Enidemiology (DUBE)
424		study Lancet 2015-386(0000)-266-73 doi: 10.1016/S0140-6736(14)62000-6
425	28	Roberts HC Denicon HI Martin HI Datel HD Syddall H Cooper C et al. A review of the measurement
420	20.	of grin strength in clinical and enidemiological studies: towards a standardised approach. Age Ageing
427		2011.40/4).422 0. doi: 10.1002/agoing/afr0E1
420	20	2011,40(4).425-9. UOI. 10.1095/dgellig/di1051.
429	29.	Lefit DH, JL, Alotabi W, Almutali W, Singi H. Normative Reference values and valuaty for the So-
430	20	Second Chail-Stahu Test in Healthy foung Adults. Int J Sports Phys Ther. 2022,17(5):507-14.
431	50.	brooks D, Davis Alvi, Naglie G. The reasibility of six-initiate and two-initiate waik tests in in-patient
432	24	genatric renabilitation. Can't Aging. 2007;26(2):159-62. doi: 10.3138/Cja.26.2.009.
433	31.	Bonannon RW. Comfortable and maximum walking speed of adults aged 20-79 years: reference
434		values and determinants. Age Ageing. 1997;26(1):15-9. doi: 10.1093/ageing/26.1.15.
435	32.	Springer BA, Marin R, Cyhan T, Roberts H, Gill NW. Normative values for the unipedal stance test with
436		eyes open and closed. J Geriatr Phys Ther. 2007;30(1):8-15. doi: 10.1519/00139143-200704000-
437		00003. doi: 10.1519/00139143-200/04000-00003.
438	33.	Romero-Ortuno R, Walsh CD, Lawlor BA & Kenny RA. A frailty instrument for primary care: findings
439		from the Survey of Health, Ageing and Retirement in Europe (SHARE). BMC Geriatr, 2010.10: p. 57.
440	~ .	doi: 10.1186/1471-2318-10-57.
441	34.	Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain
442		(VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form
443		McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain
444		Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). Arthritis
445		Care Res (Hoboken). 2011;63 Suppl 11:S240-52. doi: 10.1002/acr.20543.
446	35.	Ware J, Jr., Kosinski M, Keller SD. A 12-item Short-Form Health Survey: construction of scales and
44/		preliminary tests of reliability and validity. Med Care. 1996;34(3):220-33. doi: 10.1097/00005650-
448	20	199003000-00003.
449	30.	Guigoz Y, Veilas B, Garry PJ. Assessing the nutritional status of the elderly: The Mini Nutritional
450		Assessment as part of the geriatric evaluation. Nutr Rev. 1996;54(1 Pt 2):559-65. doi: 10.1111/j.1753-
451		4887.1930.tDU3793.X.

452	37.	Kaiser MJ, Bauer JM, Ramsch C, Uter W, Guigoz Y, Cederholm T, et al. Validation of the Mini
453		Nutritional Assessment short-form (MNA-SF): a practical tool for identification of nutritional status. J
454		Nutr Health Aging. 2009;13(9):782-8. doi: 10.1007/s12603-009-0214-7.
455	38.	Steiber N. Strong or Weak Handgrip? Normative Reference Values for the German Population across
456		the Life Course Stratified by Sex, Age, and Body Height. PLoS One. 2016;11(10):e0163917.
457 458	39.	Rolland Y, Lauwers-Cances V, Cournot M, Nourhashémi F, Reynish W, Rivière D, et al. Sarcopenia, calf circumference, and physical function of elderly women: a cross-sectional study. J Am Geriatr Soc.
459		2003:51(8):1120-4. doi: 10.1046/i.1532-5415.2003.51362.x.
460	40.	Gonzalez MC, Mehrnezhad A, Razaviarab N, Barbosa-Silva TG, Heymsfield SB. Calf circumference:
461		cutoff values from the NHANES 1999-2006. Am J Clin Nutr. 2021;113(6):1679-87. doi:
462		10.1093/ajcn/nqab029.
463	41.	Cano NJ, Miolane-Debouit M, Léger J, Heng AE. Assessment of body protein: energy status in chronic
464		kidney disease. Semin Nephrol. 2009;29(1):59-66. doi: 10.1016/j.semnephrol.2008.10.008.
465	42.	Chen B, Mitchell A, Tran D. "Step up for foot care": addressing podiatric care needs in a sample
466		homeless population. J Am Podiatr Med Assoc. 2014;104(3):269-76.
467		doi: 10.1371/journal.pone.0167463
468	43.	Tang AM, Chung M, Dong KR, Bahwere P, Bose K, Chakraborty R, et al. Determining a global mid-
469		upper arm circumference cut-off to assess underweight in adults (men and non-pregnant women).
470		Public Health Nutr. 2020;23(17):3104-13. doi: 10.1017/S1368980020000397.
471	44.	Borg GA. Psychophysical bases of perceived exertion. Med Sci Sports Exerc. 1982;14(5):377-81.
472	45.	Rosenthal M. Qualitative research methods: Why, when, and how to conduct interviews and focus
473		groups in pharmacy research. Currents in Pharmacy Teaching and Learning. 2016;8(4):509-16.
474		https://doi.org/10.1016/j.cptl.2016.03.021.
475	46.	Althubaiti A. Sample size determination: A practical guide for health researchers. J Gen Fam Med.
476		2023;24(2):72-78.doi: 10.1002/jgf2.600.
4//	47.	Lancaster GA, Dodd S, Williamson PR. Design and analysis of pilot studies: recommendations for good
4/8		practice. J Eval Clin Practice 2004;10:307-312. doi: 10.1111/j2002.384.doc.x.
479	48.	Braun V, Clarke V. What can "thematic analysis" offer health and wellbeing researchers? Int J Qual
480	40	Stud Health Well-being. 2014;9:26152. doi:10.3402/qhw.v9.26152.
481	49.	Rogans-Watson RS, C. Lewer, D. Armstrong, M and Hudson B. Premature fraility, geriatric conditions
482		and multimorbialty among people experiencing nomelessness: a cross-sectional study in a London
403 101		nostei. Housing Care and Support.2020.23:77-91. DOI: 10.1108/HCS-05-2020-0007.
404		

485

486 Supporting information

487 S1 Fig. Exit Interview Schedule