

# Learning from the COVID-19 pandemic: a systematic review of mathematical vaccine prioritization models

Gilberto Gonzalez-Parra<sup>a,b,\*</sup>, Md Shahriar Mahmud<sup>c</sup>, Claus Kadelka<sup>c,\*</sup>

<sup>a</sup>*Instituto de Matemática Multidisciplinar, Universitat Politècnica de València, València, Spain*

<sup>b</sup>*Department of Mathematics, New Mexico Tech, 801 Leroy Place, Socorro, 87801, NM, USA*

<sup>c</sup>*Department of Mathematics, Iowa State University, 411 Morrill Rd, Ames, 50011, IA, USA*

---

## Abstract

As the world becomes ever more connected, the chance of pandemics increases as well. The recent COVID-19 pandemic and the concurrent global mass vaccine roll-out provides an ideal setting to learn from and refine our understanding of infectious disease models for better future preparedness. In this review, we systematically analyze and categorize mathematical models that have been developed to design optimal vaccine prioritization strategies of an initially limited vaccine. As older individuals are disproportionately affected by COVID-19, the focus is on models that take age explicitly into account. The lower mobility and activity level of older individuals gives rise to non-trivial trade-offs. Secondary research questions concern the optimal time interval between vaccine doses and spatial vaccine distribution. This review showcases the effect of various modeling assumptions on model outcomes. A solid understanding of these relationships yields better infectious disease models and thus public health decisions during the next pandemic.

*Keywords:* Review, mathematical model, age, COVID-19, vaccine allocation, vaccine roll-out

---

\*Contributed equally, gilberto.gonzalezparra@nmt.edu and ckadelka@iastate.edu

## 1. Introduction

In December 31, 2019, the World Health Organization (WHO) was informed of several cases of a pneumonia of unknown cause occurring in Wuhan, China ([Centers for Disease Control \(CDC\)](#)). Only 71 days later, the WHO declared - after 118,000 cases in 114 countries and 4,291 deaths - COVID-19, caused by the Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV-2), a pandemic. Despite the widespread implementation of numerous types of non-pharmaceutical interventions (NPIs), aimed at curbing virus spread and keeping hospitals functional, COVID-19 continued to spread rapidly in the absence of a vaccine ([Odusanya et al., 2020](#)). Recent advances in mRNA vaccine technology enabled rapid development of highly effective vaccines ([Zhang et al., 2019](#)). Since the globalized world had never experienced a pandemic, nor a mass vaccine roll-out at this scale, various formerly mainly theoretical questions related to vaccine access and prioritization became all of a sudden very important. These included: Who should be vaccinated first? Should the second vaccine dose be delayed in order to provide a first dose to more people? What parameters must be taken into account to accurately determine the best prioritization strategy? Should high-income countries share some of their limited vaccine with poorer countries? For ethical reasons only, or does a more equitable vaccine coverage have even epidemiological benefits? In response to these quickly emerging questions, scientists from many fields started to collaborate and suggest answers.

Globally, by the end of 2023, there have been over 770 million confirmed COVID-19 cases and over 7 million deaths, reported to the WHO ([World Health Organization \(WHO\)](#)). Despite the development of highly effective vaccines and 13.6 billion administered COVID-19 vaccine doses, with over 72 percent of the world population having received at least one dose ([New York Times](#)), the disease still surges in waves around the world in early 2024. While the infection fatality rate is now substantially lower than in the beginning of the pandemic ([Sorensen et al., 2022](#)) and most NPIs have disappeared, large COVID-19 outbreaks and community spread still appear around the world, causing, for example, numerous individuals to suffer from so-called long COVID symptoms that can linger for years post infection ([Sudre et al., 2021](#)). Reasons COVID-19 has not disappeared after sufficient production of vaccines include the ongoing emergence of SARS-CoV-2 variants, partial vaccine escape by some variants ([Chakraborty et al., 2022](#); [Wang et al., 2021](#)), issues related to vaccine access and distribution specifically in low-income

38 countries (Sheikh et al., 2021), as well as vaccine hesitancy and wariness  
39 driven by rampant misinformation (Sallam, 2021). Learning from mistakes  
40 made during the COVID-19 pandemic and the first global mass vaccine roll-  
41 out is thus paramount for future pandemic preparedness.

42 This review identifies and analyzes a variety of studies related to finding  
43 the optimal vaccine allocation given a limited supply. While the specific re-  
44 search questions and settings differ from study to study, age is a crucial factor  
45 in all COVID-19 vaccine prioritizations as older people have a substantially  
46 higher COVID-19 fatality rate. The primary focus in this review is there-  
47 fore on studies that were based on a mathematical model, which takes age  
48 into consideration. Other important attributes which were used by public  
49 health decision-makers to differentiate COVID-19 vaccine access and which  
50 are investigated in some of the studies include, among others, occupation  
51 (e.g., prioritizing healthcare and essential workers) and comorbidity status  
52 (e.g., prioritizing individuals with known risk factors). Moreover, the rec-  
53 ommended two-dose vaccine regimen for most COVID-19 vaccines raised the  
54 related prioritization question whether it is beneficial to delay the second dose  
55 in order to increase initial vaccine coverage. Another related prioritization  
56 question concerns spatial aspects (e.g., the optimal distribution of limited  
57 vaccine supply between different states or countries). We summarize innova-  
58 tive and interesting mathematical model-based studies that investigate these  
59 related prioritization questions, no matter whether the models specifically  
60 consider age.

61 Given the large number of studies related to optimal COVID-19 vaccine  
62 allocations, we decided to restrict ourselves to studies that employ a math-  
63 ematical model for decision-making. The included studies employ several  
64 modeling frameworks. Most studies are based on an ordinary differential  
65 equation (ODE) model, in which the population is stratified into different  
66 compartments. The simplest model, colloquially known as SIR model and  
67 first studied nearly 100 years ago (Kermack and McKendrick, 1927), con-  
68 tains three compartments: susceptible (S), infected (I), and recovered (R).  
69 More complex models possess additional compartments for individuals that  
70 are e.g. infected but not yet infectious, asymptotically versus symptomat-  
71 ically infected, quarantined but not yet recovered, or dead. To account for  
72 different ages and possibly other attributes (e.g., occupation), the population  
73 is stratified into a finite number of sub-populations (e.g., age classes) and the  
74 compartments are duplicated for each sub-population. Each sub-population  
75 can have its own characteristics. This enables modelers to account for het-

76 erogeneity (e.g., age dependency) in contact patterns, NPI adherence, vac-  
77 cine hesitancy, susceptibility to infection, as well as various factors related  
78 to disease progression. ODE-based models implicitly make a number of as-  
79 sumptions that are inaccurate for COVID-19 disease dynamics and hard to  
80 overcome within the ODE modeling framework. Among others, they assume  
81 that (i) the population is homogeneously mixed, (ii) the time spent in each  
82 transient compartment is exponentially distributed, and (iii) disease dynam-  
83 ics are deterministic.

84 Another modeling framework, agent-based models (ABMs; also known as  
85 individual-based models), is stochastic in nature and employed by a smaller  
86 number of studies. In ABMs, individual agents (i.e., people) are modeled;  
87 agents interact with each other and possibly spread the disease through e.g.  
88 heterogeneous interaction networks. This modeling framework is highly flex-  
89 ible (e.g., each individual can have its own characteristics and decision rules)  
90 and can be adaptive (e.g., the decisions of an agent can depend on other's  
91 decisions). However, ABMs inherently rely on simulations. Their stochastic  
92 nature further increases the computational needs, rendering an exhaustive  
93 exploration of a large parameter space impossible. Lastly, a few studies  
94 employ partial differential equation (PDE) models. These studies typically  
95 focus on spatial aspects of vaccine prioritization.

96 Contrary to other review articles on this topic ([Saadi et al., 2021](#); [Liu and](#)  
97 [Lou, 2022](#); [Noh et al., 2021](#); [Thakkar and Spinardi, 2023](#)), the focus of this  
98 review is on understanding the effect of modeling assumptions and paramete-  
99 rs on policy recommendations. For example, while most studies agree that  
100 elderly and vulnerable should be vaccinated first due to their substantially  
101 higher infection fatality ratio, some studies suggest the opposite. We look in  
102 detail at which model parameters and assumptions cause these discrepancies.

103 This review includes 94 articles, which use a mathematical model to an-  
104 swer at least one of three questions related to COVID-19 vaccine prioritiza-  
105 tion:

- 106 1. How should a limited vaccine be optimally distributed among a popu-  
107 lation stratified by age (and possibly other factors)?
- 108 2. For limited vaccines with a two-dose regimen, should the second dose  
109 be delayed in order to provide more people with a first vaccine dose?
- 110 3. How should a limited vaccine be optimally distributed given spatial  
111 heterogeneity?

112 In Section 2, we briefly describe how we identified articles of interest. Sec-




Figure 1: Number of (A) investigated and (B) included studies, stratified by source.

113 tion 3 summarizes, at a high-level, the main findings of these articles related  
114 to vaccine prioritization. Section 4 puts these findings into context, particu-  
115 larly those without a clear consensus strategy. Several key COVID-19 model  
116 parameters and assumptions are introduced, with a focus on how they affect  
117 optimal vaccine prioritization strategies. Section 5 provides a brief summary  
118 of particularly interesting and noteworthy studies. Finally, Section 6 briefly  
119 presents related works that employ optimal control methods to answer ques-  
120 tions related to vaccine prioritization.

## 121 2. Methods

122 To find studies of interest, we searched PubMed, the Web of Science Core  
123 Collection and MathSciNet (all in February 2024) for research articles that  
124 contain the following keywords: 'age' AND 'model' AND ('COVID-19' OR  
125 'SARS-CoV-2') AND ('vaccine' OR 'vaccination') AND ('best' OR 'optimal'  
126 OR 'priorit\*') AND ('mathematical' OR 'computational' OR 'stochastic' OR  
127 'network'). After removing duplicates (e.g., preprints and journal articles)  
128 and non-peer-reviewed preprints, this yielded a total of 285 articles, which  
129 we manually reviewed, in addition to 43 articles known to the authors and/or  
130 referenced in one of the 285 articles (Fig. 1A). For each article, we decided if  
131 it contained a mathematical model that answers at least one of the COVID-  
132 19 vaccine prioritization questions stated above. This yielded a total of 94  
133 articles included in this review (Fig. 1B). Any article that did not assume lim-  
134 ited vaccine availability (e.g., studies looking into the epidemiological effect  
135 of boosters in high-income countries) was excluded.

136 Eighty of the included articles contain an age-stratified mathematical

137 model that provides answers to our primary research question: How should  
138 a limited vaccine be optimally distributed among a population stratified by  
139 age? Fifteen articles contain a mathematical model (not necessarily age-  
140 stratified) to answer the secondary research question: For limited vaccines  
141 with a two-dose regimen, should the second dose be delayed in order to pro-  
142 vide more people with a first vaccine dose? Finally, seven articles contain  
143 a mathematical model that considers spatial aspects of vaccine distribution.  
144 The low number of articles related to the latter two research questions is, at  
145 least partially, due to the fact that the keywords were selected to preferen-  
146 tially find articles investigating our primary research question.

### 147 **3. Summary of findings**

148 Collectively, the included articles contain models that are tailored to  
149 cities, states or countries from all continents except Antarctica (Fig. 2). A  
150 few studies tailor their model to more than one country to showcase how  
151 variability in e.g. age distributions, age-stratified contact patterns, or imple-  
152 mented NPIs can affect optimal vaccine prioritization, see e.g., [Gozzi et al. \(2021\)](#);  
153 [Liu et al. \(2022b\)](#); [Wang et al. \(2022\)](#); [Liu et al. \(2022a\)](#). Other stud-  
154 ies employ more abstract models, frequently ABMs, which are not tailored  
155 to any specific setting, see e.g., [Romero-Brufau et al. \(2021\)](#); [Grauer et al. \(2020\)](#);  
156 [Kadelka and McCombs \(2021\)](#). These models contain tuneable pa-  
157 rameters and are well-suited to reveal the qualitative dependence of optimal  
158 allocation strategies on key parameters and assumptions.

159 While the included articles employ a variety of metrics to quantify the  
160 quality of a given vaccine allocation strategy, there are several common ob-  
161 jectives. In decreasing order of use (Fig. 3), these include: minimizing deaths  
162 (used in 80 of the 94 included studies), cases/infections (56), hospitalizations  
163 (22), and years of life lost (YLL; 10). Other, less frequently used objectives  
164 include minimizing quality- and disability-adjusted life years (QALYS and  
165 DALYS, respectively; used in 6 studies), minimizing the peak number of  
166 hospitalized, as well as several equitable and economic considerations. While  
167 technically different and considered as separate objectives in at least one  
168 study ([Islam et al., 2021](#)), we do not differentiate between the objectives  
169 minimizing cases and infections. Some studies attempt, furthermore, to op-  
170 timize multiple objectives at the same time, e.g., through the use of optimal  
171 control methods (summarized in Section 6), or an analysis of Pareto-optimal  
172 allocation strategies.




Figure 2: **Number of studies that contain a model for a specific country.** Some studies include models tailored to several countries, while others are more abstract and not tailored to a specific country. Census data and age-stratified contact matrices are two examples of frequently used country-specific data.

173 When minimizing mortality is the sole objective, the majority of studies  
174 (47 out of 70) agree that vaccinating older individuals, vulnerable individuals,  
175 and - if considered - health care workers first is optimal, irrespective of  
176 the specific setting or assumptions (Fig. 3A, Table 1). There exists, however,  
177 some disagreement about the prioritization among these subpopulations.  
178 Interestingly, 23 model-based studies (32.9%) conclude that under  
179 certain circumstances a prioritization of younger people who have on average  
180 more contacts leads to lower death counts. Qualitatively, the optimal  
181 prioritization strategies do generally not shift much when minimizing other  
182 morbidity-based metrics such as YLL or hospitalizations. On the other hand,  
183 most studies (41 out of 48, 85.4%) agree that to minimize the total number  
184 of infections and/or the effective reproductive number younger individuals  
185 should be vaccinated first since they typically have more contacts and thus  
186 more chances to spread the virus (Fig. 3A).



<b>A</b>		prioritize			<b>B</b>		modify dosing interval			
		when minimizing	older	depends			younger	when minimizing	longer	depends
	cases/infections		2	5	41		cases/infections	7	1	0
	YLL		3	6	0		YLL	1	0	0
	QALYS/DALYS		3	3	0		QALYS/DALYS			
	hospitalizations		10	4	3		hospitalizations	2	0	0
	deaths		47	20	3		deaths	9	3	1

Figure 3: **High-level summary of findings.** (A) Number of studies that agree at a high-level on a given prioritization strategy (columns) when minimizing a given metric (rows). Only studies that are based on a mathematical model that considers stratifying vaccine access by age are included. Note that all studies that recommend prioritization of the oldest and most vulnerable people, possibly after vaccinating health care workers, were nevertheless counted as prioritizing older. (B) Number of studies that agree at a high-level on a dosing interval strategy (columns) when minimizing a given metric (rows). (A-B) The second column (“depends”) includes all studies that present more subtle findings where the prioritization and dosing interval depends on certain assumptions. In Table 1 and Table 2, the high-level summaries are stratified by study.



Table 1: **High-level summary of COVID-19 vaccine prioritization studies.** Each row summarizes the high-level prioritization strategy identified by a given study (white: not assessed, red: prioritize older/vulnerable population, blue: prioritize younger/high-contact population, gray: prioritization depends on model assumptions). Fig. 3A provides summary counts. Only studies that were based on a mathematical model that accounts for age were included. The models are further classified by framework, type of vaccine roll-out (as specified in 4.4), as well as considered vaccine functions.

Reference	prioritization when minimizing					modeling framework			type vaccine roll-out	vaccine-induced reduction in				
	cases/infections	YLL	QALYS/DALYS	hospitalizations	deaths	ODE	ABM	optimal control		infection	symptomatic disease	hospitalization	death	onward transmission
<a href="#">Althobaity et al. (2022)</a>	blue					gray			2	gray	gray			
<a href="#">Angelov et al. (2023)</a>					blue	gray		gray	2	gray				
<a href="#">Anupong et al. (2023)</a>	blue				red	gray			2	gray	gray			gray
<a href="#">Aruffo et al. (2022)</a>	blue					gray			4	gray				
<a href="#">Ayoub et al. (2021)</a>	red			red	red	gray			4	gray	gray			gray
<a href="#">Ben-Zuk et al. (2022)</a>	blue			red		gray	gray		2	gray				
<a href="#">Bubar et al. (2021)</a>	blue				red	gray			2	gray				
<a href="#">Buckner et al. (2021)</a>	blue	gray			gray	gray			2	gray				
<a href="#">Bushaj et al. (2023)</a>	blue			blue	gray	gray	gray		?	gray				
<a href="#">Campos et al. (2021)</a>					red	gray			5	gray				
<a href="#">Cartocci et al. (2021)</a>	blue		gray		red	gray			2	gray				
<a href="#">Cattaneo et al. (2022)</a>					red	gray	gray		5	gray	gray			
<a href="#">Chen et al. (2021)</a>	blue			gray	gray	gray	gray		5	gray				
<a href="#">Childs et al. (2022)</a>	blue					gray			2	gray		gray		
<a href="#">Choi et al. (2021)</a>	gray				red	gray			2	gray				
<a href="#">Choi and Shim (2021)</a>	gray			gray		gray	gray		2	gray	gray			
<a href="#">Conway et al. (2023)</a>	blue				blue	gray	gray		4	gray				gray
<a href="#">Ferranna et al. (2021)</a>	blue	gray			red	gray	gray		2	gray			gray	gray
<a href="#">Ferreira et al. (2022)</a>					red	gray			4	gray	gray	gray		
<a href="#">Foy et al. (2021)</a>	gray				red	gray			2	gray		gray		gray
<a href="#">Gavish and Katriel (2022)</a>	blue				red	gray			?	gray				gray
<a href="#">González-Parra et al. (2022)</a>					red	gray			2	gray				
<a href="#">Gozzi et al. (2021)</a>	blue					gray			2	gray	gray			
<a href="#">Gozzi et al. (2022)</a>	blue				red	gray			4	gray				gray
<a href="#">Grundel et al. (2021)</a>					gray	gray	gray		2	gray				
<a href="#">Han et al. (2021)</a>	blue			red	red	gray			2	gray	gray			
<a href="#">Hogan et al. (2021)</a>		red			red	gray			5	gray	gray			

**Table 1** continued

Reference	prioritization when minimizing				modeling framework			type vaccine roll-out	vaccine-induced reduction in				
	cases/infections	YLL	QALYS/DALYS	hospitalizations	deaths	ODE	ABM		optimal control	infection	symptomatic disease	hospitalization	death
<a href="#">Hong et al. (2022)</a>	■				■			5	■				
<a href="#">Hupert et al. (2022)</a>	■			■	■			5	■	■			■
<a href="#">Islam et al. (2021)</a>	■	■			■			5	■	■			■
<a href="#">Jahn et al. (2021)</a>				■	■	■		5,4					
<a href="#">Jentsch et al. (2021)</a>					■			2	■	■			
<a href="#">Kadelka and McCombs (2021)</a>					■		■	5	■				
<a href="#">Kadelka et al. (2022)</a>	■				■		■	4	■		■		■
<a href="#">Karabay et al. (2021)</a>					■		■	2			■		■
<a href="#">Kekić et al. (2023)</a>	■			■				4	■				
<a href="#">Kiem et al. (2021)</a>				■	■			4			■		■
<a href="#">Li et al. (2021)</a>					■			?	■				
<a href="#">Li et al. (2022)</a>	■				■			4	■				
<a href="#">Liu et al. (2021)</a>					■		■	4					
<a href="#">Liu et al. (2022a)</a>					■			?	■	■	■	■	■
<a href="#">Liu et al. (2022b)</a>	■		■		■			2	■		■		
<a href="#">Luangasanatip et al. (2023)</a>	■				■			2	■	■			
<a href="#">Luebben et al. (2023)</a>					■			4	■				
<a href="#">Luo et al. (2022)</a>					■		■	2	■				
<a href="#">MacIntyre et al. (2022)</a>	■				■			2	■				
<a href="#">Makhoul et al. (2020)</a>	■				■			5	■		■		
<a href="#">Mandal et al. (2021)</a>					■			5	■	■			
<a href="#">Matrajt et al. (2021b)</a>	■				■			2	■				■
<a href="#">Matrajt et al. (2021a)</a>	■				■			5	■		■		
<a href="#">McBryde et al. (2021)</a>	■	■		■	■			5	■	■	■	■	■
<a href="#">Miura et al. (2021)</a>	■			■	■			5	■				
<a href="#">Molla et al. (2022)</a>					■		■	2	■	■	■		
<a href="#">Moore et al. (2021a)</a>			■		■			5	■	■	■		
<a href="#">Moore et al. (2021b)</a>				■	■			2	■		■	■	
<a href="#">Morales-Zamora et al. (2022)</a>				■	■			2	■	■		■	
<a href="#">Nuraini et al. (2021)</a>					■			4	■				
<a href="#">Pearson et al. (2021)</a>			■		■			4	■				
<a href="#">Penn and Donnelly (2023)</a>					■			?	■				■
<a href="#">Rahmandad (2022)</a>		■			■			2	■			■	■

**Table 1** continued

Reference	prioritization when minimizing				modeling framework			type vaccine roll-out	vaccine-induced reduction in				
	cases/infections	YLL	QALYS/DALYS	hospitalizations	deaths	ODE	ABM		optimal control	infection	symptomatic disease	hospitalization	death
Rao and Brandeau (2021a)	■	■	■	■	■			?	■				
Rao and Brandeau (2021b)	■							5					
Rodriguez-Maroto et al. (2023)					■			4					
Saldaña and Scoglio (2022)	■							2					
Shim (2021)	■	■			■			5					
Stafford et al. (2023)		■					■	5	■	■	■		
Tatapudi et al. (2021)	■			■	■		■	4	■				
Tran et al. (2021)	■							2					
Trejo et al. (2024)					■			4	■	■	■		
Vo et al. (2023)	■				■			4	■				
Walker et al. (2022)					■			2			■		
Wang et al. (2022)	■				■			5	■	■		■	
Yasuda et al. (2022)	■							5	■				
Zanella et al. (2021)	■							2					
Zavrakli et al. (2023)	■						■	3					
Zhao et al. (2021b)	■				■			1	■				■
Ziarelli et al. (2023)	■			■	■		■	4				■	
Zuo et al. (2022)	■				■			2	■				

187

188 The recommended dosage for some of the most effective and initially most  
 189 widely available COVID-19 vaccines, e.g., the Pfizer-BioNTech, the Moderna,  
 190 and the AstraZeneca vaccine, was two doses. While a single dose offers some  
 191 protection, two doses, spaced out at least a few weeks, induce a substan-  
 192 tially stronger protection. Thus, a related prioritization question concerns  
 193 the optimal allocation of each individual vaccine dose. If the vaccine supply  
 194 is limited, a delay of the second dose allows for more individuals to receive a  
 195 first dose. A total of 15 studies (not necessarily age-structured) investigated  
 196 this particular prioritization question. Most studies agree that a delay of  
 197 the second dose is beneficial, irrespective of the specific objective (Fig. 3B,  
 198 Table 2). This aligns with findings from a pooled analysis of four randomised

199 trials (Voysey et al., 2021). Several studies identify the relative protection  
 200 induced by the first dose compared to the full vaccine regimen as a key pa-  
 201 rameter in this decision (Romero-Brufau et al., 2021; Matrajt et al., 2021b;  
 202 Souto Ferreira et al., 2022), highlighting the need for detailed vaccine effec-  
 203 tiveness data.

Table 2: **High-level summary of COVID-19 vaccine dosing interval studies.** Each row summarizes the high-level dosing interval recommendation identified by a given study (indexed by reference number). White: not assessed, green: delay second dose, orange: shorten dosing interval, gray: recommendation depends on model assumptions. Fig. 3B provides summary counts. Only studies that were based on a mathematical model were included. The models are further classified by framework, type of vaccine roll-out (as specified in 4.4), as well as considered vaccine functions.

Reference	dosing interval recommendation when minimizing					modeling framework			type vaccine roll-out	vaccine-induced reduction in				
	cases/infections	YLL	QALYS/DALYS	hospitalizations	deaths	ODE	ABM	optimal control		infection	symptomatic disease	hospitalization	death	onward transmission
Barpounakis et al. (2022)	■				■	■			4	■				
Childs et al. (2022)	■					■			2	■		■		
Diarra et al. (2022)					■	■			5	■		■		
Ferreira et al. (2022)					■	■			4	■		■		■
Gianatti et al. (2023)					■			■	2	■				
Jimenez-Rodriguez et al. (2022)	■				■	■			2	■				
Kobayashi and Nishiura (2022)	■				■	■			2	■				
Liu et al. (2022a)					■	■			?	■				
Mak et al. (2022)	■			■	■	■			4	■				
Matrajt et al. (2021b)					■	■			2	■		■		■
Moghadas et al. (2021)	■			■	■	■			2	■		■		
Romero-Brufau et al. (2021)					■	■			2	■				
Souto Ferreira et al. (2022)					■	■		■	4	■		■		■
Tuite et al. (2021)	■				■	■			5	■		■		
Zuo et al. (2022)	■				■	■			2	■				

204

205 Countries are spatially heterogeneous. Thus, spatial factors can affect  
 206 the optimal allocation of limited vaccine. While not the primary objective of

207 this review, we identified a number of studies that investigate spatial aspects  
 208 of vaccine distribution (Table 3). The considered questions are more diverse  
 209 than in the previous two research questions; we therefore provide most details  
 210 in Subsection 5.3. In summary, the investigated studies all agree that spa-  
 211 tial factors are important when designing deaths-minimizing optimal vaccine  
 212 prioritization plans and that non-trivial trade-offs emerge, e.g. between pri-  
 213 oritizing regions with high incidence counts whose inhabitants are on average  
 214 younger and regions with more retirees. Economic factors are also taken into  
 215 consideration by multiple studies.

Table 3: **Summary of spatial COVID-19 vaccine distribution studies.** Each row describes a study that developed a spatial vaccine prioritization model. The models are classified by framework, type of vaccine roll-out (as specified in 4.4), as well as considered vaccine functions.

Reference	modeling framework			type vaccine roll-out	vaccine-induced reduction in				
	ODE	ABM	optimal control		infection	symptomatic disease	hospitalization	death	onward transmission
<a href="#">Caga-anan et al. (2023)</a>	■		■	4	■				
<a href="#">Grauer et al. (2020)</a>		■		2	■				
<a href="#">Hong et al. (2022)</a>	■			5	■				
<a href="#">Lemaitre et al. (2022)</a>			■	2	■				
<a href="#">Molla et al. (2022)</a>	■		■	2	■		■		
<a href="#">Vo et al. (2023)</a>	■			4	■				
<a href="#">Zhou et al. (2021)</a>		■		5	■				

216

#### 217 4. Key implementation details in vaccine prioritization models

218 Modelers make many decisions - some consciously, some unconsciously  
 219 - when creating a mathematical vaccine prioritization model. Some choices  
 220 can fundamentally affect the resulting optimal vaccine allocation. In this  
 221 section, we focus on the studies that identify a dependence of the optimal

222 prioritization strategy (Fig. 3) to better understand the effect of certain mod-  
223 eling assumptions as well as the impact of setting-to-setting differences in key  
224 parameters.

#### 225 *4.1. Modeling framework*

226 The choice of modeling framework may affect outcomes of vaccine pri-  
227 oritization models. In particular, as briefly described in the introduction,  
228 the popular ODE-based compartmental models come with several implicit  
229 assumptions. The homogeneous mixing assumption can be overcome by  
230 stratifying the population into sub-populations and accurately describing  
231 heterogeneous mixing (see Subsection 4.6). Another implicit assumption of  
232 ODE-based models is that the time spent in each transient compartment is  
233 exponentially distributed. This is frequently unrealistic. For example, upon  
234 infection with SARS-CoV-2 the virus needs time to replicate before a person  
235 becomes contagious. The latent period is therefore not exponentially dis-  
236 tributed [Zhao et al. \(2021a\)](#). While most included studies ignore this issue  
237 - likely since there exists no apparent direct effect on vaccine prioritizations  
238 -, some studies stratify a single transient compartment into multiple (see  
239 e.g., [Moore et al. \(2021b\)](#); [Childs et al. \(2022\)](#)). This has the effect that the  
240 total time spent in these compartments follows an Erlang distribution (as-  
241 suming equal average time in each of the multiple compartments) rather than  
242 an exponential distribution. The Erlang distribution, as a special case of the  
243 Gamma distribution, is more flexible and can thus describe more accurately  
244 the average time an individual is e.g. latently infected with SARS-CoV-2  
245 ([Lloyd, 2001](#)).

#### 246 *4.2. Prediction horizon*

247 Public health decision makers typically operate within a defined plan-  
248 ning horizon. That is, they attempt to make decisions that yield "optimal"  
249 outcomes over the course of a given time period. Similarly, mathematical  
250 models compare outcomes (e.g., total deaths or cases under different vaccine  
251 allocation strategies) over a defined time interval whose length is known as  
252 prediction horizon. The main benefit of a short prediction horizon is reduced  
253 uncertainty since long-term disease dynamics are very difficult to predict. A  
254 German ODE-based study nicely highlights that the choice of prediction hori-  
255 zon fundamentally influences who to vaccinate first ([Grundel et al., 2021](#)).  
256 If the horizon is too short (less than 8 weeks in the study), prioritization  
257 targets may switch as the strategy suffers from shortsightedness. Another

258 study shows that vaccinating elderly is always preferred for a short prediction  
259 horizon, which may however yield sub-optimal long-term outcomes ([Campos  
260 et al., 2021](#)).

#### 261 *4.3. Vaccine eligibility*

262 Given an initially limited COVID-19 vaccine supply, people with a known  
263 history of COVID-19 infection were excluded from early vaccine access by  
264 most public health agencies. Correspondingly, most reviewed mathematical  
265 models assume that only susceptible individuals can be initially vaccinated.  
266 Some ODE-based studies with more compartments (see e.g., [Islam et al.  
267 \(2021\)](#); [Karabay et al. \(2021\)](#); [Taboe et al. \(2023\)](#); [Luo et al. \(2022\)](#); [Anupong  
268 et al. \(2023\)](#); [Grundel et al. \(2021\)](#)) allow for vaccination of any individuals  
269 without known COVID-19 history. That means pre- or asymptotically  
270 infected as well as recovered individuals without known history of infections  
271 (e.g., through positive test results or symptoms) are also eligible for early  
272 vaccination, leading to some vaccine doses being used sub-optimally. A few  
273 studies even quantify the reduction in deaths, YLL, and infections that could  
274 be achieved through the hypothetical use of seroprevalence tests prior to  
275 vaccination ([Bubar et al., 2021](#); [Ayoub et al., 2021](#)). While challenging to  
276 implement, these studies find, as expected, that vaccinating only seronegative  
277 individuals always leads to improved outcomes, with the difference being  
278 larger at higher levels of seroprevalence.

#### 279 *4.4. Vaccine roll-out*

280 When deciding who to vaccine first, public health officials must antici-  
281 pate the speed of the vaccine roll-out. In mathematical models, this results  
282 in assumptions about the daily number of vaccinations. Post-hoc analyses  
283 benefit from access to historic vaccination data and can simply ask the ques-  
284 tion: Given this number of vaccinations per day, how could these vaccines  
285 have been allocated in an optimal way? When such data is unavailable, e.g.,  
286 prior to the start of a mass vaccine roll-out, modelers typically make one of  
287 the following assumptions in ODE models. Here, let  $X = X(t)$  denote the  
288 subset of the population that is eligible for vaccination (e.g., all suscepti-  
289 bles) at time  $t$ . Then, vaccination of part of these eligible individuals can be  
290 described by

$$\frac{dX(t)}{dt} = -f(X, t). \quad (1)$$



291 Note that this equation only considers the vaccination process. The size of  
292  $X$  may also change due to natural infection, immunity waning, etc. The rate  
293 of newly vaccinated,  $f(X, t)$ , typically takes one of the following forms:

- 294 1.  $f(X, t) = \nu X(t)$ , where  $\nu \geq 0$  describes the proportion of eligible indi-  
295 viduals vaccinated per unit time. This form implies that the number  
296 of vaccinations is proportional to the size of  $X$ . Specifically, as the size  
297 of  $X$  decreases over time (due to vaccination, natural infection, etc.),  
298 the number of newly vaccinated decreases as well. Mathematically, this  
299 form guarantees that  $X(t)$  remains positive for all time. This form is  
300 used in [Zhao et al. \(2021b\)](#); [Acuña-Zegarra et al. \(2021\)](#).
- 301 2.  $f(X, t) = c$ , where  $c \geq 0$  is a constant that describes the number of  
302 vaccinations per unit time. Mathematically, this form does not guar-  
303 antee positivity of  $X(t)$  for all time, as all individuals may eventually  
304 become vaccinated (or otherwise removed from  $X$ ). This necessitates  
305 careful attention when numerically solving the ODE. Nevertheless, this  
306 form is used in many models (Table 1), likely due to its simplicity.
- 307 3.  $f(X, t) = \nu(t)X(t)$ . This form is the most complex. The proportion  
308 of eligible individuals being vaccinated may vary over time. This form  
309 has the same nice mathematical property as form 1: positivity of  $X(t)$   
310 is guaranteed for all time. Contrary to form 1 and form 2, this more  
311 complex third form allows for the rate - and also the number - of vacci-  
312 nations to increase over time, as is typically the case at the beginning  
313 of a mass vaccine roll-out. Form 1 (and form 2), on the other hand,  
314 assume that the number of vaccinations decreases (remains constant,  
315 respectively) as the number of eligible individuals decreases. This form  
316 is used in a few models that employ optimal control techniques ([Acuña-  
317 Zegarra et al., 2021](#); [Zavrakli et al., 2023](#)).
- 318 4.  $f(X, t) = c(t)$ . In this form, the number of vaccinations only depends  
319 on time but not on the size of  $X$ . This form is well-suited for post-hoc  
320 analyses, in which the number of vaccinations that were conducted per  
321 unit time (e.g., day or week) is known, see e.g. [Islam et al. \(2021\)](#);  
322 [Gozzi et al. \(2022\)](#); [Luebben et al. \(2023\)](#); [Kekić et al. \(2023\)](#); [Aruffo  
323 et al. \(2022\)](#); [Ziarelli et al. \(2023\)](#); [Ferreira et al. \(2022\)](#); [Cattaneo  
324 et al. \(2022\)](#). Mathematically, this form requires careful attention when  
325 solving the ODE numerically, to ensure  $X(t)$  remains non-negative at  
326 all time. This can be achieved by adding a number of model constraints,  
327 as in [Han et al. \(2021\)](#). One study tailors an ODE model to three

328 different Indonesian provinces and optimizes the function  $c(t)$  such that  
329 active cases remain below an acceptable threshold and total vaccination  
330 cost is minimized; interestingly, the optimal function  $c(t)$  is highly non-  
331 monotonic (Nuraini et al., 2021). Another study optimizes  $c(t)$  as well,  
332 by assuming that vaccines are produced at a constant speed but that  
333 vaccine stock needs not to be used immediately (Souto Ferreira et al.,  
334 2022).

335 5. A number of studies do not specify  $f(X, t)$ . Rather, they assume that  
336 all vaccinations have been completed prior to the simulation of the  
337 disease spread. This simplifying assumption decouples the vaccine roll-  
338 out from the disease spread. A modified version of this approach is  
339 implemented in Matrajt et al. (2021b) where the simulation of disease  
340 dynamics is stopped once a week when a specified number of (weekly)  
341 vaccinations occur. A similar approach is implemented in an ABM  
342 in Jahn et al. (2021).

343 Despite different implementations of the vaccine roll-out, model-based studies  
344 generally agree that prioritization of younger, high-contact individuals may  
345 be beneficial and even lead to fewer deaths, when the entire vaccine roll-out  
346 takes place very quickly, i.e., when vaccines for a large proportion of the pop-  
347 ulation are available quickly (Matrajt et al., 2021a; Buckner et al., 2021; Liu  
348 et al., 2022b; McBryde et al., 2021). In this case, the vulnerable population  
349 is protected indirectly, by reaching herd immunity and a stop of community  
350 spread. This strategy becomes particularly reasonable in situations with low  
351 community spread (i.e., the effective reproductive number  $R_{\text{eff}} \approx 1$ ) (Chen  
352 et al., 2021; Althobaity et al., 2022; Gozzi et al., 2021) and in which the  
353 epidemic is already in decline (i.e.,  $dR_{\text{eff}}(t)/dt < 0$ ) (Molla et al., 2022). One  
354 prominent study agrees that younger individuals should only be prioritized,  
355 when minimizing deaths, if effective reproductive numbers are low but finds  
356 that a slow roll-out (and not a fast one) is an additional requirement (Bubar  
357 et al., 2021).

#### 358 4.5. Vaccine function and efficacy

359 In theory, vaccines can improve outcomes in a variety of ways. A vac-  
360 cinated individual may be less likely (than an unvaccinated individual with  
361 same characteristics) to (i) become infected, (ii) experience symptoms when  
362 infected, (iii) require hospitalization due to severe symptoms, (iv) die. In  
363 addition, a vaccinated person (v) may be less contagious (e.g., due to a lower

364 average viral load), and (vi) may have a shorter duration of infectiousness,  
 365 i.e., faster disease progression. To illustrate how these different vaccine func-  
 366 tions are frequently included in compartmental models, consider the following  
 367 COVID-19 model, which stratifies the population by disease status (suscep-  
 368 tible (S), recently infected but not yet infectious (E=exposed), symptomati-  
 369 cally infected (I), asymptotically infected (A), severely infected/requiring  
 370 hospitalization (H), deceased from COVID-19 (D), recovered (R)) and vac-  
 371 cine status (superscript v for vaccinated)):

$$\begin{aligned} \frac{dS(t)}{dt} &= -\Lambda S, \\ \frac{dS^v(t)}{dt} &= -(1 - \epsilon_1)\Lambda S^v, \\ \frac{dE(t)}{dt} &= \Lambda S - \gamma_E E, \\ \frac{dE^v(t)}{dt} &= (1 - \epsilon_1)\Lambda S^v - \gamma_E^v E^v, \\ \frac{dI(t)}{dt} &= p_{E \rightarrow I} \gamma_E E - \gamma_I I, \\ \frac{dI^v(t)}{dt} &= (1 - \epsilon_2) p_{E \rightarrow I} \gamma_E^v E^v - \gamma_I^v I, \\ \frac{dA(t)}{dt} &= (1 - p_{E \rightarrow I}) \gamma_E E - \gamma_A A, \\ \frac{dA^v(t)}{dt} &= (1 - (1 - \epsilon_2) p_{E \rightarrow I}) \gamma_E^v E^v - \gamma_A^v A, \\ \frac{dH(t)}{dt} &= p_{I \rightarrow H} \gamma_I I - \gamma_H H, \\ \frac{dH^v(t)}{dt} &= (1 - \epsilon_3) p_{I \rightarrow H} \gamma_I^v I - \gamma_H^v H, \\ \frac{dD(t)}{dt} &= p_{H \rightarrow D} \gamma_H H, \\ \frac{dD^v(t)}{dt} &= (1 - \epsilon_4) p_{H \rightarrow D} \gamma_H^v H, \\ \frac{dR(t)}{dt} &= \gamma_A A + (1 - p_{I \rightarrow H}) \gamma_I I + (1 - p_{H \rightarrow D}) \gamma_H H, \\ \frac{dR^v(t)}{dt} &= \gamma_A^v A^v + (1 - (1 - \epsilon_3) p_{I \rightarrow H}) \gamma_I^v I^v + (1 - (1 - \epsilon_4) p_{H \rightarrow D}) \gamma_H^v H^v, \end{aligned}$$

372 where  $\Lambda = \beta(A + I + \alpha A^v + \alpha I^v)$  is the force of infection, with  $\alpha \in [0, 1]$

373 describing the vaccine-induced reduction in onward transmission. Vaccine-  
374 induced faster disease progression may be implemented by  $\gamma_x^v \geq \gamma_x$  for  $x \in$   
375  $\{E, I, A, H\}$ .

376 Three important notes: First, this model implements a so-called leaky  
377 vaccine: any vaccinated individual may still become infected, at a lower rate  
378 than unvaccinated. A leaky vaccine represents the most frequent implemen-  
379 tation of vaccine function. An alternative, also frequently observed assump-  
380 tion is an all-or-nothing vaccine. In that case,  $\epsilon_1$  determines the fraction of  
381 vaccinated individuals that are completely immune to infection, while the  
382 remaining proportion of vaccinated ( $1 - \epsilon_1$ ) are typically assumed to be as  
383 susceptible as unvaccinated individuals. In some models, their susceptibility  
384 is reduced by a certain degree. Second, a stratification by age can easily  
385 be included by duplicating all compartments for each age group, including  
386 contact patterns in the force of infection, and considering age-dependent pa-  
387 rameters. Third, the parameters  $\epsilon_2, \epsilon_3, \epsilon_4$  describe conditional probabilities.  
388 For example,  $\epsilon_2$  describes the reduction in symptomatic disease among in-  
389 fected vaccinated compared to infected unvaccinated individuals. The overall  
390 vaccine-induced reduction in symptomatic disease, measured in clinical tri-  
391 als and commonly referred to as vaccine efficacy (Halloran et al., 1997), is  
392 thus  $VE_{\text{COVID}} = 1 - (1 - \epsilon_1)(1 - \epsilon_2)$ . Similarly, the overall vaccine-induced  
393 reduction in deaths is  $VE_{\text{death}} = 1 - (1 - \epsilon_1)(1 - \epsilon_2)(1 - \epsilon_3)(1 - \epsilon_4)$ .

394 Used in 90 of the 94 investigated models (Table 1), reduction in infection  
395 ( $\epsilon_1$ , implemented either as a leaky or all-or-nothing vaccine) is the most  
396 frequently considered vaccine function, followed by reduction in symptoms  
397 ( $\epsilon_2$ , used in 28 studies), reduction in severe disease ( $\epsilon_3$ , used in 22 studies),  
398 reduction in onward transmission ( $\alpha$ , used in 17 studies), and reduction in  
399 death ( $\epsilon_4$ , used in 15 studies). Other vaccine functions considered in only a  
400 few models include a shorter period of infectiousness (Makhoul et al., 2020;  
401 Penn and Donnelly, 2023), as well as a reduced vaccine efficacy for older  
402 individuals (Bubar et al., 2021; Aruffo et al., 2022; Buckner et al., 2021) and  
403 children (Han et al., 2021). 46 out of 94 studies (48.9%) considered only  
404 one type of vaccine function, while three studies (Liu et al., 2022a; McBryde  
405 et al., 2021; Mak et al., 2022) differentiated five types (Table 1).

406 The range of parameter values considered for a given vaccine function  
407 also varied wildly. One study investigated optimal prioritization strategies  
408 for mass vaccinations with commonly used vaccines that had shown some  
409 beneficial heterologous effects against SARS-CoV-2 infection (Hupert et al.,  
410 2022). This study considered  $\epsilon_1, \epsilon_2, \epsilon_4 \in [5\%, 15\%]$ . In line with results from

411 COVID-19 vaccine clinical trials, most studies assumed relatively high levels  
412 of vaccine efficacy against symptomatic disease,  $VE_{\text{COVID}}$ , with the specific  
413 values varying based on vaccine product, number of doses and predominant  
414 virus variant. Moreover, the relative contribution of  $\epsilon_1$  and  $\epsilon_2$  differs, with  
415 some studies (see e.g., [Islam et al. \(2021\)](#); [Matrajt et al. \(2021b,a\)](#); [Han  
416 et al. \(2021\)](#); [Ayoub et al. \(2021\)](#); [Makhoul et al. \(2020\)](#); [Choi and Shim  
417 \(2021\)](#); [Hogan et al. \(2021\)](#); [Moore et al. \(2021a\)](#); [Liu et al. \(2022b\)](#); [Jahn  
418 et al. \(2021\)](#); [Kiem et al. \(2021\)](#)) contrasting optimal vaccination strategies  
419 for both extreme cases:  $\epsilon_1 = VE_{\text{COVID}}, \epsilon_2 = 0$  (sterilizing vaccine), and  
420  $\epsilon_1 = 0, \epsilon_2 = VE_{\text{COVID}}$  (non-sterilizing vaccine). These studies agree that  
421 at a fixed (overall) vaccine efficacy against symptomatic disease, higher  $\epsilon_1$   
422 (i.e., lower  $\epsilon_2$ ) leads to better outcomes. The higher  $\epsilon_2$  relative to  $\epsilon_1$ , the  
423 more important is the prioritization of older and vulnerable people when  
424 optimizing morbidity-based metrics ([Islam et al., 2021](#); [Choi and Shim, 2021](#);  
425 [Liu et al., 2022b](#); [Kiem et al., 2021](#)). Although hard to disentangle in practice,  
426 it is therefore important for optimal prioritization design to understand the  
427 relative contribution of  $\epsilon_1$  and  $\epsilon_2$  to the vaccine efficacies observed in clinical  
428 trials.

429 Studies which differentiate between single-dose and "fully" vaccinated in-  
430 dividuals include two parameters for each vaccine function. One study shows  
431 that, for a fixed  $VE_{\text{COVID}}$ , delaying second doses and thus covering a larger  
432 part of the population with first doses becomes more important at higher  $\epsilon_1$   
433 (i.e., lower  $\epsilon_2$ ) when minimizing mortality ([Matrajt et al., 2021b](#)). Another  
434 important factor is the speed of the vaccine roll-out. One study shows that  
435 a generally delayed second dose only leads to fewer deaths if the roll-out is  
436 slow ([Romero-Brufau et al., 2021](#)). An age-dependent strategy (providing  
437 two doses to people 65 and older but delaying the second dose for younger  
438 people) performs consistently well, irrespective of the speed of the roll-out.  
439 The most important parameter in determining whether a delay of second  
440 doses is beneficial is, however, the relative difference in the reduction in sus-  
441 ceptibility after one dose versus two doses. As expected, all studies agree  
442 that a delay becomes more beneficial the smaller the difference, irrespective  
443 of the optimization objective ([Romero-Brufau et al., 2021](#); [Moghadas et al.,  
444 2021](#); [Mak et al., 2022](#); [Souto Ferreira et al., 2022](#); [Matrajt et al., 2021b](#);  
445 [Childs et al., 2022](#); [Gonzalez-Parra, 2021](#); [Tuite et al., 2021](#)). One notewor-  
446 thy ABM-based study uses differential vaccine function parameters for the  
447 Moderna and Pfizer-BioNTech vaccine, and finds that to minimize cases the  
448 second Moderna dose should be delayed while a delay of the second Pfizer-

449 BioNTech dose may be detrimental if pre-existing immunity is low and if  
450 single dose-induced immunity wanes (Moghadas et al., 2021). To minimize  
451 deaths or hospitalizations, this study suggests delayed second doses, irrespec-  
452 tive of the type of vaccine.

453 The various studies differ in how the two-dose vaccination campaign is im-  
454 plemented. In compartment-based models, separate compartments for single-  
455 dose ( $V_1$ ) and “fully” vaccinated ( $V_2$ ) individuals are used. One common im-  
456 plementation employs two rates  $\nu_1(t), \nu_2(t) \geq 0$  to describe the proportion of  
457 susceptible and single-dose vaccinated that receive a vaccine dose on a given  
458 day  $t$  (see e.g., Zhao et al. (2021b); Childs et al. (2022); Liu et al. (2022a)).  
459 That is,

$$\begin{aligned}\frac{dS(t)}{dt} &= -\nu_1(t)S(t), \\ \frac{dV_1(t)}{dt} &= \nu_1(t)S(t) - \nu_2(t)V_1(t), \\ \frac{dV_2(t)}{dt} &= \nu_2(t)V_1(t),\end{aligned}$$

460 with constraints on  $\nu_1(t)$  and  $\nu_2(t)$  ensuring that only available vaccines are  
461 used. Other implementations include delay differential equations (Souto Fer-  
462 reira et al., 2022; Sepulveda et al., 2023) or weekly pulse vaccinations (Ma-  
463 trajt et al., 2021b).

#### 464 4.6. Transmission rates and heterogeneous contact patterns

465 The rate at which susceptible individuals acquire an infection, the force  
466 of infection, depends, among others, on contact rates, the community in-  
467 cidence and the infectivity of the virus. It is well-established that human  
468 interactions are age-assortative and that older individuals have on average  
469 fewer contacts (Mossong et al., 2008). A realistic account for age-specific  
470 mixing patterns is thus of paramount importance in infectious disease mod-  
471 els that guide policy-makers to prioritize either high-contact young people  
472 or lower-contact older people. A common approach to model infection of  
473 sub-population  $i, i = 1, \dots, n$  in an age-structured ODE is

$$\frac{dS_i(t)}{dt} = -\beta_i \sum_{j=1}^n C_{ij}(aA_j + I_j)S_i,$$

474 where

- 475 •  $a \geq 0$  represents the relative contagiousness of asymptomatic ( $A$ ) com-  
476 pared to symptomatic ( $I$ ) individuals. All investigated studies chose  
477  $a \in (0, 1]$ .
- 478 • The transmission rate  $\beta_i$  can account for age-dependent susceptibility  
479 and risk mitigation (e.g., mask wearing). Multiple studies assumed  
480 that older, more vulnerable individuals suffer from higher susceptibil-  
481 ity (Davies et al., 2020; Moore et al., 2021b; Jahn et al., 2021) but also  
482 engage in more risk mitigation measures (Masters et al., 2020; Kadelka  
483 and McCombs, 2021; Bushaj et al., 2023; Vo et al., 2023). The trans-  
484 mission rate may also vary over time, e.g., due to the emergence of more  
485 transmissible SARS-CoV-2 variants (Islam et al., 2021; Moore et al.,  
486 2021b), or time-varying social distancing levels (Moore et al., 2021b).  
487 It may further vary from location to location in spatially distributed  
488 models (Vo et al., 2023).
- 489 • The  $n \times n$ -matrix  $C$  describes the average number of contacts an in-  
490 dividual in sub-population  $i$  has with individuals from sub-population  
491  $j$ . Just like  $\beta_i$ , this matrix may also vary over time and by location to  
492 account for periods of school closures, work-from-home orders, etc. A  
493 reduction in activity levels of sub-population  $i$  (e.g., due to NPI adher-  
494 ence) can be implemented in two ways: (i) through a reduction in  $\beta_i$ ,  
495 or (ii) through a proportional reduction of row and column  $i$  of contact  
496 matrix  $C$ . It is very important to understand the differential effect of  
497 these choices on the model. Only the latter choice reduces both new  
498 infections of sub-population  $i$  and onward transmission by members of  
499 sub-population  $i$ . For this reason, this choice should be preferred in  
500 infectious disease models.

501 The seminal, diary-based POLYMOD study surveyed roughly one thou-  
502 sand individuals each in eight European countries and established country-  
503 specific contact matrices for a population stratified into 15 age groups (0 –  
504 4, 5 – 9, . . . , 75 – 79, 80+) (Mossong et al., 2008). Contact rates were fur-  
505 ther stratified by location (home, workplace, school, other). By combining  
506 this data with various other data sources, age-and-location-specific synthetic  
507 contact rates were obtained for 177 countries, and it was shown that syn-  
508 thetic and empirical contact matrices employed in epidemiological models  
509 yield similar findings (Prem et al., 2017, 2021).



510 Most investigated mathematical models use a country-specific contact  
511 matrix. One ODE-based study shows explicitly how the optimal prioritization  
512 strategy depends on the country-specific age pyramid and contact  
513 matrix: to minimize deaths, it is optimal to first vaccinate the oldest people  
514 in India and Italy but middle-aged people in China (Wang et al., 2022). An-  
515 other study quantifies the inter-generational mixing, which is typically lower  
516 in high-income countries (Gozzi et al., 2021). Since the population in high-  
517 income countries is on average also older, non-trivial dependencies arise when  
518 designing morbidity- or mortality-minimizing vaccine allocation strategies.

519 One commonly stated limitation of these contact matrices is that they  
520 have been derived before the COVID-19 pandemic, during which relative  
521 mixing patterns may have shifted. Two general approaches have been used  
522 to overcome this issue. First, several studies recomputed the overall contact  
523 matrix as a linear combination of the four location-specific pre-pandemic  
524 contact matrices (Matrajt et al., 2021b; Foy et al., 2021; Kiem et al., 2021;  
525 Chen et al., 2021; Moore et al., 2021b). That is,

$$C(t) = a_1(t)C_{\text{home}} + a_2(t)C_{\text{work}} + a_3(t)C_{\text{school}} + a_4(t)C_{\text{other}}.$$

526 All studies agree that during a pandemic,  $a_1(t) \geq 1$  while  $0 \leq a_2(t), a_3(t), a_4(t) <$   
527 1. Cell phone mobility data (Jentsch et al., 2021; Foy et al., 2021) as well  
528 as specific policy implementations (school closures, work-from-home orders,  
529 etc) (Jentsch et al., 2021; Karabay et al., 2021) have been used to inform  
530 the weights  $a_1(t), \dots, a_4(t)$ . The Oxford Stringency Indices are based on  
531 information on implemented government policies related to closure and con-  
532 tainment, health and economic policy, and provide means to quantify the  
533 time-varying level of NPIs in 180 countries - for several countries, even at  
534 the level of individual jurisdictions (Hale et al., 2021). Second, empirical con-  
535 tact matrices have been derived during the pandemic in several settings, and  
536 are used by a variety of studies, see e.g. (Han et al., 2021; Tran et al., 2021;  
537 Zhao et al., 2021b). Comparisons of Chinese as well as Belgian diary-based  
538 pre-pandemic contact matrices with contact matrices obtained during and  
539 after the first COVID-19 lockdowns revealed not only drastic changes in the  
540 number of contacts but also in the mixing patterns (Zhang et al., 2020, 2021;  
541 Coletti et al., 2020). Contact matrices during a pandemic - specifically dur-  
542 ing periods of strong NPI adherence, e.g., a lockdown - exhibit much lower  
543 levels of age-assortativity, likely due to school and work closures.

544 While age-assortativity decreases during a pandemic, assortative mixing  
545 with respect to other attributes, also known as homophily (McPherson et al.,

546 2001), may be high or even increase, and may profoundly influence disease  
547 dynamics. High levels of homophily with respect to COVID-19 vaccine status  
548 have been reported (Are et al., 2024). Both network- and ODE-based studies  
549 show that this may lead, at a fixed level of vaccine coverage, to more frequent  
550 outbreaks and higher attack rates (Kadelka and McCombs, 2021; Hiraoka  
551 et al., 2022; Burgio et al., 2022). Another study, employing a novel approach  
552 to include homophily with respect to binary attributes in established contact  
553 matrices (Kadelka, 2023), shows that accounting for high levels of ethnic  
554 homophily in the United States, coupled with proportionately more people  
555 of color working in high-contact jobs but fewer being of old age, leads to  
556 non-trivial trade-offs in optimal vaccine prioritization design (Kadelka et al.,  
557 2022).

558 A less frequently mentioned limitation of contact matrices is that they  
559 are by default non-reciprocal. In empirical contact matrices, elderly tend to  
560 more frequently report a brief contact (Mossong et al., 2008), and generally  
561 provide less reliable responses in surveys Perry (1982). Physical contacts,  
562 which are required for COVID-19 spread, are however reciprocal. That is,

$$N_i C_{ij} = N_j C_{ji}$$

563 should hold for all  $i, j = 1, \dots, n$ , where  $N_i$  is the size of sub-population  $i$ ;  
564 otherwise, disease dynamics are inaccurate. A common procedure to generate  
565 a reciprocal contact matrix is outlined in (Funk et al., 2019; Kadelka, 2023).  
566 Several studies employ this or a similar procedure (see e.g. Matrajt et al.  
567 (2021b); Kadelka et al. (2022); Islam et al. (2021)).

#### 568 4.7. Behavioral responses

569 The force of infection depends proportionately on the contact level. Over  
570 the course of a pandemic, implemented government policies, adherence to  
571 NPIs (e.g., social distancing), and thus contact levels differ. Variations in  
572 contact levels that affect the entire population homogeneously can be easily  
573 implemented by multiplying contact matrices with a time-varying factor, as  
574 described above. Heterogeneous behavioral responses, reported in several  
575 surveys (see e.g. Masters et al. (2020); Dryhurst et al. (2022); Pasion et al.  
576 (2020)), are often much harder to model but crucial to accurately predict,  
577 for example, the effect of a specific vaccine prioritization strategy. A variety  
578 of studies included aspects of homogeneous and heterogeneous behavioral  
579 responses.

580 Several studies assume that the population-wide contact level depends on  
581 the number of recent infections, hospitalizations and/or deaths ([Rahmandad, 2022](#);  
582 [Althobaity et al., 2022](#); [Gozzi et al., 2021](#)), the number of currently ac-  
583 tive cases ([Jentsch et al., 2021](#); [Islam et al., 2021](#)), recent changes in these  
584 numbers, or a combination of these factors. For example in [Islam et al. \(2021\)](#),  
585 the population-wide contact level has been modeled to depend on  
586 the number of active cases, using a sigmoidal function. Another study, ex-  
587 panding the ODE model by Bubar et al ([Bubar et al., 2021](#)), assumes that  
588 the contact reduction depends exponentially on the number of deaths re-  
589 ported a few days ago ([Rahmandad, 2022](#)). The authors argue that inclusion  
590 of this endogenous behavioral feedback loop provides a better model fit to  
591 data. This study further assumes that the level of contact reduction can  
592 depend on age/perceived risk, and specifically, that vaccinated individuals  
593 may engage in less contact reduction. This complicates the trade-off in vac-  
594 cine prioritization when minimizing deaths or YLL: vaccinate high-contact,  
595 less NPI-compliant individuals first or more compliant people at higher risk.  
596 This study concludes that the answer primarily depends on the speed of the  
597 vaccine roll-out, as well as the differences in NPI compliance. Several other  
598 studies come to the same or similar conclusion ([Bushaj et al., 2023](#)). An  
599 ABM-based study employs a binary stratification of NPI compliance, and  
600 quantifies how much lower the compliance level of low-risk (versus high-risk)  
601 individuals must be for them to be the optimal prioritization target ([Kadelka  
602 and McCombs, 2021](#)). It finds that the switching point depends on the vac-  
603 cine efficacy, as well as the level of homophily with respect to vaccine status  
604 and NPI adherence. An ODE-based model, first proposed in [Gozzi et al.  
605 \(2021\)](#) and then adapted to several Arab countries in [Althobaity et al. \(2022\)](#),  
606 uses the same binary classification and explicitly models dynamic shifts be-  
607 tween these two sub-populations that depend on the vaccine coverage and  
608 the number of recent deaths. These studies find that, at  $R_{\text{eff}} = 1.15$ , elderly  
609 should be prioritized when minimizing deaths as long as the vaccine roll-out  
610 is sufficiently fast; the speed at which the optimal prioritization switches  
611 is country-dependent. Some studies have been even shown that a vaccine  
612 with low effectiveness may be detrimental and yield worse outcomes than in  
613 the absence of a vaccine, due to behavior adaptation and a false belief of  
614 protection ([Kadelka and McCombs, 2021](#); [Luebben et al., 2023](#)).

#### 615 4.8. *Vaccine hesitancy*

616 Despite the availability of highly effective vaccines, a sizeable proportion  
617 of people refuses to get vaccinated against COVID-19. When determining  
618 the optimal roll-out of a limited vaccine, this factor must be taken into  
619 account, either explicitly by certain modeling assumptions or implicitly by  
620 only considering those roll-out solutions that appear feasible given levels of  
621 vaccine hesitancy. A number of survey-based studies have accessed these lev-  
622 els in different countries and at different times during the pandemic ([Sallam,](#)  
623 [2021](#); [Soares et al., 2021](#)). While vaccine hesitancy differs from country to  
624 country, it generally differs more with age. As expected, older individuals  
625 who are more at risk are also more willing to be vaccinated.

626 Many infectious disease models explicitly include vaccine hesitancy by  
627 assuming that a proportion of each sub-population cannot be vaccinated.  
628 Some models simply assume that this proportion is fixed ([Bubar et al., 2021](#);  
629 [Gavish and Katriel, 2022](#); [Islam et al., 2021](#); [Li et al., 2022](#); [Luebben et al.,](#)  
630 [2023](#); [Kadelka et al., 2022](#); [Makhoul et al., 2020](#); [Miura et al., 2021](#); [Moore](#)  
631 [et al., 2021a](#); [Rodriguez-Maroto et al., 2023](#); [Tatapudi et al., 2021](#); [Walker](#)  
632 [et al., 2022](#); [Hogan et al., 2021](#); [McBryde et al., 2021](#); [Rahmandad, 2022](#)),  
633 while others account for lower hesitancy among older individuals ([Liu et al.,](#)  
634 [2022a](#); [Moghadas et al., 2021](#); [Han et al., 2021](#); [Liu et al., 2022b](#); [Moore](#)  
635 [et al., 2021b](#); [Zavrakli et al., 2023](#)). Most models in the latter category  
636 differentiate hesitancy using a binary age threshold. A Greek model infers  
637 age-specific values from a telephone survey ([Sypsa et al., 2022](#); [Barmounakis](#)  
638 [et al., 2022](#)). One study assumes that the level of hesitancy evolves over time  
639 following an ODE formulation ([Jentsch et al., 2021](#)). An ABM-based study  
640 shows that COVID-19 outbreaks are more frequent, at a given level of vaccine  
641 coverage and NPI adherence, if those who comply with NPIs are also those  
642 who get vaccinated, as is frequently the case ([Kadelka and McCombs, 2021](#)).

#### 643 4.9. *Effective reproductive number*

644 While the basic reproductive number describes the number of secondary  
645 infections caused on average by an infected individual in a fully suscepti-  
646 ble population, the effective reproductive number varies over time and takes  
647 into account population-wide levels of immunity, NPI adherence, emergence  
648 of more infectious variants, etc. Many models have investigated how opti-  
649 mal prioritization schemes depend on this key epidemiological parameter. As  
650 described above, prioritization of high-contact, low-risk younger individuals

651 becomes a reasonable choice, when minimizing deaths, if the effective repro-  
652 ductive number is close to 1 (Chen et al., 2021; Althobaity et al., 2022; Gozzi  
653 et al., 2021; Bubar et al., 2021), especially if it is decreasing (Molla et al.,  
654 2022). A French study further clarifies that young individuals should only  
655 be prioritized if the vaccine acts almost entirely by reducing infections (that  
656 is, if  $\epsilon_1 \gg \epsilon_2$ ) (Kiem et al., 2021). In these circumstances, the vulnerable,  
657 older part of the population is indirectly protected by the vaccine through  
658 herd immunity, pending no changes in NPI adherence.

#### 659 *4.10. Variant Considerations*

660 Like most RNA viruses, SARS-CoV-2 evolves rapidly (Markov et al.,  
661 2023). Over the course of the last four years, a plethora of variants has  
662 emerged. These variants exhibit different phenotypes characterized e.g. by  
663 transmissibility, severity of disease and immune evasion. The vaccine roll-out  
664 happened in parallel to the emergence of SARS-CoV-2 variants. For exam-  
665 ple, the United States started to vaccinate individuals in December 2020. By  
666 April 2021 - when weekly vaccination counts still increased - Alpha (B.1.1.7),  
667 which was an estimated 50% more transmissible, had become the dominat-  
668 ing virus strain, followed soon after by Delta (B.1.617.2), which was even  
669 more contagious and caused also more hospitalizations and deaths (Camp-  
670 bell et al., 2021). Therefore, mathematical models used to predict optimal  
671 vaccine allocation strategies should consider the emergence of variants. In  
672 practice, the time to emergence of a new variant and its specific phenotypic  
673 characteristics can, however, not be reliably predicted. Some studies used  
674 genomic SARS-CoV-2 surveillance time-series and estimates of the pheno-  
675 typic characteristics of circulating variants to predict the future distribution  
676 of circulating virus strains (Islam et al., 2021; Childs et al., 2022). This dis-  
677 tribution can yield estimates of the time-varying transmissibility and various  
678 transition probabilities, even when the model does not account explicitly for  
679 variant-specific infections (Liu et al., 2022a). A few studies went further  
680 and even included different compartments for those infected with e.g. the  
681 wildtype and Alpha variant (Aruffo et al., 2022; Gozzi et al., 2022). Qualita-  
682 tively, accounting for the emergence of more infectious variants yields higher  
683 effective reproductive numbers (pending no changes in NPI adherence, etc)  
684 with effects as described in Subsection 4.9.

## 685 5. Summary of selected modeling studies

686 In the previous sections, we focused on key modeling assumptions, both  
687 explicit and implicit ones, and their impact on the outcomes of vaccine pri-  
688 oritization models. We highlighted different studies wherever suitable. In  
689 this section, we provide a brief summary of selected modeling studies that  
690 present interesting features. Only some features of the models and findings  
691 from the studies can be described. Subsection 5.1 summarizes studies that  
692 answer the question: How should a limited vaccine be optimally distributed  
693 among a population stratified by age? Subsection 5.2 summarizes studies  
694 that answer the question: Should the second COVID-19 vaccine dose be de-  
695 layed given limited vaccine availability? Subsection 5.3 summarizes spatial  
696 vaccine distribution studies.

### 697 5.1. Summary of studies that employ an age-structured mathematical model

#### 698 5.1.1. Differential equation and difference equation based models

699 In [Bubar et al. \(2021\)](#), the authors compared five different vaccine priori-  
700 zation strategies using an age-stratified ODE-based SEIR model. Outcomes  
701 were assessed using the number of infections, deaths and YLL. The authors  
702 found that prioritizing adults aged 20 to 49 years minimized infections at all  
703 considered values of the effective reproduction number (1.1-2). Furthermore,  
704 this same prioritization provided the best way to reduce mortality and YLL  
705 when the effective reproduction number is low ( $\leq 1.15$ ) and if the vaccine  
706 roll-out is slow. This study highlights the importance of the transmissibility  
707 of SARS-CoV-2 and the pace of vaccine roll-out on the choice of an opti-  
708 mal vaccination strategy. The authors also consider the potential benefit of  
709 seroprevalence tests prior to vaccination.

710 In [Moore et al. \(2021b\)](#), an ODE-based SEAIHR model was fitted to data  
711 from the United Kingdom. The authors show that even a vaccine as effective  
712 as those by Pfizer-BioNTech and Oxford-AstraZeneca would not suffice to  
713 contain the COVID-19 outbreak, partially due to age-varying vaccine hesi-  
714 tancy. The study further highlights that the number of deaths that appear  
715 among vaccinated will naturally increase as vaccine coverage increases. The  
716 model accounts for time-varying population-wide social distancing levels as  
717 well as the emergence of more transmissible variants.

718 In [Buckner et al. \(2021\)](#), the authors investigated optimal vaccination  
719 strategies by using an ODE-based SEPIAR model that takes into account  
720 essential workers. Stochastic nonlinear programming techniques were used



721 to find the vaccine prioritization. Assuming  $\mathcal{R}_{\text{eff}} = 2.5$ , vaccines were as-  
722 signed only to susceptible individuals and updates to the prioritization were  
723 made each month. Outcomes were assessed using the number of infections,  
724 deaths and YLL. The authors found that to minimize infections, it is opti-  
725 mal to prioritize older essential workers. However, depending on the objective  
726 and alternative model scenarios considered, younger essential workers may  
727 be prioritized to control SARS-CoV-2 spread or elderly to directly control  
728 mortality. A combination of a genetic algorithm (global) and a simulated an-  
729 nealing algorithm (local) was used to obtain the optimal vaccination strategy  
730 each month.

731 In [Foy et al. \(2021\)](#), the authors employed an age-structured ODE-based  
732 SEIARQ ( $Q$  = quarantined, not spreading) model to inform the optimal vac-  
733 cine roll-out in India. Assuming  $\mathcal{R}_{\text{eff}} = 2.4$ , four vaccine prioritizations were  
734 compared: even across the population, prioritize 20–40 year olds, 40–60 year  
735 olds, or those 60 and older. To minimize deaths, the authors found that pri-  
736 oritizing the oldest is optimal regardless of the vaccine efficacy, control mea-  
737 sures, vaccination pace, or immunity dynamics. However, this prioritization  
738 results in more symptomatic infections. To minimize infections, vaccination  
739 of 20-40 year olds should be prioritized. A faster vaccine roll-out reduces the  
740 differences between the compared vaccine prioritizations.

741 In [MacIntyre et al. \(2022\)](#), the authors employed an ODE-based SEPA-  
742 IQR model that was extended to include several additional classes such as  
743 traced, undiagnosed and highly infectious. The model was fitted to the Aus-  
744 tralian state New South Wales and age-targeted and ring vaccination pro-  
745 grams were compared. The population was stratified by occupation (health-  
746 care workers) and age. The authors found that vaccinating older people  
747 prevents more deaths and that herd immunity can only be reached by mass  
748 vaccination campaigns, and only if the vaccine is sufficiently effective and  
749 rolled out sufficiently fast.

750 In [Hogan et al. \(2021\)](#), an extended SEIR discrete-time model was used to  
751 evaluate the public health impact of vaccines using data from different coun-  
752 tries. The model uses a class of individuals with a mild infection that includes  
753 both symptomatic and asymptomatic but that does not require hospitaliza-  
754 tion. The authors identified death-minimizing vaccine allocation strategies  
755 within- and between-countries. They found if less than 20% vaccine coverage  
756 is available, it is better to prioritize the elderly. However, in less limited  
757 settings, high transmitters should be prioritized.

758 In [Moore et al. \(2021a\)](#), an ODE-based SEAIR model was used to in-



759 vestigate optimal vaccine allocations in the UK. Outcomes were assessed by  
760 deaths and loss in QALYs. For a range of model assumptions, the authors  
761 found elderly should be prioritized. However for vaccines that have low effi-  
762 cacy among the elderly ( $< 20\%$ ), other prioritizations proved more effective.

763 In Shim (2021), an ODE-based SEIAR model was calibrated South Ko-  
764 rea. The authors found that to minimize deaths (infections) older (younger)  
765 individuals should be prioritized. Interestingly, the YLL-minimizing strat-  
766 egy is sensitive to vaccine efficacy and the number of vaccine doses available.  
767 When vaccine efficacy (assuming a vaccine that only reduces infections) is  
768 relatively low ( $\leq 30\%$ ) groups with high case-fatality rates should be pri-  
769 oritized, thereby maximizing the direct benefit of vaccines. However, with  
770 vaccines that have higher efficacy, prioritization shifts toward younger age  
771 groups: 40–69 year olds at 50–70% efficacy or 30–59 year olds at 90% effi-  
772 cacy.

773 In Islam et al. (2021), a detailed ODE-based model was calibrated to  
774 evaluate the U.S. vaccine roll-out. The population was stratified by age,  
775 comorbidity status, job type and living situation. The model also accounts for  
776 time-varying population-wide social distancing levels as well as the emergence  
777 of more transmissible variants. The authors compared 17.5 million 4-phased  
778 vaccine allocation strategies and found that a strategy that prioritizes people  
779 with comorbidities in all age groups is Pareto-optimal, yielding slightly fewer  
780 deaths, infections and YLL than the strategy recommended by the Centers  
781 for Disease Control (CDC).

782 In Penn and Donnelly (2023), an ODE-based SIR model was used to  
783 study the effect of the basic reproduction number  $\mathcal{R}_0$  on the optimal vacci-  
784 nation plan. An interesting counter-intuitive result was found: It is better  
785 to prioritize 45–49 year olds than 55–59 year olds despite higher case fatality  
786 rates in the latter group. The authors explained this by the fact that the  
787 latter group has much fewer contacts with those 75 and older (as parents of  
788 those 55-59 years old have already died to a large degree). Thus, priorit-  
789 ization of 45–49 year olds substantially increases the secondary protection of  
790 the elderly. This result shows the importance of the age-stratified contact  
791 matrices.

792 In Zhao et al. (2021b), three different ODE-based SEIAR models were  
793 used to find the optimal vaccination strategy against COVID-19 in Wuhan  
794 City, China. The authors used the effective reproduction number to estimate  
795 the SARS-CoV-2 transmission between age groups. They found that, before  
796 NPIs were implemented, the highest transmissibility existed among those

797 15-44 years old. In order to control transmission, this age group should be  
798 prioritized. To minimize deaths, those  $\geq 65$  years old should be prioritized,  
799 irrespective of their lower contact rates.

800 In [Matrajt et al. \(2021a\)](#), an ODE-based model with many compartments  
801 was used to determine which age group(s) should be vaccinated assuming in-  
802 stantaneous vaccination and 10-100% vaccine coverage. The authors studied  
803 many scenarios and found that for low vaccine effectiveness (10-50%), regard-  
804 less of vaccination coverage, it is optimal to prioritize elderly people when  
805 minimizing deaths. However, for higher vaccine effectiveness and if the basic  
806 reproductive number is low, it is better to prioritize younger people, espe-  
807 cially if available vaccination coverage is  $\geq 40\%$ . The optimization routine  
808 includes a coarse global search algorithm, coupled with a fast optimizer, to  
809 explore the entire space of possible combinations of vaccine allocations.

810 In [Kadelka et al. \(2022\)](#), an age-and-ethnicity-stratified ODE-based model  
811 was used to study the optimal distribution of available vaccines in the United  
812 States to two different groups: White and Asian persons and all others. Dif-  
813 ferent levels of ethnic homophily were considered. The authors found that  
814 vaccine allocations that stratify vaccine access by ethnicity could have pre-  
815 vented a number of deaths, especially assuming high levels of ethnic ho-  
816 mophily. Moreover, the authors highlight a second trade-off when designing  
817 mortality-minimizing vaccination plans and accounting for ethnic homophily:  
818 the elderly population is predominantly White and Asian, while those em-  
819 ployed in high-contact occupations are predominantly from the other ethnic  
820 groups.

821 In [Stafford et al. \(2023\)](#), an age-and-race-stratified ODE-based model was  
822 used to study the distribution of available vaccines in the United States to  
823 two different groups: non-Hispanic White persons and all others. Several  
824 objective functions that include mortality, YLLs, measures of inequity and  
825 joint disease burden were considered. The authors found that there exists  
826 a trade-off between minimizing disease burden and minimizing inequity, es-  
827 pecially if vaccine is very limited (e.g., 10%). If vaccine coverage is  $\geq 30\%$ ,  
828 both inequity and mortality can be optimized at the same time.

829 In [Zuo et al. \(2022\)](#), an ODE-based SEIQR model was used, combined  
830 with google mobility data to modify contact matrices. This study highlights  
831 that the optimal vaccine prioritization depends on particular parameters re-  
832 lated to the transmission rates. Assuming fixed daily doses, the authors  
833 found that in a scenario with low infection rate and low vaccine availability,  
834 vaccinating first people over 60 minimizes deaths, but that with more vaccine

835 availability vaccinating first those 51-60 year old is preferable due to their  
836 higher contacts.

837 In [Gavish and Katriel \(2022\)](#), an ODE-based model is used to investigate  
838 whether children should have been vaccinated earlier. The authors found that  
839 prioritization strategies that include vaccination of children lead to Pareto-  
840 optimal outcomes regarding minimizing deaths and infections, especially if  
841 the basic reproductive number is high.

842 In [Rao and Brandeau \(2021b\)](#), an ODE-based SIR model with two age  
843 groups (with age threshold 65) was used to study which vaccine allocation  
844 minimizes the effective reproduction number. Assuming that all vaccinations  
845 take place at once, the authors found that the answer depends on available  
846 vaccine coverage, vaccination pace and the initial effective reproduction num-  
847 ber. In [Rao and Brandeau \(2021a\)](#), the same authors used the model to  
848 minimize infections, deaths, YLL and loss in QALY. They found that it is  
849 better to prioritize the young group to minimize infections, but the older in-  
850 dividuals for all other metrics. This result was obtained by simple analytical  
851 conditions that describe the optimal vaccine allocation for each objective.

852 In [Rahmandad \(2022\)](#), an ODE-based SEIR model was used to study the  
853 effects of behavioral responses to risk by means of an endogenous feedback  
854 loop. Specifically, the author assumed that population-wide social distancing  
855 levels fluctuate depending on the recently reported numbers of COVID-19  
856 deaths. The author argues that high-contact individuals should be prioritized  
857 to minimize deaths or YLLs, as long as the vaccine roll-out happens fast  
858 enough. This is because the vulnerable population is already more risk-averse  
859 and thus engages in more risk mitigation.

860 In [Han et al. \(2021\)](#), an ODE-based SIR model was used to study optimal  
861 vaccine prioritization plans in China. The authors show that a time-varying  
862 vaccination program (i.e., allocating vaccines to different target groups as  
863 the epidemic evolves) can yield much better outcomes since it is capable  
864 to simultaneously achieve different objectives (e.g., minimizing deaths and  
865 infections). In addition, a high vaccination pace in the early phase of the  
866 vaccination plan is better. In a sensitivity analysis, the authors employed a  
867 contact matrix derived from contact diaries collected in Shanghai in March  
868 2020, at a time when the lockdown was over, but severe NPIs were still in  
869 place. The “pandemic” contact matrix exhibits much less age-assortativity.

870 In [Makhoul et al. \(2020\)](#), an ODE-based model was used and the authors  
871 found that a vaccine with efficacy against infection  $\geq 70\%$  would eliminate  
872 COVID-19. Outcomes were assessed over the course of ten years and the

873 authors assumed full vaccine protection over this time course, which appears  
874 too high retrospectively. The authors studied two vaccination programs:  
875 80% coverage before the onset of the epidemic and 80% coverage within one  
876 month of the onset of the epidemic.

877 In [Campos et al. \(2021\)](#), an ODE-based SIR model was used to predict  
878 the COVID-19 dynamics and compare with out-of-sample data from Rio de  
879 Janeiro. In addition, numerical simulations were used to compare age-based  
880 vaccine allocation strategies policies. Three age groups of similar size were  
881 considered as vaccination targets. In in all the tested scenarios, prioritization  
882 should be given to either those 15-34 or 50 year and older. The optimal choice  
883 depends on the evaluation time period, vaccination schedules and efficacy of  
884 the vaccine.

885 In [Angelov et al. \(2023\)](#), a non-standard age-structured ODE-based model  
886 was proposed that differentiates between isolated and non-isolated as well as  
887 symptomatically and asymptotically infected. The model further takes  
888 into account the heterogeneity of the infected sub-population with respect to  
889 the time since infection. Solving an optimal control problem, which considers  
890 as one of the constraints the hospital capacity, the authors found that deaths  
891 in Austria are minimized if those 18-30 years old (highest transmitters) are  
892 vaccinated first, followed by those 80 and older (most at risk), followed by  
893 other age groups.

894 In [Babus et al. \(2023\)](#), a linear programming problem is solved in order  
895 to find a U.S. vaccination plan that minimizes deaths and the economic  
896 cost of a stay-at-home order. The study considered occupation-based risk  
897 exposure (454 occupations). The authors compared three different plans.  
898 Under the only considered plan without a stay-at-home order, the largest  
899 number of vaccines should be allocated to those 50–59 years old, followed by  
900 those 60–69. In general, the best plans focused on age-based risk rather than  
901 occupation-based risk exposure.

902 In [Miura et al. \(2021\)](#), the age-specific transmission intensities (i.e., the  
903 next generation matrix) are reconstructed using an approximation method.  
904 This enables the inference of the expected impact of vaccinating each sub-  
905 group from data on incidence and force of infection. This unique approach  
906 requires only routine surveillance data on the number of cases to determine  
907 the best possible allocation of vaccines, and can be employed in data-scarce  
908 environments. The method is tested with data from the Netherlands. The  
909 authors conclude that the optimal timing of changing from vaccinating one  
910 age group to another depends on the specific objective.

911 In [Cartocci et al. \(2021\)](#) an ODE-based SIR model that considers time-  
912 varying parameters and sex was used to compare Italian vaccination pro-  
913 grams, using the outcomes YLL, deaths and infections. According to the  
914 model, deaths (infections) are minimized by prioritizing elderly (younger).  
915 However, the optimal YLL-minimizing strategy depends on the effective re-  
916 productive number. If it is high, younger individuals should be prioritized.

917 In [Galli et al. \(2023\)](#) an ODE-based SIR model was used to predict  
918 COVID-19 dynamics and evaluate vaccination plans in the Southwest Shewa  
919 Zone in Ethiopia. A plan that prioritizes those 50 years and older was found  
920 to avoid more critical cases than a random vaccine allocation.

921 In [González-Parra et al. \(2022\)](#), two ODE-based SIAR models were used  
922 to study vaccine allocation strategies. Different scenarios related to the speed  
923 of the vaccine roll-out were compared. The authors found that generally  
924 those 55 years and older should be prioritized to minimize deaths. However,  
925 whenever the transmission rate is relatively high and elderly have a substan-  
926 tially lower transmission rate than younger people, the optimal prioritization  
927 switches.

928 In [Aruffo et al. \(2022\)](#), an ODE-based model with many compartments  
929 is used to study different Canadian vaccine roll-out and NPI-lifting scenar-  
930 ios. To minimize infections and shorten the time until NPIs can be lifted,  
931 those 20-59 years old should be prioritized. Different reopening scenarios  
932 and strategies were compared, with total cases and deaths depending on the  
933 timing of lifting NPIs.

### 934 *5.1.2. Agent-based models (ABMs)*

935 ABMs offer more flexibility and potential realism than ODE-based models  
936 but a proper analysis of these stochastic models requires simulations and is  
937 thus computationally expensive.

938 In [Jahn et al. \(2021\)](#), an ABM was developed to derive optimal vaccine  
939 allocation strategies for Austria. The model contains 9 million agents, one  
940 for each Austrian resident. Each agent possesses an associated state variable  
941 that describes its disease state. The model further accounts for age, occupa-  
942 tion (health care workers), testing and notification delays. The probability of  
943 an infection occurring during a single contact between an infected and a sus-  
944 ceptible was determined by calibrating the model to the number of detected  
945 Austrian COVID-19 cases in March 2020. The authors found that hospital-  
946 ization and deaths were minimized if elderly and vulnerable were prioritized,  
947 assuming very limited vaccine availability. To assign more vaccines, the au-

948 thors highlight the usefulness of a stepwise optimal allocation technique, in  
949 which small batches of vaccine are assigned at a time.

950 In [Ben-Zuk et al. \(2022\)](#), an ABM was used to derive and compare op-  
951 timal vaccine allocation strategies for two Israeli cities of similar size but  
952 with different household size and age distributions. The authors compared  
953 two strategies: vaccinate those prioritized by public health decision makers,  
954 or dynamically prioritize neighborhoods with a high estimated reproductive  
955 number. Using infections and deaths as outcomes, the study highlights that  
956 optimal vaccination plans depend on subpopulation-specific infection rates  
957 and unique demographic characteristics.

958 In [Kadelka and McCombs \(2021\)](#), an ABM was used to highlight the ef-  
959 fect of homophily and correlation between attitudes and opinions on vaccine  
960 prioritization. The authors argue that the U.S. society exhibits high levels  
961 of homophily w.r.t. to vaccine willingness and NPI adherence and that these  
962 two attributes are correlated, i.e., that people who get vaccinated are also  
963 more likely to engage in other risk mitigation. The authors found that these  
964 attributes must be taken into account to inform the optimal vaccine prior-  
965 itization strategy, as they influence at which relative contact level of older  
966 compared to younger individuals the optimal prioritization target switches.

967 In [Tatapudi et al. \(2021\)](#), an ABM that considers various NPIs was de-  
968 veloped to track the number of COVID-19 cases, hospitalized, and deaths  
969 for all age groups. 2.8 million agents were used to represent each resident  
970 in Miami Dade County, United States. Three vaccine allocation strategies  
971 were compared: (i) random allocation, (ii) prioritization by age, (iii) a minor  
972 variant of the CDC strategy, which prioritizes health care workers in addition  
973 to elderly. The authors found that a random allocation minimizes infections,  
974 while the CDC strategy minimizes deaths and YLL, although it proved only  
975 slightly better than the other two strategies.

976 In [Bushaj et al. \(2023\)](#), the Covasim ABM from [Kerr et al. \(2021\)](#) was  
977 expanded to compare a random with an age-structured vaccine allocation  
978 strategy. The authors show that a “governor Deep Reinforcement Learning  
979 agent” can learn effective strategies and suggest, based on a multi-objective  
980 reward structure, optimal ABM interventions when presented with a spe-  
981 cific epidemic situation. Moreover, the study shows that focused vaccina-  
982 tion of super-spreaders can substantially reduce infections at the expense of  
983 marginally more total deaths. The model was tested with data from the U.S.  
984 states New Jersey and Kansas.

985 In [Cattaneo et al. \(2022\)](#), the Covasim model is used to determine the



986 number of infections and deaths prevented by vaccines in the Italian region  
987 Lombardy, and to retrospectively evaluate vaccine allocation strategies. Pri-  
988 oritization of the elderly and at-risk categories, as used in Italy, was validated  
989 as the most effective in reducing deaths, however only as long as the vaccine  
990 roll-out happens fast enough.

991 *5.2. Summary of optimal COVID-19 vaccine dosing interval studies that em-  
992 ploy a mathematical model*

993 The following studies all use a mathematical model that differentiates  
994 between those vaccinated with a single dose and two doses (i.e., fully vac-  
995 cinated). The fundamental vaccine prioritization trade-off is between vacci-  
996 nating more people at lower levels of protection or inducing higher protection  
997 for fewer individuals.

998 In [Moghadas et al. \(2021\)](#), an age-structured ABM with compartments  
999 SEPIAR was used that differentiated between the Pfizer and the Moderna  
1000 vaccine. Varying rates of vaccine-induced immunity waning were consid-  
1001 ered. In addition, maximum vaccine coverage (i.e., vaccine hesitancy) was  
1002 assumed to be age-dependent. Model parameters were informed by data from  
1003 the United States. Outcomes were assessed by infections, hospitalizations,  
1004 and deaths. The authors found that a delay of the second dose of at least 9  
1005 weeks would have averted deaths and hospitalizations compared to the rec-  
1006 ommended 4-week interval. For infections, the results differed for the two  
1007 considered vaccines: while a delay of the second dose of Moderna vaccines  
1008 would have reduced infections, delaying second doses of Pfizer vaccines may  
1009 have caused more infections if pre-existing immunity is below 30% and if  
1010 vaccine-induced one-dose immunity wanes.

1011 In [Tuite et al. \(2021\)](#), a decision analytic cohort model was used to as-  
1012 sess strategies for dose allocation (assuming a steady vaccine supply). The  
1013 authors found that variants of a flexible strategy that keeps only 10% of the  
1014 supply for second doses during the first 3 weeks are better than the fixed  
1015 strategy employed by the United States.

1016 In [Souto Ferreira et al. \(2022\)](#), an age-structured SEAIHR delay differen-  
1017 tial equation model was used to study the optimal timing between first and  
1018 second dose. A constant vaccine production rate was assumed and vaccina-  
1019 tion rates were optimized using linear programming, with outcomes assessed  
1020 by deaths. The authors found that the best strategy depends on an interplay  
1021 between the vaccine production rate and the relative efficacy of the first dose.



1022 In [Ferreira et al. \(2022\)](#), a discrete-time compartmental model, fitted  
1023 to Brazil and differentiating between three different vaccines, was used to  
1024 investigate the optimal vaccine prioritization and dosing interval, which was  
1025 varied from 8 to 12 weeks. The authors found that a shorter time interval  
1026 between first and second dose for the AstraZeneca vaccine would minimize  
1027 deaths. However, in their analysis, it appears that the vaccine availability is  
1028 not fixed, i.e., a shorter time interval corresponds to more available vaccine,  
1029 which is obviously beneficial. Moreover, the authors assumed large differences  
1030 in vaccine efficacy between the first and the second doses, contrary to many  
1031 other studies.

1032 In [Zuo et al. \(2022\)](#), an ODE-based SEIQR model was fitted to South  
1033 Africa and used to answer questions related to vaccine prioritization and  
1034 delay of the second dose. The authors found that, assuming limited vaccine  
1035 availability, a delay of second doses leads to fewer severe COVID-19 cases.

1036 In [Gianatti et al. \(2023\)](#), a discrete-time model with compartments SEPIHR  
1037 (no age groups) is fitted to data from the city of Tandil, Argentina. Assuming  
1038 constant numbers of daily available vaccines, different fixed delays between  
1039 the vaccine doses (28, 42, 72 days) were compared using death as the out-  
1040 come metric. An optimal control problem was solved to determine the best  
1041 way to administrate the available vaccines, by considering two controls that  
1042 represent the number of first and second doses applied each day. The authors  
1043 found that delaying the second dose as long as possible (72 days in the study)  
1044 was optimal.

1045 In [Mak et al. \(2022\)](#), an ODE-based SEPAIHR model was used to in-  
1046 vestigate three different policies related to vaccine roll-out: holding back  
1047 second doses, releasing second doses, and delaying the time between doses.  
1048 The authors found that releasing second doses reduces infections. However,  
1049 stretching the between-dose time flattens the infection curve and reduces  
1050 both hospitalizations and mortality compared with a strategy that releases  
1051 second doses. The model includes details related to the inventory dynamics  
1052 of the vaccine roll-out process not found in other models. The authors further  
1053 conduct extensive sensitivity analyses related to age composition, risk-based  
1054 prioritization, supply disruptions, and disease transmissibility.

1055 In [Romero-Brufau et al. \(2021\)](#), an age-structured ABM was used to in-  
1056 vestigate the effect of a delayed second dose on deaths, infections and hospi-  
1057 talizations. A total of 100k agents interact in 3 types of networks (occupation,  
1058 family and random) over a period of six months. In all compared vaccina-  
1059 tion plans, the allocation started with the oldest group and proceeded by

1060 decreasing age. The authors found that a delayed second dose yields lower  
1061 deaths as long as the first dose is sufficiently effective ( $\geq 80\%$ ) but that a  
1062 delay does not affect the YLL and infections much.

1063 In [Diarra et al. \(2022\)](#), an ODE-based SEIARQ model, an adaptation  
1064 of the CoMo model ([Aguas et al., 2020](#)), was used to study vaccination  
1065 strategies in Senegal. Three particular vaccination strategies were evaluated,  
1066 using deaths as outcome metric. The authors found the second dose should  
1067 be delayed for those 40 years or younger.

1068 In [Childs et al. \(2022\)](#), an age-structured ODE-based SEIS model that  
1069 considers reinfections and immunity was used to determine questions related  
1070 to vaccine prioritization and delay of the second dose in Canada. The authors  
1071 found that a delay, as well as earlier vaccination of 15-19 year olds would both  
1072 yield lower infections numbers.

### 1073 *5.3. Summary of spatial vaccine distribution studies that employ a mathe-* 1074 *matical model*

1075 In [Grauer et al. \(2020\)](#), a computational model with Brownian agents  
1076 moving randomly through a continuous square space with periodic bound-  
1077 ary conditions was introduced. Each agent has an internal state variable  
1078 describing its disease state (e.g., S, I, or R). A statistical mean-field model  
1079 was applied to study three vaccine allocation strategies: (i) distribution of  
1080 vaccines proportional to population density, (ii) an “infection weighted strat-  
1081 egy” that distributes vaccines proportional to the quantitative value of the  
1082 bi-linear incidence rate  $\beta SI$ , and (iii) a “focusing strategy” that distributes  
1083 the vaccines sequentially by prioritizing the regions with the highest inci-  
1084 dence rate. The authors found that the last strategy minimized deaths; age  
1085 was however not considered.

1086 In [Molla et al. \(2022\)](#), a spatial ODE-based model was developed to model  
1087 COVID-19 disease dynamics in five different Finnish regions. The authors  
1088 combined age-specific contact data with geographic movement data to in-  
1089 vestigate the optimal vaccination strategies. Using optimal control methods,  
1090 the authors found that allocating vaccines demographically and in an age-  
1091 descending order is not optimal for minimizing deaths or infection cases.  
1092 Instead, it proved optimal to prioritize high-incidence regions and allocate  
1093 vaccines at the same time to different age groups.

1094 In [Zhou et al. \(2021\)](#), the authors used cell phone data from a Chinese  
1095 city to develop a spatial ABM for a realistic urban scenario. To compare  
1096 seven different scenarios related to vaccine allocation, the authors assigned

1097 the vaccines by fulfilling the priority group before advancing to the next pri-  
1098 ority group. The authors found that the vaccine coverage to reach herd im-  
1099 munity varies strongly across locations, highlighting the immense usefulness  
1100 of knowledge of the spatial heterogeneity when designing vaccine allocation  
1101 strategies.

1102 In [Lemaitre et al. \(2022\)](#), an ODE-based spatial model of the 107 Italian  
1103 provinces, originally developed in [Gatto et al. \(2020\)](#), was used to study opti-  
1104 mal vaccine distribution across space. Google Community Mobility Reports  
1105 was used to estimate the variations in mobility across provinces and as a  
1106 proxy for changes in social contacts. The authors developed a novel optimal  
1107 control framework that yields the best vaccination strategy under realistic  
1108 supply and logistics constraints. The identified optimal strategy, which sub-  
1109 stantially outperforms standard strategies, has a complex structure: while  
1110 mainly dependent on the projected incidence of each province, it also takes  
1111 into account the spatial connectivity between provinces.

1112 In [Vo et al. \(2023\)](#), an age-stratified ODE-based spatial SEIR model of  
1113 the 50 U.S. states was used to illustrate the utility of mechanistic expressions  
1114 for the basic and effective reproductive number, as well as to compare two  
1115 vaccine prioritization strategies: a uniform allocation and an allocation along  
1116 the gradient of the effective reproductive number. The authors showed that  
1117 the latter approach yields fewer infections but they acknowledged that this  
1118 would come at the expense of more hospitalizations and deaths.

## 1119 **6. Related studies that employ optimal control methods**

1120 The majority of studies included in this review identified vaccine alloca-  
1121 tion strategies that optimize a given metric, e.g. minimizing deaths or infec-  
1122 tions. Some studies went further and identified strategies that are Pareto-  
1123 optimal with respect to multiple objectives, see e.g. [Islam et al. \(2021\)](#); [Gav-  
1124 ish and Katriel \(2022\)](#); [Diarra et al. \(2022\)](#). A number of studies, some  
1125 already described above, went even further and employed classical opti-  
1126 mal control theory to find vaccination strategies that minimize a variety  
1127 of health and/or economic outcomes. Some of these studies even consider  
1128 age-dependent vaccine access.

1129 These studies define a functional - often a linear combination of different  
1130 metrics - that is optimized given some constraints, e.g., to account for limited  
1131 vaccine availability. A general challenge of optimal control approaches is the  
1132 high sensitivity of the resulting optimal vaccination strategy on the choice

1133 of weights in the functional. Moreover, the choice of functional itself affects  
1134 the results. Nevertheless, these studies can provide important insights as the  
1135 setup is more flexible, and we briefly describe some interesting approaches  
1136 and note that several others (e.g., [Lemaitre et al. \(2022\)](#); [Angelov et al.  
1137 \(2023\)](#)) are already summarized above.

1138 In [Acuña-Zegarra et al. \(2021\)](#), an ODE-based SEAIR model (no age  
1139 structure) was used to show that the optimal vaccination strategy depends  
1140 on the speed of the vaccine roll-out and the length of natural immunity.  
1141 The transmission contact rates and proportion of symptomatic cases were  
1142 estimated by calibrating the model to observed death counts. The basic  
1143 reproductive number was estimated to be in the range of [3.30, 4.84]. The  
1144 authors minimized a functional that was a linear combination of YLL and  
1145 Years Lost due to Disability. The authors found that varying the number of  
1146 doses during the vaccine roll-out (if supply allows) yields to better outcomes  
1147 than an approach with fixed number of vaccinations per day.

1148 In [Tu et al. \(2023\)](#), the authors proposed a reaction-diffusion COVID-19  
1149 model (no age structure) to investigate how different vaccination-isolation  
1150 strategies impact the COVID-19 pandemic. The functional included three  
1151 metrics: social cost, social benefit, and the basic reproduction number. The  
1152 authors found that for a given social cost or benefit, there are many Pareto-  
1153 optimal vaccination-isolation strategies. The proposed model considered also  
1154 a spatial variable, in addition to parameters related to social distancing and  
1155 vaccination.

1156 In [Olivares and Staffetti \(2021a\)](#), two control variables, vaccination and  
1157 testing, were used to find the optimal strategy that minimizes a functional  
1158 that accounts for the number of infected people with life-threatening symp-  
1159 toms and the number of deaths. The underlying model is ODE-based with  
1160 a variety of compartments. Several optimal control problems were solved  
1161 for different scenarios. Among others, the authors found that it is optimal  
1162 to roll-out a vaccine as fast as possible. In [Olivares and Staffetti \(2021b\)](#),  
1163 the same authors studied scenarios with different vaccine availability. The  
1164 functional here depends on the number of symptomatic and asymptomatic  
1165 infectious. The authors found again that early implementation of vaccina-  
1166 tion and testing reduces the number of symptomatically infected the most.  
1167 However, if vaccine availability increases gradually, the optimal vaccination  
1168 strategy differs quite strongly from other scenarios. Finally in [Olivares and  
1169 Staffetti \(2021c\)](#), the same authors considered a mass vaccination plan, and  
1170 polynomial chaos expansion was used to assess the uncertainty of the mod-

1171 eling outcomes.

1172 In [Ziarelli et al. \(2023\)](#), an age-stratified two-dose ODE-based SIR model  
1173 was calibrated to death counts from Italy, and several optimal control prob-  
1174 lems were solved, minimizing deaths, infections and hospitalizations inde-  
1175 pendently. In each problem, the total number of vaccine doses was fixed  
1176 but the distribution of the available doses among susceptibles and those who  
1177 already received their first dose was optimized. The authors found that the  
1178 deaths-minimizing strategy prioritized those 80 years and older, followed,  
1179 interestingly, by those 20-39 years old. On the other hand, the infections-  
1180 minimizing vaccination strategy prioritizes the 20-39 and 40-59 age groups  
1181 but not children and teenagers despite them having the most contacts. This  
1182 work nicely highlights the complexities of designing optimal age-based vac-  
1183 cine prioritization strategies.

1184 In [Choi and Shim \(2021\)](#), an age-structured ODE-based model for South  
1185 Korea was developed. Solving an optimal control problem with a functional  
1186 that considers the cost of vaccination, as well as the cost of symptomatic  
1187 and hospitalized infected, the authors found that the optimal vaccination  
1188 strategy depends on the way the vaccine functions. While “susceptibility-  
1189 reducing” vaccines should be allocated relatively evenly . On the other hand,  
1190 “symptom-reducing” vaccines should, surprisingly, be allocated to those 20-  
1191 29 and 50 and older but not to those 30-49 years old. The impact of vaccine  
1192 function proved particular strong if the roll-out was assumed to be fast.

1193 In [Libotte et al. \(2020\)](#), an SIR model was calibrated to data from China.  
1194 An inverse problem was solved to determine the transmission rate, infectious  
1195 period, initial number of infecteds and basic reproduction number ( $\mathcal{R}_0$ ). The  
1196 authors developed a multi-objective optimal control problem, in which the  
1197 number of vaccines and the total number of infected are simultanenously  
1198 minimized. This problem is solved using Differential Evolution, yielding a  
1199 set of Pareto-optimal vaccination strategies.

1200 In [Zhang et al. \(2024\)](#), an optimal control problem was solved with the  
1201 aim of minimizing deaths and conserving vaccines at the same time. The  
1202 population was divided into four subpopulations: health workers, young in-  
1203 dividuals, middle-aged individuals, and the elderly. The authors found that  
1204 the optimal vaccination strategy substantially improved upon a proportional  
1205 vaccine roll-out.

1206 There exist numerous other works that use classical optimal control to  
1207 identify optimal COVID-19 vaccination strategies, most of them minimizing  
1208 an objective functional which accounts for infected cases, deaths or the num-

1209 ber of vaccines [Agossou et al. \(2021\)](#); [Al-arydah \(2023\)](#); [Salcedo-Varela et al.](#)  
1210 [\(2023\)](#); [Shen et al. \(2021\)](#); [Zaitri et al. \(2022\)](#). In particular, some works  
1211 have combined optimal control with age-structured models to find the opti-  
1212 mal vaccination allocation [Avci and Yurtoğlu \(2023\)](#); [Chhetri et al. \(2022\)](#);  
1213 [Kumar et al. \(2021\)](#). Optimal control employed on infectious disease models  
1214 represents a powerful tool to identify optimal vaccine allocation strategies.  
1215 However, setting up the optimal control problem including the constraints  
1216 regarding vaccine availability is crucial but it is challenging to restrict the  
1217 search to vaccination programs that can be implemented in the real world.  
1218 The choice of the functional to be minimized is also crucial, as strongly affects  
1219 the optimal outcomes, see e.g., [Ledzewicz and Schättler \(2020\)](#).

## 1220 7. Conclusion

1221 The COVID-19 pandemic constitutes one of the worst pandemics hu-  
1222 mankind has ever endured, both in terms of lives lost and economic reper-  
1223 cussions. It is also the first pandemic in a globalized world. The rapid  
1224 spread of the disease around the world was enabled by high levels of connec-  
1225 tion, transport and travel between distant parts of the world. This is not  
1226 going to change, which is why the world will eventually face another pan-  
1227 demic. Whether this will be caused by a highly transmissible SARS-CoV-2  
1228 that has evolved to evade immune defenses or by an entirely novel pathogen  
1229 cannot be predicted. However, we can learn from mistakes made during the  
1230 COVID-19 pandemic to ensure better preparedness for a future pandemic.  
1231 On the mathematical modeling front, this includes fully understanding the  
1232 effect realistic human behavior and social processes have on the outcomes in  
1233 infectious disease models. Specifically for models designed to inform priori-  
1234 tization strategies for a vaccine that will initially always be limited, we need  
1235 to look beyond the details of specific models and understand the greater  
1236 connections behind explicit and implicit model assumptions and outcomes.  
1237 This is what we attempted in this systematic review of mathematical models  
1238 designed to find optimal COVID-19 vaccine prioritization strategies.

## 1239 Acknowledgments

1240 G.G.-P. acknowledges funding from Maria Zambrano (UPV, Ministry of  
1241 Universities of Spain by the European Union-Next generation EU) and an



1242 Institutional Development Award (IDeA) from the National Institute of Gen-  
1243 eral Medical Sciences of the National Institutes of Health under grant number  
1244 P20GM103451. C.K. was partially supported by a travel grant (712537) from  
1245 the Simons Foundation.

#### 1246 **Conflict of interest**

1247 The authors declare there is no conflict of interest.

#### 1248 **References**

- 1249 Acuña-Zegarra, M.A., Díaz-Infante, S., Baca-Carrasco, D., Olmos-Liceaga,  
1250 D., 2021. COVID-19 optimal vaccination policies: a modeling study on  
1251 efficacy, natural and vaccine-induced immunity responses. *Mathematical*  
1252 *Biosciences* 337, 108614.
- 1253 Agossou, O., Atchadé, M.N., Djibril, A.M., 2021. Modeling the effects of  
1254 preventive measures and vaccination on the COVID-19 spread in Benin  
1255 Republic with optimal control. *Results in Physics* 31, 104969.
- 1256 Aguas, R., White, L., Hupert, N., Shretta, R., Pan-Ngum, W., Celhay, O.,  
1257 Moldokmatova, A., Arifi, F., Mirzazadeh, A., Sharifi, H., et al., 2020. Mod-  
1258 elling the COVID-19 pandemic in context: an international participatory  
1259 approach. *BMJ global health* 5, e003126.
- 1260 Al-arydah, M., 2023. Mathematical modeling and optimal control for  
1261 COVID-19 with population behavior. *Mathematical Methods in the Ap-*  
1262 *plied Sciences* 46, 19184–19198.
- 1263 Althobaity, Y., Wu, J., Tildesley, M.J., 2022. Non-pharmaceutical interven-  
1264 tions and their relevance in the COVID-19 vaccine rollout in Saudi Arabia  
1265 and Arab Gulf countries. *Infectious Disease Modelling* 7, 545–560.
- 1266 Angelov, G., Kovacevic, R., Stilianakis, N.I., Veliov, V.M., 2023. Optimal  
1267 vaccination strategies using a distributed model applied to COVID-19.  
1268 *Central Europe an Journal of Operations Research* 31, 499–521.
- 1269 Anupong, S., Chantanasaro, T., Wilasang, C., Jitsuk, N.C., Sararat,  
1270 C., Sornbundit, K., Pattanasiri, B., Wannigama, D.L., Amarasiri, M.,  
1271 Chadsuthi, S., et al., 2023. Modeling vaccination strategies with limited



- 1272 early COVID-19 vaccine access in low-and middle-income countries: A case  
1273 study of Thailand. *Infectious Disease Modelling* 8, 1177–1189.
- 1274 Are, E.B., Card, K.G., Colijn, C., 2024. The role of vaccine status homophily  
1275 in the COVID-19 pandemic: a cross-sectional survey with modelling. *BMC*  
1276 *Public Health* 24, 1–16.
- 1277 Aruffo, E., Yuan, P., Tan, Y., Gatov, E., Moyles, I., Bélair, J., Watmough, J.,  
1278 Collier, S., Arino, J., Zhu, H., 2022. Mathematical modelling of vaccination  
1279 rollout and NPIs lifting on COVID-19 transmission with VOC: a case study  
1280 in Toronto, Canada. *BMC public health* 22, 1–12.
- 1281 Avcı, D., Yurtoğlu, M., 2023. An optimal vaccination scenario for COVID-19  
1282 transmission between children and adults, in: *Mathematical Modeling and*  
1283 *Intelligent Control for Combating Pandemics*. Springer, pp. 93–108.
- 1284 Ayoub, H.H., Chemaitelly, H., Makhoul, M., Al Kanaani, Z., Al Kuwari, E.,  
1285 Butt, A.A., Coyle, P., Jeremijenko, A., Kaleeckal, A.H., Latif, A.N., et al.,  
1286 2021. Epidemiological impact of prioritising SARS-CoV-2 vaccination by  
1287 antibody status: mathematical modelling analyses. *BMJ innovations* 7.
- 1288 Babus, A., Das, S., Lee, S., 2023. The optimal allocation of COVID-19  
1289 vaccines. *Economics Letters* 224, 111008.
- 1290 Barmounakis, P., Demiris, N., Kontoyiannis, I., Pavlakis, G.N., Sypsa, V.,  
1291 2022. Evaluating the effects of second-dose vaccine-delay policies in Europe  
1292 an countries: A simulation study based on data from Greece. *PLOS One*  
1293 17, e0263977.
- 1294 Ben-Zuk, N., Daon, Y., Sasson, A., Ben-Adi, D., Huppert, A., Nevo, D.,  
1295 Obolski, U., 2022. Assessing COVID-19 vaccination strategies in varied  
1296 demographics using an individual-based model. *Frontiers in public health*  
1297 10, 966756.
- 1298 Bubar, K.M., Reinholt, K., Kissler, S.M., Lipsitch, M., Cobey, S., Grad,  
1299 Y.H., Larremore, D.B., 2021. Model-informed COVID-19 vaccine prioritization  
1300 strategies by age and serostatus. *Science* 371, 916–921.
- 1301 Buckner, J.H., Chowell, G., Springborn, M.R., 2021. Dynamic prioritization  
1302 of COVID-19 vaccines when social distancing is limited for essential work-  
1303 ers. *Proceedings of the National Academy of Sciences* 118, e2025786118.

- 1304 Burgio, G., Steinegger, B., Arenas, A., 2022. Homophily impacts the success  
1305 of vaccine roll-outs. *Communications Physics* 5, 70.
- 1306 Bushaj, S., Yin, X., Beqiri, A., Andrews, D., Büyüктаhtakın, I.E., 2023.  
1307 A simulation-deep reinforcement learning (SiRL) approach for epidemic  
1308 control optimization. *Annals of Operations Research* 328, 245–277.
- 1309 Caga-anan, R.L., Macalisang, J.M., Dalisay, J.L.M., Raza, M.N., Martinez,  
1310 J.G.T., Arcede, J.P., 2023. Optimal vaccination control for COVID-19 in  
1311 a metapopulation model: a case of the philippines. *Frontiers in Applied  
1312 Mathematics and Statistics* 9, 1154634.
- 1313 Campbell, F., Archer, B., Laurenson-Schafer, H., Jinnai, Y., Konings, F.,  
1314 Batra, N., Pavlin, B., Vandemaele, K., Van Kerkhove, M.D., Jombart, T.,  
1315 et al., 2021. Increased transmissibility and global spread of SARS-CoV-2  
1316 variants of concern as at June 2021. *Eurosurveillance* 26, 2100509.
- 1317 Campos, E.L., Cysne, R.P., Madureira, A.L., Mendes, G.L., 2021. Multi-  
1318 generational SIR modeling: Determination of parameters, epidemiologi-  
1319 cal forecasting and age-dependent vaccination policies. *Infectious Disease  
1320 Modelling* 6, 751–765.
- 1321 Cartocci, A., Cevenini, G., Barbini, P., 2021. A compartment modeling  
1322 approach to reconstruct and analyze gender and age-grouped COVID-19  
1323 Italian data for decision-making strategies. *Journal of Biomedical Informatics* 118, 103793.
- 1325 Cattaneo, A., Vitali, A., Mazzoleni, M., Previdi, F., 2022. An agent-based  
1326 model to assess large-scale COVID-19 vaccination campaigns for the Italian  
1327 territory: The case study of Lombardy region. *Computer Methods and  
1328 Programs in Biomedicine* 224, 107029.
- 1329 Centers for Disease Control (CDC), . <https://www.cdc.gov/museum/timeline/covid19.html>  
1330 [accessed: 01/14/2024].
- 1331 Chakraborty, C., Sharma, A.R., Bhattacharya, M., Lee, S.S., 2022. A de-  
1332 tailed overview of immune escape, antibody escape, partial vaccine escape  
1333 of SARS-CoV-2 and their emerging variants with escape mutations. *Frontiers in Immunology* 13, 801522.
- 1334

- 1335 Chen, X., Zhu, G., Zhang, L., Fang, Y., Guo, L., Chen, X., 2021. Design and  
1336 analysis of network behaviors for optimizing network energy efficiency in 5g  
1337 mmwave systems. *IEEE Transactions on Network Science and Engineering*  
1338 8, 1862–1872.
- 1339 Chhetri, B., Vamsi, D., Prakash, D.B., Balasubramanian, S., Sanjeevi, C.B.,  
1340 2022. Age structured mathematical modeling studies on COVID-19 with  
1341 respect to combined vaccination and medical treatment strategies. *Com-  
1342 putational and Mathematical Biophysics* 10, 281–303.
- 1343 Childs, L., Dick, D.W., Feng, Z., Heffernan, J.M., Li, J., Röst, G., 2022.  
1344 Modeling waning and boosting of COVID-19 in Canada with vaccination.  
1345 *Epidemics* 39, 100583.
- 1346 Choi, W., Shim, E., 2021. Vaccine effects on susceptibility and symptoma-  
1347 tology can change the optimal allocation of COVID-19 vaccines: South  
1348 Korea as an example. *Journal of Clinical Medicine* 10, 2813.
- 1349 Choi, Y., Kim, J.S., Kim, J.E., Choi, H., Lee, C.H., 2021. Vaccination  
1350 prioritization strategies for COVID-19 in Korea: a mathematical modeling  
1351 approach. *International Journal of Environmental Research and Public  
1352 Health* 18, 4240.
- 1353 Coletti, P., Wambua, J., Gimma, A., Willem, L., Vercruyse, S., Vanhoutte,  
1354 B., Jarvis, C.I., Van Zandvoort, K., Edmunds, J., Beutels, P., et al., 2020.  
1355 CoMix: comparing mixing patterns in the belgian population during and  
1356 after lockdown. *Scientific reports* 10, 21885.
- 1357 Conway, E., Walker, C.R., Baker, C., Lydeamore, M.J., Ryan, G.E., Camp-  
1358 bell, T., Miller, J.C., Rebuli, N., Yeung, M., Kabashima, G., et al., 2023.  
1359 COVID-19 vaccine coverage targets to inform reopening plans in a low  
1360 incidence setting. *Proceedings of the Royal Society B* 290, 20231437.
- 1361 Davies, N.G., Klepac, P., Liu, Y., Prem, K., Jit, M., Eggo, R.M., 2020. Age-  
1362 dependent effects in the transmission and control of COVID-19 epidemics.  
1363 *Nature medicine* 26, 1205–1211.
- 1364 Diarra, M., Kebir, A., Talla, C., Barry, A., Faye, J., Louati, D., Opatowski,  
1365 L., Diop, M., White, L.J., Loucoubar, C., et al., 2022. Non-pharmaceutical  
1366 interventions and COVID-19 vaccination strategies in Senegal: a modelling  
1367 study. *BMJ global health* 7, e007236.

- 1368 Dryhurst, S., Schneider, C.R., Kerr, J., Freeman, A.L., Recchia, G., Van  
1369 Der Bles, A.M., Spiegelhalter, D., Van Der Linden, S., 2022. Risk per-  
1370 ceptions of COVID-19 around the world, in: COVID-19. Routledge, pp.  
1371 162–174.
- 1372 Ferranna, M., Cadarette, D., Bloom, D.E., 2021. COVID-19 vaccine allo-  
1373 cation: Modeling health outcomes and equity implications of alternative  
1374 strategies. *Engineering* 7, 924–935.
- 1375 Ferreira, L.S., de Almeida, G.B., Borges, M.E., Simon, L.M., Poloni, S.,  
1376 Bagattini, Á.M., da Rosa, M.Q.M., Diniz Filho, J.A.F., de Souza Kuchen-  
1377 becker, R., Camey, S.A., et al., 2022. Modelling optimal vaccination strate-  
1378 gies against COVID-19 in a context of Gamma variant predominance in  
1379 Brazil. *Vaccine* 40, 6616–6624.
- 1380 Foy, B.H., Wahl, B., Mehta, K., Shet, A., Menon, G.I., Britto, C., 2021.  
1381 Comparing COVID-19 vaccine allocation strategies in India: A mathe-  
1382 matical modelling study. *International Journal of Infectious Diseases* 103,  
1383 431–438.
- 1384 Funk, S., Knapp, J.K., Lebo, E., Reef, S.E., Dabbagh, A.J., Kretsinger, K.,  
1385 Jit, M., Edmunds, W.J., Strebel, P.M., 2019. Combining serological and  
1386 contact data to derive target immunity levels for achieving and maintaining  
1387 measles elimination. *BMC medicine* 17, 1–12.
- 1388 Galli, M., Zardini, A., Gamshie, W.N., Santini, S., Tsegaye, A., Trentini,  
1389 F., Marziano, V., Guzzetta, G., Manica, M., d’Andrea, V., et al., 2023.  
1390 Priority age targets for COVID-19 vaccination in Ethiopia under limited  
1391 vaccine supply. *Scientific Reports* 13, 5586.
- 1392 Gatto, M., Bertuzzo, E., Mari, L., Miccoli, S., Carraro, L., Casagrandi, R.,  
1393 Rinaldo, A., 2020. Spread and dynamics of the COVID-19 epidemic in  
1394 Italy: Effects of emergency containment measures. *Proceedings of the*  
1395 *National Academy of Sciences* 117, 10484–10491.
- 1396 Gavish, N., Katriel, G., 2022. The role of childrens’ vaccination for COVID-  
1397 19—Pareto-optimal allocations of vaccines. *PLoS Computational Biology*  
1398 18, e1009872.

- 1399 Gianatti, J., Lotito, P., Neder, J., Núñez, P., Parente, L., 2023. Optimal  
1400 vaccination policies for covid-19 considering vaccine doses delays. *Trends*  
1401 in Computational and Applied Mathematics 24, 121–139.
- 1402 Gonzalez-Parra, G., 2021. Analysis of delayed vaccination regimens: A math-  
1403 ematical modeling approach. *Epidemiologia* 2, 271–293.
- 1404 González-Parra, G., Cogollo, M.R., Arenas, A.J., 2022. Mathematical mod-  
1405 eling to study optimal allocation of vaccines against COVID-19 using an  
1406 age-structured population. *Axioms* 11, 109.
- 1407 Gozzi, N., Bajardi, P., Perra, N., 2021. The importance of non-  
1408 pharmaceutical interventions during the COVID-19 vaccine rollout. *PLoS*  
1409 *computational biology* 17, e1009346.
- 1410 Gozzi, N., Chinazzi, M., Davis, J.T., Mu, K., Pastore y Piontti, A., Ajelli,  
1411 M., Perra, N., Vespignani, A., 2022. Anatomy of the first six months of  
1412 COVID-19 vaccination campaign in Italy. *PLoS Computational Biology*  
1413 18, e1010146.
- 1414 Grauer, J., Löwen, H., Liebchen, B., 2020. Strategic spatiotemporal vaccine  
1415 distribution increases the survival rate in an infectious disease like COVID-  
1416 19. *Scientific reports* 10, 1–10.
- 1417 Grundel, S.M., Heyder, S., Hotz, T., Ritschel, T.K., Sauerteig, P., Worth-  
1418 mann, K., 2021. How to coordinate vaccination and social distancing to  
1419 mitigate SARS-CoV-2 outbreaks. *SIAM Journal on Applied Dynamical*  
1420 *Systems* 20, 1135–1157.
- 1421 Hale, T., Angrist, N., Goldszmidt, R., Kira, B., Petherick, A., Phillips, T.,  
1422 Webster, S., Cameron-Blake, E., Hallas, L., Majumdar, S., et al., 2021.  
1423 A global panel database of pandemic policies (Oxford COVID-19 Govern-  
1424 ment Response Tracker). *Nature human behaviour* 5, 529–538.
- 1425 Halloran, M.E., Struchiner, C.J., Longini Jr, I.M., 1997. Study designs for  
1426 evaluating different efficacy and effectiveness aspects of vaccines. *American*  
1427 *journal of epidemiology* 146, 789–803.
- 1428 Han, S., Cai, J., Yang, J., Zhang, J., Wu, Q., Zheng, W., Shi, H., Ajelli,  
1429 M., Zhou, X.H., Yu, H., 2021. Time-varying optimization of COVID-19

- 1430 vaccine prioritization in the context of limited vaccination capacity. *Nature*  
1431 *communications* 12, 4673.
- 1432 Hiraoka, T., Rizi, A.K., Kivelä, M., Saramäki, J., 2022. Herd immunity and  
1433 epidemic size in networks with vaccination homophily. *Physical Review E*  
1434 105, L052301.
- 1435 Hogan, A.B., Winskill, P., Watson, O.J., Walker, P.G., Whittaker, C.,  
1436 Baguelin, M., Brazeau, N.F., Charles, G.D., Gaythorpe, K.A., Hamlet,  
1437 A., et al., 2021. Within-country age-based prioritisation, global allocation,  
1438 and public health impact of a vaccine against SARS-CoV-2: A mathematical  
1439 modelling analysis. *Vaccine* 39, 2995–3006.
- 1440 Hong, Z., Li, Y., Gong, Y., Chen, W., 2022. A data-driven spatially-specific  
1441 vaccine allocation framework for COVID-19. *Annals of Operations Re-*  
1442 *search* , 1–24.
- 1443 Hupert, N., Marín-Hernández, D., Gao, B., Águas, R., Nixon, D.F., 2022.  
1444 Heterologous vaccination interventions to reduce pandemic morbidity and  
1445 mortality: Modeling the US winter 2020 COVID-19 wave. *Proceedings of*  
1446 *the National Academy of Sciences* 119, e2025448119.
- 1447 Islam, M.R., Oraby, T., McCombs, A., Chowdhury, M.M., Al-Mamun, M.,  
1448 Tyshenko, M.G., Kadelka, C., 2021. Evaluation of the United States  
1449 COVID-19 vaccine allocation strategy. *PLOS One* 16, e0259700.
- 1450 Jahn, B., Sroczynski, G., Bicher, M., Rippinger, C., Mühlberger, N., Santa-  
1451 maria, J., Urach, C., Schomaker, M., Stojkov, I., Schmid, D., et al., 2021.  
1452 Targeted COVID-19 Vaccination (TAV-COVID) Considering Limited Vac-  
1453 cination Capacities—An Agent-Based Modeling Evaluation. *Vaccines* 9,  
1454 434.
- 1455 Jentsch, P.C., Anand, M., Bauch, C.T., 2021. Prioritising COVID-19 vacci-  
1456 nation in changing social and epidemiological landscapes: a mathematical  
1457 modelling study. *The Lancet Infectious Diseases* 21, 1097–1106.
- 1458 Jimenez-Rodriguez, P., Munoz-Fernandez, G.A., Rodrigo-Chocano, J.C.,  
1459 Seoane-Sepulveda, J.B., Weber, A., 2022. A population structure-sensitive  
1460 mathematical model assessing the effects of vaccination during the third  
1461 surge of COVID-19 in Italy. *Journal of Mathematical Analysis and Appli-*  
1462 *cations* 514, 125975.

- 1463 Kadelka, C., 2023. Projecting social contact matrices to populations stratified  
1464 by binary attributes with known homophily. *Mathematical biosciences and*  
1465 *engineering: MBE* 20, 3282–3300.
- 1466 Kadelka, C., Islam, M.R., McCombs, A., Alston, J., Morton, N., 2022. Ethnic  
1467 homophily affects vaccine prioritization strategies. *Journal of Theoretical*  
1468 *Biology* 555, 111295.
- 1469 Kadelka, C., McCombs, A., 2021. Effect of homophily and correlation of  
1470 beliefs on COVID-19 and general infectious disease outbreaks. *PLOS One*  
1471 16, e0260973.
- 1472 Karabay, A., Kuzdeuov, A., Ospanova, S., Lewis, M., Varol, H.A., 2021.  
1473 A vaccination simulator for COVID-19: Effective and sterilizing immu-  
1474 nization cases. *IEEE Journal of Biomedical and Health Informatics* 25,  
1475 4317–4327.
- 1476 Kekić, A., Dehning, J., Gresele, L., von Kügelgen, J., Priesemann, V.,  
1477 Schölkopf, B., 2023. Evaluating vaccine allocation strategies using  
1478 simulation-assisted causal modeling. *Patterns* 4, 100739.
- 1479 Kermack, W.O., McKendrick, A.G., 1927. A contribution to the mathe-  
1480 matical theory of epidemics. *Proceedings of the Royal Society of London.*  
1481 *Series A, Containing papers of a mathematical and physical character* 115,  
1482 700–721.
- 1483 Kerr, C.C., Stuart, R.M., Mistry, D., Abeysuriya, R.G., Rosenfeld, K., Hart,  
1484 G.R., Núñez, R.C., Cohen, J.A., Selvaraj, P., Hagedorn, B., et al., 2021.  
1485 Covasim: an agent-based model of COVID-19 dynamics and interventions.  
1486 *PLOS Computational Biology* 17, e1009149.
- 1487 Kiem, C.T., Massonnaud, C.R., Levy-Bruhl, D., Poletto, C., Colizza, V.,  
1488 Bosetti, P., Fontanet, A., Gabet, A., Olié, V., Zanetti, L., et al., 2021.  
1489 A modelling study investigating short and medium-term challenges for  
1490 COVID-19 vaccination: From prioritisation to the relaxation of measures.  
1491 *EclinicalMedicine* 38.
- 1492 Kobayashi, T., Nishiura, H., 2022. Prioritizing COVID-19 vaccination. part  
1493 2: Real-time comparison between single-dose and double-dose in Japan.  
1494 *Math. Biosci. Eng* 19, 7410–7424.



- 1495 Kumar, A., Viswakarma, N.K., Adlakha, A., Mukherjee, K., 2021. How  
1496 successful have the lockdowns been in controlling the (COVID-19/SARS-  
1497 CoV-2) pandemic—A simulation-based analysis. *International Journal of*  
1498 *Modeling, Simulation, and Scientific Computing*, 2041002.
- 1499 Ledzewicz, U., Schättler, H., 2020. On the role of the objective in the op-  
1500 timization of compartmental models for biomedical therapies. *Journal of*  
1501 *optimization theory and applications* 187, 305–335.
- 1502 Lemaitre, J.C., Pasetto, D., Zanon, M., Bertuzzo, E., Mari, L., Miccoli, S.,  
1503 Casagrandi, R., Gatto, M., Rinaldo, A., 2022. Optimal control of the  
1504 spatial allocation of COVID-19 vaccines: Italy as a case study. *PLoS*  
1505 *computational biology* 18, e1010237.
- 1506 Li, M., Zu, J., Zhang, Y., Ma, L., Shen, M., Li, Z., Ji, F., 2022. COVID-19  
1507 epidemic in New York City: development of an age group-specific math-  
1508 ematical model to predict the outcome of various vaccination strategies.  
1509 *Virology Journal* 19, 1–13.
- 1510 Li, R., Bjørnstad, O.N., Stenseth, N.C., 2021. Prioritizing vaccination by  
1511 age and social activity to advance societal health benefits in Norway: a  
1512 modelling study. *The Lancet Regional Health–Europe* 10.
- 1513 Libotte, G.B., Lobato, F.S., Platt, G.M., Neto, A.J.S., 2020. Determination  
1514 of an optimal control strategy for vaccine administration in COVID-19  
1515 pandemic treatment. *Computer methods and programs in biomedicine*  
1516 196, 105664.
- 1517 Liu, K., Lou, Y., 2022. Optimizing COVID-19 vaccination programs during  
1518 vaccine shortages. *Infectious Disease Modelling* 7, 286–298.
- 1519 Liu, Y., Pearson, C.A., Sandmann, F.G., Barnard, R.C., Kim, J.H., Flasche,  
1520 S., Jit, M., Abbas, K., 2022a. Dosing interval strategies for two-dose  
1521 COVID-19 vaccination in 13 middle-income countries of Europe: Health  
1522 impact modelling and benefit-risk analysis. *The Lancet Regional Health–*  
1523 *Europe* 17.
- 1524 Liu, Y., Sandmann, F.G., Barnard, R.C., Pearson, C.A., Pastore, R., Pebody,  
1525 R., Flasche, S., Jit, M., 2022b. Optimising health and economic impacts of  
1526 COVID-19 vaccine prioritisation strategies in the WHO European Region:  
1527 a mathematical modelling study. *The Lancet Regional Health–Europe* 12.

- 1528 Liu, Z., Omayrat, M., Stursberg, O., 2021. A study on model-based optimization of vaccination strategies against epidemic virus spread., in: ICINCO,  
1529 pp. 630–637.  
1530
- 1531 Lloyd, A.L., 2001. Realistic distributions of infectious periods in epidemic  
1532 models: changing patterns of persistence and dynamics. *Theoretical popu-*  
1533 *lation biology* 60, 59–71.
- 1534 Luangasanatip, N., Painter, C., Pan-ngum, W., Saralamba, S., Wichaita,  
1535 T., White, L., Aguas, R., Clapham, H., Wang, Y., Isaranuwachai, W.,  
1536 et al., 2023. How to model the impact of vaccines for policymaking when  
1537 the characteristics are uncertain: A case study in Thailand prior to the  
1538 vaccine rollout during the COVID-19 pandemic. *Vaccine* 41, 4854–4860.
- 1539 Luebben, G., González-Parra, G., Cervantes, B., 2023. Study of optimal vac-  
1540 cination strategies for early COVID-19 pandemic using an age-structured  
1541 mathematical model: A case study of the USA. *Mathematical Biosciences*  
1542 *and Engineering* 20, 10828–10865.
- 1543 Luo, Q., Weightman, R., Mcquade, S., Díaz, M., Trélat, E., Barbour, W.,  
1544 Work, D., Samaranyake, S., Piccoli, B., 2022. Optimization of vaccination  
1545 for COVID-19 in the midst of a pandemic. *Networks and Heterogeneous*  
1546 *Media* 17, 443–466.
- 1547 MacIntyre, C.R., Costantino, V., Trent, M., 2022. Modelling of COVID-19  
1548 vaccination strategies and herd immunity, in scenarios of limited and full  
1549 vaccine supply in NSW, Australia. *Vaccine* 40, 2506–2513.
- 1550 Mak, H.Y., Dai, T., Tang, C.S., 2022. Managing two-dose COVID-19 vaccine  
1551 rollouts with limited supply: Operations strategies for distributing time-  
1552 sensitive resources. *Production and Operations Management* 31, 4424–  
1553 4442.
- 1554 Makhoul, M., Ayoub, H.H., Chemaitelly, H., Seedat, S., Mumtaz, G.R.,  
1555 Al-Omari, S., Abu-Raddad, L.J., 2020. Epidemiological impact of SARS-  
1556 CoV-2 vaccination: Mathematical modeling analyses. *Vaccines* 8, 668.
- 1557 Mandal, S., Arinaminpathy, N., Bhargava, B., Panda, S., 2021. India’s prag-  
1558 matic vaccination strategy against COVID-19: a mathematical modelling-  
1559 based analysis. *BMJ open* 11, e048874.

- 1560 Markov, P.V., Ghafari, M., Beer, M., Lythgoe, K., Simmonds, P., Stilianakis,  
1561 N.I., Katzourakis, A., 2023. The evolution of SARS-CoV-2. *Nature Re-*  
1562 *views Microbiology* 21, 361–379.
- 1563 Masters, N.B., Shih, S.F., Bukoff, A., Akel, K.B., Kobayashi, L.C., Miller,  
1564 A.L., Harapan, H., Lu, Y., Wagner, A.L., 2020. Social distancing in re-  
1565 sponse to the novel coronavirus (COVID-19) in the United States. *PLOS*  
1566 *One* 15, e0239025.
- 1567 Matrajt, L., Eaton, J., Leung, T., Brown, E.R., 2021a. Vaccine optimization  
1568 for COVID-19: Who to vaccinate first? *Science Advances* 7, eabf1374.
- 1569 Matrajt, L., Eaton, J., Leung, T., Dimitrov, D., Schiffer, J.T., Swan, D.A.,  
1570 Janes, H., 2021b. Optimizing vaccine allocation for COVID-19 vaccines  
1571 shows the potential role of single-dose vaccination. *Nature communications*  
1572 12, 3449.
- 1573 McBryde, E.S., Meehan, M.T., Caldwell, J.M., Adekunle, A.I., Ogunlade,  
1574 S.T., Kuddus, M.A., Ragonnet, R., Jayasundara, P., Trauer, J.M., Cope,  
1575 R.C., 2021. Modelling direct and herd protection effects of vaccination  
1576 against the SARS-CoV-2 delta variant in Australia. *Medical Journal of*  
1577 *Australia* 215, 427–432.
- 1578 McPherson, M., Smith-Lovin, L., Cook, J.M., 2001. Birds of a feather:  
1579 Homophily in social networks. *Annual review of sociology* 27, 415–444.
- 1580 Miura, F., Leung, K.Y., Klinkenberg, D., Ainslie, K.E., Wallinga, J., 2021.  
1581 Optimal vaccine allocation for COVID-19 in the Netherlands: A data-  
1582 driven prioritization. *PLoS computational biology* 17, e1009697.
- 1583 Moghadas, S.M., Vilches, T.N., Zhang, K., Nourbakhsh, S., Sah, P., Fitz-  
1584 patrick, M.C., Galvani, A.P., 2021. Evaluation of COVID-19 vaccination  
1585 strategies with a delayed second dose. *PLoS Biology* 19, e3001211.
- 1586 Molla, J., Ponce de León Chávez, A., Hiraoka, T., Ala-Nissila, T., Kivelä,  
1587 M., Leskelä, L., 2022. Adaptive and optimized COVID-19 vaccination  
1588 strategies across geographical regions and age groups. *PLoS computational*  
1589 *biology* 18, e1009974.

- 1590 Moore, S., Hill, E.M., Dyson, L., Tildesley, M.J., Keeling, M.J., 2021a. Mod-  
1591 elling optimal vaccination strategy for SARS-CoV-2 in the UK. *PLoS*  
1592 *computational biology* 17, e1008849.
- 1593 Moore, S., Hill, E.M., Tildesley, M.J., Dyson, L., Keeling, M.J., 2021b. Vac-  
1594 cination and non-pharmaceutical interventions for COVID-19: a mathe-  
1595 matical modelling study. *The Lancet Infectious Diseases* 21, 793–802.
- 1596 Morales-Zamora, G., Espinosa, O., Puertas, E., Fernández, J.C., Hernández,  
1597 J., Zakzuk, V., Cepeda, M., Alvis-Gúzman, N., Castañeda-Orjuela, C.,  
1598 Paternina-Caicedo, A., 2022. Cost-effectiveness analysis of strategies of  
1599 COVID-19 vaccination in Colombia: comparison of high-risk prioritization  
1600 and no prioritization strategies with the absence of a vaccination plan.  
1601 *Value in Health Regional Issues* 31, 101–110.
- 1602 Mossong, J., Hens, N., Jit, M., Beutels, P., Auranen, K., Mikolajczyk, R.,  
1603 Massari, M., Salmaso, S., Tomba, G.S., Wallinga, J., et al., 2008. Social  
1604 contacts and mixing patterns relevant to the spread of infectious diseases.  
1605 *PLoS medicine* 5, e74.
- 1606 New York Times, . [https://www.nytimes.com/interactive/2021/world/covid-](https://www.nytimes.com/interactive/2021/world/covid-vaccinations-tracker.html)  
1607 [vaccinations-tracker.html](https://www.nytimes.com/interactive/2021/world/covid-vaccinations-tracker.html) [accessed: 01/14/2024].
- 1608 Noh, E.B., Nam, H.K., Lee, H., 2021. Which group should be vaccinated  
1609 first?: A systematic review. *Infection & chemotherapy* 53, 261–270.
- 1610 Nuraini, N., Sukandar, K., Hadisoemarto, P., Susanto, H., Hasan, A.,  
1611 Sumarti, N., 2021. Mathematical models for assessing vaccination sce-  
1612 narios in several provinces in Indonesia. *Infectious Disease Modelling* 6,  
1613 1236–1258.
- 1614 Odusanya, O.O., Odugbemi, B.A., Odugbemi, T.O., Ajisegiri, W.S., 2020.  
1615 COVID-19: A review of the effectiveness of non-pharmacological interven-  
1616 tions. *Nigerian Postgraduate Medical Journal* 27, 261–267.
- 1617 Olivares, A., Staffetti, E., 2021a. Optimal control applied to vaccination and  
1618 testing policies for COVID-19. *Mathematics* 9, 3100.
- 1619 Olivares, A., Staffetti, E., 2021b. Optimal control-based vaccination and  
1620 testing strategies for COVID-19. *Computer Methods and Programs in*  
1621 *Biomedicine* 211, 106411.

- 1622 Olivares, A., Staffetti, E., 2021c. Uncertainty quantification of a mathe-  
1623 matical model of COVID-19 transmission dynamics with mass vaccination  
1624 strategy. *Chaos, Solitons & Fractals* 146, 110895.
- 1625 Pasion, R., Paiva, T.O., Fernandes, C., Barbosa, F., 2020. The age ef-  
1626 fect on protective behaviors during the COVID-19 outbreak: Sociodemo-  
1627 graphic, perceptions and psychological accounts. *Frontiers in psychology*  
1628 11, 561785.
- 1629 Pearson, C.A., Bozzani, F., Procter, S.R., Davies, N.G., Huda, M., Jensen,  
1630 H.T., Keogh-Brown, M., Khalid, M., Sweeney, S., Torres-Rueda, S., et al.,  
1631 2021. COVID-19 vaccination in Sindh Province, Pakistan: A modelling  
1632 study of health impact and cost-effectiveness. *PLoS Medicine* 18, e1003815.
- 1633 Penn, M.J., Donnelly, C.A., 2023. Asymptotic analysis of optimal vaccination  
1634 policies. *Bulletin of Mathematical Biology* 85, 15.
- 1635 Perry, B.C., 1982. Validity and reliability of responses of the aged to surveys  
1636 and questionnaires. *The Journal of Family Practice* 15, 182–183.
- 1637 Prem, K., Cook, A.R., Jit, M., 2017. Projecting social contact matrices in  
1638 152 countries using contact surveys and demographic data. *PLoS compu-  
1639 tational biology* 13, e1005697.
- 1640 Prem, K., Zandvoort, K.v., Klepac, P., Eggo, R.M., Davies, N.G., Centre  
1641 for the Mathematical Modelling of Infectious Diseases COVID-19 Working  
1642 Group and Cook, A.R., Jit, M., 2021. Projecting contact matrices in 177  
1643 geographical regions: an update and comparison with empirical data for  
1644 the covid-19 era. *PLoS computational biology* 17, e1009098.
- 1645 Rahmandad, H., 2022. Behavioral responses to risk promote vaccinating  
1646 high-contact individuals first. *System Dynamics Review* 38, 246–263.
- 1647 Rao, I.J., Brandeau, M.L., 2021a. Optimal allocation of limited vaccine to  
1648 control an infectious disease: Simple analytical conditions. *Mathematical  
1649 biosciences* 337, 108621.
- 1650 Rao, I.J., Brandeau, M.L., 2021b. Optimal allocation of limited vaccine to  
1651 minimize the effective reproduction number. *Mathematical Biosciences*  
1652 339, 108654.

- 1653 Rodriguez-Maroto, G., Atienza-Diez, I., Ares, S., Manrubia, S., 2023. Vacci-  
1654 nation strategies in structured populations under partial immunity and re-  
1655 infection. *Journal of Physics A: Mathematical and Theoretical* 56, 204003.
- 1656 Romero-Brufau, S., Chopra, A., Ryu, A.J., Gel, E., Raskar, R., Kremers, W.,  
1657 Anderson, K.S., Subramanian, J., Krishnamurthy, B., Singh, A., et al.,  
1658 2021. Public health impact of delaying second dose of BNT162b2 or  
1659 mRNA-1273 COVID-19 vaccine: simulation agent based modeling study.  
1660 *bmj* 373.
- 1661 Saadi, N., Chi, Y.L., Ghosh, S., Eggo, R.M., McCarthy, C.V., Quaipe, M.,  
1662 Dawa, J., Jit, M., Vassall, A., 2021. Models of COVID-19 vaccine priorit-  
1663 isation: a systematic literature search and narrative review. *BMC medicine*  
1664 19, 1–11.
- 1665 Salcedo-Varela, G.A., Peñuñuri, F., González-Sánchez, D., Díaz-Infante, S.,  
1666 2023. Synchronizing lockdown and vaccination policies for COVID-19: An  
1667 optimal control approach based on piecewise constant strategies. *Optimal*  
1668 *Control Applications and Methods* .
- 1669 Saldaña, J., Scoglio, C., 2022. Influence of heterogeneous age-group contact  
1670 patterns on critical vaccination rates for herd immunity to SARS-CoV-2.  
1671 *Scientific Reports* 12, 2640.
- 1672 Sallam, M., 2021. COVID-19 vaccine hesitancy worldwide: a concise system-  
1673 atic review of vaccine acceptance rates. *Vaccines* 9, 160.
- 1674 Sepulveda, G., Arenas, A.J., González-Parra, G., 2023. Mathematical model-  
1675 ing of COVID-19 dynamics under two vaccination doses and delay effects.  
1676 *Mathematics* 11, 369.
- 1677 Sheikh, A.B., Pal, S., Javed, N., Shekhar, R., 2021. COVID-19 vaccination in  
1678 developing nations: challenges and opportunities for innovation. *Infectious*  
1679 *disease reports* 13, 429–436.
- 1680 Shen, Z.H., Chu, Y.M., Khan, M.A., Muhammad, S., Al-Hartomy, O.A.,  
1681 Higazy, M., 2021. Mathematical modeling and optimal control of the  
1682 COVID-19 dynamics. *Results in Physics* 31, 105028.
- 1683 Shim, E., 2021. Optimal allocation of the limited COVID-19 vaccine supply  
1684 in South Korea. *Journal of clinical medicine* 10, 591.

- 1685 Soares, P., Rocha, J.V., Moniz, M., Gama, A., Laires, P.A., Pedro, A.R.,  
1686 Dias, S., Leite, A., Nunes, C., 2021. Factors associated with COVID-19  
1687 vaccine hesitancy. *Vaccines* 9, 300.
- 1688 Sorensen, R., Barber, R., Pigott, D., Carter, A., Spencer, C., Ostroff, S.,  
1689 Reiner, R., Abbafati, C., Adolph, C., Allorant, A., et al., 2022. Variation in  
1690 the COVID-19 infection-fatality ratio by age, time, and geography during  
1691 the pre-vaccine era: A systematic analysis. *The Lancet* 399, 1469–1488.
- 1692 Souto Ferreira, L., Canton, O., da Silva, R.L.P., Poloni, S., Sudbrack, V.,  
1693 Borges, M.E., Franco, C., Marquitti, F.M.D., de Moraes, J.C., Veras,  
1694 M.A.d.S.M., et al., 2022. Assessing the best time interval between doses in  
1695 a two-dose vaccination regimen to reduce the number of deaths in an ongo-  
1696 ing epidemic of SARS-CoV-2. *PLoS computational biology* 18, e1009978.
- 1697 Stafford, E., Dimitrov, D., Ceballos, R., Campelia, G., Matrajt, L., 2023.  
1698 Retrospective analysis of equity-based optimization for COVID-19 vaccine  
1699 allocation. *PNAS Nexus* 2, pgad283.
- 1700 Sudre, C.H., Murray, B., Varsavsky, T., Graham, M.S., Penfold, R.S.,  
1701 Bowyer, R.C., Pujol, J.C., Klaser, K., Antonelli, M., Canas, L.S., et al.,  
1702 2021. Attributes and predictors of long COVID. *Nature medicine* 27,  
1703 626–631.
- 1704 Sypsa, V., Roussos, S., Engeli, V., Paraskevis, D., Tsiodras, S., Hatzakis, A.,  
1705 2022. Trends in COVID-19 vaccination intent, determinants and reasons  
1706 for vaccine hesitancy: Results from repeated cross-sectional surveys in  
1707 the adult general population of Greece during November 2020–June 2021.  
1708 *Vaccines* 10, 470.
- 1709 Taboe, H.B., Asare-Baah, M., Iboi, E.A., Ngonghala, C.N., 2023. Critical  
1710 assessment of the impact of vaccine-type and immunity on the burden of  
1711 COVID-19. *Mathematical Biosciences* , 108981.
- 1712 Tatapudi, H., Das, R., Das, T.K., 2021. Impact of vaccine prioritization  
1713 strategies on mitigating COVID-19: an agent-based simulation study using  
1714 an urban region in the United States. *BMC medical research methodology*  
1715 21, 1–14.



- 1716 Thakkar, K., Spinardi, J.R., 2023. Impact of vaccination and non-  
1717 pharmacological interventions on covid-19: a review of simulation mod-  
1718 eling studies in asia. *Frontiers in Public Health* 11, 1252719.
- 1719 Tran, T.N.A., Wikle, N.B., Albert, E., Inam, H., Strong, E., Brinda, K.,  
1720 Leighow, S.M., Yang, F., Hossain, S., Pritchard, J.R., et al., 2021. Optimal  
1721 SARS-CoV-2 vaccine allocation using real-time attack-rate estimates in  
1722 Rhode Island and Massachusetts. *BMC medicine* 19, 1–14.
- 1723 Trejo, I., Hung, P.Y., Matrajt, L., 2024. Covid19Vaxplorer: a free, online,  
1724 user-friendly COVID-19 vaccine allocation comparison tool. *PLOS Global  
1725 Public Health* 4, e0002136.
- 1726 Tu, Y., Hayat, T., Hobiny, A., Meng, X., 2023. Modeling and multi-objective  
1727 optimal control of reaction-diffusion COVID-19 system due to vaccination  
1728 and patient isolation. *Applied Mathematical Modelling* 118, 556–591.
- 1729 Tuite, A.R., Zhu, L., Fisman, D.N., Salomon, J.A., 2021. Alternative dose al-  
1730 location strategies to increase benefits from constrained COVID-19 vaccine  
1731 supply. *Annals of internal medicine* .
- 1732 Vo, M., Feng, Z., Glasser, J.W., Clarke, K.E., Jones, J.N., 2023. Analysis of  
1733 metapopulation models of the transmission of SARS-CoV-2 in the United  
1734 States. *Journal of Mathematical Biology* 87, 24.
- 1735 Voysey, M., Clemens, S.A.C., Madhi, S.A., Weckx, L.Y., Folegatti, P.M.,  
1736 Aley, P.K., Angus, B., Baillie, V.L., Barnabas, S.L., Bhorat, Q.E., et al.,  
1737 2021. Single-dose administration and the influence of the timing of  
1738 the booster dose on immunogenicity and efficacy of ChAdOx1 nCoV-1  
1739 (AZD1222) vaccine: a pooled analysis of four randomised trials. *The  
1740 Lancet* 397, 881–891.
- 1741 Walker, J., Paul, P., Dooling, K., Oliver, S., Prasad, P., Steele, M.,  
1742 Gastañaduy, P.A., Johansson, M.A., Biggerstaff, M., Slayton, R.B., 2022.  
1743 Modeling strategies for the allocation of SARS-CoV-2 vaccines in the  
1744 United States. *Vaccine* 40, 2134–2139.
- 1745 Wang, R., Chen, J., Gao, K., Wei, G.W., 2021. Vaccine-escape and fast-  
1746 growing mutations in the United Kingdom, the United States, Singapore,  
1747 Spain, India, and other COVID-19-devastated countries. *Genomics* 113,  
1748 2158–2170.

- 1749 Wang, X., Wu, H., Tang, S., 2022. Assessing age-specific vaccination strate-  
1750 gies and post-vaccination reopening policies for COVID-19 control using  
1751 SEIR modeling approach. *Bulletin of Mathematical Biology* 84, 108.
- 1752 World Health Organization (WHO), . <https://data.who.int/dashboards/covid19/cases>  
1753 [accessed: 01/14/2024].
- 1754 Yasuda, H., Ito, F., Hanaki, K.i., Suzuki, K., 2022. COVID-19 pandemic  
1755 vaccination strategies of early 2021 based on behavioral differences between  
1756 residents of Tokyo and Osaka, Japan. *Archives of Public Health* 80, 180.
- 1757 Zaitri, M.A., Bibi, M.O., Torres, D.F., 2022. Transport and optimal control  
1758 of vaccination dynamics for COVID-19, in: *Mathematical Analysis of*  
1759 *Infectious Diseases*. Elsevier, pp. 27–39.
- 1760 Zanella, M., Bardelli, C., Dimarco, G., Deandrea, S., Perotti, P., Azzi, M.,  
1761 Figini, S., Toscani, G., 2021. A data-driven epidemic model with social  
1762 structure for understanding the COVID-19 infection on a heavily affected  
1763 Italian province. *Mathematical Models and Methods in Applied Sciences*  
1764 31, 2533–2570.
- 1765 Zavrakli, E., Parnell, A., Malone, D., Duffy, K., Dey, S., 2023. Optimal  
1766 age-specific vaccination control for COVID-19: An Irish case study. *PLOS*  
1767 *One* 18, e0290974.
- 1768 Zhang, C., Maruggi, G., Shan, H., Li, J., 2019. Advances in mRNA vaccines  
1769 for infectious diseases. *Frontiers in immunology* 10, 594.
- 1770 Zhang, J., Litvinova, M., Liang, Y., Wang, Y., Wang, W., Zhao, S., Wu, Q.,  
1771 Merler, S., Viboud, C., Vespignani, A., et al., 2020. Changes in contact  
1772 patterns shape the dynamics of the COVID-19 outbreak in China. *Science*  
1773 368, 1481–1486.
- 1774 Zhang, J., Litvinova, M., Liang, Y., Zheng, W., Shi, H., Vespignani, A., Vi-  
1775 boud, C., Ajelli, M., Yu, H., 2021. The impact of relaxing interventions on  
1776 human contact patterns and SARS-CoV-2 transmission in China. *Science*  
1777 *Advances* 7, eabe2584.
- 1778 Zhang, J., Wang, X., Rong, L., Pan, Q., Bao, C., Zheng, Q., 2024. Plan-  
1779 ning for the optimal vaccination sequence in the context of a population-  
1780 stratified model. *Socio-Economic Planning Sciences* , 101847.

- 1781 Zhao, S., Tang, B., Musa, S.S., Ma, S., Zhang, J., Zeng, M., Yun, Q., Guo,  
1782 W., Zheng, Y., Yang, Z., et al., 2021a. Estimating the generation interval  
1783 and inferring the latent period of COVID-19 from the contact tracing data.  
1784 *Epidemics* 36, 100482.
- 1785 Zhao, Z.y., Niu, Y., Luo, L., Hu, Q.q., Yang, T.l., Chu, M.j., Chen, Q.p.,  
1786 Lei, Z., Rui, J., Song, C.l., et al., 2021b. The optimal vaccination strategy  
1787 to control COVID-19: a modeling study in Wuhan City, China. *Infectious*  
1788 *diseases of poverty* 10, 48–73.
- 1789 Zhou, S., Zhou, S., Zheng, Z., Lu, J., 2021. Optimizing spatial allocation of  
1790 COVID-19 vaccine by agent-based spatiotemporal simulations. *GeoHealth*  
1791 5, e2021GH000427.
- 1792 Ziarelli, G., Parolini, N., Verani, M., Quarteroni, A., et al., 2023. Optimized  
1793 numerical solutions of SIRDVW multiage model controlling SARS-CoV-2  
1794 vaccine roll out: An application to the Italian scenario. *Infectious Disease*  
1795 *Modelling* .
- 1796 Zuo, C., Meng, Z., Zhu, F., Zheng, Y., Ling, Y., 2022. Assessing vacci-  
1797 nation prioritization strategies for COVID-19 in South Africa based on  
1798 age-specific compartment model. *Frontiers in public health* 10.