1	Single-cell RNA sequencing unraveled the expression heterogeneity of
2	hematopoietic stem and progenitor cells and lymphoid cell development
3	dysregulation in childhood asthma
4	Danying Zhu ^{*1,2} , Guang Li ^{*,3} , Lang Yuan ^{*,1} , Zeyu Zeng ^{1,2} , Na Dong ¹ , Chao Wang ¹ , Ming
5	Chen ¹ , Lijian Xie ⁵ , Guohui Ding ^{#,4} , Libing Shen ^{#,4} , Xiaoyan Dong ^{#,1,2}
6	
7	¹ Department of Pulmonology, Shanghai Children's Hospital, School of medicine, Shanghai
8	Jiao Tong University, Shanghai, China
9	² Institute of Pediatric Infection, Immunity, and Critical Care Medicine, Shanghai Jiao Tong
10	University School of Medicine, Shanghai, China.
11	³ Daozhi Precision Medicine Technology (Shanghai) Co., Ltd; Shanghai, China.
12	⁴ Institute for Digital Health, International Human Phenome Institutes (Shanghai), Shanghai,
13	China.
14	⁵ Department of Pediatrics, Jinshan Hospital, Fudan University, Shanghai, 201508, China.
15	
16	*These authors contributed equally: Danying Zhu, Guang Li, Lang Yuan
17	
18	#Corresponding authors
19	Xiaoyan Dong, Ph.D.
20	Department of Respiratory, Shanghai Children's Hospital, School of medicine, Shanghai Jiao
21	Tong University, Shanghai, China.
22	Institute of Pediatric Infection, Immunity, and Critical Care Medicine, Shanghai Jiao Tong
23	University School of Medicine, Shanghai, China.
24	No.355, Luding Road, Shanghai 200062, China.
25	Tel: 18930734642
26	E-mail: dongxy@shchildren.com.cn
27	
28	Libing Shen, Ph.D
29	Institute for Digital Health, International Human Phenome Institutes (Shanghai), Shanghai,
30	China.

- 31 12/F, No.258 Guoxia Road, Yangpu District, Shanghai, China.
- 32 Tel: 13482391370
- 33 E-mail: <u>shenlibing@ihup.org.cn</u>
- 34
- 35 Guohui Ding, *Ph.D.*
- 36 Institute for Digital Health, International Human Phenome Institutes (Shanghai), Shanghai,
- 37 China.
- 38 12/F, No.258 Guoxia Road, Yangpu District, Shanghai, China.
- 39 <u>Tel:17721236391</u>
- 40 Email: dingguohui@ihup.org.cn
- 41
- 42
- 43

44 Abstract

45 Asthma is a long-term inflammatory disease affecting airways and lungs with usual 46 onset in childhood. Its cause is not fully understood up to now. Here, using single-cell 47 RNA sequencing, we profile peripheral blood mononuclear cells (PBMCs) from three 48 pediatric patients with onset asthma and four age-matched healthy controls to 49 investigate the cellular etiology of childhood asthma. The overall expression features 50 among three asthma patients' PBMCs demonstrate that innate immunity is commonly 51 upregulated while adaptive immunity is commonly downregulated in childhood 52 asthma, but each patient has different molecular phenotypes. The analyses of the 53 expression profiles of hematopoietic stem and progenitor cells (HSPCs) further show 54 that the HSPCs of asthma patients have heterogeneous expression backgrounds with 55 more specific differentially expressed genes (DEGs) in each patient than common 56 DEGs and a common feature of low S100 protein binding gene expression. S100A8, 57 S100A9, S100A12, and RETN are universally upregulated in various cell types of 58 asthma patients. The cell developmental trajectories in three asthma cases exhibit an 59 abnormal immune cell development pattern compared to that in health control. The 60 dysregulated lymphoid lineage development is observed in all 3 patients, but there is 61 no identical abnormal pattern for each patient. The pseudo-time analyses of gene 62 expression show that the expression dynamics of two proto-oncogenes, JUN and SPI1, 63 and six inflammatory response related genes (S100A8, S100A9, S100A12, IL7R, 64 IL32, CCL5) are relevant to abnormal immune cell development in asthma patients. 65 The cell-cell communication analyses reveal the contribution of incoming annexin 66 signal towards dendritic cells and the outgoing resistin signal from dendritic cells to 67 asthma heterogeneity. Interestingly, the plasma blast cells of asthma patient 3 with 68 severe symptoms exhibit dual cell identities of both plasma blast cells and T cells. Our 69 scRNA-Seq analyses for three asthma patients reveal a complex cellular etiology for 70 childhood asthma and provide a new research direction for the comprehensive and 71 systematic understanding of key molecular mechanisms of childhood asthma.

72

74 Introduction

75 Asthma, a persistent respiratory condition, is marked by inflammation of the 76 airways, heightened bronchial responsiveness, and variable airway obstruction. This 77 complex disease involves a multitude of cellular participants, including eosinophils, T 78 and B lymphocytes, macrophages, neutrophils, and epithelial cells, which interact 79 with environmental factors to induce chronic inflammation, reversible airflow 80 limitations, and airway remodeling in patients¹. The involvement of various cell types and their mediators is a key aspect of asthma's pathogenesis². This diversity in asthma 81 82 phenotypes and the intricacy of its pathogenesis are further complicated by the fact 83 that respiratory infections and allergen exposures are known to trigger asthma exacerbations¹. While asthma can affect individuals across all age groups, the 84 85 manifestation, underlying pathophysiology, and response to treatment can vary significantly between children and adults. Recognizing the molecular and cellular 86 87 distinctions between childhood and adult asthma is crucial for crafting age-tailored 88 diagnostic and therapeutic approaches.

89 childhood asthma, in particular, presents with distinct clinical and immunological 90 characteristics compared to its adult counterpart. Asthma in children often manifests 91 as allergic phenotypes with a higher incidence of eosinophilic inflammation, whereas 92 adults may display a broader range of subtypes, including non-eosinophilic variants. 93 The immature and plastic nature of the immune system in children can lead to unique 94 cellular responses and immune dysregulation. Therefore, there are some differences in 95 the treatment by unique age groups³. Single-cell sequencing offers a powerful tool to 96 explore age-specific alterations in immune cell populations, their activation states, and 97 functional attributes, providing a deeper understanding of the mechanisms driving 98 childhood asthma.

99 Previous research has predominantly focused on the single-cell sequencing of adult asthma^{4, 5}, whereas this study applies single-cell technology to childhood asthma, 100 101 aiming to uncover the role of immune abnormalities in the pathogenesis of asthma in 102 children. Through single-cell sequencing, specific cell subpopulations can be 103 identified, and their roles in the inflammatory responses of children with asthma can 104 be understood, as well as their responses to current treatment methods. These findings 105 may help explain why childhood asthma exhibits differences in clinical presentation 106 and treatment response compared to adult asthma, and they provide new perspectives 107 for future research and clinical practice.

108 Method

109 Patients

110 During the initial hospital stay, before any treatment, we collected three fresh 111 peripheral blood samples from patients experiencing acute asthma. The diagnosis of 112 asthma was confirmed using the criteria outlined in the 2023 Global Initiative for 113 Asthma (GINA) report, which can be accessed at https://ginasthma.org/2023-gina-114 main-report/. Atopic status was determined by a total IgE level of 200 IU/ml or higher. 115 For single-cell RNA sequencing (scRNA-seq), we enrolled four healthy individuals 116 who were participating in routine physical examinations and had no recent history of 117 fever, infection, or immunization. Data from three of these individuals have 118 previously been published⁶. All participants were recruited from Shanghai Children's 119 Hospital between December 2019 and February 2023. Exclusion criteria for the study 120 included: Participants who were undergoing immunotherapy, such as anti-IgE 121 treatment, or had received such therapy within the last three months. Individuals with 122 significant abnormalities identified in their complete blood count. Subjects who were 123 currently enrolled in an asthma-related pharmaceutical or interventional study, or had 124 been part of such a study within the last four weeks. Patients with concurrent medical conditions necessitate systemic corticosteroids or other immunomodulatory 125 126 treatments. The study was approved by the Ethics Committee of Shanghai Children's 127 Hospital (Protocol Numbers: 2019R081, 2022R029-F-01). Informed consent was 128 obtained from all participants and their guardians.

129

130 Single-cell preparation and sequencing

131 A total of 2 milliliters of venous blood was collected from each participant using 132 tubes containing EDTA as an anticoagulant. These samples were processed within 133 four hours to preserve cell integrity. Peripheral blood mononuclear cells (PBMCs) 134 were separated from the collected blood using Ficoll-Paque medium and density 135 gradient centrifugation. The viability of the isolated PBMCs was assessed using 136 trypan blue staining, and only samples with more than 90% viable cells were selected 137 for further analysis. A calculated volume of the cell suspension, estimated to contain 138 around 12,000 cells per sample, was prepared to ensure an adequate cell count for 139 subsequent capture and sequencing processes. The Chromium Next GEM Single Cell

140 V(D)J Reagent Kits v1.1 from 10x Genomics were utilized for capturing single cells 141 and constructing the corresponding libraries. The cell suspension, along with 142 barcoded gel beads and partitioning oil, were loaded onto the $10 \times$ Genomics 143 Chromium Chip to create single-cell Gel Beads-in-Emulsion (GEMs). Inside each 144 GEM, cells were lysed, and their transcripts were barcoded through reverse 145 transcription. After reverse transcription, the cDNA, now carrying cell barcodes, 146 underwent PCR amplification to ensure a sufficient quantity for sequencing. The 147 construction of single-cell RNA sequencing (scRNA-seq) libraries was accomplished 148 using the 5' Library Kits. The prepared libraries were sequenced on an Illumina 149 NovaSeq platform, producing paired-end reads with a length of 2×150 base pairs. 150 This sequencing step generates the raw data necessary for subsequent data analysis.

151

152 scRNA-seq data analysis

153 The demultiplexed reads were then aligned to the GRCh38 reference genome 154 using the Cell Ranger software suite. This gene-barcode matrix was subsequently 155 analyzed using Seurat version 3.0.2 for various analytical steps, including quality 156 control, data normalization, dimensionality reduction, batch effect correction, 157 clustering, and data visualization. The specific procedures carried out with Seurat 158 included: Applying quality control criteria for the majority of samples, which 159 involved ensuring a total UMI count ranging from 200 to 6000 and limiting the 160 percentages of mitochondrial, hemoglobin, and ribosome genes to below 10%, 0.1%, 161 and 3%, respectively. Integrating samples collected during the initial phase of the 162 study (C1-C4) to eliminate batch effects using anchor. Scaling the integrated matrix 163 and employing the principal components derived from principal component analysis 164 (PCA) for uniform manifold approximation and projection (UMAP). Utilizing UMAP 165 to visualize cells in a two-dimensional space, highlighted their gene expression 166 pattern similarities. Conducting shared nearest neighbor graph-based clustering on the 167 PCA-reduced data to identify the primary cell types within the PBMCs. Examining 168 the expression of canonical marker genes to refine cell cluster annotations. 169 Determining cell identities through multimodal reference mapping with the 170 SeuratDisk package.

172 Differential expression, functional enrichment analysis, and pseudo-time analysis

173 The marker genes for each cluster were calculated with the FindAllMarkers function 174 within the Seurat package. The clusterProfiler package (version 3.16.0) was used for 175 function over-representation analysis of the differentially expressed genes (DEGs) 176 with a false discovery rate (FDR) threshold of <0.05. Gene Ontology (GO), KEGG 177 pathways, and hallmark gene sets from the MSigDB database (version 7.1) were used 178 as gene function databases to assess the enrichment of specific biological functions or 179 pathways among the DEGs. Pseudo-time analysis of cell differentiation trajectories 180 for each sample dataset was performed with R package Monocle 27. The expression 181 feature and inferred cell type for each sample dataset from the Seurat result were used 182 to annotate the cell dataset for the Monocle analysis pipeline. We used the Monocle 183 built-in approach named "dpFeature" to detect the variable genes that define a cell's 184 differentiation. Its advantages are needing no prior biological knowledge and 185 discovering important ordering genes from the data itself. Dimension reduction was 186 performed with 2 max components and the "DDRTree" method. Cell-cell 187 communication analysis was performed with the R package CellChat 1.6.1⁸.

188

189 Human serum resistin Enzyme-Linked Immunosorbent Assay (ELISA) analysis

EDTA-anticoagulated whole blood was transferred to the laboratory and
processed immediately after collection. Centrifuge samples for 15 minutes at 1000 x g
at 4 □ within 30 minutes of collection and stored at -80 □. We followed a human
resistin ELISA kit (Signalway Antibody, Maryland, USA, Catalog No: EK2351)
protocol to analyze peripheral blood serums.

195

196 **Results**

197 Characteristics of the study subjects for asthma patients and health controls

In our single-cell RNA analysis study, we included a cohort of 7 childhood subjects, comprising 3 children experiencing asthma exacerbations and 4 healthy individuals without asthma. T The mean age of the participants ranges from 1 to 9 years old. An additional group of 14 asthma patients and health controls ranging from 1 to 13 years old, were included in the study. Allergic testing in asthmatic children revealed sensitization to house dust mites. Allergic testing in asthmatic children revealed sensitization to house dust mites.

205

206 Single-cell RNA profiling of PBMCs in health and asthma patients

We collected the peripheral blood samples derived from 3 patients with onset asthma (A1-A3). The patients were diagnosed according to the GINA 2023 (https://ginasthma.org/2023-gina-main-report/). For each patient, the blood sample was taken on the first days before therapy. We also collected fresh peripheral blood samples from four age-matched healthy donors as controls (C1–C4).

212 We used the 10× Genomics platform for scRNA-seq of PBMCs isolated from the 213 samples. PBMCs were loaded onto the platform and about 5000 - 14000 cells per 214 sample could be recovered from the sequencing data. The total number of detected 215 cells passing quality control was 34,354 for 3 asthma patients and 22,398 cells from 216 healthy controls. Based on the scRNA-seq profiles, we clustered the cells across 217 samples with Seurat 4.0 and visualized them in two-dimensional space (Fig. 1a). The PBMCs in our study could be clustered into 12 major cell types including B cells, 218 219 CD4 T cells, CD8 T cells, CD14 monocytes (CD14 mono), CD16 monocytes (CD16 220 mono), dendritic cells (DC), mucosa-associated invariant T cells (MAIT), gamma-221 delta cells (gdT), hematopoietic stem and progenitor cells (HSPC), natural killer cells 222 (NK), plasma blast cells (Plasmablast), and platelets. Each cell type was verified with 223 PBMC multimodal reference object and canonical gene markers (Supplementary 224 Figure 1a and 1b). The percentage of major cell types for each sample is shown in 225 Figure 1b. The more detailed classification of cell types excluding CD4 T and CD8 T 226 cells shows that CD14 monocytes are significantly increased and CD16 monocytes 227 are significantly decreased in asthma patients (Figure 1c, *P*-value < 0.05). The 228 detailed classification of T cell types in each sample is shown in Supplementary 229 Figure 1c.

230

231 Expression features of all cells and HSPCs in asthma patients

We first compared the gene expression level of all cells in each asthma patient to controls. The overall differentially expressed genes (DEGs) were identified for each asthma patient. There are 174 upregulated genes in A1 (Asthma1), 29 upregulated genes in A2 (Asthma2), and 246 upregulated genes in A3 (Asthma3) (Figure 2a). There are 21 common upregulated genes among three asthma patients. GO analysis shows that these commonly upregulated genes are involved in the extracellular region,

238 secretion, and antimicrobial function (Figure 2b). GO result also shows that innate 239 immunity plays a major role in childhood asthma which manifests as a defense 240 response to bacterium and inflammatory response. There are 57 specific upregulated 241 genes in A1, 3 specific upregulated genes in A2, and 128 specifically upregulated 242 genes in A3. GO analyses of the specifically upregulated genes in A1 and A3 show 243 that these two patients have different molecular phenotypes (Supplemental Tables 1 244 and 2). A1 exhibits the specific expression of signal sequences (usually 20-30 amino 245 acids long) and positive regulation of T cell migration while A3 has an excessive 246 expression of immunoglobulin. 3 specific upregulated genes in A2 are IGHV3-73, 247 CDK6 and NELL2 which are not sufficient for GO analysis. We also find 33 248 downregulated genes in A1 (Asthma1), 80 downregulated genes in A2 (Asthma2), and 249 66 downregulated genes in A3 (Asthma3) (Figure 2c). There are 9 common 250 upregulated genes among three asthma patients. GO analysis shows that these 251 common downregulated genes are involved in adaptive immunity and T cell receptor 252 (Figure 2d). There are 13 specific downregulated genes in A1, 37 specific 253 downregulated genes in A2, and 20 specific downregulated genes in A3. A1 exhibits 254 the specific repression of the external side of the plasma membrane and B cell 255 receptor signaling pathway expression (Supplemental Table 3). A2 exhibits the 256 specific repression of the immunoglobulin complex (Supplemental Table 4). A3 257 exhibits the specific repression of the T cell receptor complex (Supplemental Table 5). 258

259 We further examined the expression features of HSPCs in asthma patients, because 260 the other types of cells are all derived from them. There are 176 upregulated HSPC 261 genes in A1 (Asthma1), 211 upregulated HSPC genes in A2 (Asthma2), and 199 262 HSPC upregulated genes in A3 (Asthma3) (Figure 3a). There are 81 common 263 upregulated HSPC genes among three asthma patients. GO analysis shows that these 264 commonly upregulated HSPC genes are mainly involved in the cellular components 265 of extracellular exosome and ER to Golgi transport vesicle membrane (Figure 3b). 266 There are 29 specifically upregulated HSPC genes in A1, 72 specifically upregulated 267 HSPC genes in A2, and 55 specifically upregulated HSPC genes in A3. No significant 268 GO result is found for 29 specifically upregulated HSPC genes in A1. HSPCs in A2 269 exhibit a specific expression of extracellular exosome related genes and HSPCs in A3 270 exhibit a specific expression of endoplasmic reticulum related genes (Supplemental 271 Tables 6 and 7). There are 179 downregulated HSPC genes found in A1 (Asthma1),

272 203 downregulated HSPC genes found in A2 (Asthma2), and 119 HSPC 273 downregulated genes found in A3 (Asthma3) (Figure 3c). GO analysis shows that 274 these common downregulated HSPC genes are mainly involved in S100 protein 275 binding (Figure 3d). There are 85 specifically downregulated HSPC genes in A1, 113 276 specifically downregulated HSPC genes in A2, and 53 specifically downregulated 277 HSPC genes in A3. HSPCs in A1 exhibit a specific repression of adaptive immunity 278 and T cell receptor complex expression (Supplemental Table 8). HSPCs in A2 exhibit 279 a specific repression of immune response related genes' expression (Supplemental 280 Table 9). HSPCs in A3 exhibit a specific repression of extracellular exosome and 281 immunity relate genes' expression (Supplemental Table 10). Interestingly, we find that 282 T cell receptor genes are repressed in all three asthma patients, although they are not 283 statically significant in the HSPCs of A2 and A3 (Supplemental Tables 9 and 10).

284

In expression feature analyses, we find that S100A8, S100A9, S100A12, and RETN are universally upregulated in various cell types of asthma patients, especially S100A8 and S100A9 (Figure 4). S100A8, S100A9, and S100A12 are known for participating in inflammatory response while RETN has antibacterial activity⁹⁻¹¹.

289

290 Single-cell trajectory reconstructions of PBMCs in health controls and 3 asthma

291 patients

292 The expression features of asthma patients' HSPCs demonstrate that the expression of 293 immunity and T cell receptor genes are abnormally repressed in them. Since HSPCs 294 give rise to the other cells, it is worth investigating the cell developmental states in 295 asthma patients. Single-cell trajectory analysis could help us reconstruct the cell 296 developmental paths of PBMCs based on featured variable genes. The analyses show 297 that the PBMCs in healthy controls can be divided into 5 developmental states while 298 the PBMCs in asthma patients have 7, 3, and 5 developmental states, differently 299 (Figure 5a, 5c, 5e, and 5g). All cell types are plotted on each trajectory in each sample 300 (Figure 5b, 5d, 5f, and 5h).

301

302 Based on the major cell types and numbers, state 1 in healthy controls can be 303 classified as B lineage, state 2 can be classified as monocyte lineage, state 3 is a 304 mixed lineage (B and T), state 4 can be classified as T lineage, and state 5 can be

305 classified as T-NK lineage (Figure 3b and Supplement Figure 2a). Myeloid lineage 306 (monocytes) and two lymphoid lineages (B cells and T cells) are clearly differentiated 307 in the PBMCs of healthy controls. However, the clear cell development patterns 308 disappear in 3 asthma patients. Each patient exhibits unique cell developmental 309 patterns different from healthy controls. A1 has 7 states and 4 of them are T lineages 310 according to the major cell numbers (state 1, 2, 3, and 6, Figure 5d and Supplement 311 Figure 2b). A2 has only 3 states and a part of B cells are mixed with T cells (state 1, 312 Figure 5f and Supplement Figure 2c). A3 has 5 states, but B cells are mixed with T 313 cells in state 3 (state 1, Figure 5h and Supplement Figure 2d). Although different 314 childhood asthma patient has a different cell developmental state, abnormal lymphoid 315 lineage development is observed in all 3 patients (multiple T lineage states or no clear 316 B lineage state). This result is consistent with the expression feature analyses showing 317 that adaptive immunity is repressed in asthma patients. T cells and B cells are a major 318 constitution of adaptive immune system and their poor development naturally leads to 319 insufficient the adaptive immunity.

320

321 Pseudo-time expression dynamic analyses of proto-oncogenes and inflammatory

322 response related genes

323 The abnormal cell developmental trajectories in asthma patients should be 324 accompanied by abnormal gene expression patterns. JUN gene is a transcription factor 325 regulated by various extracellular stimuli including peptide growth factors, proinflammatory cytokines, and even UV irradiation¹². SPI1 gene is also a transcription 326 factor regulating gene expression during myeloid and B-lymphoid cell development¹³. 327 Both of them are proto-oncogenes essential for cellular differentiation^{14, 15}. We further 328 329 include six inflammatory response related genes in pseudo-time expression dynamic 330 analyses. Their express dynamics were plotted along cell developmental trajectories 331 in both health control and three asthma patients.

332

In health controls, we observed the onset expression of JUN in the early pseudo-time state and a peak expression of six inflammatory response related genes in the earlymiddle pseudo-time state. SPI1 is expressed in the late pseudo-time state (Figure 6a). In A1 and A2, the expression dynamics of JUN, SPI1, and six inflammatory response related genes are almost the same (Figure 6b and 6c). JUN's onset expression is

338 repressed while SPI1 exhibits an onset expression in the early pseudo-time state. 339 S100A8, S100A9, and S100A12 show an early onset expression rather than a peak 340 expression in the early-middle pseudo-time state. IL7R, IL32, and CCL5 start to 341 express in the early-middle pseudo-time state as health control, but they show a 342 persistent expression level instead of an obvious expression peak. Notably, IL7R, 343 IL32, and CCL5 all participate in the biological process of negative regulation of the 344 T cell apoptotic process (GO:0070233). It implies that the T cells have prolonged cell 345 survival in A1 and A2. A3 shows a different expression dynamic pattern from health 346 control, A1, and A2 (Figure 6d). In A3, JUN is also repressed while the expression 347 patterns of IL7R, IL32, and CCL5 are more similar to health control. S100A8, 348 S100A9, S100A12, and SPI1 show a persistent expression level without decline. 349 Since SPI1 activates gene expression during myeloid and B-lymphoid cell 350 development and S100 protein family plays a role in cell growth and differentiation, their persistent expression pattern indicates an abnormal lymphoid cell development 351 352 fate in A3.

353

354 DCs function as a signaling hub in cell communication and play a role in asthma 355 heterogeneity

356 We performed cell communication analyses for both health control and asthma 357 patients to investigate their cell-cell interactions. The number of interactions and 358 interaction strength for health control and three asthma patients are shown in Figure 7. 359 Generally, asthma patients have a weaker interaction strength than health control and 360 A3 has the smallest number of interactions and the weakest interaction strength 361 among four samples. Among the major immune cell types, we find that dendritic cells 362 largest number of both incoming and outgoing interaction have the 363 numbers(Supplemental Figure 3a, 3b, 3c, and 3d). DCs are known to be the messenger cells communicating between innate and adaptive immune systems¹⁶. They 364 365 are also known as the antigen presenting cells which interact with T cells and B cells to activate and regulate the adaptive immune response¹⁷. Although DCs don't have the 366 367 strongest interaction strength among the major immune cell types (Supplemental 368 Figure 4a, 4b, 4c, and 4d), they surely function as a signaling hub in cell 369 communication based on cell communication analyses.

370

371 We further examined the DCs' signaling differences between health control and

372 asthma patients. Using health control as a signaling background, the signaling 373 changes were detected in three asthma patients. The DCs of A1 has a specific 374 incoming signal of annexin (Supplemental Figure 5a). The DCs of A2 has a specific 375 outgoing signal of resistin (Supplemental Figure 5b). The DCs of A3 has a specific 376 incoming signal of annexin and a specific outgoing signal of resistin (Supplemental 377 Figure 5c). Annexin A1, A2 and A5 have a role in the regulation of inflammation activation and may serve as the potential biomarkers for asthma¹⁸⁻²⁰. They all show an 378 379 elevated expression tendency in asthma cases (Supplemental Figure 6). Resistin also serves a potential biomarker for asthma and is related to inflammation as well^{21, 22}. 380 381 Our results show that annexin and resistin are actually two opposite directional signals 382 for the DCs of asthma patients. One of them is a sufficient predictor of asthma and 383 both of them indicate a severe symptom of asthma, since A3 is an acute severe asthma 384 patient. Both annexin and resistin show a self-reinforce trend in the CD14 monocytes 385 of asthma patients (Figure 8a and 8b). The upregulated signaling interactions in 386 asthma are mainly centered on the molecular component of external side of plasma 387 membrane and cell adhesion molecules (GO:0009897 and hsa04514, Supplemental 388 Figure 7a, 7b, and 7c). Furthermore, we found that CLEC2B-KLRB1 and LGALS9-389 CD44 signaling interactions are downregulated in all three asthma patients 390 (Supplemental Figure 8a, 8b, and 8c). CLEC2B-KLRB1 interaction mediates the activation of NK cells and monocytes²³. LGALS9-CD44 interaction enhances the 391 stability and function of adaptive regulatory T cells²⁴. The down-regulated CLEC2B-392 393 KLRB1 and LGALS9-CD44 interactions proposes a weakened functions of NK cells, 394 monocytes and T cells in asthma patients.

395

396 Plasma blast cells in A3 express T cell receptor genes

Patient A3, a severe asthma sufferer who has been admitted to the pediatric intensive
care unit, exhibits a unique group of cells not present in healthy controls A1 and A2
(Figure 1a). This group predominantly consists of plasma blast cells along with a
mixed cluster comprising plasma blast cells (35.2%), erythrocytes (26.5%), CD14
monocytes (14.4%), CD4 T cells (11.2%), CD8 T cells (7.7%), and other cells (5%).

402

Upon comparing the gene expression profile of A3's plasma blast cells with those of
the healthy controls, A1 and A2, we discovered an unexpected finding: the plasma
blast cells in A3 express a range of T cell receptor genes, including TRBV20-1, in

addition to immunoglobulins (Figure 9a). Further investigation into the expression
levels of classical markers for plasma blast cells (CD38, CD19, CD27), CD4 T cells
(CD8A, CD27), CD8 T cells (CD40LG, CD27), and NK cells (GNLY, NKG7)
revealed through a violin plot that all these markers are indeed expressed in A3's
plasma blast cells (Figure 9b). This suggests that the plasma blast cells in A3 likely
possess dual cellular identities, exhibiting characteristics of both plasma blast cells
and T cells, which accounts for the expression of T cell receptor genes.

413

414 The expression level of RETN in lymphoid linage cells

The feature plot shows that the RETN expression is not evenly distributed among different cell clusters. CD14 mono, CD16 mono, and dendritic cells have an especially high expression level of RETN. The violin plot further shows that RETN expression level in these four clusters is always higher in 3 asthma cases than in healthy controls. RETN encodes resistin, then we tested the level of resistin in plasma between children with asthma and healthy controls, we found that resistin in the serum increased significantly in the asthma group than in controls (Figure 10).

422

423 Discussions

In individuals with allergic airway disease, such as allergic asthma, the airways are already sensitized to specific allergens. This means that the immune system has developed an exaggerated response to these allergens, leading to chronic inflammation and airway hyperresponsiveness. During an asthma exacerbation, this preexisting allergic airway disease is further aggravated by additional inflammatory stimuli or triggers. Nevertheless, the etiology of childhood asthma is so far poorly understood²⁵.

431

432 The analyses of PBMCs from three childhood asthma patients show that childhood 433 asthma is a complex disease with distinct individualities at the cellular level. The 434 overall expression features among three asthma patients' PBMCs demonstrate that 435 innate immunity is commonly upregulated while adaptive immunity is commonly 436 downregulated in childhood asthma, but each patient has different molecular 437 phenotypes. The common upregulated genes in the HSPCs of asthma patient are 438 related to the cellular components of extracellular exosome and ER to Golgi transport 439 vesicle membrane and the common downregulated genes in the HSPCs of asthma

440 patients are related to S100 protein binding. The specific downregulated genes in the 441 HSPCs of each asthma patients converge on immunity and T cell receptor related biological entities. Because HSPCs give the rise to the other cells, they are a cellular 442 443 source of immunity. The expression abnormality in HSPCs would surely influence 444 their descendant cells. The downregulation of S100 protein binding in the HSPCs 445 indicates that S100 proteins are in an active form in the HSPCs of asthma patients. 446 The downregulation of immunity related genes in HSPCs is a sign of hypo-immune 447 tendency started from the cellular source in asthma cases. The active S100 protein and 448 hyo-immune tendency in the HSPCs of asthma patients seem to lead to over innate 449 immunity and low adaptive immunity for all cells of asthma patients. Although 450 childhood asthma has common pathophysiological features at the cellular level, there 451 are more specific differentially expressed genes (DEGs) in each patient than common 452 DEGs, which proposes that the genetic causes of asthma vary across different children 453 or different allergens can trigger different immune genes in a pediatric patient's 454 immune system. These two explanations are mutually inclusive. Future studies 455 includes bone marrow samples might give us much clearer views for this question, 456 since HSPCs are generated from bone marrow.

457

458 The single-cell trajectory analyses further demonstrate the cell developmental 459 heterogeneities among asthma patients. Different asthma patient has different 460 developmental states. The abnormal lymphoid lineage development is observed in all 3 patients. A1 has multiple T cell states while A2 and A3 have a poorly developed 461 462 lymphoid state of mixed B and T cells. The expression of T cell receptor genes is 463 repressed in the HSPCs of all three asthma patients, although this repression is not 464 statically significant in A2 and A3. It could partly explain the abnormal cell 465 developmental states in asthma patients and the developmental heterogeneities among 466 them. The delayed expression of T cell receptor genes could lead to the HSPC's 467 descendant cells having difficulty receiving the lymphoid development signals. In A1, 468 such deficiency is responded to the development of multiple T cell states by A1's 469 immune system, which seems to compensate for the lack of a proper T cell 470 developmental state. In A2 and A3, the delayed expression of T cell receptor genes is 471 less severe than that in A1, which leads to the underdevelopment of lymphoid lineage. 472 The abnormal single-cell developmental trajectories in asthma patients are also 473 accompanied by the key genes' aberrant expression dynamics. JUN and SPI1 are two

proto-oncogenes, both of which have multiple functions in cell fate decisions^{15, 26}. The 474 475 expression of JUN is repressed in the early stage of cell trajectories for all three 476 asthma patients. It indicates that the cell developmental fate of asthma patients has 477 trouble at the very beginning, which is consistent with the expression abnormality in 478 the HSPCs of asthma patients. In A1 and A2, the repression of JUN seems to be 479 replaced by the expression of SPI1. In A3, the beginning of the expression of SPI1 is 480 consistent with that in health control, but the expression of SPI1 shows no decline like 481 in health control. It implies that the developmental fates of distal cells in A3 could be 482 influenced by SPI1.

483

484 The cell-cell communication analyses further revealed the possible mechanisms 485 underlying the etiology of asthma. Although the number of cell-cell interactions is 486 similar in health control and asthma cases, the strength of cell-cell interactions is 487 much weaker in asthma patients, especially in A3. The overall weak signaling strength 488 is a clear sign of cell-cell communication disorder in asthma. It is in accordance with 489 the abnormal cell developmental trajectories of asthma. DCs induce primary immune 490 responses by processing and presenting antigen material on the cell surface of T 491 cells²⁷. They function as the signaling hub communicating between innate and 492 adaptive immune systems. Two aberrant signals, annexin, and resistin, are found for 493 the DCs of asthma, both of which play a part in inflammatory response. Inflammation 494 is a generic immune response and is considered as a mechanism of innate immunity. It 495 explains why innate immunity is hyperactive in asthma patients. In our cases, each 496 asthma patient has his or her aberrant signaling pattern, annexin for A1, resistin for A2, 497 and both annexin and resistin for A3. It further explains the heterogeneities among 498 asthma patients. Resistin as an outgoing signal from DCs has been reported as a predictor of asthma²¹. Our ELISA results show that its expression can be detected in 499 500 12 enlisted childhood asthma patients.

501

A3 is a special case in our study for the patient with severe symptoms compared to A1 and A2 whose symptoms are mild. Severe asthma means strong inflammation in patient's airway which further causes breathlessness and continuous cough. The profiling of A3's PBMCs first presents the patient's severity in our study. A3 has two extra cell clusters compared with health control, A1 and A2. These two extra cell clusters are annotated as mixed cells and plasma blast cells. The UMAP plot shows

that the mixed cells and plasma blast cells in A3 are stemmed from CD4 T cells. The expression features of A3's plasma blast cells demonstrate that they express T cell receptor genes as well. It is noticed that SPI1 has a persistent expression dynamic in A3. That A3's plasma blast cells show a duality of T cell identity could be linked to the SPI1's aberrant expression pattern. SPI1 might influence the developmental fate of plasma blast cells in A3 and further exacerbate the patient's symptoms.

514

515 Conclusions

516 Our research substantiates that childhood asthma is a multifaceted disease 517 characterized by an upregulation of the innate immune system and a downregulation 518 of the adaptive immune response. Single-cell RNA profiling of PBMCs from asthma 519 patients has uncovered a variety of expression patterns among HSPCs, along with 520 aberrant cell developmental trajectories and diminished cell-cell interaction signaling. 521 While significant heterogeneity exists among different cases of asthma, the 522 pathological features converge on certain commonalities, particularly the abnormal 523 development of the lymphoid lineage and an overabundance of expression in genes 524 related to \$100 protein binding and immune response. Notably, \$100A8, \$100A9, 525 S100A12, and RETN, along with ANXA1, ANXA2, and ANXA5, have emerged as 526 pivotal molecular markers in asthma, directly implicated in the inflammatory response.

527

528 Contributions

529 Xiaoyan Dong and Guohui Ding designed the study. Danying Zhu wrote the 530 manuscript. Libing Shen analysis the data. Zeyu Zeng did literature search. Lang Yuan 531 and Na Dong contributed clinical samples. Chao Wang and Ming Chen collected the 532 clincal data. Lijian Xie and Xiaoyan Dong funded the single cell sequencing.

533

534 **Declaration of interests**

535 The authors declare no competing interests.

536

537 Acknowledgements

538 This study was funded by Jinshan Distinct key medical specialty project 539 (JSZK2023A04) and Institute of Pediatric Infection, Immunity, and Critical Care

540 Medicine Cultivate Research Project.

541

542 Data Sharing Statement

- 543 All data generated or analyzed during this study are included in this article. Further
- 544 enquiries can be directed to the corresponding author.

546 **References**

- 547 1. Tang W, Li M, Teng F, Cui J, Dong J, Wang W. Single-cell RNA-sequencing in asthma research.
- 548 Front Immunol. 2022;13:988573. doi:10.3389/fimmu.2022.988573
- 549 2. Victor JR, Lezmi G, Leite-de-Moraes M. New Insights into Asthma Inflammation: Focus on iNKT,
- 550 MAIT, and gammadeltaT Cells. Clin Rev Allergy Immunol. Dec 2020;59(3):371-381.
- 551 doi:10.1007/s12016-020-08784-8
- 552 3. Reddel HK, Bacharier LB, Bateman ED, et al. Global Initiative for Asthma Strategy 2021:
- Executive Summary and Rationale for Key Changes. Am J Respir Crit Care Med. Jan 1
 2022;205(1):17-35. doi:10.1164/rccm.202109-2205PP
- 4. Alladina J, Smith NP, Kooistra T, et al. A human model of asthma exacerbation reveals
 transcriptional programs and cell circuits specific to allergic asthma. *Sci Immunol.* May 12
 2023;8(83):eabq6352. doi:10.1126/sciimmunol.abq6352
- 5. Li H, Wang H, Sokulsky L, et al. Single-cell transcriptomic analysis reveals key immune cell
 phenotypes in the lungs of patients with asthma exacerbation. *J Allergy Clin Immunol*. Mar
 2021;147(3):941-954. doi:10.1016/j.jaci.2020.09.032
- 561 6. Wang Z, Xie L, Ding G, et al. Single-cell RNA sequencing of peripheral blood mononuclear cells
 562 from acute Kawasaki disease patients. *Nat Commun.* Sep 14 2021;12(1):5444. doi:10.1038/s41467563 021-25771-5
- 564 7. Qiu X, Mao Q, Tang Y, et al. Reversed graph embedding resolves complex single-cell trajectories.
 565 *Nat Methods*. Oct 2017;14(10):979-982. doi:10.1038/nmeth.4402
- 566 8. Jin S, Guerrero-Juarez CF, Zhang L, et al. Inference and analysis of cell-cell communication using
- 567 CellChat. Nat Commun. Feb 17 2021;12(1):1088. doi:10.1038/s41467-021-21246-9
- 568 9. Shah RD, Xue C, Zhang H, et al. Expression of Calgranulin Genes S100A8, S100A9 and
- 569 S100A12 Is Modulated by n-3 PUFA during Inflammation in Adipose Tissue and Mononuclear Cells.
- 570 PLoS One. 2017;12(1):e0169614. doi:10.1371/journal.pone.0169614
- 571 10. Harris TA, Gattu S, Propheter DC, et al. Resistin-like Molecule alpha Provides Vitamin-A-
- 572 Dependent Antimicrobial Protection in the Skin. Cell Host Microbe. Jun 12 2019;25(6):777-788 e8.
- 573 doi:10.1016/j.chom.2019.04.004
- 574 11. Propheter DC, Chara AL, Harris TA, Ruhn KA, Hooper LV. Resistin-like molecule beta is a
- 575 bactericidal protein that promotes spatial segregation of the microbiota and the colonic epithelium.

- 576 Proc Natl Acad Sci U S A. Oct 17 2017;114(42):11027-11033. doi:10.1073/pnas.1711395114
- 577 12. Wisdom R, Johnson RS, Moore C. c-Jun regulates cell cycle progression and apoptosis by distinct
- 578 mechanisms. *Embo j.* Jan 4 1999;18(1):188-97. doi:10.1093/emboj/18.1.188
- 579 13. Burda P, Laslo P, Stopka T. The role of PU.1 and GATA-1 transcription factors during normal and
- 580 leukemogenic hematopoiesis. *Leukemia*. Jul 2010;24(7):1249-57. doi:10.1038/leu.2010.104
- 581 14. Angel P, Hattori K, Smeal T, Karin M. The jun proto-oncogene is positively autoregulated by its
- 582 product, Jun/AP-1. Cell. Dec 2 1988;55(5):875-85. doi:10.1016/0092-8674(88)90143-2
- 583 15. Ray D, Culine S, Tavitain A, Moreau-Gachelin F. The human homologue of the putative proto-
- oncogene Spi-1: characterization and expression in tumors. *Oncogene*. May 1990;5(5):663-8.
- 585 16. Monga I, Kaur K, Dhanda SK. Revisiting hematopoiesis: applications of the bulk and single-cell
- 586 transcriptomics dissecting transcriptional heterogeneity in hematopoietic stem cells. Brief Funct
- 587 Genomics. May 21 2022;21(3):159-176. doi:10.1093/bfgp/elac002
- 588 17. Banchereau J, Steinman RM. Dendritic cells and the control of immunity. Nature. Mar 19
- 589 1998;392(6673):245-52. doi:10.1038/32588
- 590 18. Lee SH, Lee PH, Kim BG, et al. Annexin A1 in plasma from patients with bronchial asthma: its
- 591 association with lung function. BMC Pulm Med. Jan 4 2018;18(1):1. doi:10.1186/s12890-017-0557-5
- 592 19. Dallacasagrande V, Hajjar KA. Annexin A2 in Inflammation and Host Defense. Cells. Jun 19
- 593 2020;9(6)doi:10.3390/cells9061499
- 20. Lee SH, Lee PH, Kim BG, Hong J, Jang AS. Annexin A5 Protein as a Potential Biomarker for the
- 595 Diagnosis of Asthma. Lung. Dec 2018;196(6):681-689. doi:10.1007/s00408-018-0159-x
- 596 21. Ballantyne D, Scott H, MacDonald-Wicks L, Gibson PG, Wood LG. Resistin is a predictor of
- 597 asthma risk and resistin: adiponectin ratio is a negative predictor of lung function in asthma. *Clin Exp*
- 598 *Allergy*. Aug 2016;46(8):1056-65. doi:10.1111/cea.12742
- 599 22. Małyszko J, Małyszko JS, Pawlak K, Mysliwiec M. Resistin, a new adipokine, is related to
 600 inflammation and renal function in kidney allograft recipients. *Transplant Proc.* Dec
 601 2006;38(10):3434-6. doi:10.1016/j.transproceed.2006.10.140
- 602 23. Welte S, Kuttruff S, Waldhauer I, Steinle A. Mutual activation of natural killer cells and
 603 monocytes mediated by NKp80-AICL interaction. *Nat Immunol.* Dec 2006;7(12):1334-42.
 604 doi:10.1038/ni1402
- 605 24. Wu C, Thalhamer T, Franca RF, et al. Galectin-9-CD44 interaction enhances stability and function

- 606 of adaptive regulatory T cells. Immunity. Aug 21 2014;41(2):270-82.
- 607 doi:10.1016/j.immuni.2014.06.011
- 608 25. Martinez FD. Genes, environments, development and asthma: a reappraisal. Eur Respir J. Jan
- 609 2007;29(1):179-84. doi:10.1183/09031936.00087906
- 610 26. Vogt PK. Fortuitous convergences: the beginnings of JUN. Nat Rev Cancer. Jun 2002;2(6):465-9.
- 611 doi:10.1038/nrc818
- 612 27. Kelly RR. Microcomputers in education: age of romance, or age of reason? Am Ann Deaf. Sep
- 613 1982;127(5):693-7. doi:10.1353/aad.2012.1147
- 614





616

Figure 1. Single-cell profiling of PBMCs in six samples. A. The integration singlecell profiling analysis of six samples including 4 healthy controls, asthma patient 1, asthma patient 2, and asthma patient 3. B. The percentage bar chart shows the continent of different cell types in each sample. C. The percentage bar chart shows the different cell types except T cells in each sample.

622



624

Figure 2. The analysis of upregulated and downregulated genes in asthma patients. A. Venn gram of the upregulated genes in 3 asthma patients. B. GO term enrichment analysis of the specific upregulated genes in 3 asthma patients. C. Venn gram of the downregulated genes in 3 asthma patients. D. GO term enrichment analysis of the specific downregulated genes in 3 asthma patients.



Figure 3. The analysis of upregulated and downregulated genes in asthma patients'
HSPC. A. Venn gram of the upregulated genes in 3 asthma patients' HSPC. B. GO
term enrichment analysis of the specific upregulated genes in 3 asthma patients'
HSPC. C. Venn gram of the downregulated genes in 3 asthma patients' HSPC. D. GO
term enrichment analysis of the specific downregulated genes in 3 asthma patients'
HSPC. C. Venn gram of the specific downregulated genes in 3 asthma patients'
HSPC. D. GO
term enrichment analysis of the specific downregulated genes in 3 asthma patients'



Figure 4. The violin plot of S100A9, S100A8, S100A12 and RETN gene expression
levels across different cell types for four sample groups.



641 Figure 5. Pseudo-time analysis of all cells in four sample groups. A. The 642 differentiation trajectory of all cells in healthy controls by state. B. The differentiation 643 trajectory of all cells in healthy controls by cell types. C. The differentiation trajectory 644 of all cells in asthma patient 1 (Ast1) by state. D. The differentiation trajectory of all 645 cells in asthma patient 1 (Ast1) by cell types. E. The differentiation trajectory of all 646 cells in asthma patient 2 (Ast2) by state. F. The differentiation trajectory of all cells in 647 asthma patient 2 (Ast2) by cell types. G. The differentiation trajectory of all cells in 648 asthma patient 3 (Ast3) by state. H. The differentiation trajectory of all cells in asthma 649 patient 3 (Ast3) by cell types.



Figure 6. Pseudo-time analyses of expression dynamics of two cell-fate and six inflammation related genes in four sample groups. A. Expression dynamics two cellfate and six inflammation related genes in health control. B. Expression dynamics two cell-fate and six inflammation related genes in health control in asthma patient 1 (Ast1). C. Expression dynamics two cell-fate and six inflammation related genes in health control in asthma patient 2 (Ast2). D. Expression dynamics two cell-fate and six inflammation related genes in health control in asthma patient 3 (Ast3).



Figure 7. The number of cell-cell interactions and interaction strength for four sample

659 groups.



Figure 8. The annexin and resistin signaling pathway network of four sample groups.
The annexin signaling pathway network is shown for health control, asthma patient 1
(Ast1) and asthma patient 3 (Ast3). The resistin signaling pathway network is shown
for health control, asthma patient 1 (Ast1) and asthma patient 2 (Ast2).



666 Figure 9. The dual cell identity of the plasma blast cells in asthma patient 3. A. The 667 expression heatmap of immunoglobulin (IGLV or IGHV) genes and T cell receptor 668 (TRBV or TRAV) genes in the plasma blast cells of four sample groups. B. The violin 669 plot of CD38, CD19, CD27, CD8A, CD40LG, GNLY and NKG7 gene expression 670 levels across CD4+ T cell, CD8+ T cell, NK cell and plasma blast cells among four 671 sample groups. CD38 and CD19 for the markers for plasma blast cells. CD27 is the 672 marker for both plasma blast cells and T cells. CD8A, CD40LG, GNLY and NKG7 673 are the markers for T cells.



- **Figure 10.** The expression of RETN gene in four sample groups. The level of serum
- 675 resistin between asthma patients and health controls(***p < 0.001).
- 676

677 Supplemental Figures



Supplemental Figure 1. Integrated single-cell profiling of PBMCs for four sample groups. A. The predicted cell types for the integrated single-cell profiling of PBMCs for four sample groups B. Expression of canonical gene markers for each cell type in four sample groups based on integration analysis. C. The detailed classification and proportion of T cells for all seven samples in our study.



Supplemental Figure 2. Pseudo-time analysis of all cells in four sample groups. A. The proportion of major cell types in each state for healthy controls. B. The proportion of major cell type in each state for asthma patient 1 (Ast1). C. The proportion of major cell type in each state for asthma patient 2 (Ast2). D. The proportion of major cell type in each state for asthma patient 3 (Ast3).



Supplemental Figure 3. The number of incoming and outgoing interactions for each cell type in four sample groups. A. The number of incoming and outgoing interactions in health control. B. The number of incoming and outgoing interactions in asthma patient 1 (Ast1). C. The number of incoming and outgoing interactions in asthma patient 2 (Ast2). D. The number of incoming and outgoing interactions in asthma patient 3 (Ast3).



Supplemental Figure 4. The number of incoming and outgoing interaction strength for each cell type in four sample groups. A. The number of incoming and outgoing interaction strength in health control. B. The number of incoming and outgoing interaction strength in asthma patient 1 (Ast1). C. The number of incoming and outgoing interaction strength in asthma patient 2 (Ast2). D. The number of incoming and outgoing interaction strength in asthma patient 3 (Ast3).



Supplemental Figure 5. The signaling changes of DCs in three asthma patients
(health control as the backgroup). A. The signaling changes of DCs in asthma patient
1 (Ast1). B. The signaling changes of DCs in asthma patient 2 (Ast2). C. The
signaling changes of DCs in asthma patient 3 (Ast3).



Supplemental Figure 6. The violin plot of ANXA1, ANXA2 and ANXA2 gene
expression levels across different cell types for four sample groups.



711 **Supplemental Figure 7.** The upregulated signaling from DCs to the other cell types

712 in three asthma patients. A. The upregulated signaling from DCs to the other cell

types in asthma patient 1 (Ast1). B. The upregulated signaling from DCs to the other

cell types in asthma patient 2 (Ast2). C. The upregulated signaling from DCs to the

715 other cell types in asthma patient 3 (Ast3).



~ ^ ~

- 718 Supplemental Figure 8. The downregulated signaling from DCs to the other cell
- 719 types in three asthma patients. A. The downregulated signaling from DCs to the other
- cell types in asthma patient 1 (Ast1). B. The downregulated signaling from DCs to the
- 721 other cell types in asthma patient 2 (Ast2). C. The downregulated signaling from DCs
- to the other cell types in asthma patient 3 (Ast3).