| 1 | |
|----|----------------------------------------------------------------------------------------------------------------------------------------|
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| | |
| 6 | |
| 7 | Patient Preferences for Diagnostic Imaging |
| 8 | Services: Decentralize or not? |
| 9 | |
| 10 | Decentralizing Imaging Services |
| 11 | |
| 12 | |
| 13 | |
| 14 | Eline M. van den Broek-Altenburg, Ph.D. ¹ *; Jamie S. Benson, BA ² ; Adam J. Atherly, Ph.D. ³ ; |
| 15 | Kristen K. DeStigter, MD ¹ |
| 16 | |
| 17 | ¹ University of Vermont, Larner College of Medicine |
| 18 | ² University of Pennsylvania, Perelman School of Medicine |
| 19 | ³ Virginia Commonwealth University, College of Health Professions |
| 20 | |
| 21 | *Corresponding author |
| 22 | 111 Colchester Avenue |
| 23 | Burlington, VT 05401 |
| 24 | <u>Eline.altenburg@med.uvm.edu</u> |
| 25 | |
| 26 | |
| 27 | |
| 28 | |
| 29 | NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice. |

30 Abstract (250)

- 31 The objective of this study was to identify patient preferences for outpatient diagnostic
- 32 imaging services and analyze how patients make trade-offs between attributes of these
- 33 services using a discrete choice experiment (DCE).
- 34 We used a DCE with 14 choice questions asking which imaging locations patients would
- 35 prefer. We used latent class analysis to analyze preference heterogeneity between different
- 36 patient groups and to estimate the relative value they assign to different attributes of
- 37 imaging services.
- 38 Our analysis showed that the "Experienced Patients" subgroup generally value diagnostic
- 39 imaging services in both acute and chronic situations and had a strong preference for
- 40 hospital outpatient radiology departments (HORD) that would provide services at lower
- 41 costs, where their images would be interpreted by a specialty radiologist, the clinic would be
- 42 recommended by their PCP, online scheduling would be available, service rating were higher,
- 43 and travel and wait times would be shorter. New Patients significantly valued the service
- 44 rating of the hospital outpatient radiology departments (HORD) and online scheduling.
- 45 HORDs can be more competitive by providing services that live up to expectations better
- 46 than available retail radiology clinics (RRCs). Most RRCs do not currently offer online
- 47 scheduling so ease of use may also steer patients towards HORDs. HORDs have the
- 48 advantage of being linked to the main medical center which has the reputation of having
- 49 clinical expertise and more sophisticated technology. We conclude that there is room for
- 50 medical centers to build HORDs that provide an appealing and competitive alternative to
- 51 current RRC.
- 52
- 53
- 54

55 Introduction

An important challenge for many hospitals and clinics is addressing the escalating diagnostic imaging demands for hospital outpatient radiology departments (HORDs)¹. HORDs have struggled to keep up with the increasing need for imaging, with MRI and CT imaging rates for adults more than doubling from 2006 to 2016². This escalating demand for services, coupled with an increase in patient complexity not only leads to increased costs, but also reduced accessibility and quality. Delays in imaging at HORDs can also lead to delayed care delivery, which can negatively impact patient outcomes.

63

64 Retail radiology clinics (RRCs) emerged from the general retail health clinics' market boom as 65 a potential solution to the imaging bottleneck effect of HORDs. RRC offer a variety of 66 diagnostic imaging services, typically at lower cost, at a decentralized clinic ³⁻⁵. RRCs have 67 gained popularity by cutting wait-times, increasing accessibility, and reducing costs 3-5. 68 Despite the increase in RRC utilization, price variability and unreliable quality of the technology used, imaging interpretation and reporting of images have led RRCs to be labeled 69 as low-value substitutes to HORD services 6-8. For example, one study found significant 70 71 variability in diagnostic reports from 10 different RRCs performing an MRI scan of the lower 72 back on the same patient within a 3-week period 5. Another study reported that radiology 73 departments in tertiary care centers are frequently asked to perform secondary interpretations 74 of imaging studies, finding that discrepancy rates vary widely ⁶.

75

In parallel, hospital networks are struggling with care integration and coordination, as well as maintaining quality standards across all departments and campuses within their network. Some hospital systems are now looking at options to develop their own RRCs to provide convenient and efficient locations, without sacrificing the technological or interpretation quality, while leveraging the existing trust patients may have with their hospital system. This is a potentially valuable innovation for patients, as they would be able to receive the "specialty read" quality at locations with shorter wait times and lower out-of-pocket costs.

To successfully decentralize imaging services at a larger scale, however, it is important to define value not just from the perspective of the healthcare system, but also from the perspective of the patient. Despite widespread encouragement to incorporate the patient voice in healthcare, developments such as RRCs are usually done without meaningful input from patients ⁷. In the push for patient-centeredness, CMS recommends hospital departments incorporate patient insight into their service quality measures ⁸.

90

Existing research on patient preferences in diagnostic imaging services is sparse, however, and what exists is often contradictory ⁹⁻¹⁵. Though survey studies have explored from whom patients prefer to learn their results and how (e.g., by phone, e-mail, or in-person), findings to date have been inconsistent ^{9-11,16}. For example, one study found a majority preference for learning results in-person and from a radiologist¹⁰ while another found that most patients prefer to learn results from their primary care provider, over the phone ¹¹.

97

98 The objective of this study is to identify patient preferences for outpatient diagnostic imaging 99 services and analyze how patients make trade-offs between attributes of these services using 100 a stated choice experiment. In this study, we analyze patient's individual preference 101 heterogeneity for these services and study how preferences vary among and within different 102 patient populations.

103

104 Methods

105 Discrete Choice Experiment

In this study, we used a Discrete Choice Experiment (DCE) to explore which attributes of imaging services matter to patients and how they make trade-offs between these attributes. A DCE provides the opportunity to estimate pair-wise choices and analyze marginal values for the total value of a health service or good ¹⁷. The unique contribution of a DCE is that it allows researchers to analyze the trade-offs that patients are willing to make, including options that

111 may not currently exist but could in the future. We do this by using a survey in which patients 112 answer a repeated set of choice tasks. In these choice tasks, they are asked to choose their 113 preferred option of three locations offering imaging services. The imaging locations vary by 114 different attributes and levels of those attributes. We explain this further in the paragraph 115 about the experimental design.

116

With the DCE, we can estimate the utilities that patients assign to different attributes of imaging services. Utilities represent the total satisfaction or benefit derived from consuming a good or service and the capacity of a service to satisfy patients' wants. To estimate the relative utilities patients assign to different aspects of services, the DCE included 14 choice tasks. The experimental design of the DCE was based on "prior" estimates of the utilities for attributes of the choice which are calculated by using expectations on what the model parameters will be. These numbers are based on sample data from a pilot and from the literature.

124

125 Qualitative Research

126 To understand which aspects or attributes of imaging services are important to patients and 127 should therefore be included in the choice experiment, we first performed qualitative research 128 to identify key aspects of value to patients. This qualitative work defined the most important 129 attributes of imaging services to patients and provided the base for a DCE. We performed two 130 focus groups, each with 12 patients over the age of 18 who had received outpatient imaging 131 services in 2019 from a medium-sized academic medical center, at the main campus of a six-132 hospital, two-state network in the Northeastern part of the United States. A question guide 133 aimed at understanding how patients perceive their radiology experiences led participants 134 through their focus group session. Questions were intended to be exploratory and were 135 sometimes followed by probes to allow differences between patient insights and experiences 136 to emerge. From the thematic analysis of the focus groups, we learned that important themes 137 included: overarching trust in system and referring providers; preference for transparent and

138 informative communication; personal interaction and compassionate bedside manner from 139 staff; and accessibility of radiology services and the facility. We also performed qualitative 140 research among leaders in the radiology department of the medical center who were asked 141 what they considered to be important to patients. Their suggestions provided additional input 142 for the stated choice design, such as the potential importance of wait time to be seen and 143 whether or not the location for getting imaging was recommended by patient's primary care 144 provider.

145

146 Experimental Design

147 Two discrete choice experiments were designed to represent an acute patient's need (X-ray 148 service), and a sub-acute need using Magnetic Resonance Imaging (MRI service). Participants 149 were asked to imagine a situation where they were hurt and needed imaging services. 150 Participants were randomized between two arms with the following descriptions of the choice 151 situation:

• Situation 1): "For the purpose of this study, suppose you hurt your arm and

153 your primary care provider wants to send you for an X-ray. You have three

154 options of locations where you can have your imaging done."

• Situation 2): "For the purpose of this study, suppose you hurt your back a

156 while ago, and are having persistent pain. Your primary care provider wants to

157 send you for an MRI. You have three options of locations where you can have

158 your imaging done."

159

160 The DCE included 14 choice tasks where patients were asked to choose the imaging clinic they 161 preferred. Each choice task had three different clinics which varied by 10 different attributes, 162 which were determined by prior qualitative research. These selected attributes included: 163 whether the interpreting radiologist is a general or sub-specialty radiologist; whether the clinic 164 was recommended by their primary care physician (PCP); time to results; out of pocket cost; wait time to an appointment; travel time to the clinic; parking costs; parking accessibility;
service; and whether or not online scheduling is available. Service is a multifactorial attribute
(e.g., staff attentiveness and facility amenities) combined into a star rating. The rating scale is
between one and five stars, with a five-star rating representing an excellent service and a onestar rating suggesting a poor service, as rated by other hypothetical patients. Figure 1 shows
an example of a choice task.

171

Following the DCE choice tasks, we asked patients attitudinal questions on a 1–5-point Likert scale from "strongly disagree" to "strongly agree". These questions were borrowed from and validated by the national Medical Expenditures Panel Survey (MEPS) ¹⁹, a set of large-scale surveys of families and individuals, their medical providers, and employers across the United States. The questions focused on perceived need for healthcare, need for insurance, riskaversion, and perceived personal health status.

178

179 Data Source, Participants and Study Size Estimation

Our data were sampled from an online Centiment panel between April 11, 2021 through November 19, 2021, with a second sample between May 16, 2022, and June 24, 2022. Centiment contacted approximately 472 individuals in the catchment area of a moderate size academic medical center in a rural Northeastern part of the United States. Participants completed written consent before continuing to the online survey questions. All data were fully anonymized before shared with the study team. The Institutional Review Board at the University of Vermont reviewed the study and determined it was exempt from full review.

Of the total sample, 268 finished the survey and met initial inclusion criteria (Age >= 188 18): 134 were assigned to the arm X-ray group, 134 to the back MRI group. We excluded 98 189 subjects for failing consistency criteria, qualifying for a closed quota, failing a bot-behavior 190 check, or failing an attention (response quality) check, leaving a final sample of 170: 84 in X-191 ray, 86 in MRI. As each subject answered 14 choice tasks, we obtained an effective sample size of n=2,380 for modeling. We used NGene 1.2.1 (ChoiceMetrics, 2018) to estimate the
minimum sample size required for this study.

194

195 Statistical Methods

196 We used a mixed logit model (ML) to estimate the probability of a choice alternative being 197 chosen, depending on the characteristics of the choice (attributes and levels) and the 198 characteristics of the chooser. We used latent class analysis (LCA) which addresses the issue 199 of unobserved preferences of patients by probabilistically segmenting a sample population 200 into different groups or "classes" based on a latent variable²⁴. Class membership is first defined 201 by a membership function including the indicator variables, after which the utility functions 202 of different classes can be estimated. All statistical analyses were conducted using Stata 18 203 (StataCorp LLC, 2023).

204

205 Latent Class Analysis

The latent class conditional logit (LCL) model further extends the ML model by incorporating a discrete representation of unobserved preference heterogeneity within groups in a population. The LCL model might explain, for example, that patients who had previous imaging services are more likely to fall into the class that is more sensitive to appointment wait time, while older patients might be more likely to fall into the class that is more sensitive to PCP recommendation.

212

213 **Results**

214 Descriptive Results

215 A total of 84 patients answered questions about preferences for attributes of an X-ray; 86

216 people responded to the MRI choice questions. The summary statistics are reported in Table

1. On average, patients answering choice questions on X-ray tended to be female (60%),

218 white (91%) and live in rural areas (58%); about half had private insurance (46%) and

relatively few had met their insurance deductible (21%). Patients answering the choice
questions on the MRI were generally similar, with 67% female, 96% white, 69% in rural
areas, 45% with private insurance, 22% had met their insurance deductible. Patients
receiving the X-ray choice questions had had an average of 4.8 previous images while MRI
patients had had 2.9.

224

225 Table 1: Descriptive Statistics226

| | X-ray | | MRI | |
|---------------------------------------------------------------|----------|------------|----------|------------|
| | | | | |
| Age (mean, sd) | 46.0 | (16.1) | 47.8 | (16.9) |
| Income (USD) (mean, sd) | \$48,425 | (\$40,939) | \$47,444 | (\$47,881) |
| Number of Previous Images (mean, sd) | 4.8 | (1.5) | 2.9 | (1.7) |
| Agreement: I don't need insurance (mean, sd) | 1.5 | (0.9) | 1.7 | (1.2) |
| Agreement: Insurance isn't worth the cost (mean, sd) | 2.3 | (1.3) | 2.4 | (1.4) |
| Agreement: I take more risks than others (mean, sd) | 2.3 | (1.2) | 2.3 | (1.2) |
| Agreement: I can fight illness without a doctor (mean, sd) | 2.5 | (1.3) | 2.4 | (1.2) |
| Agreement: I am healthier than others (mean, sd) | 2.7 | (1.1) | 2.7 | (1.1) |
| Gender Identity (%, n) | | | | |
| Male | 38.1% | 32 | 32.6% | 28 |
| Female | 59.5% | 50 | 67.4% | 58 |
| Non-Binary | 2.4% | 2 | 0.0% | 0 |
| Race (%, n) | | | | |
| White | 91.7% | 77 | 96.5% | 83 |
| Black | 1.2% | 1 | 0.0% | 0 |
| Asian | 2.4% | 2 | 2.3% | 2 |
| AIAN | 3.6% | 3 | 1.2% | 1 |
| Multiple | 1.2% | 1 | 0.0% | 0 |

| Ethnicity (%, n) | | | | |
|------------------------------|-------|----|-------|----|
| Not Hispanic | 92.9% | 78 | 96.5% | 83 |
| Hispanic | 7.1% | 6 | 3.5% | 3 |
| Rurality (%, n) | | | | |
| Urban | 14.3% | 12 | 14.0% | 12 |
| Suburban | 27.4% | 23 | 17.4% | 15 |
| Rural | 58.3% | 49 | 68.6% | 59 |
| Education Level (%, n) | | | | |
| Less than High School | 2.4% | 2 | 1.2% | 1 |
| High school | 27.4% | 23 | 23.3% | 20 |
| Some college, no degree | 25.0% | 21 | 17.4% | 15 |
| Associate's: vocational | 0.0% | 0 | 3.5% | 3 |
| Associate's: academic | 6.0% | 5 | 2.3% | 2 |
| Bachelor's (BA, AB, BS, BBA) | 23.8% | 20 | 32.6% | 28 |
| Master's | 11.9% | 10 | 15.1% | 13 |
| Professional, Doctoral | 3.6% | 3 | 4.7% | 4 |
| Insurance Status (%, n) | | | | |
| Insured - Private | 46.4% | 39 | 45.3% | 39 |
| Insured - Medicare | 21.4% | 18 | 24.4% | 21 |
| Insured - Medicaid | 26.2% | 22 | 23.3% | 20 |
| Uninsured | 6.0% | 5 | 7.0% | 6 |
| Deductible Status (%, n) | | | | |
| Met | 21.4% | 18 | 22.1% | 19 |
| Unmet | 44.0% | 37 | 45.3% | 39 |
| No Deductible | 34.5% | 29 | 32.6% | 28 |
| Health Status (%, n) | | | | |
| No Chronic Condition | 50.0% | 42 | 47.7% | 41 |
| Chronic Condition | 50.0% | 42 | 52.3% | 45 |
| | | | | |

227

228 Mixed logit Results

The results of the mixed logit model are shown in Table 2 where we separated results for Xray and MRI. We found that out-of-pocket costs, interpreting doctor specialty, whether or not the clinic was recommended by the primary care provider, the wait time to results, the clinic service rating and online scheduling were all statistically significant and had the expected signs for both MRI and X-ray. Patients were less likely to choose a clinic if the out-of-pocket costs were higher and the wait time to results was longer, but more likely to choose it if images were interpreted by a specialty radiologist, the clinic was recommended to them by their primary care provider, the service rating was higher and if online scheduling was available. For X-ray, free parking was associated with a higher probability of choosing a clinic.

| | (MRI) | (MRI) | (MRI) | (X-ray) | (X-ray) | (X-ray) |
|--------------------------------------|-----------|---------|---------------|---------------------------|---------|-------------------|
| | В | SE | 95% CI | В | SE | 95% CI |
| Out of Pocket Cost | -0.004*** | (0.001) | -0.005,-0.003 | -0.006* | (0.002) | -0.010,-0.00 |
| Interpreting Doctor Specialty | 0.732*** | (0.116) | 0.504,0.960 | 0.374** | (0.122) | 0.135,0.614 |
| Clinic Recommended by PCP | 0.626*** | (0.112) | 0.407,0.845 | 0.678*** | (0.155) | 0.374,0.982 |
| Wait Time to Results | -0.012* | (0.006) | -0.024,-0.001 | - 0.013 ^{***} | (0.004) | -0.020,- 0.006 |
| Travel Time to Clinic | -0.007 | (0.006) | -0.004,-0.018 | -0.018* | (0.008) | 0.001,0.034 |
| Wait Time to Appointment | -0.014 | (0.018) | -0.050,-0.022 | -0.023* | (0.012) | 0.000,0.047 |
| Clinic Service Rating | 0.520*** | (0.096) | 0.332,0.708 | 0.451*** | (0.126) | 0.205,0.698 |
| Free Parking | 0.150 | (0.097) | -0.040,0.339 | 0.669*** | (0.108) | 0.457,0.881 |
| Clinic Parking Lot Distance | -0.041 | (0.023) | -0.087,0.005 | -0.011 | (0.007) | -0.025,0.002 |
| Online Scheduling Availability | 0.288* | (0.112) | 0.068,0.508 | 0.375*** | (0.089) | 0.201,0.550 |

239 Table 2: Results of the MMNL Model240

The attributes that mattered the most to patients for both MRI and X-ray were specialty radiologist reading (0.732, standard error= 0.116 for MRI; 0.374, se =0.122) for X-ray); recommendation by the primary care provider reading (0.626, se=0.112 for MRI; 0.678, se=0.155 for X-ray) and the clinic's service rating reading (0.520, se=0.096 for MRI; 0.451, se=0.126 for X-ray).

248

249 Latent Class Analysis

250 Table 3 shows the results of the Latent Class model where age, gender, education, income, 251 whether someone had more than two previous scans, whether they considered themselves 252 healthier than others, whether they had private insurance and whether they were more likely 253 to take risks than others were the indicator variables estimating the probability of class 254 membership. We found that 54.7 percent of patients were in class 1 and 45.3 percent in class 255 2. We found that females, older patients, whether they had at least two previous scans and 256 those who were less likely to take risks than others were highly predictive of being in class 2 257 (p<0.02) which we therefore labeled as the "Experienced Patients" class. Those who had fewer 258 than two previous scans were more likely to be in class 2 (2.7111, p<0.01) as were male patients 259 (-0.8479, p=0.02) and younger patients (-1.7197, p=0.02), which we labeled "New Patients". 260 The results show that someone who had a 1-point increase in the Likert scale for "more likely 261 to take risks than others" was significantly more likely to be in class 2 (-3.1354.7277, p=0.01). 262 they offer at the centralized location.

- 263 Table 3: Results of the Latent Class Analysis
- 264
- 265

| | Class membership | Standard error |
|-----------------------------------------|------------------|----------------|
| Male | 84793** | (.3791) |
| Income (ref: lowest income category) | 0.1477 | (.3979) |
| Age (years) | 1.7196** | (.7559) |
| More than 2 previous scans | 2.7110*** | (.9873) |
| Education (ref: no high school) | -12.9818 | (3.6767) |
| More likely to take risk than others | -3.1353*** | (1.2389) |

| Private insurance | -1.0671 | (1.1982) (10.5153 | |
|-----------------------|----------------------|----------------------------|--|
| Healthier than others | -14.4012 | | |
| | Class 1 (Experienced | Class 2 (New Users) B (SE) | |
| | Patients) B (SE) | | |
| Costs > current | -0.1774*** (0.0367) | -0.2783 (0. 1764) | |
| Specialty read | 0.35123*** (0.0419) | 0.0525 (0. 2753) | |
| PCP recommendation | 0.3890*** (0.0432) | 0.1641 (0.2178) | |
| Online Scheduling | 0.1240*** (0.0419) | 4.7541*** (0.4703) | |
| Service rating | 0.3033***(0.0509) | 0.1148*** (0.2636) | |
| Time to Results | -0.0225 (0.0340) | -0.0289 (0.1763) | |
| Travel Time | -0.1473*** (0.0324) | -0.2649 (0.1618) | |
| Wait Time | -0.1165*** (0.0447) | -0.1271 (0.1968) | |
| Access to parking | -0.0308 (0.0254) | -0.0509 (0.1829) | |

266 267

267 **** **p<.01**, *** **p<.05**, * **p<.10** 268

When decentralizing some services away from the main hospital, the focus should be on passing on the medical center's high service rating to the HORD and offering online scheduling. Indeed, patients value easier access, shorter wait times and lower out-ofpocket costs which the growing popularity of RRCs has shown. HORDs can win in popularity by making sure they attain and retain high star ratings while offering better access than RRCs. Primary care providers can potentially play an important role in directing patients to HORDs for their diagnostic imaging services.

276

277 For patients in class 1 ("Experienced Patients"), costs (-0.1774), specialty read (0.3512), PCP 278 recommendation (0.3890), travel time (-0.1474), wait time (-0.1165), service (0.3033) and 279 online scheduling (0.1240) all had significant effects (p<0.01) and had the expected signs. For 280 example, patients in the "Experienced Patients" class will be more likely to choose a clinic 281 location if it is lower cost, takes less travel time, has shorter wait times for results, if results 282 are read by a specialty radiologists, the clinic is recommended by their PCP, service is better 283 and online scheduling is available. Patients in the "New Patients" class only cared about online 284 scheduling (4.7542) and service rating (1.1149), but the effect size was relatively high. None of 285 the other attributes of the service would affect their choice.

286 Marginal Rates of Substitution

287 Table 4 shows an analysis of the trade-offs patients were willing to make, known as the 288 marginal rates of substitution. We found that for patients in the Experienced Patient class, 289 even though most attributes significantly affected their choice for clinic, the effect size was 290 considerably smaller than for the New Patients class. Experienced Patients were willing to 291 pay: \$2 more than what they currently pay (out-of-pocket) to have their images read by a 292 radiologist specialist; \$2 more to go to a clinic that was recommended by their PCP; \$0.70 293 more for online scheduling; \$1.70 more for a 1-point higher star rating; \$0.80 more to have a clinic that would be 1 minutes closer than their current one; \$0.65 more for a clinic that had a 294 295 1-hour shorter wait and \$0.17 more for a clinic that would decrease their walk-up time by 1 296 minute. The New Patients were willing to pay \$17 more for online scheduling and \$4 more for 297 a higher star rating.

298

299 Table 4: Willingness-to-Pay (MRS)

300

| Class 1 (Experienced Patients) | Class 2 (New Patients) |
|-----------------------------------|-----------------------------------------------------------------------------|
| 1.980 | 0.189 |
| 2.192 | 0.590 |
| -0.127 | -0.104 |
| -0.831 | -0.952 |
| -0.657 | -0.457 |
| 1.709 | 4.005 |
| -0.174 | -0.183 |
| 0.699 | 17.079 |
| | (Experienced Patients) 1.980 2.192 -0.127 -0.831 -0.657 1.709 -0.174 |

301 302 303

304 Attribute Importance

305 Patients were also asked to rank-order the attributes in how they prioritized them when 306 making a choice for a clinic. This is usually consistent with the magnitude of the coefficients 307 in the choice models, but also allows us to see if there was any attribute non-attendance. This 308 means that when processing the attributes, some patients may not consider particular 309 attributes at all. The violin plot in Figure 2 shows the results of the question on attribute 310 importance. We see that the results are consistent with the ML results: the most important 311 attributes included interpreting doctor specialty level, PCP recommendation, and costs. There 312 was a small difference between the X-ray and MRI arms: among X-ray patients, service rating 313 and parking access also ranked highly.

314

315 **Discussion**

In this paper, we sought to identify patient preferences for outpatient diagnostic imaging services. We analyzed how patients make trade-offs between attributes of services using a discrete choice experiment. We explored patient's individual preference heterogeneity for these services and reported how preferences vary among and within different groups of patients.

321

322 In our base analysis, we found that specialty reading of images, PCP recommendation, lower 323 costs, travel time and wait time, as well as higher star rating (representing better service or 324 reputation) and online scheduling are all significant predictors of choice regarding where to 325 get diagnostic imaging services. However, when we segmented the sample population 326 deterministically, we found that males, younger people and more likely to take risks than 327 others only We termed this group "New Patients" as they had significantly fewer previous 328 scans and generally do not highly value health services. Insurance status, health status and 329 chronic conditions, education and income did not define class membership.

331 These results suggest that decision-makers looking to decentralize imaging services while 332 incorporating patient preferences for attributes of those services should differentiate between 333 the different patient sub-populations they are serving. This requires careful consideration of 334 patient characteristics as well as preferences. Overall, we found in this study that New Patients 335 care about the reputation or star rating of a clinic and online scheduling availability. But 336 additionally, Experienced Users – who will be the majority of radiology users – are focused on 337 wait time, price and recommendations from primary care providers. This is a concrete 338 message to medical centers seeking to decentralize their services away from the main hospital: 339 patients will want to know that the new location offers the same service level in addition to 340 convenience and will be relying on their primary care providers for advice, suggesting outreach 341 to primary care providers will be important for success.

342

343 Limitations

Although Centiment used a quota sampling approach, the gender balance may not be reflective of the total population, although we do not believe this is a major threat to external validity. However, while our sample is largely representative of the larger population, we measure intention for hypothetical choices and cannot say for sure that these consistently translate to real-life behavioral trade-offs, especially in acute situations. More work needs to be done to further explore factors that affect decision-making and preferences in these circumstances.

350

351 Conclusions

In this study, we analyzed the trade-offs patients make between attributes of radiology services to inform decision-making around designing optimal HORDs. Our analysis showed that a patient population can and should be segmented into subgroups that evaluate the value of imaging services differently. The "Experienced Patients" subgroup generally value diagnostic imaging services in both acute and more chronic situations and they had a strong preference for a HORD that would provide services at lower costs, where their images would be interpreted by a specialty radiologist, the clinic would be recommended by their PCP, online scheduling would be available, service rating were higher, and travel and wait times would be
shorter. HORDs can therefore be more competitive by providing services that live up to these
expectations better than available RRCs.

362

New Patients significantly valued the service rating of the HORD and online scheduling. Most 363 364 RRCs do not currently offer online scheduling so ease of use may also steer these potential 365 future patients towards HORDs. More importantly, HORDs have the advantage of being linked to the main medical center which has the reputation of having clinical expertise and 366 367 more sophisticated technology. We conclude that there is room for medical centers to build 368 HORDs that provide an appealing and competitive alternative to the current RRC. RRCs have 369 the risk of unreliable quality in the technology and imaging interpretation and it is therefore 370 desirable that HORDs provide the same or more benefits while maintaining the quality care

372 **References**

373

374 Levin DC, Parker L, Rao VM. Recent trends in imaging use in hospital settings: 1. 375 Implications for future planning. J Am Coll Radiol Internet. Elsevier Inc; 376 2017;14:331–6. Available from: http://dx.doi.org/10.1016/j.jacr.2016.08.025 377 Smith-Bindman R, Kwan ML, Marlow EC, Theis MK, Bolch W, Cheng SY, et al. 2. 378 Trends in use of medical imaging in US health care systems and in Ontario, Canada, 379 2000-2016. JAMA. 2019;322:843-56. 380 Boland GWL. Diagnostic imaging centers for hospitals: A different business 3. 381 proposition for outpatient radiology. J Am Coll Radiol. 2007;4:581-3. 382 Iglehart JK. The new era of medical imaging - Progress and pitfalls. N Engl J Med. 4. 383 2006;354:2822-8. 384 Scott Ashwood J, Reid RO, Setodji CM, Weber E, Gaynor M, Mehrotra A. Trends in 5. 385 Retail Clinic Use Among The Commercially Insured Internet. Available from: 386 www.aimc.com 387 Herzog R, Elgort DR, Flanders AE, Moley PJ. Variability in diagnostic error rates of 6. 388 10 MRI centers performing lumbar spine MRI examinations on the same patient 389 within a 3-week period. Spine J Internet. Elsevier Inc.; 2017;17:554–61. Available 390 from: http://dx.doi.org/10.1016/j.spinee.2016.11.009 391 Kostrubiak, D.E., DeHay, P.W., Ali, N., D'Agostino, R., Keating, D.P., Tam, J.K. and 7. 392 Akselrod, D.G., 2020. Body MRI subspecialty reinterpretations at a tertiary care 393 center: discrepancy rates and error types. American Journal of Roentgenology, 394 215(6), pp.1384-1388. 395 8. Johnson FR, Beusterien K, Özdemir S, Wilson L. Giving patients a meaningful voice 396 in United States regulatory decision making: The role for health preference research. 397 Patient. 2017;10:523-6. 398 Centers for Medicare & Medicaid Services. CMS quality measure development plan: 9. 399 Supporting the transition to the Merit-based Incentive Payment System (MIPS) and 400 Alternative Payment Models (APMs) Internet. 2016. Available from: 401 https://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-402 Instruments/Value-Based-Programs/MACRA-MIPS-and-APMs/Final-MDP.pdf 403 Basu PA, Ruiz-Wibbelsmann JA, Spielman SB, Van Dalsem VF, Rosenberg JK, Glazer 10. 404 GM. Creating a patient-centered imaging service: Determining what patients want. 405 Am J Roentgenol. 2011;196:605-10. Pahade J, Couto C, Davis RB, Patel P, Siewert B, Rosen MP. Reviewing imaging 406 11. 407 examination results with a radiologist immediately after study completion: Patient 408 preferences and assessment of feasibility in an academic department. Am J 409 Roentgenol. 2012;199:844-51. 410 Cabarrus M, Naeger DM, Rybkin A, Qayyum A. Patients prefer results from the 12. 411 ordering provider and access to their radiology reports. J Am Coll Radiol Internet. 412 Elsevier Inc; 2015;12:556–62. Available from: 413 http://dx.doi.org/10.1016/j.jacr.2014.12.009 414 Mangano MD, Bennett SE, Gunn AJ, Sahani D V., Choy G. Creating a patient-13. 415 centered radiology practice through the establishment of a diagnostic radiology 416 consultation clinic. Am J Roentgenol. 2015;205:95-9. 417 Woolen S, Kazerooni EA, Wall A, Parent K, Cahalan S, Alameddine M, et al. Waiting 14. 418 for radiology test results: Patient expectations and emotional disutility. J Am Coll 419 Radiol Internet. Elsevier Inc; 2018;15:274–81. Available from: 420 https://doi.org/10.1016/j.jacr.2017.09.017 421 15. Kleij KS, Tangermann U, Amelung VE, Krauth C. Patients' preferences for primary 422 health care - A systematic literature review of discrete choice experiments. BMC 423 Health Serv Res. BMC Health Services Research; 2017;17:1-12. 424 Mühlbacher AC, Bethge S, Reed SD, Schulman KA. Patient preferences for features of 16. 425 health care delivery systems: A discrete choice experiment. Health Serv Res. 426 2016;51:704-27.


- 427 17. van den Broek-Altenburg, E. and Atherly, A., 2020. Using discrete choice
 428 experiments to measure preferences for hard to observe choice attributes to inform
 429 health policy decisions. Health economics review, 10(1), pp.1-8.
- 430 18. AA. Hess, S., Rose, J.M. Should Reference Alternatives in Pivot Design SC Surveys be
 431 Treated Differently?. Environ Resource Econ 42, 297–317 (2009).
- 432 19. Cohen, J.W., Cohen, S.B. and Banthin, J.S., 2009. The medical expenditure panel
 433 survey: a national information resource to support healthcare cost research and
 434 inform policy and practice. Medical care, pp.S44-S50.
- 435 20. McFadden, Daniel. "Conditional logit analysis of qualitative choice behavior." (1973).
- 436 21. Hensher DA, Greene WH. The mixed Logit model: the state of practice.
 437 Transportation. 2003;30(2):133-76.
- 438 22. McFadden D, Train K. Mixed MNL models for discrete response. J Appl Econ.
 439 2000;15(5):447-70.
- van den Broek-Altenburg, E., et al., Exploring Heterogeneity in Moral Terminology in
 Palliative Care Consultations. BMC palliative care, 2020.
- 442 24. Yoo HI. lclogit2: An enhanced command to fit latent class conditional logit models.
 443 The Stata Journal. 2020 Jun;20(2):405-25.
- 444 25. Train KE. EM algorithms for nonparametric estimation of mixing distributions.
- 445 Journal of Choice Modelling. 2008 Jan 1;1(1):40-69.

Figure 1 Example of a Choice Task in the DCE

| | Clinic 1 | Clinic 2 | Clinic 3 |
|-----------------------------|---------------------|-----------------------|---------------------|
| Interpreting Radiologist | General Radiologist | Specialty Radiologist | General Radiologist |
| Primary Care Recommendation | Yas | No | Yes |
| Time to Results | 18 hours | 24 hours | 30 hours |
| Cost | \$50 | \$40 | \$50 |
| Walt Time | 10 days | 10 days | 8 days |
| Travel Time | 22 minutes | 37 minutes | 22 minutes |
| Parking Cost | Paid | Free | Paid |
| Parking Accessibility | 7.5 minutes | 12.5 minutes | 7.5 minutes |
| Service | 3 Star. *** | 4 Star: **** | 2 Star. ** |
| Online Scheduling | Not Available | Available | Not Available |
| Which would you choose? | Clinic 1 | Clinic 2 | Clinic 3 |

Figure

Figure 2: Attribute Importance



Figure