

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

A brain vital sign study of subconcussion in female and male high school student athletes

Ryan C.N. D’Arcy^{1,2,3*}, David McCarthy⁴, Derek Harrison⁴, Zander Levenberg⁴, Julian Wan⁴, Aidan Hepburn⁴, Eric D. Kirby^{1,5}, Tanja Yardley⁶, Nikita Yamada-Bagg², Shaun D. Fickling^{1,2}, Thayne A. Munce⁷, David W. Dodick^{8,9}, Christopher Ahmad^{10,11}, & Ken Shubin Stein^{9,12}

- 1) BrainNET, Health and Technology District, Surrey, BC, Canada;
- 2) Centre for Neurology Studies, HealthTech Connex, Surrey, BC, Canada;
- 3) Faculty of Engineering Science, Simon Fraser University, Burnaby, BC, Canada;
- 4) Brentwood College School, BRAIN team, Mill Bay, BC, Canada;
- 5) Faculty of Individualized Interdisciplinary Studies, Simon Fraser University, Burnaby, BC, Canada;
- 6) Surrey Neuroplasticity Clinic, HealthTech Connex, Surrey, BC, Canada;
- 7) Environmental Influences on Health & Disease Group, Sanford Research, Sioux Falls, SD, USA;
- 8) Mayo Clinic College of Medicine, Rochester, MN, USA;
- 9) Atria Academy of Science and Medicine, New York City, NY, USA;
- 10) New York Presbyterian Hospital, New York City, NY, USA;
- 11) Columbia University Irving Medical Center, New York City, NY, USA;
- 12) Hospital for Special Surgery, New York City, NY, USA.

*Corresponding author: Ryan C.N. D’Arcy, PhD, PIEng, Centre for Neurology Studies, HealthTech Connex, Surrey, BC Canada ryan@healthtechconnex.com

BRENTWOOD Research Study
Manuscript

35 **Abstract**

36 Emerging evidence from neurophysiological brain vital sign studies show repeatable sensitivity
37 to cumulative subconcussive impairments over a season of contact sports. The current study addressed
38 the need for a large prospective study comparing a low-contact control group to high-contact group.
39 Importantly, the study also expanded the scope of neurophysiological changes related to repetitive head
40 impacts to include female athletes in addition to male athletes. In total, 89 high school student athletes
41 underwent 231 brain vital sign scans over a full calendar year. The results replicated prior subconcussive
42 cognitive impairments (N400 delays) and sensory impairments (N100 amplitude reductions) in male
43 athletes and demonstrated similar subconcussive impairments for the first time in female athletes.
44 While there was no significant subconcussive difference between female and male athletes, female
45 athletes show overall larger responses in general. The findings demonstrated that subconcussive
46 impairments are detectable in a controlled experimental comparison for both female and male high
47 school athletes. The study highlights the opportunity to monitor subconcussive changes in cognitive
48 processing for both female and male athletes to help advance prevention, mitigation and management
49 efforts aimed at reducing athletes' risk of potential long-term negative health outcomes related to
50 cumulative exposure to repetitive head impacts.

51

52 **Introduction**

53

54 **Background**

55 Head impact exposure from participating in contact or collision sports is commonly associated
56 with concussion; however, subconcussion is a growing concern for athletes as well as a developing
57 research area in parallel (Smith et al., 2018; 2019). Subconcussion has been increasingly implicated with
58 potential short-term consequences on the developing brains and longer-term concerns related to chronic
59 traumatic encephalopathy (McKee et al. 2009; Guskiewicz et al. 2005; Dioso et al. 2022). Concussion
60 research has also historically been primarily focused on males (Valera et al. 2021; Snedaker et al. 2022).
61 Therefore, emerging subconcussive research must be inclusive of both female and male athletes. This is a
62 timely priority as new intervention options begin to emerge from the clinical trials literature (Breuer et al.
63 2023).

64 Operationally defined, a subconcussive impact is a mechanical force transmitted to the brain
65 below the threshold for a diagnosis of an acute concussive injury. The effects of these repetitive head
66 impacts (RHI) may neither be detectable to players nor to observers; however, it has been shown that RHI
67 over time can result in both microstructural alterations and functional brain impairment (Mainwaring et
68 al. 2018). The extent of impairment is highly correlated with the frequency of exposure to these impacts
69 (Fickling, Smith, et al. 2021; Fickling, Poel, et al. 2021; Hirad et al. 2019; Saghafi et al. 2018; Fickling et al.
70 2019).

71 Recent research has increasingly focused on the role of cumulative RHI exposure rather than a
72 concussion as a singular, acute traumatic event. Evidence suggests that even a single practice session
73 involving head contact, such as heading a ball in soccer, can result in impairment (Nowak et al. 2020).
74 Magnetic resonance imaging (MRI) studies have demonstrated that players exposed to RHI over the

BRENTWOOD Research Study Manuscript

75 course of a season of sport show significant structural abnormalities compared to controls over the same
76 time period (Hirad et al. 2019; Saghafi et al. 2018; Veksler et al. 2020). Both Hirad et al. (2019) and Saghafi
77 et al. (2018) used diffusion-tensor imaging, an MRI-derived technique, to show abnormalities in the white
78 matter of a group of football athletes were related to different measures of head impact exposure.
79 Interestingly, while detecting differences in permeability and white matter microstructure in the brains of
80 contact sport athletes, Veksler et al. (2020) did not find any statistical differences in blood-brain barrier
81 permeability between non-contact athletes and non-athlete healthy controls, leading the authors to
82 merge the groups into a single control group.

83 A major challenge for current medical imaging approaches, including MRI, relates to the need for
84 accessible, objective, and sensitive measures of subconcussive impacts at points-of-care (PoC). Portable
85 electroencephalography (EEG) provides the evaluation of individual neurophysiological event-related
86 potentials (ERPs). These have recently been translated into a brain vital signs framework, facilitating
87 access to sensitive measures for various PoC applications (Ghosh Hajra et al. 2016). Three well-established
88 ERP responses are extracted within the brain vital signs framework: 1) the N100 for auditory sensation
89 (Davis 1939); 2) the P300 for basic attention (Sutton et al. 1967); and 3) the N400 for cognitive processing
90 (Kutas and Federmeier 2011). Brain vital signs have been successfully implemented in the evaluation of
91 both concussion and subconcussion (Fickling et al. 2019; Fickling, Smith, et al. 2021; Fickling, Poel, et al.
92 2021).

93 Identification of subconcussive cognitive impairments over a season of contact sports was initially
94 reported in male Junior-A ice hockey players (Fickling et al. 2019, N=23). Subsequent studies (Fickling,
95 Smith, et al. 2021, N=23a; Fickling, Poel, et al. 2021, N=15b; Breuer et al. 2023, N=30) replicated and
96 confirmed identification of accumulated subconcussive exposure in male athletes for different age ranges
97 (12-14, 16-21, & 21-28, respectively) and different contact sports, including ice hockey, tackle football,
98 and mixed martial arts. Importantly, this work demonstrated strong, significant, and linearly predictive
99 relationships between brain vital sign changes and exposure to subconcussive impacts (Fickling, Smith, et
100 al. 2021; Fickling, Poel, et al. 2021). These relationships were demonstrated both when subconcussive
101 impacts were quantified directly by impact sensors or indirectly by the number of games and practices
102 played. A pattern of subconcussive changes have emerged, identified by changes in the N400 (cognitive
103 processing) and N100 (sensory processing). The N400 and N100 components have been implicated in the
104 rate of information processing and synchronous pyramidal neural activation (Luck 2014): the two are
105 interdependent factors reflecting an association with cognitive processing (Kutas and Federmeier 2011)
106 and auditory sensation (Davis 1939). While the replications of N400 and N100 subconcussive changes in
107 male athletes are noteworthy, a single prospective study with a large sample size including female athletes
108 is critical.


109

110 Objectives & Hypothesis

111 The study was conducted at Brentwood College School (a Grade 8-12 high school in Mill Bay,
112 British Columbia, Canada) by student researchers from the Brentwood Research, Action, and Innovation
113 in Neuroscience (BRAIN) team. Brain vital signs were monitored in 89 female and male high school
114 students (15-17 years of age) participating in high-contact (e.g. rugby, ice hockey, and soccer) and low-
115 contact sports (e.g. rowing, climbing, and tennis) over three academic terms across the school year (Figure

BRENTWOOD Research Study Manuscript

116 1). With respect to matching participants in high-contact versus low-contact sports, Brentwood is a private
117 boarding school, in which all students follow a structured program with commonly matched ages, daily
118 schedules, diets, academics, athletics, and sleep routines. The study used a mixed-model, repeated
119 measures design evaluated across 3 terms before and after each of the three school year terms as well as
120 after the summer break, resulting in 7 possible scan timepoints. The objectives were to replicate prior
121 subconcussion findings in male athletes, as well as in female athletes, with the primary hypothesis
122 predicting that brain vital sign subconcussive differences would be detectable in the comparisons for high-
123 contact sports relative to low-contact sports. Subsequent analyses were conducted to explore specific
124 differences between female and male subconcussive and overall neurophysiological results.
125
126



127
128 **Figure 1:** Overview of the experimental design and study timeline (N: Number of participants; n: brain
129 vital signs scans completed).
130
131

132 Results


133
134 In the current study, 89 female and male student athletes were scanned a total of 231 times
135 before and after varying terms of contact or non-contact sports. All three brain vital signs, the N100,
136 P300, and N400, were verified at both the group- and individual- levels. Peak amplitude and latency
137 values were examined in the multivariate analysis of variance (MANOVA), which were first examined
138 with univariate tests of normality and Levene's test of equality of error variances. N100 latency and
139 N400 amplitude failed tests of normality ($p < 0.05$), while all amplitude and latency values passed
140 Levene's test ($p > 0.05$). Therefore, non-parametric Kruskal-Wallis test results are also included in
141 Supplemental Data 1 to demonstrate the robustness of the following MANOVA results.
142

BRENTWOOD Research Study Manuscript

143 High-contact versus low-contact effects:

144 Figure 2a shows brain vital sign differences in the N400 latency and N100 amplitude, while
145 figure 2b provides the underlying waveform differences. The findings confirmed the consistent
146 replication of subconcussive changes in the N400 and N100 at the overall group level, with a significant
147 delay in N400 peak latency and a significant increase in N100 peak amplitude. Specifically, the MANOVA
148 showed a significant main effect of condition between high- versus low-contact groups [$F(6,104) = 7.95$,
149 $p < 0.001$, $\eta_p^2 = 0.314$] (Table 1). Separate univariate between-subjects tests on the outcome variables
150 revealed significant effects of group on the N400 latency, [$F(1) = 35.79$, $p < 0.001$, $\eta_p^2 = 0.247$], and N100
151 amplitude, [$F(1) = 7.88$, $p = 0.006$, $\eta_p^2 = 0.067$] (Figure 2, Table 2). Figure 3 provides the individual scatter
152 plot data for both the N400 latency and N100 amplitude effects, including means and standard
153 deviations. The current results addressed the planned high- versus low-contact group comparison to
154 build on prior pre- versus post- season differences in male athletes. Specific pre- versus post-
155 comparisons are provided as Supplemental data for comparison (Supplemental Figure 1).

156
157



158
159 **Figure 2: (A):** Overall group (female and male combined) brain vital sign radar plot comparing high-
160 versus low-contact groups. **(B)** Group-level grand average ERP waveforms from tones (top) and
161 word pairs (bottom) for overall group comparing high- versus low-contact group differences. Deviant tones
162 and incongruent word pair waveforms are plotted, with standard error shaded. N100, P300, and N400
163 brain vital sign components of the high-contact group are highlighted with labels and arrows. Asterisks
164 denote significant effects in both the brain vital sign and waveform results.

165

166 **Table 1:** MANOVA multivariate effect results for high- and low-contact scans.

BRENTWOOD Research Study
Manuscript

MANOVA	DF	PILLAI	F-VALUE	P-VALUE
GROUP	6,104	0.314	7.947	<0.001*
SEX	6, 104	0.148	3.020	0.009*
GROUP X SEX	6, 104	0.003	0.047	1.000


167

168 **Table 2:** MANOVA results for the group effect on each brain vital sign component.

Brain vital sign	High-contact mean (standard deviation)	Low-contact mean (standard deviation)	F-value	p-value
N100 Amplitude	-4.60 (2.82)	-3.15 (2.11)	7.883	0.006*
N100 Latency	100.78 (16.58)	100.78 (25.64)	0.750	0.388
P300 Amplitude	4.52 (2.84)	3.86 (3.26)	0.508	0.477
P300 Latency	274.77 (40.03)	270.14 (35.22)	0.304	0.583
N400 Amplitude	-3.49 (2.85)	-3.35 (2.26)	0.001	0.977
N400 Latency	433.50 (70.42)	359.16 (57.86)	35.786	<0.001*

169

170



171

172 **Figure 3:** Individual scatter plot data, including group means and standard deviations (middle) showing a
173 significant N400 latency and N100 amplitude differences for high-contact (red) versus low-contact
174 (blue).

175

176 Female compared to male subconcussive effects:

177

178

179

180

181

182


183

184

There was no significant sex by contact group interaction present [$F(6,104) = 0.047, p = 1.00, \eta_p^2 = 0.003$] (Table 1), indicating female and male athletes experienced similar subconcussive effects. Figure 4a shows brain vital sign radar plots for female and male athletes for high- versus low-contact and figure 4b provides the corresponding waveform results. The MANOVA confirmed a significant main effect for female athletes compared to male athletes [$F(6, 104) = 3.02, p = 0.009, \eta_p^2 = 0.148$] (Table 1) across these scans. Separate univariate between-subjects tests revealed significant effects for the P300 amplitude [$F(1) = 7.45, p = 0.007, \eta_p^2 = 0.064$] and N400 amplitude [$F(1) = 4.71, p = 0.032, \eta_p^2 = 0.041$] (Table 3). Note in Figure 4b that the vertical y-axis (Amplitude) for female athletes and male athletes is

BRENTWOOD Research Study Manuscript

185 fixed, highlighting the overall increased response amplitude in female responses relative to male
186 responses across all peaks.
187



188
189 **Figure 4:** (A) Radar plots comparing individual-level brain vital signs of the high-contact and low-contact
190 groups for both female and male athletes. (B) Group-level grand average ERP waveforms from tones
191 (top) and word pairs (bottom) for female athletes (left, dark shading) and male athletes (right, light
192 shading) comparing high- versus low-contact group differences. Deviant tones and incongruent word
193 pair waveforms are plotted (with standard error shading). N100, P300, and N400 brain vital sign
194 components of the high-contact group are highlighted with labels and arrows. Note in Figure 4b that the
195 vertical y-axis (Amplitude) for female athletes and male athletes is fixed, highlighting the overall
196 increased response amplitude in female responses relative to male responses across all peaks.

197
198
199 **Table 3:** MANOVA results for the main effect of sex on each brain vital sign component (high- and low-
200 contact scans).


BRENTWOOD Research Study
Manuscript

Brain vital sign component	Female athlete mean (standard deviation)	Male athlete mean (standard deviation)	F-value	p-value
N100 Amplitude	-4.29 (2.47)	-3.36 (2.47)	2.427	0.122
N100 Latency	94.28 (15.32)	101.86 (25.74)	2.597	0.110
P300 Amplitude	5.18 (3.24)	3.47 (2.85)	7.447	0.007*
P300 Latency	273.21 (37.50)	271.17 (37.04)	0.025	0.874
N400 Amplitude	-4.07 (2.96)	-3.00 (2.08)	4.707	0.032*
N400 Latency	391.67 (78.34)	385.91 (69.232)	0.053	0.819

201
202
203
204
205
206
207
208
209
210
211
212
213
214

Female compared to male effects:


To further characterize brain vital sign profiles, the multivariate main effect of sex, and the univariate results above (Table 3) for female and male athletes, contact condition was collapsed across high- and low-contact scans to create Figure 5. P300 and N400 amplitude differences accounts for a significant main effect with a near significant effect in the N100 amplitude. The trends driving the main effect can be seen in Figure 6, which showed the individual scatter plots across female athletes and male athletes, including means and standard deviations. To further characterize female and male effects with a larger and closely weighted group number, Figure 7 provides radar plot and waveform results from female and male athletes for all collected scans (N=202 scans). The purpose of this comparison was to include all scans as an evaluation of robust differences between female and male athletes. Corresponding statistics are in Supplemental Data 3.



215


BRENTWOOD Research Study Manuscript

216 **Figure 5:** Comparison of all female and male high- and low-contact scans showing amplitude differences.
217 **(A)** Radar plot showing overall female and male brain vital sign difference. **(B)** Group-level waveforms
218 from deviant tones (top) and incongruent word pairs (bottom) are plotted (standard error shading) for
219 female athletes (dark blue) and male athletes (light blue). N100, P300, and N400 components are
220 highlighted with arrows.
221
222



223 **Figure 6:** Scatter plots of individual-level data points for all scans of the amplitudes (top) and latencies
224 (bottom) for all brain vital signs components in female athletes (left) and male athletes (right).
225 Significant differences are highlighted by the asterisk, all p -values are reported as well (Bonferroni
226 adjusted).
227
228
229

BRENTWOOD Research Study Manuscript



230
231 **Figure 7:** Comparison of all female and male scans. **(A)** Radar plot showing individual-level brain vital
232 sign differences across all six components. **(B)** Group-level waveforms from deviant tones (top) and
233 incongruent word pairs (bottom) are plotted (standard error shading) confirming lower amplitudes in
234 male athletes (highlighted with arrows).

235
236 The current analyses examined high- versus low-contact, using equivalent exposure durations to
237 prior studies. For additional analyses (N=202 scans), contact exposure levels were compared between
238 low-contact exposure and both a single term and two or more terms of high-contact exposure. These
239 results were closely consistent with both subconscious and sex effects above and are provided as
240 Supplemental Data 4.

241 242 Discussion

243
244 The current findings replicated prior studies of subconscious changes in brain vital signs. The
245 study used a large, prospective, repeated measures design to compare high- versus low-contact exposure
246 across both female and male athletes. The findings supported the primary hypothesis that prior
247 subconscious differences in the N400 and N100 would be detectable in a high- versus low- contact
248 comparison (Figure 2), not only for male athletes but also in female athletes (Figure 4). Additional
249 comparisons of female and male results showed significantly larger P300 and N400 responses for female
250 athletes (Figure 6).

251
252 *Subconcussive analyses*

BRENTWOOD Research Study Manuscript

253 Subconcussive N400 changes replicate four previous studies on concussion (Fickling et al.
254 2019; Fickling, Smith, et al. 2021; Fickling, Poel, et al. 2021; Breuer et al. 2023). The N400 is commonly
255 associated with high-order cognitive semantic processing (D'Arcy et al. 2004; Kutas and Federmeier 2011)
256 and measures changes in semantic processing, suggesting the initial demonstration of cognitive changes
257 due to contact (Connolly and D'Arcy 2000; D'Arcy et al. 2003; Gawryluk et al. 2010; Ghosh Hajra et al.
258 2018). It is noteworthy that these replications have spanned across different sports and age ranges in
259 male athletes. However, the specific N400 changes have varied with respect to latency delays and/or
260 amplitude reductions, with the former directly measuring the rate of information processing and the latter
261 related to synchronous pyramidal neural activation (Luck 2014). The two are interdependent factors
262 reflecting a relative impairment in cognitive processing. While the initial results for female athletes
263 showed comparable N400 latency delays, a specific pattern of change in N400 latency versus amplitude
264 remains to be characterized.

265 Similarly, subconcussive changes in the N100 have been replicated across a number of prior
266 studies. While the studies have reported only N100 amplitude changes, the direction of the effects has
267 varied (i.e., decreases, increases, and correlated variability (Fickling et al. 2019; Fickling, Smith, et al. 2021;
268 Fickling, Poel, et al. 2021). In prior linear regression analyses of the relationship between brain vital signs
269 and helmet mounted accelerometers, inclusion of the N100 with the N400 provided additional
270 explanatory variance (Fickling, Poel, et al. 2021). Consequently, N100 changes suggest that subconcussive
271 impairments can span from lower-level sensory processing to higher level cognitive processing. As with
272 the N400 above, the common preliminary results showing N100 changes in both female and male athletes
273 do not rule out the possibility of specific differences. Indeed, examination of the female and male
274 waveforms shows early sensory and attention differences in other ERP components (see below).

275 Of note, the current findings are the first to compare brain vital signs between high- and low-
276 contact groups using a between-subjects design. To anchor to prior studies, a direct evaluation of the
277 within-subjects pre- versus post- design has also demonstrated a significant delay in the N400 latency
278 (Supplemental Data 2). The contribution of a control group is an important step, with the interpretation
279 further underpinned by the nature of the closely matched comparison (i.e., related to questions about
280 whether the effects are indeed attributable to subconcussive impact exposure). The student athletes in
281 this study attended a private school with full time residence. Accordingly, all students follow a highly
282 structured daily and weekly schedule, matched across programs, activities, diet, sleep, physical fitness,
283 rest, and related factors.

284 Another novel aspect of the current study was the ability to begin exploring differences between
285 a single term versus two or more terms of exposure to subconcussive impacts (Supplemental Data 4).
286 While preliminary, the findings suggested that there may be a graded effect of exposure on the N400 peak
287 latency (Supplemental Data 4). These results highlighted important questions related to tracking the
288 extent of exposure with increased temporal resolution to characterize the changes over time. As reflected
289 in the study design (Figure 1), a large cohort of these students were also re-evaluated after the summer
290 break to determine whether this duration of rest resulted in a return to pre- baseline levels following post-
291 subconcussive changes. Evaluation of these results are currently being analyzed as a follow-up study.
292 While the emerging subconcussive findings are concerning, the demonstrated utility of using brain vital
293 signs to objectively identify acute cognitive changes associated with repeated impact exposure has

BRENTWOOD Research Study Manuscript

294 enabled clinical trials on potential interventions. A recent study on subconcussion focused on a
295 prospective randomized placebo-controlled trial evaluating the effect of a dietary supplement effects
296 created to reduce neuroinflammation and support energy production, on brain structure and function in
297 Junior-A ice hockey players (Breuer et al. 2023). Breuer et al. (2023) randomized players into two groups:
298 a group that took a supplement containing a combination of non-pharmacological ingredients shown to
299 support brain health (Synaquell™ - Thorne Research Inc., NY, USA) daily over the season and a placebo
300 control group who followed the same protocol without supplementation. Multivariate analyses showed
301 significant neurocognitive pre- to post- season changes in only the control group, with significant negative
302 impacts on the N100 and N400 along with reduced saccadic eye movement (King-Devick Test) and
303 neurofilament light chain measures.

304

305 *Female and male comparison*

306 Robust P300 and N400 differences between female and male athletes were shown with and
307 without subconcussive contact exposure as a factor. While the significant P300 and N400 differences
308 demonstrated larger amplitudes for female athletes, examination of the overall waveforms showed this
309 finding generally across all ERP responses. There are at least two potential factors related to interpretation
310 of the difference between female and male athletes. One factor may relate to general attention difference
311 during the approximately 6-minute test time, whereby female athletes showed increased sustained
312 attention on average relative to male athletes. In this case, improved sustained attention to the task
313 would result in a higher signal-to-noise ratio in the signal averaged ERPs and consequently higher response
314 amplitudes. Alternatively, another factor may relate to a fundamental difference in ERP response size
315 between female and male individuals. While these two factors are not mutually exclusive, the notable
316 differences in baseline variance suggested sustained attention during the test time is likely a factor (Figure
317 5). Differences in attention between female and male participants has been noted in prior literature, but
318 these results are still mixed (Davidson, Cave, and Sellner 2000; Solianik, Brazaitis, and Skurvydas 2016;
319 Riley et al. 2016).

320 As mentioned above, despite the subconcussive similarities between female and male athletes, it
321 is too early to exclude differences, as an “absence of evidence is not evidence of absence.” For instance,
322 examination of the ERP waveforms (Figure 4) shows noteworthy subconcussive differences in responses
323 outside of the N100, P300, and N400. In particular, there appears to be a subconcussive reduction in the
324 N200-P200 difference in female athletes, which is not present in male athletes. The N200-P200 is
325 associated with early perceptual-attentional processing (Portella et al. 2012; D’Arcy, Connolly, and
326 Crocker 2000; Patel and Azzam 2005), and further research should explore this in the context of sex-
327 specific changes in subconcussion and head-impact exposure. With respect to this, the clinical importance
328 of even small attentional impairments can consequently have a larger impact on high-level processing for
329 individuals in attention demanding professions. For example, we identified early on in clinical concussion
330 evaluation that these changes can have a compounding effect on high demand jobs such as air traffic
331 controllers relative to less attentionally-demanding positions (Mateer and D’Arcy 2000). A comparable
332 impact may be a consideration for enhanced sustained attention processing in female athletes. Given
333 recent advances in understanding functional neuroanatomical differences (Andrushko et al. 2023), further
334 studies needed to better characterize female- versus male- specific subconcussive profiles.

BRENTWOOD Research Study Manuscript

335

336 *Limitations and Future Directions*

337 The current study included data across a large sample made up of smaller subsets across different
338 sports, with differing contact levels. While this was beneficial to create a large sample size for this study,
339 larger subsets of groups pertaining to each specific sport would help elucidate sport-specific differences.
340 Similarly, history of high-contact sport participation and concussion are important variables that may
341 affect the current state of participants. Nonetheless, the current study is the first in a continuing
342 investigation that will subsequently focus on specific sports and recovery times following subconcussive
343 impairment in contact sport. Future studies will expand to focus on recovery effects and differences
344 following summer break. Also, the current study combined subjects in grades 10-12 in a highly structured
345 school environment; however, previous work has shown differing neurological developmental rates
346 between female and male adolescents (Paus 2010; Kaczurkin, Raznahan, and Satterthwaite 2019; Gur
347 and Gur 2016), which may have been a factor for female and male-specific profiles. Future work will
348 include larger age subset sizes and longitudinal analyses, as well as controlling for age as a co-variate in
349 future models, to have a better understanding of developmental differences between female and male
350 individuals. Finally, the current study did not track important biological factors that differ between female
351 and male athletes. These factors include but are not limited to menstrual cycle phase at time of exposure
352 or in the case of multiple exposures, when the majority of exposures occurred (menstrual phase has been
353 a predictor of mild TBI outcome (Wunderle et al. 2014) and progesterone shows neuroprotective as a TBI
354 treatment (Wright et al. 2007)); body fat percentages given recent understanding of fat being a major
355 endocrine organ (Musi and Guardado-Mendoza 2014); and diet given the understanding of diet impact
356 on inflammation and brain function (Patel et al. 2023; Markovic et al. 2021). With clear neurophysiological
357 differences identified between females and males, future work must include expanded characterization
358 of these key biological factors to understand change that may affect results at the time of evaluation.
359 Lastly, to better understand the resulting effect of these subconcussive neurophysiological changes and
360 differences, additional study variables should include heart rate variability data to understand clinical and
361 sub-clinical autonomic dysfunction and time to recovery, as well as changes in sleep and endocrine levels
362 due to RHI, as TBI may lead to endocrine changes, specifically in developing adolescents (Richmond and
363 Rogol 2014).

364

365 **Conclusion**

366 With increasing attention on subconcussive impairments and potential interventions to limit
367 future risks (e.g., long-term neuropsychiatric impairment, chronic traumatic encephalopathy – CTE), the
368 current findings provide a large sample demonstration of subconcussive changes between closely
369 matched high- versus low-contact student athletes that replicate previous investigations. The findings
370 provide strong support for continued focus on the role of subconcussive injuries in sport-related brain
371 health research. Importantly, as this field of study has historically focused on male athletes, inclusion of
372 female athletes in the current findings represents a critical milestone towards understanding specific
373 neurophysiological differences before and after exposure to subconcussive impacts. Initial results strongly
374 support continued investigations that span beyond male athletes in helmeted contact sports.

375

BRENTWOOD Research Study
Manuscript

376 **Methods**

377
378 A total of 89 high school students were scanned 231 times over the year (34 females and 55
379 males). In total, 8 scans were removed due to attrition, 12 scans had a data collection error, failing to
380 produce scan data, 5 due to dropping all epochs for a condition (i.e. deviant tones), and 4 due to excess
381 noise (N=202 scans). For analysis and results we ended with N=81, n=202 (33 females and 48 males). The
382 current study focused on a scan-based between contact groups analysis, instead of a pre-post subject
383 based within-subjects analysis. I.e. results from a subject scan after a season of low contact sport would
384 be characterized as a low contact data point (not a benchmark or pre data point), while if that same
385 subject had a separate scan later in the school year after 2 terms of high contact, that scan would be
386 characterized as a high contact data point. This was done to maximize the total data point number for
387 analyses and provide more equal weighting across groups, as opposed to a repeated measure subject-
388 based analysis including only subjects that had not had any contact immediately prior to the study.

389 All healthy participants were in grades 10 through 12 (age range: 15-17), fluent in English, no
390 history of concussion in the past 6-months, no self-reported problems with brain function, and had no
391 hearing problems. All participants were recruited from Brentwood College School athletic programs,
392 which involve either high-contact (e.g., rugby, ice-hockey, soccer, field hockey, jujitsu, and basketball) or
393 no-contact/low-contact sports (e.g., rowing, tennis, and climbing) over three distinct three-month terms
394 in the school year (Fig. 1). As described above, Brentwood is a private boarding school in which all students
395 follow a structured daily schedule and a similar diet, exercise, and sleep schedule throughout the
396 academic year. Accordingly, participants were closely matched across a wide number of typical lifestyle
397 factors. The study had Advarra institutional research ethics board approval and all participants along with
398 their guardians provided informed consent.

399
400 **Data Collection**

401 The brain vital sign framework (Ghosh Hajra et al. 2016) was employed with the NeuroCatch®
402 Platform (Version 1.1), which uses EEG to rapidly extract the N100, P300, and N400 ERPs. All three ERPs
403 were stimulated using a compressed and standardized auditory stimulus sequence (approximately 6
404 minutes) consisting of sets of tones and spoken word pairs. The brain vital signs framework utilizes an
405 oddball paradigm, involving frequent standard tones with infrequent oddball tones, which elicit the N100
406 and P300 components. The tone stimuli precede word pairs, which are either matched or mismatched
407 semantically-related word-pair primes (e.g., match: bread-butter; mismatch: bread-window) to elicit the
408 N400 component. Participants passively listened to the stimuli and decided if the word pairs matched or
409 mismatched.

410 Participants were fitted with an elasticized cloth cap containing an 8-channel EEG amplifier with
411 standardized electrode locations (Fz, Cz, & Pz). Three additional electrodes were placed on the
412 participant's forehead to record the ground (GND) and electrooculogram (vEOG, hEOG) signals. A
413 reference electrode was clipped to the participant's right earlobe. Standard EEG skin preparation (70%
414 isopropyl alcohol wipes) and conductive gel products (SignaGel) were used to ensure appropriate contact
415 for all EEG sensors. Distractions were mitigated by performing the scans in a quiet and closed classroom.

BRENTWOOD Research Study Manuscript

416 The same environment was used for all scans. Participants were asked to listen attentively to the stimuli,
417 but no active response was required. To reduce motor and ocular artifacts, participants were instructed
418 to sit motionlessly, and maintain visual fixation on a cross positioned at eye level 2 m away.

419
420

421 Data Processing

422 The NeuroCatch® Platform was calibrated for any trigger latency delays. The most common delay
423 was identified and applied to shift all trigger data. EEG data were processed using a fourth order
424 Butterworth filter (0.5-10 Hz) and a Notch filter (60 Hz). Adaptive filtering using electrooculogram
425 channels was used to correct for ocular artifacts. EEG was epoched based on stimulus events using an
426 epoch from -100ms pre-stimulus to 1000ms post-stimulus. All epochs were linearly detrended and any
427 epochs with noise exceeding $\pm 75\mu\text{V}$ were rejected. N100, P300, and N400 peaks were calculated using
428 the maximal peak within a standard temporal window and manually verified. Amplitude and latency
429 metrics from these peaks were then linearly transformed into standardized scores on a scale from 0 to
430 100, derived from entire group means (Fickling et al. 2019). All preprocessing was completed in Python
431 using the Scipy and MNE libraries.

432
433

434 Statistical Analysis

435 All individual-level statistical analyses of peak amplitude and latencies were performed using
436 SPSS (Version 29.0.0, IBM, NY, USA). In the current study, individual-level can be defined as amplitudes
437 and latencies from each individual scan data, not derived from the group-level average waveforms. To
438 assess differences in brain vital signs between contact groups and sex, brain vital sign scores were
439 compared using a multivariate analysis of variance (MANOVA) (**Between:** Group: low-contact and high-
440 contact; Sex: Female and Male) on N100, P300, and N400 amplitudes and latencies for only scans
441 characterized as low-contact or high-contact for two or more terms. To further assess differences in
442 brain vital signs between sex in a larger number of scans, brain vital sign scores were compared using a
443 multivariate analysis of variance (MANOVA) (**Between:** Sex: Female and Male) on N100, P300, and N400
444 amplitudes and latencies for all 202 scans collected. All univariate results are adjusted for multiple
445 comparisons using Bonferroni adjustment. All statistics were also rerun only with subjects with less than
446 50% of epochs dropped (12 of 202 scans dropped) due to the $75\mu\text{V}$ thresholding measures mentioned
447 above. To keep our participant number large and because the analysis resulted in very similar outcomes,
448 the current manuscript reports the statistics from the full group data, including subjects with more than
449 50% of epochs dropped.

450
451

452 Acknowledgements

453

BRENTWOOD Research Study Manuscript

454 The authors would like to acknowledge the important leadership of Dr. Aynsley Smith (Mayo
455 Clinic, retired), whose passionate leadership for research advances in sports-related concussion and
456 ever-lasting priority to translate research advances to young females and males served as an important
457 inspiration for this study. The authors would also like to acknowledge the technical assistance by Joshua
458 Ighalo and the assistance in data collection by Laila Reed.
459

460 Funding

461 This work has been supported in part by funding from the Natural Sciences and Engineering
462 Research Council of Canada (Discovery Grant, PI R.C.N. D'Arcy; CGS M, E.D. Kirby).
463

464 Competing interests

465 Some of the authors (EDK, TY, NYB, SDF, RCND) have financial and/or business interests in
466 HealthTech Connex Inc., which may be affected by the research reported in this paper. Over the past 36
467 months, DWD reports the following competing interests: Consulting: Amgen, Atria, CapiThera Ltd.,
468 Cerecin, Ceruvia Lifesciences LLC, CoolTech, Ctrl M, Allergan, AbbVie, Biohaven, GlaxoSmithKline,
469 Lundbeck, Eli Lilly, Novartis, Impel, Satsuma, Theranica, WL Gore, Genentech, Nocira, Perfood, Praxis,
470 AYYA Biosciences, Revance, Pfizer. Honoraria: American Academy of Neurology, Headache Cooperative
471 of the Pacific, Canadian Headache Society, MF Med Ed Research, Biopharm Communications, CEA Group
472 Holding Company (Clinical Education Alliance LLC), Teva (speaking), Amgen (speaking), Eli Lilly
473 (speaking), Lundbeck (speaking), Pfizer (speaking), Vector Psychometric Group, Clinical Care Solutions,
474 CME Outfitters, Curry Rockefeller Group, DeepBench, Global Access Meetings, KLJ Associates, Academy
475 for Continued Healthcare Learning, Majallin LLC, Medlogix Communications, Medica Communications
476 LLC, MJH Lifesciences, Miller Medical Communications, WebMD Health/Medscape, Wolters Kluwer,
477 Oxford University Press, Cambridge University Press. Non-profit board membership: American Brain
478 Foundation, American Migraine Foundation, ONE Neurology, Precon Health Foundation, International
479 Headache Society Global Patient Advocacy Coalition, Atria Health Collaborative, Arizona Brain Injury
480 Alliance, Domestic Violence HOPE Foundation/Panfila. Research support: Department of Defense,
481 National Institutes of Health, Henry Jackson Foundation, Sperling Foundation, American Migraine
482 Foundation, Henry Jackson Foundation, Patient Centered Outcomes Research Institute (PCORI). Stock
483 options/shareholder/patents/board of directors: Aural analytics (options), Axon Therapeutics, ExSano
484 (options), Palion (options), Man and Science, Healint (options), Theranica (options), Second
485 Opinion/Mobile Health (options), Epien (options/board), Nocira (options), Matterhorn (shares/board),
486 Ontologics (shares/board), King-Devick Technologies (options/board), Precon Health (options/board),
487 AYYA Biosciences (options), Axon Therapeutics (options/board), Cephalgia Group (options/board), Atria
488 Health (options/employee). Patent 17189376.1-1466:vTitle: Onabotulinum Toxin Dosage Regimen for
489 Chronic Migraine Prophylaxis (Non-royalty bearing). Patent application submitted: Synaquell® (Precon
490 Health).
491

492 Author Contributions

493

BRENTWOOD Research Study
Manuscript

494 RCND assisted with project study design, manuscript drafting, and acted as lab PI. DM
495 contributed with data collection and project study design. DH contributed with data collection, data
496 processing, and manuscript editing. ZL contributed with data collection and manuscript editing. JW
497 contributed with data collection and manuscript editing. AH contributed with data collection, statistical
498 analysis, and manuscript editing. EDK contributed with data processing, statistical analysis, and
499 manuscript drafting. TY, NYB, SDF, TAM, DD, CA, and KSS contributed with manuscript editing.

500

501 References

502

- 503 Andrushko, Justin W., Shie Rinat, Eric D. Kirby, Julia Dahlby, Chelsea Ekstrand, and Lara A.
504 Boyd. 2023. "Females Exhibit Smaller Volumes of Brain Activation and Lower Inter-
505 Subject Variability during Motor Tasks." *Scientific Reports* 13 (1): 17698.
506 <https://doi.org/10.1038/s41598-023-44871-4>.
- 507 Breuer, Logan T., Shaun D. Fickling, David A. Krause, Ryan C. N. D'Arcy, Tory O. Frizzell,
508 Tania Gendron, and Michael J. Stuart. 2023. "The Effects of a Dietary Supplement on
509 Brain Function and Structure in Junior A Ice Hockey Players: A Prospective Randomized
510 Trial." medRxiv. <https://doi.org/10.1101/2023.04.03.23288079>.
- 511 Connolly, John F, and Ryan C. N D'Arcy. 2000. "Innovations in Neuropsychological Assessment
512 Using Event-Related Brain Potentials." *International Journal of Psychophysiology* 37 (1):
513 31–47. [https://doi.org/10.1016/S0167-8760\(00\)00093-3](https://doi.org/10.1016/S0167-8760(00)00093-3).
- 514 D'Arcy, Ryan C. N, John F Connolly, and Susan F Crocker. 2000. "Latency Shifts in the N2b
515 Component Track Phonological Deviations in Spoken Words." *Clinical Neurophysiology*
516 111 (1): 40–44. [https://doi.org/10.1016/S1388-2457\(99\)00210-2](https://doi.org/10.1016/S1388-2457(99)00210-2).
- 517 D'Arcy, Ryan C. N, Yannick Marchand, Gail A Eskes, Edmund R Harrison, Stephen J Phillips,
518 Alma Major, and John F Connolly. 2003. "Electrophysiological Assessment of Language
519 Function Following Stroke." *Clinical Neurophysiology* 114 (4): 662–72.
520 [https://doi.org/10.1016/S1388-2457\(03\)00007-5](https://doi.org/10.1016/S1388-2457(03)00007-5).
- 521 D'Arcy, Ryan C.N., John F. Connolly, Elisabet Service, Colin S. Hawco, and Michael E.
522 Houlihan. 2004. "Separating Phonological and Semantic Processing in Auditory
523 Sentence Processing: A High-Resolution Event-Related Brain Potential Study." *Human*
524 *Brain Mapping*. <https://doi.org/10.1002/hbm.20008>.
- 525 Davidson, Heather, Kyle R Cave, and Daniela Sellner. 2000. "Differences in Visual Attention
526 and Task Interference between Males and Females Reflect Differences in Brain
527 Laterality." *Neuropsychologia* 38 (4): 508–19. [https://doi.org/10.1016/S0028-3932\(99\)00084-6](https://doi.org/10.1016/S0028-3932(99)00084-6).
- 529 Davis, P. A. 1939. "Effects of Acoustic Stimuli on the Waking Human Brain." *Journal of*
530 *Neurophysiology* 2 (6): 494–99. <https://doi.org/10.1152/jn.1939.2.6.494>.
- 531 Dioso, Emma, John Cerillo, Mohammed Azab, Devon Foster, Isaac Smith, Owen Leary, Michael
532 Goutnik, and Brandon Lucke-Wold. 2022. "Subconcussion, Concussion, and Cognitive
533 Decline: The Impact of Sports Related Collisions." *Journal of Medical Research and*
534 *Surgery* 3 (4): 54–63. <https://doi.org/10.52916/jmrs224081>.
- 535 Fickling, Shaun D, Daniel N Poel, Jason C Dorman, Ryan C N D'Arcy, and Thayne A Munce.
536 2021. "Subconcussive Changes in Youth Football Players: Objective Evidence Using
537 Brain Vital Signs and Instrumented Accelerometers." *Brain Communications* 4 (2):
538 fcab286. <https://doi.org/10.1093/braincomms/fcab286>.
- 539 Fickling, Shaun D., Aynsley M. Smith, Gabriela Pawlowski, Sujoy Ghosh Hajra, Careesa C. Liu,
540 Kyle Farrell, Janelle Jorgensen, Xiaowei Song, Michael J. Stuart, and Ryan C.N. D'Arcy.

BRENTWOOD Research Study

Manuscript

- 541 2019. "Brain Vital Signs Detect Concussion-Related Neurophysiological Impairments in
542 Ice Hockey." *Brain* 142 (2): 255–62. <https://doi.org/10.1093/brain/awy317>.
- 543 Fickling, Shaun D, Aynsley M Smith, Michael J Stuart, David W Dodick, Kyle Farrell, Sara C
544 Pender, and Ryan C N D'Arcy. 2021. "Subconcussive Brain Vital Signs Changes Predict
545 Head-Impact Exposure in Ice Hockey Players." *Brain Communications* 3 (2): fcab019.
546 <https://doi.org/10.1093/braincomms/fcab019>.
- 547 Gawryluk, Jodie R., Ryan C.N. D'Arcy, John F. Connolly, and Donald F. Weaver. 2010.
548 "Improving the Clinical Assessment of Consciousness with Advances in
549 Electrophysiological and Neuroimaging Techniques." *BMC Neurology*.
550 <https://doi.org/10.1186/1471-2377-10-11>.
- 551 Ghosh Hajra, Sujoy, Careesa C. Liu, Xiaowei Song, Shaun D. Fickling, Teresa P. L. Cheung,
552 and Ryan C. N. D'Arcy. 2018. "Multimodal Characterization of the Semantic N400
553 Response within a Rapid Evaluation Brain Vital Sign Framework." *Journal of
554 Translational Medicine* 16 (June): 151. <https://doi.org/10.1186/s12967-018-1527-2>.
- 555 Ghosh Hajra, Sujoy, Careesa C. Liu, Xiaowei Song, Shaun Fickling, Luke E. Liu, Gabriela
556 Pawlowski, Janelle K. Jorgensen, et al. 2016. "Developing Brain Vital Signs: Initial
557 Framework for Monitoring Brain Function Changes Over Time." *Frontiers in
558 Neuroscience* 10. <https://www.frontiersin.org/article/10.3389/fnins.2016.00211>.
- 559 Gur, Raquel E., and Ruben C. Gur. 2016. "Sex Differences in Brain and Behavior in
560 Adolescence: Findings from the Philadelphia Neurodevelopmental Cohort."
561 *Neuroscience & Biobehavioral Reviews*, The Adolescent Brain, 70 (November): 159–70.
562 <https://doi.org/10.1016/j.neubiorev.2016.07.035>.
- 563 Guskiewicz, Kevin M., Stephen W. Marshall, Julian Bailes, Michael McCrea, Robert C. Cantu,
564 Christopher Randolph, and Barry D. Jordan. 2005. "Association between Recurrent
565 Concussion and Late-Life Cognitive Impairment in Retired Professional Football
566 Players." *Neurosurgery* 57 (4): 719.
567 <https://doi.org/10.1227/01.NEU.0000175725.75780.DD>.
- 568 Hirad, Adnan A., Jeffrey J. Bazarian, Kian Merchant-Borna, Frank E. Garcea, Sarah
569 Heilbronner, David Paul, Eric B. Hintz, et al. 2019. "A Common Neural Signature of Brain
570 Injury in Concussion and Subconcussion." *Science Advances* 5 (8): eaau3460.
571 <https://doi.org/10.1126/sciadv.aau3460>.
- 572 Kaczurkin, Antonia N., Armin Raznahan, and Theodore D. Satterthwaite. 2019. "Sex
573 Differences in the Developing Brain: Insights from Multimodal Neuroimaging."
574 *Neuropsychopharmacology* 44 (1): 71–85. <https://doi.org/10.1038/s41386-018-0111-z>.
- 575 Kutas, Marta, and Kara D. Federmeier. 2011. "Thirty Years and Counting: Finding Meaning in
576 the N400 Component of the Event-Related Brain Potential (ERP)." *Annual Review of
577 Psychology*. <https://doi.org/10.1146/annurev.psych.093008.131123>.
- 578 Luck, Steven J. 2014. *An Introduction to the Event-Related Potential Technique, Second
579 Edition*. MIT Press.
- 580 Mainwaring, Lynda, Kaleigh M. Ferdinand Pennock, Sandhya Mylabathula, and Benjamin Z.
581 Alavie. 2018. "Subconcussive Head Impacts in Sport: A Systematic Review of the
582 Evidence." *International Journal of Psychophysiology*, Long-term outcomes of sport-
583 related brain injuries: A psychophysiological perspective, 132 (October): 39–54.
584 <https://doi.org/10.1016/j.ijpsycho.2018.01.007>.
- 585 Markovic, Shaun J., Melinda Fitzgerald, Jeremiah J. Peiffer, Brendan R. Scott, Stephanie R.
586 Rainey-Smith, Hamid R. Sohrabi, and Belinda M. Brown. 2021. "The Impact of Exercise,
587 Sleep, and Diet on Neurocognitive Recovery from Mild Traumatic Brain Injury in Older
588 Adults: A Narrative Review." *Ageing Research Reviews* 68 (July): 101322.
589 <https://doi.org/10.1016/j.arr.2021.101322>.

BRENTWOOD Research Study

Manuscript

- 590 Mateer, Catherine A., and Ryan C. N. D'Arcy. 2000. "Current Concepts in Assessment and
591 Management." In *Neuropsychological Management of Mild Traumatic Brain Injury*, edited
592 by Sarah A. Raskin and Catherine A. Mateer. New York, NY, USA: Oxford University
593 Press.
- 594 McKee, Ann C., Robert C. Cantu, Christopher J. Nowinski, E. Tessa Hedley-Whyte, Brandon E.
595 Gavett, Andrew E. Budson, Veronica E. Santini, Hyo-Soon Lee, Caroline A. Kubilus, and
596 Robert A. Stern. 2009. "Chronic Traumatic Encephalopathy in Athletes: Progressive
597 Tauopathy After Repetitive Head Injury." *Journal of Neuropathology & Experimental*
598 *Neurology* 68 (7): 709–35. <https://doi.org/10.1097/NEN.0b013e3181a9d503>.
- 599 Miller, Jeff, Rolf Ulrich, and Wolfgang Schwarz. 2009. "Why Jackknifing Yields Good Latency
600 Estimates." *Psychophysiology* 46 (2): 300–312. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8986.2008.00761.x)
601 [8986.2008.00761.x](https://doi.org/10.1111/j.1469-8986.2008.00761.x).
- 602 Musi, Nicolas, and Rodolfo Guardado-Mendoza. 2014. "Chapter 14 - Adipose Tissue as an
603 Endocrine Organ." In *Cellular Endocrinology in Health and Disease*, edited by Alfredo
604 Ulloa-Aguirre and P. Michael Conn, 229–37. Boston: Academic Press.
605 <https://doi.org/10.1016/B978-0-12-408134-5.00014-7>.
- 606 Nowak, Madeleine K., Zachary W. Bevilacqua, Keisuke Ejima, Megan E. Huibregtse, Zhongxue
607 Chen, Timothy D. Mickleborough, Sharlene D. Newman, and Keisuke Kawata. 2020.
608 "Neuro-Ophthalmologic Response to Repetitive Subconcussive Head Impacts: A
609 Randomized Clinical Trial." *JAMA Ophthalmology* 138 (4): 350–57.
610 <https://doi.org/10.1001/jamaophthalmol.2019.6128>.
- 611 Patel, Palak R., Patrick Armistead-Jehle, Nicholas R. Eltman, Kelly M. Heath, David X. Cifu, and
612 Randel L. Swanson. 2023. "Brain Injury: How Dietary Patterns Impact Long-Term
613 Outcomes." *Current Physical Medicine and Rehabilitation Reports* 11 (3): 367–76.
614 <https://doi.org/10.1007/s40141-023-00413-7>.
- 615 Patel, Salil H., and Pierre N. Azzam. 2005. "Characterization of N200 and P300: Selected
616 Studies of the Event-Related Potential." *International Journal of Medical Sciences* 2 (4):
617 147–54.
- 618 Paus, Tomáš. 2010. "Chapter 2 - Sex Differences in the Human Brain: A Developmental
619 Perspective." In *Progress in Brain Research*, edited by Ivanka Savic, 186:13–28. Sex
620 Differences in the Human Brain, Their Underpinnings and Implications. Elsevier.
621 <https://doi.org/10.1016/B978-0-444-53630-3.00002-6>.
- 622 Portella, Claudio, Sergio Machado, Oscar Arias-Carrión, Alexander T. Sack, Julio Guilherme
623 Silva, Marco Orsini, Marco Antonio Araujo Leite, et al. 2012. "Relationship Between
624 Early and Late Stages of Information Processing: An Event-Related Potential Study."
625 *Neurology International* 4 (3): e16. <https://doi.org/10.4081/ni.2012.e16>.
- 626 Richmond, Erick, and Alan D. Rogol. 2014. "Traumatic Brain Injury: Endocrine Consequences in
627 Children and Adults." *Endocrine* 45 (1): 3–8. <https://doi.org/10.1007/s12020-013-0049-1>.
- 628 Riley, Elizabeth, Hidefusa Okabe, Laura Germine, Jeremy Wilmer, Michael Esterman, and
629 Joseph DeGutis. 2016. "Gender Differences in Sustained Attentional Control Relate to
630 Gender Inequality across Countries." *PLoS ONE* 11 (11): e0165100.
631 <https://doi.org/10.1371/journal.pone.0165100>.
- 632 Saghafi, Behrouz, Gowtham Murugesan, Elizabeth Davenport, Ben Wagner, Jillian Urban,
633 Mireille Kelley, Derek Jones, et al. 2018. "Quantifying the Association between White
634 Matter Integrity Changes and Subconcussive Head Impact Exposure from a Single
635 Season of Youth and High School Football Using 3D Convolutional Neural Networks."
636 *Proceedings of SPIE--the International Society for Optical Engineering* 10575
637 (February): 105750E. <https://doi.org/10.1117/12.2293023>.
- 638 Smulders, Fren T. Y. 2010. "Simplifying Jackknifing of ERPs and Getting More out of It:
639 Retrieving Estimates of Participants' Latencies." *Psychophysiology* 47 (2): 387–92.

BRENTWOOD Research Study

Manuscript

- 640 <https://doi.org/10.1111/j.1469-8986.2009.00934.x>.
- 641 Snedaker, Katherine Price, Jennifer P. Lundine, Angela H. Ciccio, Mohammad Nadir Haider,
642 and Katy H. O'Brien. 2022. "Gaps in Concussion Management across School-Aged
643 Children." *Brain Injury* 36 (6): 714–21. <https://doi.org/10.1080/02699052.2022.2034954>.
- 644 Solianik, Rima, Marius Brazaitis, and Albertas Skurvydas. 2016. "Sex-Related Differences in
645 Attention and Memory." *Medicina* 52 (6): 372–77.
646 <https://doi.org/10.1016/j.medic.2016.11.007>.
- 647 Sutton, Samuel, Patricia Tueting, Joseph Zubin, and E. R. John. 1967. "Information Delivery
648 and the Sensory Evoked Potential." *Science* 155 (3768): 1436–39.
649 <https://doi.org/10.1126/science.155.3768.1436>.
- 650 Ulrich, Rolf, and Jeff Miller. 2001. "Using the Jackknife-Based Scoring Method for Measuring
651 LRP Onset Effects in Factorial Designs." *Psychophysiology* 38 (5): 816–27.
652 <https://doi.org/10.1111/1469-8986.3850816>.
- 653 Valera, Eve M., Annie-Lori C. Joseph, Katherine Snedaker, Matthew J. Breiding, Courtney L.
654 Robertson, Angela Colantonio, Harvey Levin, et al. 2021. "Understanding Traumatic
655 Brain Injury in Females: A State-of-the-Art Summary and Future Directions." *The Journal*
656 *of Head Trauma Rehabilitation* 36 (1): E1–17.
657 <https://doi.org/10.1097/HTR.0000000000000652>.
- 658 Veksler, Ronel, Udi Vazana, Yonatan Serlin, Ofer Prager, Jonathan Ofer, Nofar Shemen,
659 Andrew M Fisher, et al. 2020. "Slow Blood-to-Brain Transport Underlies Enduring Barrier
660 Dysfunction in American Football Players." *Brain* 143 (6): 1826–42.
661 <https://doi.org/10.1093/brain/awaa140>. Markovic, Shaun J., Melinda Fitzgerald, Jeremiah
662 J. Peiffer, Brendan R. Scott, Stephanie R. Rainey-Smith, Hamid R. Sohrabi, and Belinda
663 M. Brown. 2021. "The Impact of Exercise, Sleep, and Diet on Neurocognitive Recovery
664 from Mild Traumatic Brain Injury in Older Adults: A Narrative Review." *Ageing Research*
665 *Reviews* 68 (July): 101322. <https://doi.org/10.1016/j.arr.2021.101322>.
- 666 Wright, David W., Arthur L. Kellermann, Vicki S. Hertzberg, Pamela L. Clark, Michael Frankel,
667 Felicia C. Goldstein, Jeffrey P. Salomone, et al. 2007. "ProTECT: A Randomized Clinical
668 Trial of Progesterone for Acute Traumatic Brain Injury." *Annals of Emergency Medicine*
669 49 (4): 391-402.e2. <https://doi.org/10.1016/j.annemergmed.2006.07.932>.
- 670 Wunderle, Kathryn, Kathleen M. Hoeger, Erin Wasserman, and Jeffrey J. Bazarian. 2014.
671 "Menstrual Phase as Predictor of Outcome After Mild Traumatic Brain Injury in Women."
672 *The Journal of Head Trauma Rehabilitation* 29 (5): E1.
673 <https://doi.org/10.1097/HTR.0000000000000006>.
- 674