

**Title:** Development of Prediction Models of COVID-19 Vaccine Uptake among Lebanese and Syrians in a district of Beirut, Lebanon: a population-based study

**Authors:** Marie-Elizabeth Ragi MSc<sup>1</sup>, Hala Ghattas PhD<sup>1,2</sup>, Hazar Shamas MSc<sup>1</sup>, Jocelyn DeJong PhD<sup>3</sup>, Nada M. Melhem PhD<sup>4</sup>, Stephen J. McCall DPhil<sup>1\*</sup> with the CAEP Study Group<sup>¶</sup>

### **Affiliations**

<sup>1</sup>Center for Research on Population and Health, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon

<sup>2</sup>Department of Health Promotion, Education, and Behavior, University of South Carolina, Columbia, South Carolina, USA

<sup>3</sup>Department of Epidemiology and Population Health, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon

<sup>4</sup>Medical Laboratory Sciences Program, Division of Health Professions, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon

\*Corresponding author: Stephen J McCall, Center for Research on Population and Health, Faculty of Health Sciences, American University of Beirut, Beirut 1107 2020, Lebanon  
sm227@aub.edu.lb

¶ Membership of the CAEP Study Group is listed in the Acknowledgments.

**Word count:** 3043

**Keywords:** COVID-19, Vaccination, Prediction model, Health inequalities

## Abstract

**Introduction:** Vaccines are essential to prevent infection and reduce morbidity of infectious diseases. Previous evidence has shown that migrants and refugees are particularly vulnerable to exclusion and discrimination, and low COVID-19 vaccine intention and uptake were observed among refugees globally. This study aimed to develop and internally validate prediction models of COVID-19 vaccine uptake by nationality.

**Methods:** This is a nested prognostic population-based cross-sectional analysis. Data was collected between June and October 2022 in Sin-El-Fil, a district of Beirut, Lebanon. All Syrian adults and a random sample of adults from low-socioeconomic status neighborhoods were invited to participate in a telephone survey. The main outcome was uptake of COVID-19 vaccine. Predictors of COVID-19 vaccine uptake were assessed using LASSO regression for Lebanese and Syrian nationalities, respectively.

**Results:** Of 2,045 participants, 79% were Lebanese, 18% Syrians and 3% of other nationalities. COVID-19 vaccination uptake was higher among Lebanese (85% (95%CI:82-86)) compared to Syrians (47% (95% CI:43-51)) ( $P < 0.001$ ); adjusted odds ratio (aOR) 6.8 (95% CI:5.5-8.4). Predictors of uptake of one or more vaccine dose for Lebanese were older age, presence of an older adult in the household, higher education, greater asset-based wealth index, private healthcare coverage, feeling susceptible to COVID-19, belief in the safety and efficacy of vaccines and previous receipt of flu vaccine. For Syrians they were older age, male, completing school or higher education, receipt of cash assistance, presence of comorbidities, belief in the safety and efficacy of vaccines, previous receipt of flu vaccine, and legal residency status in Lebanon.

**Conclusions:** These findings indicate barriers for vaccine uptake in Syrian migrants and refugees, including legal residency status. They call for urgent action to enable equitable access to vaccines by raising awareness about the importance of vaccination and the targeting of migrant and refugee populations through vaccination campaigns.

## **Key Messages**

### **What is already known on this topic**

Vaccines are essential to prevent infection and reduce morbidity of infectious diseases, and vulnerable populations may lack access to vaccination campaigns.

### **What this study adds**

To the best of our knowledge, no studies have examined compared predictors of COVID-19 vaccine uptake and measured the rate of vaccination among between Syrian migrants and refugees and their Lebanese host communities. This study illustrates a clear difference in vaccine uptake between nationalities and developed prediction models among the Syrian and Lebanese that identified differential predictors of COVID-19 vaccine uptake for each population.

### **How this study might affect research, practice or policy**

These findings indicate barriers for vaccine uptake in Syrian migrants and refugees, including legal status, and calls for urgent action to enable access to vaccines by raising awareness about the importance of vaccination against COVID-19 in vulnerable groups and targeting migrant and refugee populations through vaccination campaigns.

## Introduction

Vaccines were an essential public health tool to mitigate the impact of the COVID-19 pandemic. In Lebanon, vaccines were available free of charge to all residents starting February 2021 following the WHO SAGE risk-and age-based approach for prioritization (1-4). The COVID-19 national vaccination plan aimed to reach community immunization levels of at least 70-80% by the end of 2022 (2-4), yet there were subpopulations that remained unvaccinated. A survey amongst older Syrian refugees in Lebanon showed that only 42.5% were vaccinated against COVID-19 by March 2022 (5), while the national average for all residents of Lebanon was 50% by October 2022 (6).

Lebanon hosts the largest number of refugees per capita in the world, including Syrian, Palestinian and Iraqi refugees, that account for approximately one third of the population, many of which are dispersed in urban residential areas living amidst Lebanese communities (4, 7). Migrants and refugees are particularly vulnerable to exclusion, stigma and discrimination that may limit access to vaccination campaigns (7). Required renewal of residency permits per year at significant economic costs adds additional burden (8). Studies in other settings have shown that refugees and non-native ethnic groups were less willing to get vaccinated against COVID-19 as compared to native populations (9-13), with racial discrimination and lack of trust in authorities' facilities and systems being identified as determinants of COVID-19 vaccine hesitancy (12, 13). Thus, the factors that impact vaccine uptake are likely to differ between national and non-national populations, and understanding demographic variations and predictors of vaccination status between these subpopulations remains important to inform the response to future outbreaks of COVID-19 and future pandemics.

This study aims to determine whether there are differences in COVID-19 vaccine uptake by nationality in a suburb of Beirut, Lebanon, and to develop and internally validate prediction models that examine context-specific differential predictors of COVID-19 vaccine uptake by nationality.

## **Methods**

### ***Study design and setting***

This is a nested cross-sectional analysis within an ongoing multi-wave longitudinal study that aims to track vulnerabilities to COVID-19 and other health emergencies in Sin-EI-Fil, a suburb of Beirut, Lebanon. The parent study focused on subpopulations with vulnerabilities that may increase risk of COVID-19 infection, morbidity or mortality. These included 1) adults aged 60 years or older, 2) pregnant women, 3) adults living in areas of low socioeconomic status, and 4) Syrian migrants and refugees. The study protocol was reviewed and approved by the American University of Beirut Social and Behavioral Sciences Institutional Review Board (reference: SBS-2021-0268).

### ***Sampling and study population***

This was a multi-stage stratified sampling design using an area-based sampling (14). The suburb of Sin-EI-Fil was stratified into areas of low and high socioeconomic status, with boundaries defined geographically through stakeholder consultation with the Sin-EI-Fil municipality and NGOs. A household listing exercise was carried out in April 2022 whereby all households in Sin-EI-Fil were enumerated face-to-face to complete an eligibility screening survey, which identified households and individuals with the specific vulnerability criteria mentioned above (15). Pre-consent was obtained from all households with eligible individuals and phone numbers were collected from consenting households. For the study sample selection, a listing of all eligible household members was generated. A sample of older adults and a sample of adults between 18-60 years old were randomly selected using proportionate allocation. Several of the vulnerability groups were small so these group were oversampled; this included the selection of all Syrian adults and pregnant women. For the present analysis, the study population included all participants living in areas of low socioeconomic status who completed the first wave of data collection conducted between June and October 2022 (n=2,045).

Respondents were contacted to complete a computer assisted telephone survey conducted by a trained data collector and data were entered on SurveyCTO software (Dobility Inc., Cambridge, MA, USA). Verbal informed consent to participate in an oral telephone interview was obtained from selected respondents and those aged 60 years or older were furthermore assessed for

capacity to participate using 5 modified items from the University of California, San Diego, Brief Assessment of Capacity to Consent (16).

### ***Data sources***

The questionnaire was developed using existing validated scales and questionnaire modules, contextualized questions, and community-identified priorities. The survey was created in collaboration with representatives from the municipality of Sin-EI-Fil, the Ministry of Public Health (MOPH), and Non-Governmental Organizations (NGOs) operating in Sin-EI-Fil. The survey tool was drafted in English, translated to Arabic and tested for comprehension prior to deployment. Data were frequently monitored in parallel with data collection for quality assurance, with 5% call back checks conducted to ensure accuracy of the collected data.

### ***Outcome measures***

COVID-19 vaccine uptake of at least one dose was the primary outcome of interest. Each participant was asked the following questions: “Have you received the COVID-19 vaccine?” and “If yes, how many doses did you receive?”. The outcome is presented as binary: “received the COVID-19 vaccine” and “did not receive the COVID-19 vaccine”. Additionally, unvaccinated participants were asked if they planned to take the COVID-19 vaccine and the reasons why not.

### ***Possible predictors***

Based on the literature on COVID-19 vaccine acceptance and hesitancy, several possible predictors for vaccine uptake were included. For Lebanese, these were: age, sex, education, household assets wealth index, presence of an older adult in the household, presence of chronic illness, healthcare coverage, COVID-19 knowledge and risk perceptions (susceptibility to COVID-19), general perceptions of vaccine safety and effectiveness, and receipt of the flu vaccine. The household assets-based wealth index was generated through Principal Component Analysis (PCA) from variables on presence in the household of functional transportation, communication, home technology and cooking assets. The chronic illnesses assessed included hypertension, type II diabetes, vascular diseases, dyslipidemia, chronic respiratory diseases, rheumatoid arthritis, chronic kidney diseases, cancer. For Syrians, these were age, sex, education, receipt of cash assistance and legal residency status in the country that are context specific to Syrian migrants and refugees, presence of chronic illness, healthcare coverage,

COVID-19 knowledge and risk perceptions, general perceptions of vaccine safety and effectiveness, and receipt of the flu vaccine. The missing values were assumed to be missing at random. The variable with the highest amount of missing values was at 6.3% and a complete case analysis was performed (17).

### *Statistical analysis*

Frequencies and percentages are presented for categorical variables, and medians with their interquartile range (IQR) for continuous variables. The analysis accounted for the complex design and non-response. Using unadjusted logistic regression analyses, the association between COVID-19 vaccine uptake and the possible predictors was examined among the different nationalities, unadjusted odds ratios (ORs) along with their 95% confidence intervals (CIs) are reported. The outcome was a binary variable for uptake of the COVID-19 vaccine. All variables were categorical except age, which was found to have a linear association with COVID-19 vaccine uptake.

A Least Absolute Shrinkage and Selection Operator (LASSO) logistic regression model was used to identify the predictors of vaccine uptake for each nationality (Lebanese and Syrian), and all corresponding candidate predictors were entered into each model. The final models were internally validated using cross-validation (10-folds). Multicollinearity was assessed using correlation matrices where a variance inflation factor (VIF) greater than five indicated collinearity. The final models' performances were assessed through their discrimination capabilities using the Area Under the Receiver Operating Curve (AUC), which ranges from 0.5 (representing a discriminative ability equal to chance), to 1.0 (reflecting a perfect discriminative ability between those with and without the outcome). The evaluation of the models' calibration, which determines the agreement between observed outcomes and predictive probabilities, was performed by analyzing calibration plots and slope. An ideal calibration is illustrated by a diagonal line on the graph with an intercept of 0 and a slope of 1. However, a slope less than 1 suggests overfitting in the model, meaning that respondents with high risk of the outcome have overestimated risk predictions while those with low risk of the outcome have underestimated risk predictions (18). Additionally, calibration-in-the-large (CITL) was assessed to identify the overall difference between the observed number of events and the average predictive risk. LASSO for inference analysis was conducted, which allowed the reporting of ORs and CIs. A

secondary analysis among Syrian migrants and refugees explored the association of regularized residency (possession of legal documents) on vaccination by logistic regression adjusted for confounders, including sex, education and years of living in Lebanon. All analyses were conducted using Stata/SE statistical software version 18 (STATA Corp).

The study followed the TRIPOD reporting guidelines and STROBE reporting guidelines for prediction modelling (19, 20).

## Results

A total of 2,045 participants who completed the first wave of data collection were included in this analysis (Supp. Figure 1), of whom 79% were Lebanese (n=1,322), 18% Syrians (n=664) and 3% of other nationalities (n=59), while 0.8% (n=17) had a missing outcome. Median age of the study sample was 47 years (IQR 32-62, range 18-98) and 45.6% were males.

As of October 2022, COVID-19 vaccine uptake among study participants was 77%. There were differences in vaccination status between nationalities (Figure 1) with half (53%) of Syrians not having received any doses of the COVID-19 vaccine followed by other nationalities (39%) while 15% of Lebanese had not received any doses ( $P<0.001$ ). In addition, the Lebanese population in this study had received a greater number of vaccine doses with 82% vaccinated having received 2 doses or more (vs. 39% and 55% for Syrians and other nationalities, respectively). Most common reasons reported for lack of vaccine uptake were: (1) not believing the vaccine is essential, and (2) preferring to follow other precautions - by both Lebanese (51% and 25%, respectively) and Syrians (45% and 23%, respectively). For those vaccinated with only one dose (3% of Lebanese and 8% of Syrians), the main reason was not wanting to receive the second dose for fear of side effects for Lebanese (44%), while among Syrian migrants and refugees the highest proportion stated they were still waiting for a second dose (41%) (Supp. Table 1).

For Lebanese adults, older age, the presence of an older adult in the household, the highest tertile of asset-based wealth index, the presence of at least one multi-comorbidity, having private or public funds healthcare coverage, belief that COVID-19 is a serious infection, considering themselves susceptible to COVID-19, belief in the safety and efficacy of vaccines in general, and receipt of the flu vaccine were all associated with increased odds of receiving the COVID-19 vaccine (Table 1).



For Syrian migrants and refugees, similarly to Lebanese, older age, belief that COVID-19 is a serious infection, considering themselves susceptible to COVID-19, belief in the safety and efficacy of vaccines in general, and receipt of the flu vaccine were associated with increased odds of receiving the COVID-19 vaccine; while being male and having regularized residency were additional variables found to increase the odds of vaccination amongst Syrian migrants and refugees (Table 2). In a sub-group analysis amongst Syrian migrants and refugees, having legal residency status in Lebanon, a context specific factor to Syrian migrants and refugees, doubled the odds of COVID-19 vaccine uptake (adjusted OR:1.7 (95%CI:1.1-2.8)). An unadjusted bivariate analysis for other nationalities found COVID-19 vaccine uptake associated with attitudes towards COVID-19 and general vaccination (Supp. Table 2).

The final models retained eight predictors of COVID-19 vaccine uptake for both the Lebanese and Syrian populations respectively. For Lebanese, identified predictors were older age, presence of an older adult in the household, higher education, greater asset-based wealth index, private healthcare coverage (i.e., having private or mutual insurance), feeling susceptible to COVID-19, belief in the safety and efficacy of vaccines, and previously received the flu vaccine (Table 3). While among Syrian migrants and refugees, the predictors were greater age, male sex, having completed school or higher education, receipt of cash assistance, presence of co-morbidities, belief in the safety and efficacy of vaccines, previously received the flu vaccine, and legal residency status in Lebanon (Table 4).

Both models showed good discrimination with an AUC of 0.823 (95%CI:0.786-0.859) for the Lebanese sample and 0.768 (95%CI:0.730-0.806) for the Syrian sample. The model for the Lebanese population had good calibration with a calibration slope of 1.061 (95%CI:0.905-1.217) and CITL of 0.004 (Supp. Figure 2). The model for the Syrian population had underfitting with a calibration slope of 1.199 (95%CI:0.955-1.442) and CITL of -0.012 (Supp. Figure 3).

To illustrate the model amongst Syrian migrants and refugees, a 40-year-old male Syrian with no legal documentation, who completed school, is not receiving cash assistance, does not have a co-morbidity, does not believe in the safety and efficacy of vaccines, and has never taken the flu vaccine has a 4% chance of having taken the COVID-19 vaccine.

## Discussion

This study revealed inequalities in COVID-19 vaccination status between Syrian migrants and refugees and Lebanese adults residing in a low socioeconomic area in Beirut. Overall, 53% of Syrian migrant and refugee adults had not received any doses of the COVID-19 vaccine compared to 15% among Lebanese adults. This study also developed and internally validated a prediction model that identified eight predictors of COVID-19 vaccine uptake of at least one dose for the Lebanese and Syrian populations, respectively. This study found socioeconomic, demographic and health risk perceptions to predict vaccine uptake for both Syrians and Lebanese. In addition, legal residency status in the country and sex were found to lead to differentials in vaccine uptake within the Syrian population.

The proportion of vaccinated Syrian migrants and refugees in this sample was similar to another study examining COVID-19 vaccination in Syrian refugees aged 50 years or older in Lebanon, which reported 42.5% having received at least one dose of the COVID-19 vaccine by mid-March 2022 (5). Furthermore, vaccination status in Syrian migrants and refugees in this sample was slightly lower than the 50% national COVID-19 vaccination average recorded at the time of data collection (6). In contrast, the proportion of vaccinated Lebanese reported in this study at 85% exceeds this national prevalence, and the majority (82%) were vaccinated with 2 doses or more which reached the national immunization target of 70% (3). These findings demonstrate large differentials in vaccination uptake by nationality, which indicate barriers to vaccination uptake amongst Syrian migrants and refugees. Thus, understanding the reasons for not taking the vaccine remains important to design contextualized public health interventions.

A previous study conducted on older Syrian refugees living in Lebanon found similar predictors, bar the direction of the association of age and vaccine uptake; likely due to the larger age range included in the present study (5). To the best of our knowledge, there are no prediction models for COVID-19 vaccine uptake among Lebanese adults or younger Syrian migrants and refugees.

Some of the identified predictors of COVID-19 vaccine uptake in this study, older age and education level, were common for both Lebanese (national) and Syrian (non-national) populations, which was consistent with studies conducted in the MENA region (21, 22), the UK (10), the US (11, 23, 24), and a global meta-analysis (25). In general, older adults have physiological vulnerabilities and co-morbidities, and thus may perceive themselves at higher risk

of infection and severe complications (21, 23, 25). Higher education level was associated with higher likelihood of vaccine uptake (5) and higher willingness to receive the vaccine (10, 21-26). This may be explained by those with a higher educational level being more aware of the risks of the disease, the importance of vaccination and having the ability to critically appraise vaccine misinformation (12, 23). Health and COVID-19 risk beliefs and perceptions, as well as trust in vaccination effectiveness following positive past experiences, also played a role in vaccine acceptance and uptake in line with other studies (11, 21, 23-25, 27).

The socio-economic indicators identified, where both Lebanese in the highest tertile of wealth and Syrian migrants and refugees receiving cash assistance had a higher likelihood of vaccine uptake, were in accordance with studies in low and middle-income countries reported low socio-economic status as a determinant of low willingness to take the COVID-19 vaccine (26). Studies in other context also align, as in the US low income was associated with lower vaccine uptake and greater vaccine hesitancy (23, 24). Despite the COVID-19 vaccine being available free of charge to all residents (3), this highlights that there may be other barriers to attaining the vaccine specifically for Syrian migrants and refugees.

Legal residency status (having a legal residency permit in Lebanon) was found as a context specific predictor of COVID-19 vaccine uptake for Syrian migrants and refugees. The majority of the Syrian migrants and refugees (79%) reported not having a legal residency permit; this was likely due to the high costs and requirements relating to obtaining and renewing annual residency permits (8). Previous studies in other countries have also shown that lack of legal residency permit among migrants impacts health care access (28). Consequently, lack of legal residency may have impacted vaccination uptake as residents of any nationality in Lebanon were required to register on an online government platform (3), thus Syrian migrants and refugees, particularly those without documentation, may have worried that registering may have led to arrest or deportation (8, 28, 29).

### ***Strengths and Limitations***

This study has several strengths and limitations. The study outcome was self-reported which may be prone to information bias; however, recall of vaccinations amongst adults was shown to be relatively accurate (30). The study sample may not be representative of all adults in Lebanon, yet it is representative of this low socio-economic suburb in Beirut. The model was slightly

underfitted for Syrians and a larger sample may help prevent this in the future, whilst this study had a large enough sample size of Lebanese adults to be able to develop a well calibrated model.

### ***Conclusion***

By October 2022, more than half the Syrian migrants and refugees living in a low socio-economic area of a suburb of Beirut were still not vaccinated against COVID-19 despite the availability of vaccines free of charge to all residents including non-nationals in Lebanon, as opposed to the majority of Lebanese from the same area having received the vaccine. Predictors for both populations include older age, educational level, socio-economic status, and general attitude toward vaccination. The findings indicate barriers for vaccine uptake in Syrian migrants and refugees, including legal status. This calls for urgent action to eliminate barriers and enable equitable access to vaccines for all vulnerable population in outbreaks such as the COVID-19 pandemic.

## **Contributors**

HG, JD and SJM conceptualized the study; SJM, HG and MER designed the survey, contribution to study methodology and investigation, and oversaw the data collection. The formal analysis and literature search was conducted by MER. MER and SJM drafted the manuscript. SJM supervised MER throughout the project. MER, HG, HS, JD, NM and SJM contributed to the interpretation of the results, and have edited drafts and approved the final version of the Article. MER, HG and SJM had full access to and verified the raw data. All authors had access to the study data and had final responsibility for the decision to submit the manuscript for publication.

## **Funding**

This work was funded by the International Development Research Centre (IDRC) – Canada (grant number: 103964; project number: 25941).

## **Declaration of interests**

The authors declare no competing interests.

## **Data sharing**

The anonymized data can be obtained upon reasonable request from the Center for Research on Population and Health at the American University of Beirut ([crph@aub.edu.lb](mailto:crph@aub.edu.lb)).

## **Acknowledgments**

We would like to thank the “Community Action for Equity in Pandemic preparedness and control” CAEP Study Group for supporting this work (Aline Germani, Fadi El-Jardali). We thank Berthe Abi Zeid for her support and expertise in data analysis. We acknowledge BOT (Bridge. Outsource. Transform) for their assistance in collecting the data required for the success of this study and the study participants for their participation.

## References

1. Impact Lebanese Government Platform. COVID-19 vaccination open data [Available from: <https://impact.gov.lb/home?dashboardName=vaccine-breakdown>].
2. Republic of Lebanon Ministry of Public Health. Lebanon National Deployment and Vaccination Plan for COVID-19 Vaccines. 2021.
3. Abou Hassan FF, Hamdan MB, Ali F, Melhem NM. Response to COVID-19 in Lebanon: update, challenges and lessons learned. *Epidemiology & Infection*. 2023;151:e23.
4. Jawad NK, Taweeleh LA, Elharake JA, et al. Refugee access to COVID-19 vaccines in Lebanon. *The Lancet*. 2021;397(10288):1884.
5. Abi Zeid B, El Khoury T, Ghattas H, et al. Predictors and barriers to vaccination among older Syrian refugees in Lebanon: a cross-sectional analysis of a multi-wave longitudinal study. *The Lancet Healthy Longevity*. 2023;4(5):e219-e27.
6. Mathieu E, Ritchie H, Ortiz-Ospina E, et al. Coronavirus Pandemic (COVID-19) OurWorldInData.org2020 [Available from: <https://ourworldindata.org/covid-vaccinations>].
7. Fouad FM, McCall SJ, Ayoub H, Abu-Raddad LJ, Mumtaz GR. Vulnerability of Syrian refugees in Lebanon to COVID-19: quantitative insights. *Conflict and Health*. 2021;15(1):1-6.
8. Kikano F. Informality versus short-term regularization of the Syrian refugees' situation in Lebanon. *Enhancing Disaster Preparedness: Elsevier*; 2021. p. 209-22.
9. Shaw J, Anderson KB, Fabi RE, et al. COVID-19 vaccination intention and behavior in a large, diverse, US refugee population. *Vaccine*. 2022;40(9):1231-7.
10. Robertson E, Reeve KS, Niedzwiedz CL, et al. Predictors of COVID-19 vaccine hesitancy in the UK household longitudinal study. *Brain, behavior, and immunity*. 2021;94:41-50.
11. Gerretsen P, Kim J, Caravaggio F, et al. Individual determinants of COVID-19 vaccine hesitancy. *PloS one*. 2021;16(11):e0258462.
12. Biswas MR, Alzubaidi MS, Shah U, Abd-Alrazaq AA, Shah Z. A scoping review to find out worldwide COVID-19 vaccine hesitancy and its underlying determinants. *Vaccines*. 2021;9(11):1243.
13. Savoia E, Piltch-Loeb R, Goldberg B, et al. Predictors of COVID-19 vaccine hesitancy: socio-demographics, co-morbidity, and past experience of racial discrimination. *Vaccines*. 2021;9(7):767.
14. Qader SH, Lefebvre V, Tatem AJ, et al. Using gridded population and quadtree sampling units to support survey sample design in low-income settings. *International Journal of Health Geographics*. 2020;19:1-16.
15. Kalton G. Methods for oversampling rare subpopulations in social surveys. *Survey methodology*. 2009;35(2):125-41.
16. Jeste DV, Palmer BW, Appelbaum PS, et al. A new brief instrument for assessing decisional capacity for clinical research. *Archives of general psychiatry*. 2007;64(8):966-74.
17. Harrell FE. *Regression modeling strategies: with applications to linear models, logistic regression, and survival analysis*: Springer; 2001.
18. Steyerberg EW, Steyerberg EW. Overfitting and optimism in prediction models. *Clinical prediction models: A practical approach to development, validation, and updating*. 2019:95-112.
19. Collins GS, Reitsma JB, Altman DG, Moons KG. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD) the TRIPOD statement. *Circulation*. 2015;131(2):211-9.

20. Vandembroucke JP, Elm Ev, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Annals of internal medicine*. 2007;147(8):W-163-W-94.
21. Dadras O, SeyedAlinaghi S, Karimi A, et al. Public acceptability of COVID-19 vaccines and its predictors in Middle Eastern/North African (MENA) countries: a systematic review. *Human vaccines & immunotherapeutics*. 2022;18(5):2043719.
22. Al-Mohaithef M, Padhi BK. Determinants of COVID-19 vaccine acceptance in Saudi Arabia: a web-based national survey. *Journal of multidisciplinary healthcare*. 2020:1657-63.
23. Roberts HA, Clark DA, Kalina C, et al. To vax or not to vax: Predictors of anti-vax attitudes and COVID-19 vaccine hesitancy prior to widespread vaccine availability. *PloS one*. 2022;17(2):e0264019.
24. Wang Y, Liu Y. Multilevel determinants of COVID-19 vaccination hesitancy in the United States: A rapid systematic review. *Preventive medicine reports*. 2022;25:101673.
25. Wang Q, Yang L, Jin H, Lin L. Vaccination against COVID-19: A systematic review and meta-analysis of acceptability and its predictors. *Preventive medicine*. 2021;150:106694.
26. Moola S, Gudi N, Nambiar D, et al. A rapid review of evidence on the determinants of and strategies for COVID-19 vaccine acceptance in low-and middle-income countries. *Journal of global health*. 2021;11.
27. Salibi N, Abdulrahim S, El Haddad M, et al. COVID-19 vaccine acceptance in older Syrian refugees: Preliminary findings from an ongoing study. *Preventive medicine reports*. 2021;24:101606.
28. Matlin SA, Smith AC, Merone J, et al. The challenge of reaching undocumented migrants with COVID-19 vaccination. *International Journal of Environmental Research and Public Health*. 2022;19(16):9973.
29. Kaloti R, Fouad FM. The politics of COVID-19 vaccine equity among refugee populations in Lebanon. *Journal of Global Health Economics and Policy*. 2022;2:e2022003.
30. Tjaden AH, Fette LM, Edelstein SL, et al. Self-reported SARS-CoV-2 vaccination is consistent with electronic health record data among the COVID-19 community research partnership. *Vaccines*. 2022;10(7):1016.

## Tables and Figures

### **Figure 1 - COVID-19 vaccine uptake among study participants.**

*\*Uptake of one dose of COVID-19 vaccine was 4% for the total sample, 3% for Lebanese, 8% for Syrians, and 6% for other nationalities.*



**Table 1 - Characteristics of the Lebanese population in our study area and associations with COVID-19 vaccine uptake**

	Total (n=1,313)		Not vaccinated (n=214)		Vaccinated (n=1,099)		COVID-19 vaccine uptake
	n	Weighted %	n	Weighted %	n	Weighted %	Weighted Unadjusted OR [95% CI]
<b>Age (years) Median(IQR)</b>	1304	52 (35-64)	212	42 (33-57)	1092	54 (36-65)	1.02 [1.01-1.03]
Missing	9		2		7		
<b>Sex</b>							1
Male	588	(45.0)	93	(43.8)	495	(45.2)	
Female	725	(55.0)	121	(56.2)	604	(54.8)	0.95 [0.70-1.28]
Missing	0		0		0		
<b>Presence of an older adult in the household</b>							1
No	565	(40.3)	125	(55.7)	440	(37.5)	
Yes	743	(59.7)	89	(44.3)	654	(62.5)	2.10 [1.55-2.84]
Missing	5		0		5		
<b>Education</b>							1
School not attended	103	(8.8)	24	(12.6)	79	(8.1)	
School not completed	578	(47.6)	89	(43.8)	489	(48.3)	1.72 [1.03-2.87]
School completed	174	(14.0)	30	(14.5)	144	(13.9)	1.49 [0.81-2.75]
Vocational	117	(9.3)	21	(10.5)	96	(9.1)	1.35 [0.70-2.61]
Higher Education	251	(20.3)	38	(18.6)	213	(20.6)	1.73 [0.97-3.08]
Missing	90		12		78		
<b>Assets-based wealth index</b>							1
1	137	(10.5)	31	(14.1)	106	(9.8)	
2	558	(43.4)	100	(48.1)	458	(42.6)	1.27 [0.80-2.03]
3	587	(46.1)	77	(37.8)	510	(47.6)	1.81 [1.12-2.92]
Missing	31		6		25		
<b>Presence of chronic illnesses</b>							1
No	811	(60.5)	153	(70.3)	658	(58.7)	
Yes	502	(39.5)	61	(29.7)	441	(41.3)	1.66 [1.20-2.30]
Missing	0		0		0		
<b>Healthcare coverage</b>							1
Nothing	669	(53.8)	145	(69.3)	524	(50.9)	
Private and mutual funds	78	(6.3)	6	(3.2)	72	(6.9)	2.90 [1.23-6.84]

Public funds and NGOs/UN agencies	487	(39.9)	55	(27.5)	432	(42.2)	2.08 [1.48-2.93]
Missing	79		8		71		
<b>Believing COVID-19 is a serious infection</b>							
True	1121	(86.6)	168	(79.9)	953	(87.8)	1
False	172	(13.4)	41	(20.1)	131	(12.2)	0.55 [0.37-0.82]
Missing	20		5		15		
<b>Consider themselves susceptible to a COVID-19 infection</b>							
No	310	(23.5)	75	(34.9)	235	(21.4)	1
Yes	997	(76.5)	136	(65.1)	861	(78.6)	1.97 [1.42-2.71]
Missing	6		3		3		
<b>Vaccines are safe and/or effective</b>							
Agree	1023	(78.3)	86	(39.0)	937	(85.4)	1
Neither agree or disagree	182	(13.7)	56	(26.8)	126	(11.4)	0.19 [0.13-0.29]
Disagree	106	(8.0)	71	(34.2)	35	(3.2)	0.04 [0.03-0.07]
Missing	2		1		1		
<b>Previous receipt of the flu vaccine</b>							
No	915	(73.7)	169	(83.4)	746	(71.9)	1
Yes	328	(26.3)	34	(16.6)	294	(28.1)	1.96 [1.31-2.93]
Missing	70		11		59		

Data are n and weighted % or median (IQR), OR=odds ratio, CI=confidence interval.

Age was a continuous variable; increasing age was associated with increased odds of receiving the COVID-19 vaccine.

The wealth from assets index was generated from the presence in the household of functional transportation, communication, home technology and cooking assets. The chronic illnesses assessed included hypertension, type II diabetes, vascular diseases, dyslipidemia, chronic respiratory diseases, rheumatoid arthritis, chronic kidney diseases, cancer.

**Table 2 - Characteristics of the Syrian population in our study area and associations with COVID-19 vaccine uptake**

	Total (n=658)		Not vaccinated (n=355)		Vaccinated (n=303)		COVID-19 vaccine uptake
	n	Weighted %	n	Weighted %	n	Weighted %	Weighted Unadjusted OR [95% CI]
<b>Age (years) Median(IQR)</b>	656	35 (26-42)	353	32 (24-41)	303	37 (28-45)	1.02 [1.01-1.04]
Missing	2		2		0		
<b>Sex</b>							1
Male	336	(50.3)	154	(41.9)	182	(59.6)	
Female	322	(49.7)	201	(58.1)	121	(40.4)	0.49 [0.35-0.68]
Missing	0		0		0		
<b>Education</b>							1
School not attended	93	(14.9)	58	(16.7)	35	(12.8)	
School not completed	450	(71.2)	248	(72.5)	202	(69.7)	1.25 [0.76-2.04]
School completed	51	(8.5)	20	(6.4)	31	(10.8)	2.21 [1.03-4.72]
Vocational	17	(2.7)	10	(2.9)	7	(2.5)	1.11 [0.36-3.42]
Higher Education	17	(2.7)	6	(1.5)	11	(4.2)	3.45 [1.12-10.61]
Missing	30		13		17		
<b>Receiving cash assistance</b>							1
No	357	(56.5)	203	(58.9)	154	(53.9)	
Yes	281	(43.5)	139	(41.1)	142	(46.1)	1.23 [0.88-1.71]
Missing	20		13		7		
<b>Presence of chronic illness</b>							1
No	533	(79.8)	289	(79.3)	244	(80.4)	
Yes	125	(20.2)	66	(20.7)	59	(19.6)	0.93 [0.61-1.42]
Missing	0		0		0		
<b>Healthcare coverage</b>							1
Nothing	500	(77.4)	266	(77.2)	234	(77.6)	
Private, mutual and public funds	3	(0.4)	2	(0.5)	1	(0.3)	0.56 [0.05-6.29]
NGOs/UN agencies	143	(22.2)	79	(22.3)	64	(22.1)	0.98 [0.66-1.46]
Missing	12		8		4		
<b>Believing COVID-19 is a serious infection</b>							1
True	588	(90.3)	307	(87.7)	281	(93.2)	
False	60	(9.7)	41	(12.3)	19	(6.8)	0.52 [0.28-0.97]
Missing	10		7		3		

<b>Consider themselves susceptible to a COVID-19 infection</b>							
No	161	(23.9)	98	(27.5)	63	(20.0)	1
Yes	495	(76.1)	255	(72.5)	240	(80.0)	1.51 [1.03-2.23]
Missing	2		2		0		
<b>Vaccines are safe and/or effective</b>							
Agree	444	(68.4)	183	(51.8)	261	(86.8)	1
Neither agree or disagree	142	(20.8)	106	(29.0)	36	(11.8)	0.24 [0.16-0.38]
Disagree	69	(10.8)	64	(19.2)	5	(1.4)	0.04 [0.02-0.11]
Missing	3		2		1		
<b>Previous receipt of the flu vaccine</b>							
No	449	(72.5)	260	(78.4)	189	(65.9)	1
Yes	184	(27.5)	80	(21.6)	104	(34.1)	1.88 [1.30-2.71]
Missing	25		15		10		
<b>Have regularized residency</b>							
No	518	(78.9)	299	(84.4)	219	(72.9)	1
Yes	133	(21.1)	52	(15.6)	81	(27.1)	2.00 [1.31-3.05]
Missing	7		4		3		

Data are n and weighted % or median (IQR), OR=odds ratio, CI=confidence interval.

Age was a continuous variable; increasing age was associated with increased odds of receiving the COVID-19 vaccine.

The chronic illnesses assessed included hypertension, type II diabetes, vascular diseases, dyslipidemia, chronic respiratory diseases, rheumatoid arthritis, chronic kidney diseases, cancer.

**Table 3 - Predictors of vaccine acceptance among Lebanese adults in a district of Beirut**

	Codes	Coefficients*	OR [95%CI]^
<b>Age (years)</b>		0.02	1.03 [1.01-1.05]
<b>Presence of an older adult in the household</b>			
No	0		Ref.
Yes	1	0.46	1.73 [1.15-2.60]
<b>Education</b>			
School not attended	1	-0.62	Ref.
School not completed	2	-	2.42 [1.16-5.05]
School completed	3	-	2.51 [0.93-6.77]
Vocational	4	-	2.67 [1.06-6.74]
Higher Education	5	0.02	3.02 [1.26-7.21]
<b>Assets-based wealth index</b>			
1	1	-0.08	Ref.
2	2	-	1.13 [0.64-1.99]
3	3	0.46	1.97 [1.09-3.55]
<b>Healthcare coverage</b>			
Nothing	0	-0.84	Ref.
Private and mutual funds	1	0.19	4.12 [1.40-12.13]
Public funds and NGOs/UN agencies	2	-	2.55 [1.62-3.99]
<b>Consider themselves susceptible to a COVID-19 infection</b>			
No	0		Ref.
Yes	1	0.22	1.33 [0.87-2.02]
<b>Vaccines are safe and/or effective</b>			
Agree	1	1.57	Ref.
Neither agree or disagree	2	-	0.19 [0.12-0.30]
Disagree	3	-1.50	0.04 [0.02-0.07]
<b>Previous receipt of the flu vaccine</b>			
No	0		Ref.
Yes	1	0.04	1.17 [0.60-1.96]
<b>Intercept</b>		-0.25	

<b>AUC</b>			0.823 [0.786-0.859]
<b>C-Slope</b>			1.061 [0.905-1.217]
<b>CITL</b>			0.004

*AUC=Area Under the Receiver Operating Curve, C-Slope=Calibration-slope, CITL=calibration-in-the-large.*

*OR=odds ratio, CI=confidence interval.*

*\*Obtained from the LASSO logistic regression analysis. ^Obtained from the LASSO for inference analysis.*

**Table 4 - Predictors of vaccine acceptance among Syrian adults in a district of Beirut**

	Codes	Coefficients*	OR [95%CI]^
<b>Age (years)</b>		0.02	1.03 [1.01-1.05]
<b>Sex</b>			
Male	0	-	Ref.
Female	1	-0.51	0.51 [0.34-0.74]
<b>Education</b>			
School not attended	1	-0.03	Ref.
School not completed	2	-	1.25 [0.68-2.30]
School completed	3	0.43	2.49 [1.00-6.16]
Vocational	4	-	1.25 [0.38-4.16]
Higher Education	5	0.39	3.02 [0.64-14.29]
<b>Receiving cash assistance</b>			
No	0	-	Ref.
Yes	1	0.19	1.52 [1.00-2.30]
<b>Presence of chronic illness</b>			
No	0	-	Ref.
Yes	1	0.02	1.21 [0.73-2.02]
<b>Vaccines are safe and/or effective</b>			
Agree	1	1.26	Ref.
Neither agree or disagree	2	-	0.25 [0.15-0.41]
Disagree	3	-1.12	0.05 [0.01-0.14]
<b>Previous receipt of flu vaccine</b>			
No	0	-	Ref.
Yes	1	0.35	1.68 [1.10-2.56]
<b>Have regularized residency</b>			
No	0	-	Ref.
Yes	1	0.51	2.07 [1.21-3.56]
<b>Intercept</b>		-1.77	
<b>AUC</b>			0.768 [0.730-0.806]
<b>C-Slope</b>			1.199 [0.955-1.442]

<b>CITL</b>				-0.012
-------------	--	--	--	--------

*AUC=Area Under the Receiver Operating Curve, C-Slope=Calibration-slope, CITL=calibration-in-the-large.  
OR=odds ratio, CI=confidence interval.*

*\*Obtained from the LASSO logistic regression analysis. ^Obtained from the LASSO for inference analysis.*



All (n=2028)

77%  
vaccinated

23% Not vaccinated

\*

44%  
2 doses

29%  
3 doses  
or more

Lebanese (n=1313)

85%  
vaccinated

15% Not vaccinated

\*

46%  
2 doses

36%  
3 doses  
or more

Syrian (n=658)

47%  
vaccinated

53% Not vaccinated

\*

35%  
2 doses

4%  
3 doses

Other  
nationalities  
(n=57)

61%  
vaccinated

39% Not vaccinated

\*

45%  
2 doses

10%  
3 doses

