1	The iCARE feasibility non-experimental design study: An integrated collection of education
2	modules for fall and fracture prevention for healthcare providers in long term care
3	
4	Isabel B. Rodrigues ^{1,2} , George Ioannidis ¹ , Lauren Kane ¹ , Loretta M. Hillier ³ , Caitlin McArthur ⁴ ,
5	Jonathan Adachi ¹ , Lehana Thabane ^{1,3,5,6,7} , George Heckman ^{8,9} , Jayna Holroyd-Leduc ¹⁰ , Susan
6	Jaglal ¹¹ , Sharon Kaasalainen ¹² , Sharon Straus ^{2,13} , Momina Abbas ^{3,5} , Jean-Eric Tarride ⁵ , Sharon
7	Marr ² , John Hirdes ⁹ , Arthur N. Lau ¹ , Andrew Costa ⁵ , Alexandra Papaioannou ^{1,3,5*}
8	¹ McMaster University, Faculty of Health Sciences, Department of Medicine, Hamilton, ON
9	² Unity Health Toronto St. Michael's Hospital, Knowledge Translation Program, Toronto, ON
10	³ Geras Centre for Aging Research, Hamilton, ON
11	⁴ Dalhousie University, Faculty of Health, School of Physiotherapy, Halifax, NS
12 13	⁵ McMaster University, Faculty of Health Sciences, Department of Health Research Methods, Evidence, and Impact, Hamilton, ON
14	⁶ Research Institute of St Joes Hamilton, St Joseph's Healthcare Hamilton, Hamilton ON
15	⁷ Faculty of Health Sciences, University of Johannesburg, Johannesburg, South Africa
16	⁸ Schlegel-UW Research Institute for Aging, Waterloo, ON
17	⁹ University of Waterloo, School of Public Health Sciences, Waterloo, Ontario
18 19	¹⁰ University of Calgary, Departments of Medicine and Community Health Sciences, Calgary, AB
20	¹¹ University of Toronto, Department of Physical Therapy, Toronto, ON
21	¹² McMaster University, School of Nursing, Hamilton, ON
22 23	¹³ University of Toronto, Department of Medicine, Toronto, ON
24	*Corresponding author
25	Email: papaioannou@hhsc.ca

26 ABSTRACT

Falls and hip fractures are a major health concern among older adults in long term care (LTC) with 27 almost 50% of residents experiencing a fall annually. Hip fractures are one of the most important 28 and frequent fall-related injuries in LTC. The purpose of this study was to determine the feasibility 29 (recruitment rate and adaptations) of implementing the PREVENT (Person-centred Routine 30 Fracture PreEVENTion) model in practice, with a subobjective to understand facilitators and 31 barriers. The model includes a multifactorial intervention on diet, exercise, environmental 32 adaptations, hip protectors, medications (including calcium and vitamin D), and medication 33 reviews to treat residents at high risk of fracture. Our secondary outcomes aimed to assess change 34 in knowledge uptake of the guidelines among healthcare providers and in the proportion of fracture 35 prevention prescriptions post-intervention. We conducted a mixed-methods non-experimental 36 design study in three LTC homes across southern Ontario. A local champion was selected to guide 37 the implementation. We reported recruitment rates using descriptive statistics and adaptations 38 39 using content analysis. We reported changes in knowledge uptake using the paired sample t-test and the percentage of osteoporosis medications prescriptions using absolute change. Within five 40 months, we recruited three LTC homes. We required two months to identify and train the local 41 champion over three 1.5-hour train-the-trainer sessions, and the champion required three months 42 43 to deliver the intervention to the healthcare team. We identified several facilitators, barriers, and adaptations. Benefits of the model include easy access to the Fracture Risk Scale, clear and 44 45 succinct educational material catered to each healthcare professional, and an accredited educational module for physicians and nurses. Challenges included misperceptions between the 46 47 differences in fall and fracture prevention strategies, fear of perceived side effects associated with fracture prevention medications, and time barriers with completing the audit report. Our study did 48 49 not increase knowledge uptake of the guidelines, but there was an increase in the proportion of osteoporosis medication post-intervention. 50

51

52 INTRODUCTION

Falls and hip fractures are a major health concern among older adults (1). An estimated 53 one in three individuals aged 65 years and older fall each year, and almost half of individuals living 54 in long term care (LTC) fall annually (2). Hip fractures in residents living in LTC are devastating 55 and are likely to increase with current demographic trends in aging (3,4). Each year, almost 50% 56 of residents in LTC facilities fall at least once, and 40% of residents fall twice or more (3,5). Of 57 those LTC residents that fall, 10% to 25% of falls are associated with serious injuries requiring 58 medical treatment with 2% to 6% resulting in fractures of the hip, wrist, or vertebra (6-8). Hip 59 fractures are particularly life altering as both mobility and the capacity to live independently are 60 affected (3,6–8). Moreover, about half of the residents in LTC who experience a hip fracture may 61 die within the next year or develop total dependence within six months of their hip fracture (3,6-62 8). Thus, interventions to reduce falls and fractures in LTC are important to decrease morbidity 63 and mortality. 64

65 In 2015, the Canadian recommendations for preventing fractures in LTC were released to facilitate an evidence-based decision-making process (9). Indeed, guidelines have the capacity to 66 promote high quality practice informed by evidence, enable appropriate resource allocation, and 67 advance research by identifying knowledge gaps (10); however, the existence of guidelines alone 68 is not enough to change practice (10,11). The lack of guideline uptake in practice is evident with 69 the ongoing high morbidity and mortality associated with fractures in LTC (12–14). The major 70 gap is the limited evidence on effective knowledge translation strategies to prevent falls and low 71 trauma fractures in residents who are at high risk of fractures (12,15,16). Active strategies that 72 utilize an integrated knowledge translation approach may help to uptake guidelines into practice 73 (12). Current methods to uptake guidelines involve co-developing models with end-users and 74 stakeholders and piloting their implementation under real-world conditions (17). 75

PREVENT (Person-centred <u>R</u>outine Fracture Pre<u>EVENT</u>ion) is an educational outreach model for delivering education on fall and fracture prevention to healthcare providers working in LTC. The model includes a multifactorial intervention on diet, exercise, environmental adaptations to reduce falls, hip protectors, medication (including calcium and vitamin D), and medication reviews. The model, herein known as PREVENT, aligns with the 2015 Canadian evidence-based recommendation (9) and with recommendations from Australia and USA (18). PREVENT utilizes

a multifactorial knowledge mobilization strategy that engages the entire multi-disciplinary LTC
team (e.g., physicians, nurses, physiotherapists, dietitians, personal support workers) and builds
automated identification of high-risk fractures into routine care processes.

PREVENT may have several benefits including reducing falls and fractures and improving 85 quality of life of residents in LTC. Before a trial can be implemented to determine the effectiveness 86 of PREVENT in practice, we need to determine the feasibility of implementing it under real-world 87 conditions. The purpose of the iCare (integrated collection of education modules for fall and 88 fracture prevention) study was to determine the feasibility of implementing PREVENT to 89 healthcare providers in LTC. Our secondary outcomes were to determine if PREVENT improved 90 91 knowledge uptake of the Canadian recommendations (9) and increased the proportion of osteoporosis medication prescriptions post-intervention. We selected the proportion of change in 92 93 osteoporosis medications since these medications currently have the highest certainty of evidence to reduce the risk and the rate of fractures (19,20) and are the easiest outcome to measure in 94 95 practice.

96 METHODS

97 Study design

We conducted a pre-post non-experimental design study in three LTC homes across Ontario. We conducted our study in accordance with the STROBE 2007 guidelines (21) and TIDieR 2014 checklist (22). We registered our study on clinicaltrials.gov on July 6, 2022 (NCT05445336); we updated the registry on December 21, 2022. We received ethics approval from the Hamilton Integrated Research Ethics Board (Project ID 14463).

103 Study setting

Between January to June 2022, we sent emails to five large LTC corporate offices detailing the iCARE study. Each corporation owned between three to 30 homes across Ontario. Two LTC corporations were willing to participate in the feasibility study. We recruited one for-profit and two not-for profit LTC homes, herein known as Home A (120 beds), Home B (425 beds), and Home C (240 beds). All three LTC homes are in southern Ontario, Canada in large metropolitan cities.

110 Knowledge translation model

111 PREVENT is an organizational-level, service provision approach that embeds falls and 112 fracture education and management strategies in LTC. The model utilizes several intervention

- functions and behaviour change techniques based on the Behaviour Change Wheel (Table 1)
- 114 (23,24). The specific PREVENT components are detailed in S1.

Intervention	Definition of	Rehaviour change	Rehaviour change
function	intervention function	technique	technique example
Behaviour Change Wheel			
Education	Improving knowledge and understanding of roles and responsibilities for healthcare professionals responsible for service provision and improving knowledge through a community of practice (24)	 Feedback on behaviour Prompts/cues Self-monitoring of behaviour 	• Audit and Feedback Report
Training	Providing key professionals with core skills in management and leadership (24)	 Demonstration of the behaviour Instruction on how to perform a behaviour 	 Train the trainer Educational videos
Modelling	Discussion of identified successful strategies for change within network (24)	• Demonstration of the behaviour	 Train the trainer Educational videos Training manual Fracture and Fall Prevention Guide

115 Table 1: Intervention functions and behaviour change techniques used in PREVENT

PREVENT was designed to be integrated into regularly scheduled healthcare sessions (e.g., Professional Advisory Committee, Falls Committee), which usually occur quarterly in LTC. We identified a local champion to help guide the implementation of the model and continue to promote best practices (25). The standardized model includes: 1) train-the-trainer sessions with the local champion (26); 2) a multidisciplinary educational session with fracture prevention care recommendations and real-world case studies (27); 3) orientation to the Fracture Prevention

Toolkit developed by the lead researcher (IBR) that is provided to each home with the Fracture 122 Risk Scale (FRS) which identifies residents at high risk of fracture (score between 4 to 8) and low 123 124 risk of fracture (score between 1 to 3) (28), templates, standard order sets, and quick reference guide to the 2015 Canadian LTC fracture prevention guidelines (29); 4) an audit and feedback 125 report developed by the local champion for physicians and healthcare providers regarding the 126 number of residents at high versus low risk of fracture, number of residents who have experienced 127 a fragility fracture in the last six months, and the number of high risk of fracture residents on 128 osteoporosis medications (30); and 5) implementation intentions that includes an implementation 129 proposal template, care planning template, and Process Indicator Checklist (home-level policies 130 that support and sustain fracture prevention best practices such as training and education of front-131 line staff including personal support workers) (31). We utilized a three-step process to implement 132 PREVENT. In step one (train-the-trainer), the research team worked with each home to identify a 133 local champion. We sought champions who the home's management team considered trustworthy 134 and influential, acted as a role model for behaviour change, supported and legitimized the work, 135 provided a mechanism of communication between the research team and healthcare providers in 136 137 LTC, and served to sustain gains in the long term (25). The lead investigator (IBR) met with each local champion using an online platform (i.e., Zoom) and in-person at the LTC home to review 138 139 how to deliver PREVENT, which included reviewing how to use the FRS to identify residents at high and low risk of fracture (28), a training manual (see S2), Fracture and Fall Prevention Guide 140 141 (see S3), educational module, and other point of care tools. Our processes were guided by the Getting to Outcomes Framework, which includes a 10-step program for implementing, evaluating, 142 143 and continuously improving prevention programs (32). In step two, the local champion adapted the educational meeting to the LTC home and developed the audit and feedback report. During the 144 145 third step, the local champion presented the adapted PREVENT material during the leadership team meeting which included physicians, nurse practitioners, registered nurses, dietitians, 146 physiotherapists, kinesiologists, personal support workers, and interRAI coordinators. InterRAI 147 coordinators are part of the interRAI network and are responsible for collecting health outcomes 148 in LTC (https://interrai.org/about-interrai/). During the meeting, the leadership team also 149 brainstormed and implemented intentions and care plans to identify and treat residents at high risk 150 of fracture. LTC homes were remunerated for their time and participation in the iCARE study. 151

152 Intervention timeline

	Intervention Timeline
Month 1 and Month 2	 LTC home recruitment Identify local champion
Month 3	1. Train-the-trainer to prepare audit and feedback report and adapt/deliver the educational module
Month 4	Interactive Session 11. Local champion leads education module with the leadership team2. Audit and feedback report review3. Develop action and care plan
Months 4 to Month 6	1. LTC home implements action and care plan
Month 7	Interactive Session 21. Focus group with local champion and leadership team to understand barriers, facilitators, and adaptations2. One-on-one interview with local champion to understand barriers, facilitators, and adaptations

153 Inclusion/exclusion criteria

As this is a pragmatic study, our inclusion and exclusion criteria were broad and generalizable to represent real world practice. Our organizational level inclusion criterion was to include homes with a minimum of 70 beds, which is the sample size for our larger clinical trial (i.e., PREVENT trial).

158 Outcomes

159 Primary outcome

The primary outcome was feasibility of implementation defined by the number of LTC homes recruited within two months, length of time for the local champion to deliver the PREVENT program, and the length of time it took the local champion to develop the audit and feedback report (Table 2) (33). The lead investigator (IBR) developed a checklist prior to the start of the study to assess fidelity; our method to develop the checklist was defined using a five-step guide proposed by Michie and colleagues (34,35). The fidelity checklist assessed domains related to preparing the educational session (e.g., audit reporting, scheduling a date and time for the leadership team to

meet), presenting the educational session (e.g., reviewing the FRS and guidelines, developing, and 167 reviewing the audit and feedback report), and leading discussions on the action and care plan 168 169 processes. The fidelity checklist included 23 items, and each item was rated from 0 (action was not completed), 0.25 (action was 25% completed), 0.50 (50% completed), 0.75, and 1.00. A total 170 score of 85% (20/23) was considered good fidelity, moderate between 50% to 84%, and poor <50%171 (34,35). The fidelity checklist was completed by the lead investigator (IBR) during the education 172 session. To understand adaptations after implementing the intervention, we held focus groups with 173 the local champion and the leadership team. Focus groups were recorded and transcribed verbatim. 174 We also held informal one-on-one interviews using open-ended questions with each local 175 champion to identify adaptations to the model; these interviews were held three months after the 176 educational session and led by the lead researcher. The one-on-one interview with the local 177 178 champion was not recorded, but detailed field notes were taken by the research coordinator (LK).

Tuble 2. I customety outcomes and criteria for	Success (55)
Feasibility outcomes	Criteria for success
Recruitment rate (organizational level –	Recruit three LTC homes in Ontario
length of time to recruit three LTC homes)	within 2 months
Recruitment rate (participant level - length of	• Two months to identify, meet, and train
time to identify and train local champion)	local champion
Determine challenges to deliver the	• $\geq 80\%$ on the fidelity checklist
PREVENT model and program	• Identify facilitators of and barriers to
	implementation during the focus group
Determine challenges in linking the data for	• Identify facilitators of and barriers to
the audit and feedback report	developing the audit and feedback report

179	Table 2: Feasibility outcomes and criteria for success ((33))
-----	--	------	---

180 Secondary outcomes

Our secondary outcomes were to determine if there was a change in knowledge uptake and 181 in the proportion of osteoporosis medications before and after the intervention. We assessed 182 183 change in knowledge uptake using a pre-post test. First, we developed a case study of a typical resident living in LTC. Next, healthcare providers were asked to complete a series of multiple-184 choice questions about identifying the resident's FRS, prescribing medications, supplements, and 185 exercise, conducting an environmental hazard scan, and using hip protectors. Immediately after 186 the education session, we asked healthcare providers to repeat the multiple-choice questionnaire 187 188 on the same case study. We collected data on information on osteoporosis medications at baseline (six months prior to implementing the intervention) and post-intervention (three to four months 189 after the intervention). We collected information on the number of residents at high risk of fracture 190

or low risk of fracture using the FRS, the number of residents on an osteoporosis medication (e.g., bisphosphonates, denosumab), and the number of osteoporotic fractures at baseline and postintervention. We defined fractures as any major osteoporotic fracture of the hip, pelvis, vertebrae, or distal radius (36,37). The local champion pulled data on FRS scores, fractures, and medications directly from all residents' charts in the electronic medical record or the Ontario Drug Benefit database. A list of osteoporosis medications is provided in S4 table.

197 Analysis

We reported our recruitment rates and fidelity scores using descriptive statistics as a total 198 score or value. We reported the results of the focus groups using content analysis (descriptive 199 qualitative analysis); two reviewers (IBR and LK or LH) coded the data independently using 200 NVivo, version 14 (QSR International Pty Ltd, Doncaster, VIC, Australia) (38). The reviewers 201 202 then met to discuss and compare the codes and group the codes into categories (38). Disagreements were resolved by the third reviewer (38). We collected demographic characteristics 203 of the local champion and leadership team, which were reported using the number of individuals 204 205 and a percentage. To assess change in knowledge uptake, we conducted a paired sample t-test (two-sided p). To report changes in osteoporosis medications, we conducted two data pulls at 206 baseline and post-intervention; we included residents who "moved-in" (i.e., moved into the home 207 during the post-intervention phase) and "moved-out" (i.e., passed away during the post-208 209 intervention phase) in the analysis. We reported the results using absolute change. Quantitative results were analyzed using IBM SPSS Statistics for Windows, version 28 (Armonk, NY: IBM 210 211 Corp).

212 **Results**

213 Feasibility

We recruited three LTC homes between October 2022 and February 2023; our recruitment process was affected by residual COVID-19 complications. After we identified a home, the home's administrative team took approximately a month to identify a potential local champion. Two of the local champions were nurse practitioners, and the third champion was a kinesiologist. Subsequently, our research team met with the local champion to complete the train-the-trainer sessions. Our research team completed two, one-hour train-the-trainer sessions with Home A;

220 however, based on feedback from Home A, we added a third train-the-trainer session. The local champion led the third session where they practiced delivering the educational module to the 221 222 research team. Home B and C received three sessions and each session was 1.5 hours in length. After the train-the-trainer sessions, the local champion required at least two months to review the 223 educational materials, adapt the material to their home, and develop the audit and feedback report. 224 All leadership team meetings were held in-person at a convenient time at their respective LTC 225 home. The characteristics of the local champion and leadership team are presented in Table 3. 226 PREVENT required at least 10 months per home to recruit, train, and implement. 227

	Home A	Home B	Home C
	(k=10)	(k=19)	(k=11)
Local champion			
Highest level of education	College	University/post	University/post-
-	_	-graduate	graduate
Fidelity score (total score 23 points)	19.75/23	20.25/23 (88%)	20.00/23 (87%)
	(86%)		
Leadership team			·
Gender (female), n (%)	8 (80%)	8 (42%)	10 (91%)
Level of Education, n (%)			
College	5 (50%)	4 (21%)	1 (9%)
University	5 (50%)	8 (42%)	9 (82%)
Prefer not to answer	0 (0%)	7 (37%)	1 (9%)
Role in LTC home, n(%)			
Physician	0 (0%)	4 (21%)	1 (9%)
Pharmacist	0 (0%)	1 (5%)	1 (9%)
Administrator	1 (10%)	1 (5%)	0 (0%)
Director of care/Nursing director	1 (10%)	2 (11%)	1 (9%)
Neighbourhood coordinator	2 (20%)	4 (21%)	2 (18%)
Dietitian	0 (0%)	1 (5%)	1 (9%)
Nurse practitioner	0 (0%)	1 (5%)	2 (18%)
Nursing student	1 (10%)	0 (0%)	0 (0%)
Registered nurse	1 (10%)	0 (0%)	0 (0%)
Rehabilitation	2 (20%)	2 (11%)	2 (18%)
specialist/Physiotherapist			
Personal support worker	2 (20%)	3 (16%)	1 (9%)
Length of time worked within current			
profession in LTC, n (%)			
< 1 year	2 (20%)	1 (5%)	1 (9%)
1 to 2 years	0 (0%)	2 (11%)	1 (9%)
3 to 5 years	2 (20%)	1 (5%)	1 (9%)
> 5 years	5 (50%)	8 (42%)	7 (64%)
Prefer not to answer	1 (10%)	7 (37%)	1 (9%)

	228	Table 3: Local	champion and	l leadership team	characteristics ((n = 3)
--	-----	----------------	--------------	-------------------	-------------------	---------

Length of time providing care to residents at			
risk of falls and fractures, n (%)			
< 1 year	2 (20%)	1 (5%)	1 (9%)
1 to 3 years	1 (10%)	0 (0%)	1 (9%)
3 to 5 years	6 (60%)	10(53%)	8 (73%)
> 5 years	1 (10%)	1 (5%)	0 (0%)
Prefer not to answer	0 (0%)	7 (37%)	1 (9%)

We had a diverse group of healthcare professionals on the leadership team (Table 3). In Home A, the local champion recruited ten healthcare professionals to the leadership team, in Home B, 19 healthcare professionals, and in Home C, 12 healthcare professionals. The attendance rate during the leadership team for Home A was 90% (9/10), Home B, 84% (16/19), and Home C, 90% (10/11). Fidelity scores ranged from moderate to good (Table 3).

234 Challenges, successes, and adaptations

During the focus group, each home identified similar barriers to and facilitators of implementing PREVENT in practice.

237 Challenges

Challenges to implementing PREVENT included having an unclear sustainability strategy, 238 misperceptions between fall and fracture prevention strategies, fear of prescribing osteoporosis 239 240 medications, and time barriers with completing the audit report. During the focus group, the local champion expressed that it was unclear who would sustain the role of implementing PREVENT 241 after the research team had left. The local champion cited lack of time and other commitments as 242 a barrier to sustaining the model; all three champions noted that a major barrier was dedicating 243 244 time to the audit and feedback report. To manually extract the data and link the variables to each resident for the audit report, the local champion required at least one full day per week for five 245 weeks (i.e., 40 hours total). Moreover, several leadership members were unsure of the 246 247 implementation plan after identifying a resident at high risk of fracture:

248 "I found myself often sometimes ignoring the prompts for FRS <Fracture Risk Scale> unless
249 there was a specific reason. Cold calling families to say your fracture risk scale is high, and
250 that's the only reason I'm calling is weird." [Home B]

- 251 Several healthcare professionals on the leadership team felt the strategy to prevent fractures (i.e.,
- 252 PREVENT) was akin to their current fall prevention strategy. It was perceived that if a healthcare
- 253 professional could prevent a fall, it would translate to preventing a fracture.
- 254 "I think a lot of homes already are looking at all the PREVENT aspects, it's just not known as
 255 PREVENT or they're not grouping them together, but I think it's just something that's already
 256 being looked at from all the different disciplines." [Home A]
- 257 Other healthcare providers believed PREVENT was a tool to ensure the staff were performing
- their fall prevention duties, rather than a model of change to prevent fractures in LTC.
- 259 "It's worked well just as a tool to let us know if there's any missing boxes that we're not
 260 checking for fracture prevention. It's very helpful with just making sure we're looking at
 261 everything and yea, just crossing off any things we can possibly to do help the residents." [Home
 262 A]

Additionally, several healthcare providers were unsure about the ethical implications of prescribing osteoporosis medications to residents at high risk of fracture. The main barrier to prescribing medications was fear of side effects (i.e., osteonecrosis of the jaw, atypical femur fractures, hypocalcemia). Some providers were also concerned about the ethical implications of prescribing osteoporosis medications to residents who could not consent or had limited knowledge of the medication.

- "We are worried that, we weren't overly confident that would lead to prescription. So there was
 a trigger. We didn't know if they were going to have informed consent and we didn't know if our
 nursing staff knew enough about these medications to provide it" [Home B].
- 272 Facilitating factors

Despite some challenges, we also identified several facilitators to our model. Facilitators 273 274 included easy access to the FRS, clear and succinct educational material catered to each healthcare professional, and an educational module with an accredited program for physicians and nurses. All 275 276 three homes deemed it beneficial to have the educational module delivered in person. We also received positive feedback on the type of information included in our educational module. In 277 particular, healthcare providers found it helpful to review real-world case studies followed by 278 discussions and suggestions to overcome barriers to implementation (e.g., using hip protectors for 279 280 residents with incontinence).

"A visual to see the walkers and those pieces, I think even the slide with the hip protectors, like 281 how to apply them and which ones were incorrect. I think that's a nice reminder because I didn't 282 even know how to position them correctly and I didn't even have awareness of the FRS. Like I 283 kind of knew like our FRAT <Falls Risk Assessment Tool> score and our falls risk, and when I 284 go to my units I know who the falls, the higher risk of falls are but I didn't, and then I just 285 thought of them like okay, we have hip protectors for them, like what are we doing about it but I 286 didn't necessarily do a full review of like let's look at their meds and let's look at their hips, are 287 they wearing hip protectors, what are they, let's look at all these other things. I wasn't sort of 288 looking at that." [Home C]. 289

290 Adaptations

The process of developing and implementing models into practice is an iterative process. 291 292 Through this process, we recognized three adaptations to the PREVENT model including 293 identifying additional characteristics for a successful local champion, developing an automated audit report, and creating tools to facilitate conversations for prescribing osteoporosis medications. 294 It was suggested that the role of the local champion include co-championship with at least two 295 healthcare professionals to help manage the implementation process. Suggested co-championship 296 include one champion be either a registered nurse or a nurse practitioner and the second champion 297 be a registered practical nurse or resident case manager. The benefit of including a resident case 298 manager as a local champion is their established rapport with residents and caregivers and their 299 ability to link residents to other needed services (e.g., referral to a dietitian). Additionally, the local 300 champion must have a basic understanding of medications and have strong collaborations with 301 other healthcare providers including physicians, nurses, dietitians, pharmacists, and 302 physiotherapists. Other adaptations for future practice include working with the electronic medical 303 record company to automate the audit report process, develop conversational toolkits (e.g., 304 between nurses and residents) to aid dialogues about fracture prevention, and develop one-page 305 educational guides for healthcare professionals and residents about the side effects of osteoporosis 306 medications. 307

308 Audit report

We found that the educational module did not improve knowledge uptake of the LTC recommendations among healthcare providers particularly in utilizing the FRS, administrating osteoporosis medications and supplements (calcium and Vitamin D), using hip protectors, and reviewing environmental hazards (mean 0.07, 95% confidence intervals [CI] -0.02 to 0.16). However, our intervention resulted in an absolute increase in the proportion of osteoporosis

- medications by 6% (Table 4). Home A reported an absolute change of 13%; 21% of residents were
- on an osteoporosis medication at baseline while post-intervention 34% of residents were on a
- medication (odds ratio [OR] 0.25, 95% CI 0.12 to 0.48). Similarly, Home B reported an absolute
- change of 3% (31% at baseline and 34% post-intervention, while Home C had an 8% increase
- 318 (29% at baseline and 37% post-intervention).

319 Table 4: Aggregated pre-and-post audit report

	Homes A, B and C	
	Pre-	Post-
	intervention	intervention
Total number of residents in Home A, B, and C	739	734
Number and percentage of residents at high risk of	419 (57%)	430 (59%)
fracture		
Number and percentage of residents at low risk of	320 (43%)	304 (41%)
fracture		
Number and percentage of residents who sustained a	20 (3%)	25 (3%)
fragility fracture in the last six months		
Number and percentage of residents at high risk of	122 (29%)	150 (20%)
fracture on osteoporosis medications (unspecified		
amount of time)		

320 Note: The number of residents pre-and-post intervention are different because of the "moved-in"

321 (i.e., moved into the LTC home) and "moved-out" (i.e., passed away).

322 **Discussion**

Falls and hip fractures are a major health concern among older adults living in LTC (3.6-323 324 8). The purpose of the iCARE study was to determine the feasibility (recruitment rate, facilitators, barriers, adaptations) of implementing PREVENT in LTC homes in Canada. The model includes 325 a multifactorial intervention on diet and supplements, exercise, environmental adaptations, hip 326 protectors, and medication reviews to treat residents at high risk of fracture (9). The standardized 327 328 model utilizes local champions, educational outreach methods, audit and feedback reports, and implementation intentions to change behaviour. Although we did not meet our recruitment 329 criterion to recruit three homes in two months, our process was affected by COVID-19. 330 Nevertheless, it is possible to recruit LTC homes by accounting for a longer recruitment period 331 (i.e., three homes over five months). We did meet our criterion to recruit a local champion and 332 333 provide training within our timeframe. Our local champions successfully delivered PREVENT with a fidelity score ranging from moderate to good. We also identified several facilitators, 334

barriers, and adaptations to PREVENT. Benefits of the model include easy access to the FRS, clear 335 and succinct educational material catered to each healthcare professionals, and an educational 336 337 module with an accredited program for physicians and nurses. We also identified a few challenges such as misperceptions between the differences in fall and fracture prevention strategies, fear of 338 prescribing osteoporosis medications, and time barriers with completing the audit report. Our study 339 did not find a statistically significant increase in knowledge uptake regarding recommendations 340 for preventing falls and fractures; however, there was an increase in the proportion of osteoporosis 341 medication prescriptions post-intervention. The development and implementation of knowledge 342 translation models is an iterative process. This study identified additional changes to PREVENT 343 including co-champions, automated audit report, and tools to facilitate conversations when 344 prescribing osteoporosis medications. Our next step will be to lead a large, pragmatic randomized 345 346 controlled trial to implement PREVENT in 122 homes across Ontario, Canada.

Effective and sustainable knowledge translation models are essential to encourage the 347 348 uptake of evidence-based practices (11). The greatest limitation is that there is limited evidence on effective and sustainable models in practice and such models may not be generalizable from one 349 350 setting to another (16). Compared with community or acute care settings, there has also been limited implementation models for LTC, especially focused on fracture prevention. PREVENT is 351 352 a scalable educational outreach model that utilizes a multifactorial intervention to reduce falls and fractures among residents in LTC (39). Our study provides preliminary data that suggests 353 354 PREVENT may be successfully implemented within the LTC setting, particularly when structured to fit the unique practice environment and organizational structure. Most knowledge translation 355 356 models have been evaluated in non-LTC settings (39). A 2009 Cochrane review reported improvements in care that ranged from 4% to 12% using educational meetings, educational 357 outreach programs, local champions, audit and feedback reports, and computerized reminders (39). 358 When designing PREVENT, we utilized such techniques (e.g., educational meetings, local 359 champion, audit and feedback) as a method to change behaviour (25,27,30,31,39). We observed 360 improvements similar to the Cochrane review ranging from 3% to 13% in the proportion of 361 osteoporosis medications before and after the intervention. Four months was not enough time to 362 implement change, especially in homes with over 200 beds. We saw greater improvements in 363 prescriptions being filled in homes that had a lower rate of medication use. We specifically focused 364 365 on osteoporosis medications as the highest certainty of evidence to reduce the rate and risk of

fractures is using such medications (19,20) and is the easiest outcome to measure. We hypothesize
that an increased uptake of osteoporosis medications will translate to lower fracture rates in LTC;
however, a large trial is needed to confirm the hypothesis.

Similar models to prevent fractures in LTC include the Bavarian Fall and Fracture 369 Prevention model in Germany that focuses on exercise, documentation of falls, environmental 370 371 adaptations, medication reviews, vitamin D, hip protectors, and education among healthcare providers (40,41). The Bavarian Fall and Fracture Prevention study is one of the few large studies 372 of an educational outreach program to implement fracture prevention guidelines into LTC (40.41); 373 the authors implemented a multifactorial educational outreach program and after one-year found 374 an 18% reduction in hip fracture in residents from the intervention home (hazard rate ratio 0.82, 375 95% CI 0.72-0.93) (40,41). While the model was effective, it was a non-randomized, quasi-376 experimental design that utilized insurance company staff to facilitate and disseminate knowledge 377 on fall and fracture prevention. The sustainability or scalability of this model may not be applicable 378 in the Canada healthcare system. Thus, the next steps will be to conduct a large randomized 379 controlled trial of the adapted PREVENT model in several Canadian LTC homes to determine if 380 381 the model reduces the rate of hip fractures in residents at high risk of fracture.

Our PREVENT model attempted to leverage the facilitators and limit barriers to 382 383 implementation as described in other studies. A recent systematic review by our group identified several barriers to implementing guidelines into LTC settings including time constraints and 384 385 inadequate staffing, cost and lack of resources, knowledge gaps, and lack of teamwork and organizational support (14). Our model addressed such barriers including time constraints, lack of 386 387 resources, and knowledge gaps by developing a multi-modal approach to embed a validated computer decision support tool and the FRS into already established care pathways. PREVENT 388 389 also provides an accredited Continuing Medical Education module to physicians and nurses as a 390 method to disseminate evidence-based knowledge. We worked with the home's management team to identify potential local champions who could help lead the educational module and lead best 391 practices after the research team left; from the literature we identified characteristics for an 392 393 effective local champion (25,42). Through the iCARE study, we identified additional 394 characteristics to identify a local champion including having co-championship to manage the implementation process and developing conversational toolkits to aid dialogues between 395 healthcare providers and residents on osteoporosis medications. We are also working with a 396

Canadian electronic medical record company to develop an automated audit and feedback report to address the time barrier. We suggest that future knowledge translation and implementation science researchers look to our adapted PREVENT model as there may be important changes to help facilitate the effective translation of evidence into practice.

The iCARE study had several methodological strengths including the recruitment of homes 401 that were geographically diverse and located in communities of varied population sizes; we also 402 included not-for-profit and for-profit homes. Our study implemented a multifactorial intervention 403 that engaged a wide variety of healthcare providers. In addition, our study builds on previous 404 knowledge of implementing models into practice and was designed to leverage facilitators and 405 address common barriers to implementation. To date, there are very few models on chronic disease 406 management especially in LTC (43,44), and so, more studies are needed to identified challenges 407 to implementing knowledge translation models into practice. Nonetheless, our study is not without 408 limitations. Given that our intervention was multifaceted, it was challenging to identify the most 409 significant components (e.g., educational outreach, local champion, audit and feedback report) that 410 would result in effective implementation of PREVENT. Moreover, measures of organizational 411 412 context such as work culture and leadership style, were not captured in the iCARE study which could have a significant impact on our model's components. Healthcare sustainability remains a 413 414 multi-dimension problem, and so it is unclear if local champions or co-championship is sustainable (45). Lastly, for this study, we did not involve residents or their family caregivers during the 415 416 implementation process. We acknowledge that all residents have complex needs and multiple healthcare issues, and their preferences for care should be incorporated in the decision-making 417 process. Future adaptations for the PREVENT model will be to include end-users on the research 418 team to aid the implementation process to consider how resident's preferences can be considered. 419 420 Before we implement the PREVENT model as a randomized controlled trial, we will consult and 421 collaborate with knowledge users on our implementation process.

422 **Conclusion**

PREVENT is a multifactorial educational outreach model to implement education on diet,
 exercise, environmental adaptations, hip protectors, medications (including calcium and vitamin
 D) and medication reviews to treat residents at high risk of fracture. The model utilizes educational
 meetings, local champions, and audit and feedback as methods to promote behaviour change. The

model requires ten months to recruit, train, and implement PREVENT, with five of the ten months 427 dedicated toward recruitment of LTC homes. We suggest co-championship when delivering 428 429 multifactorial models in LTC and developing tools that can be easily embedded into routine practice. Our study did not increase knowledge uptake of the guidelines; however, we did observe 430 an increase in the proportion of osteoporosis medication prescriptions post-intervention. Our next 431 steps will be to consult and collaborate with knowledge users on PREVENT and then lead a large, 432 randomized controlled trial to implement PREVENT to determine if the model reduces the risk 433 and rate of hip fractures in residents living in LTC. 434

435 **List of abbreviations**

436	CI	Confidence Intervals

- 437FRSFracture Risk Scale
- 438 iCARE Integrated collection of education modules for <u>fall</u> and <u>fracture</u> prevention
- 439 LTC Long term care
- 440 PREVENT <u>Person-centred Routine Fracture PreEVENT</u>ion
- 441 OR Odds Ratio

442 **Supporting information**

- 443 S1 File. PREVENT Components
- 444 S2 File. Training Manual for Local Champions
- 445 S3 File. Fracture & Fall Prevention Guide for Long Term Care
- 446 S4 Table. Osteoporosis Drugs
- 447 S5 Table. STROBE Checklist
- 448 S6 Table. TIDieR Checklist
- 449 S7 File. iCARE protocol

451 **References**

- Bergen G, Stevens M, Burns E. Falls and Fall Injuries Among Adults Aged ≥65 Years United States. MMWR Morb Mortal Wkly Rep. 2014;65(37):993–8.
- O'Loughlin JL, Yvonne Robitaille JFB, Samy Suissa. Incidence of and Risk Factors for
 Falls and Injurious Falls among the Community-dwelling Elderly. American journal of
 epidemiology . 1993;137(3):342–54.
- 457 3. Neuman MD, Silber JH, Magaziner JS, Passarella MA, Mehta S, Werner RM. Survival
 458 and functional outcomes after hip fracture among nursing home residents. JAMA Intern
 459 Med. 2014;174(8):1273–80.
- 4. Rapp K, Becker C, Lamb SE, Icks A, Klenk J. Hip fractures in institutionalized elderly
 people: Incidence rates and excess mortality. Journal of Bone and Mineral Research. 2008
 Nov;23(11):1825–31.
- 463 5. Zimmerman S, Chandler JM, Hawkes W, Sloane PD, Hebel ; J Richard, Magaziner J, et
 464 al. Effect of Fracture on the Health Care Use of Nursing Home Residents [Internet].
 465 Available from: https://jamanetwork.com/
- 6. Chandler JM, Zimmerman SI, Girman CJ, Martin AR, Hawkes W, Hebel JR, et al. Low
 Bone Mineral Density and Risk of Fracture in White Female Nursing Home Residents
 [Internet]. Available from: http://jama.jamanetwork.com/
- 469 7. Ooms M, Vlasman P, Lips, Nauta J, Bouter L, Valkenburg H. The incidence of hip
 470 fractures in independent and institutionalized elderly people. Osteoporosis International.
 471 1994;4(1):6–10.
- Weller I, Schatzker J. Hip fractures and Alzheimer's disease in elderly institutionalized
 Canadians. Ann Epidemiol . 2004;14(5):319–24.
- Papaioannou A, Santesso N, Morin SN, Feldman S, Adachi JD, Crilly R, et al.
 Recommendations for preventing fracture in long-term care. CMAJ [Internet].
 2015;187(15). Available from: www.cmaj.ca/lookup/
- 477 10. Graham ID, Tetroe J, Gagnon M. Knowledge Translation in Healthcare. Second Edi. Sons
 478 JW&, editor. 2013. 75–92 p.
- Lavis JN, Robertson D, Woodside JM, McLeod CB, Abelson J, Group KTS. How Can
 Research Organizations More Effectively Transfer Research Knowledge to Decision
 Makers? Milbank Q. 2003;81(2):221–48.
- Roigk P, Becker C, Schulz C, König HH, Rapp K. Long-term evaluation of the
 implementation of a large fall and fracture prevention program in long-term care facilities.
 BMC Geriatr. 2018 Oct 1;18(1).

Bai Y, McArthur C, Ioannidis G, Giangregorio L, Straus S, Papaioannou A. Strategies for
the implementation of an electronic fracture risk assessment tool in long term care: a
qualitative study. BMC Geriatr. 2021 Dec 1;21(1).

- McArthur C, Bai Y, Hewston P, Giangregorio L, Straus S, Papaioannou A. Barriers and
 facilitators to implementing evidence-based guidelines in long-term care: a qualitative
 evidence synthesis. Vol. 16, Implementation Science. BioMed Central Ltd; 2021.
- Tomasone JR, Kauffeldt KD, Chaudhary R, Brouwers MC. Effectiveness of guideline
 dissemination and implementation strategies on health care professionals' behaviour and
 patient outcomes in the cancer care context: A systematic review. Vol. 15, Implementation
 Science. BioMed Central Ltd; 2020.
- Alamri SH, Kennedy CC, Marr S, Lohfeld L, Skidmore CJ, Papaioannou A. Strategies to overcome barriers to implementing osteoporosis and fracture prevention guidelines in long-term care: A qualitative analysis of action plans suggested by front line staff in Ontario, Canada. BMC Geriatr. 2015 Aug 1;15(1).
- 499 17. Cargo M, Mercer SL. The value and challenges of participatory research: Strengthening
 500 its practice. In: Annual Review of Public Health. 2008. p. 325–50.
- Niznik JD, Gilliam MA, Colón-Emeric C, Thorpe CT, Lund JL, Berry SD, et al.
 Controversies in Osteoporosis Treatment of Nursing Home Residents. J Am Med Dir
 Assoc. 2022 Dec 1;23(12):1928–34.
- S04 19. Cummings SR, Lui LY, Eastell R, Allen IE. Association between Drug Treatments for
 Patients with Osteoporosis and Overall Mortality Rates: A Meta-analysis. JAMA Intern
 Med. 2019 Nov 1;179(11):1491–500.
- 507 20. Händel MN, Cardoso I, Von Bülow C, Rohde JF, Ussing A, Nielsen SM, et al. Fracture
 508 risk reduction and safety by osteoporosis treatment compared with placebo or active
 509 comparator in postmenopausal women: systematic review, network meta-analysis, and
 510 meta-regression analysis of randomised clinical trials. BMJ. 2023;
- 511 21. Vandenbroucke JP, von Elm E, Altman DG, Gotzsche PC, Mulrow CD, Pocock SJ, et al.
 512 Strengthening the Reporting of Observational Studies in Epidemiology (STROBE):
 513 Explanation and Elaboration. PLOS Med. 20017;16(4).
- 514 22. Hoffmann TC, Glasziou PP, Barbour V, Macdonald H. Better reporting of interventions : 515 template for intervention description and replication (TIDieR) checklist and guide. 516 2014;1687(March):1–12. Available from: http://dx.doi.org/doi:10.1136/bmj.g1687
- 517 23. Michie S, Atkins L, West R. The behaviour change wheel: a guide to designing
 518 interventions. 2nd edn. Silverback Publishing, Sutton; 2014. 59–60 p.

519 24. Michie S, van Stralen M, West R. The behaviour change wheel: A new method for 520 characterising and designing behaviour change interventions. Implementation Science. 521 2011;6(42).

522 523 524	25.	Flodgren G, O'brien MA, Parmelli E, Grimshaw JM. Local opinion leaders: Effects on professional practice and healthcare outcomes. Vol. 2019, Cochrane Database of Systematic Reviews. John Wiley and Sons Ltd; 2019.
525 526 527	26.	Yarber L, Brownson CA, Jacob RR, Baker EA, Jones E, Baumann C, et al. Evaluating a train-the-trainer approach for improving capacity for evidence-based decision making in public health. BMC Health Serv Res. 2015 Dec 12;15(1).
528 529 530	27.	O'Brien M, Oxman A, Davis D, Haynes R, Freemantle N, Harvey E. Educational outreach visits: effects on professional practice and health care outcomes. In: Cochrane Database of Systematic Reviews. John Wiley & Sons, Ltd; 1997.
531 532 533 534	28.	Ioannidis G, Jantzi M, Bucek J, Adachi JD, Giangregorio L, Hirdes J, et al. Development and validation of the Fracture Risk Scale (FRS) that predicts fracture over a 1-year time period in institutionalised frail older people living in Canada: An electronic record-linked longitudinal cohort study. BMJ Open. 2017 Sep 1;7(9).
535 536 537	29.	Papaioannou A, Ioannidis G, McArthur C, Hillier LM, Feldman S, Giangregorio L, et al. Preventing Fractures in Long-Term Care: Translating Recommendations to Clinical Practice. J Am Med Dir Assoc. 2021 Jan 1;22(1):36–42.
538 539 540	30.	Ivers N, Jamtvedt G, Flottorp S, Young JM, Odgaard-Jensen J, French SD, et al. Audit and feedback: Effects on professional practice and healthcare outcomes. Cochrane Database of Systematic Reviews. 2012;2012(6).
541 542 543	31.	Shojania KG, Jennings A, Mayhew A, Ramsay CR, Eccles MP, Grimshaw J. The effects of on-screen, point of care computer reminders on processes and outcomes of care. Cochrane Database of Systematic Reviews. John Wiley and Sons Ltd; 2009.
544 545	32.	Wandersman A, Pamela I, Matthew C, Shakeh K. Getting to outcomes: a results-based approach to accountability. Eval Program Plann. 2000;23(3):389395.
546 547 548	33.	Eldridge SM, Lancaster GA, Campbell MJ, Thabane L, Hopewell S, Coleman CL, et al. Defining feasibility and pilot studies in preparation for randomised controlled trials: development of a conceptual framework. PLoS One. 2016;11(3).
549 550 551	34.	Walton H, Spector A, Williamson M, Tombor I, Michie S. Developing quality fidelity and engagement measures for complex health interventions. Br J Health Psychol. 2020 Feb 1;25(1):39–60.
552 553	35.	Breitenstein SM, Gross D, Garvey CA, Hill C, Fogg L, Resnick B. Implementation fidelity in community-based interventions. Res Nurs Health. 2010 Apr;33(2):164–73.
554	36.	Kanis JA. Diagnosis of osteoporosis and assessment of fracture risk. 2002;359:1929–36.
555 556	37.	Kanis JA, Melton LJ, Christiansen C, Johnston CC, Khaltaev N. The Diagnosis of Osteoporosis. 1994;9(8):1137–41.

557 558	38.	White MD, Marsh EE, White MD, Marsh EE. Content Analysis : A Flexible Methodology Content Analysis : A Flexible Methodology. 2017;55(1):22–45.
559 560	39.	Grimshaw J, Eccles M, Lavis J, Hill S, Squires J. Knowledge translation of research findings. Implementation Science. 2012;7(50).
561 562 563	40.	Becker C, Cameron ID, Klenk J, Lindemann U, Heinrich S, König HH, et al. Reduction of femoral fractures in long-term care facilities: The Bavarian fracture prevention study. PLoS One. 2011;6(8).
564 565 566 567	41.	Schulz C, Lindlbauer I, Rapp K, Becker C, König HH. Long-Term Effectiveness of a Multifactorial Fall and Fracture Prevention Program in Bavarian Nursing Homes: An Analysis Based on Health Insurance Claims Data. J Am Med Dir Assoc. 2017 Jun 1;18(6):552.e7-552.e17.
568 569 570	42.	Bonawitz K, Wetmore M, Heisler M, Dalton VK, Damschroder LJ, Forman J, et al. Champions in context: Which attributes matter for change efforts in healthcare? Implementation Science. 2020 Aug 6;15(1).
571 572 573 574	43.	Boscart V, Davey M, Crutchlow L, Heyer M, Johnson K, Taucar LS, et al. Effective Chronic Disease Interventions in Nursing Homes: A Scoping Review Based on the Knowledge-to-Action Framework. Vol. 45, Clinical Gerontologist. Routledge; 2022. p. 1073–86.
575 576 577	44.	Boscart V, Crutchlow LE, Sheiban Taucar L, Johnson K, Heyer M, Davey M, et al. Chronic disease management models in nursing homes: A scoping review. Vol. 10, BMJ Open. BMJ Publishing Group; 2020.
578 579	45.	Fleiszer AR, Semenic SE, Ritchie JA, Richer MC, Denis JL. The sustainability of healthcare innovations: A concept analysis. J Adv Nurs. 2015 Jul 1;71(7):1484–98.
580		

581 Ethics approval and consent to participate

In accordance with local regulatory guidelines and standards for human participants' protection, all methods were carried out in accordance with relevant guidelines and regulations. Our study was reviewed by the Hamilton Integrated Research Ethics Board (HiREB), which is associated with McMaster University. Before enrolling in the study, all participants were briefed on the study and written informed consent was obtained. This review was registered on clinicaltrials.gov (NCT05445336).

588 **Consent for publication**

589 Not applicable.

590 Availability of data and materials

- 591 The data generated during the current study are not publicly available due to content that could
- 592 compromise research participant privacy but are available from the corresponding author on
- reasonable request for research purposes only.

594 **Competing interests**

595 The authors have no competing interests.

596 Funding

597 The authors would like to thank the following funding agents for their support with the project.

598 The lead author (IBR) was funded by the McMaster Institute for Research on Aging (MIRA),

599 AGE-WELL, the Hamilton Health Sciences New Investigator Fund, the Canadian Institutes of

600 Health Research Postdoctoral Award. The corresponding author (AP) received funding from the

- 601 Amgen Competitive Grant Program in Bone Research Award. The funders played no part in
- 602 developing the research design, collecting data collection, analyzing the results, or writing the 603 manuscript.

604



Figure 1: STROBE flow chart. STROBE, Strengthening the Reporting of Observational Studies in Epidemiology.

STROBE Flow Diagram