# (Brief Report) Title: Mental Bandwidth is Associated with HIV and Viral Suppression Among Low-Income Women in Philadelphia

Aaron Richterman,\*<sup>1,2,3</sup> Nancy Aitcheson,<sup>1,2</sup> Celeste Durnwald,<sup>4</sup> Cara Curley,<sup>4</sup> William R. Short,<sup>1,4</sup> Mirabelle Jean Louis,<sup>5</sup> Florence Momplaisir,<sup>1,2</sup> Harsha Thirumurthy<sup>2,3</sup>

**Keywords:** Behavioral Economics, Pregnancy, HIV, Poverty, Mental Bandwidth

Abbreviations: ART, antiretroviral therapy; HIV, human immunodeficiency virus; IQR, interguartile range; KDDT, Kirby Delay Discounting Task; PWH, people with HIV; PVT, psychomotor vigilance task; RPM, Raven's progressive matrices; SD, standard deviation

## Word count: 2047 Number of references: 37 Number of tables: 1 Number of figures: 0

\*Corresponding Author: Aaron Richterman, MD, MPH; Hospital of the University of Pennsylvania, 3400 Spruce Philadelphia, PA 19104: Street, e-mail: aaron.richterman@pennmedicine.upenn.edu telephone: 2674417915; ORCID: 0000-0001-7920-7191

Affiliations:

<sup>1</sup> Division of Infectious Diseases, Department of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

<sup>2</sup> Leonard Davis Institute of Health Economics, University of Pennsylvania, Philadelphia, Pennsvlvania. USA

Department of Medical Ethics and Health Policy, University of Pennsylvania, Philadelphia, Pennsylvania, USA

<sup>4</sup> Department of Obstetrics and Gynecology, University of Pennsylvania, Philadelphia, Pennsylvania, USA

Bentley University, Boston, Massachusetts, USA

#### 1 Abstract

2 Behavioral economics research suggests poverty may influence behavior by reducing 3 mental bandwidth, increasing future discounting, and increasing risk aversion. It is plausible that these decision-making processes are further impaired in the context of 4 5 HIV or pregnancy. In this cross-sectional study of 86 low-income women in 6 Philadelphia, multivariable models showed that HIV was associated with decreased 7 mental bandwidth (one of two measures) and lower risk aversion. Pregnancy was not 8 associated with any decision-making factors. Viral suppression was associated with 9 greater mental bandwidth (one of two measures), and antenatal care engagement with lower future discounting. Anti-poverty interventions may be particularly beneficial to 10 11 health behaviors of HIV. improve in the context

## 12 Introduction

Health behaviors such as lack of adherence to antiretroviral therapy (ART) and inconsistent engagement in care remain some of the greatest barriers to ending HIV epidemics. This is particularly true for pregnant women with HIV, for whom postpartum clinic attendance, ART adherence, and postpartum viral suppression rates remain unacceptably low in high- and low-income settings.<sup>1-10</sup>

40% of people with HIV (PWH) in the US live below the poverty line,<sup>11</sup> and poverty is an 18 19 important upstream contributor to poor ART adherence and engagement in care by PWH, particularly during pregnancy and in the years after giving birth.<sup>5,9</sup> Recent 20 21 behavioral economics research suggests that poverty may affect individual behavior through psychological (in addition to economic) mechanisms.<sup>12-21</sup> This may occur 22 23 through decreases in mental bandwidth (the mental resources available at a given time 24 to make complex decisions), increases in future discounting (the tendency to undervalue future outcomes), and increases in risk aversion.<sup>15,16,19-24</sup> 25

While the association between poverty and these psychological decision-making processes has been recently established, it is plausible that these processes are even further impaired in the context of HIV or pregnancy. If this were the case, it would suggest that anti-poverty interventions could be particularly beneficial to improve health behaviors in these populations. To evaluate this possibility, we measured mental bandwidth, future discounting, and risk aversion among low-income women in Philadelphia, and examined their associations with HIV and pregnancy status.

### 33 Methods

34 We conducted a cross-sectional study of low-income pregnant and non-pregnant 35 women with and without HIV receiving care at outpatient clinics in the University of Pennsylvania Health System. Eligibility criteria included (1) cisgender female, (2) age 36 37 18-40 years, (3) Philadelphia resident, (4) household income <200% federal poverty 38 level, (5) confirmed pregnancy by standard laboratory and ultrasound testing (pregnant 39 participants), (6) documented HIV seropositive (participants with HIV), (7) able to 40 provide informed consent. The study was approved by the University of Pennsylvania 41 IRB. All participants provided written informed consent.

Participants completed a one-time survey assessing sociodemographics, income, food security,<sup>25</sup> health insurance, obstetrical history, self-reported ART adherence (among PWH), and whether a pregnancy was intended (pregnant participants). We conducted chart reviews to assess for most recent HIV viral load (last 3 months) and current gestational age. For pregnant participants, we conducted a follow-up chart review after delivery to assess antenatal care engagement (proportion of scheduled obstetrics clinic appointments that were attended) and gestational age at delivery.

We assessed mental bandwidth using two measures.<sup>14</sup> The Psychomotor Vigilance 49 50 Task (PVT) assesses attentional vigilance by asking participants to press a button when 51 a stimulus appears on a screen over a period of ten minutes and measuring reaction time and accuracy.<sup>26,27</sup> After standardizing reaction time, number of false starts 52 53 (responding prior to the stimulus appearing), and number of minor lapses (correctly 54 responding but with response time >500 milliseconds) such that each variable had a 55 population mean of 0 and standard deviation (SD) of 1, we generated PVT scores to 56 synthesize reaction time and accuracy as follows: reaction time + 0.5\*(minor lapses +

false starts).<sup>28</sup> Raven's Progressive Matrices (RPM) is a non-verbal estimate of fluid intelligence, or the ability to solve novel problems or adapt to new situations.<sup>29</sup> RPM presents participants with a series of ten figures with a missing portion of a pattern and asks them to select which of six options best completes the missing pattern, with the number of correct answers representing a participant's score.

62 We assessed future discounting using the Kirby Delay-Discounting Task (KDDT), a 27-

item questionnaire that asks participants to hypothetically choose between smaller,

64 immediate monetary rewards and larger, delayed awards.<sup>30,31</sup> The KDDT generates

each participant's discounting constant (k), with higher values indicating greater

amounts of future discounting (i.e., greater impulsivity).<sup>32</sup>

67 We assessed risk-aversion using an ordered lottery selection method, asking

68 participants hypothetical questions about whether they would prefer a 100% chance of

receiving \$5, or a 50% chance of receiving \$10 and a 50% chance of receiving \$0, \$1,

\$2, \$3, or \$4, to assess the threshold (if any) at which they would choose the higher-risk
option.<sup>33,34</sup>

We used multivariable linear regression models to separately estimate associations between HIV and pregnancy (primary explanatory variables) with mental bandwidth (PVT, RPM), future discounting, and risk aversion (primary outcomes). Each of the outcomes was standardized to have a mean of 0 and standard deviation (SD) of 1, such that effect sizes were expressed in SDs of the outcome. Models controlled for age, race, ethnicity, educational attainment, income, food security, pregnancy (HIV models), and HIV (pregnancy models). These covariates were selected a priori as either known or hypothesized confounders of the relationship between the decision-making processesand HIV or pregnancy.

In secondary analyses, we used similar multivariable models to estimate associations between the same decision-making factors and viral suppression (among PWH) and antenatal care engagement (among pregnant participants). Because of smaller sample sizes, these models adjusted only for age and either HIV or pregnancy.

We had a planned sample size of at least 70 participants (approximately half with HIV, and half pregnant) to provide at least 80% power to detect similar differences in mental bandwidth as seen in prior research demonstrating the effects of poverty alleviation on mental bandwidth (0.67 SDs).<sup>15</sup> The study was approved by the University of Pennsylvania Institutional Review Board. We performed statistical analysis using SAS Version 9.4 (SAS Institute; Cary, NC).

#### 91 Results

92 We enrolled 86 participants with a median age of 28 years (IQR 25 to 32) between 93 March 2022 and August 2023. There were 69 (80%) participants who identified as non-94 Hispanic Black, 6 (7%) as non-Hispanic White, 2 (3%) as non-Hispanic Asian, 2 (3%) as 95 non-Hispanic and multiple races, and 7 (8%) as Hispanic. Sixty-seven participants 96 (78%) had a single marital status, and 76 (88%) had completed high school. 97 Participants had a median of 1 child (IQR 1 to 2). For annual household income, 25 98 participants (29%) reported ≤\$10,000, 20 (23%) reported \$10,001-\$20,000, 28 (33%) reported \$20,001-\$40,000, 10 (12%) reported \$40,001-\$60,000, and 3 (3%) reported 99

\$60,001-\$80,000. Fifty-four participants (63%) were employed, 33 (38%) had low or
very low food security, and 76 (88%) had Medicaid.

102 Thirty-five participants (41%) were living with HIV, and 36 (42%) were pregnant -26103 (30%) were non-pregnant without HIV, 24 (28%) were non-pregnant with HIV, 25 (29%) 104 were pregnant without HIV, and 11 (13%) were pregnant with HIV. PWH reported a 105 median of one (IQR 0 to 3) missed dose of ART during the last month, with 16 (46%) 106 having <95% adherence and 4 (11%) having <80% adherence. The most recent viral 107 load was undetectable for 30 (86%) of the PWH. Among pregnant participants, 9 (25%) 108 intended the current pregnancy, and the median gestational age at the time of the 109 enrollment was 25 weeks (IQR 16 to 32). In follow-up chart review, pregnant 110 participants had a median engagement in antenatal care (proportion of scheduled 111 obstetrics visits attended) of 0.86 (IQR 0.71 to 1.00), and a median gestational age at 112 delivery of 39 weeks (IQR 37 to 40).

Participants had a median number of correct answers on the RPM of 8 (IQR 7 to 9), and on the PVT they had a median reaction time of 414 milliseconds (IQR 375 to 484), median number of false starts of 1 (IQR 0 to 2), and a median number of minor lapses of 17 (IQR 8 to 38). Participants had a median log temporal discounting constant of -3.7 (IQR -4.6 to -2.7). The median risk-aversion threshold was a 50% chance of \$4 and a 50% chance of \$10 (IQR 50% chance of \$1 to no threshold [always choosing a 100% chance of \$5]).

After adjustment, HIV was associated with lower mental bandwidth on the PVT (-0.42 SD difference, 95% CI -0.84 to -0.01), and lower risk-aversion (-1.16 SD difference,

95% CI -2.05 to -0.27) (Table 2). Pregnancy was not associated with any decisionmaking factors. Among PWH, viral suppression was associated with greater mental bandwidth on the PVT (1.12 SD difference, 95% CI 0.10 to 2.13). Among pregnant participants, antenatal care engagement was associated with lower future discounting (-1.57 SD difference, 95% CI -3.08 to -0.05).

#### 127 Discussion

128 In this cross-sectional study of low-income women receiving health care in Philadelphia, 129 we found that living with HIV was associated with significantly lower mental bandwidth 130 (on one of two measures) and risk aversion, while pregnancy was not associated with 131 any psychological decision-making factors. We also found that viral suppression was 132 associated with greater mental bandwidth (on one of two measures), and antenatal care 133 engagement was associated with lower future discounting. The differences were largely similar in magnitude to those seen in prior studies of poverty alleviation.<sup>15</sup> We 134 135 hypothesize that we found associations between HIV (and viral suppression) and 136 mental bandwidth when using the PVT but not RPM because of a ceiling effect resulting 137 from high scores with little variability for the RPM, indicating that the RPM series needs 138 to be more difficult for this population in future work. This is among the first studies to 139 assess these decision-making factors among low-income PWH and pregnant women.

Our findings suggest that PWH living in poverty may be especially constrained in their ability to make decisions to promote long-term health (e.g. medication adherence), and that these constraints are associated with important health outcomes like viral suppression and engagement in care. Consequently, interventions that increase mental

144 bandwidth and decrease future discounting, such as poverty alleviation through cash 145 transfers or housing support, may lead to improvements in health behaviors and health 146 outcomes. While we did not find any specific differences associated with pregnancy, it 147 remains possible that sub-groups of pregnant women - such as those with limited social 148 support – experience greater losses in mental bandwidth. Studies with larger sample 149 sizes of low-income pregnant women could further explore such associations. Our 150 findings nonetheless remain relevant during the perinatal period since poverty often increases around the time of pregnancy.<sup>35</sup> 151

152 This study expands our understanding of poverty's impact on psychological decision-153 making processes into the context of HIV and HIV-related behaviors. The psychology 154 literature points to a dual-process model of thinking and decision-making — System 1 is 155 intuitive, and rapidly guides one's decision-making on a regular basis; System 2 is 156 deliberative and considers longer-term consequences of each decision, but requires cognitive effort.<sup>36</sup> Mental bandwidth has been defined as the mental resources available 157 158 at a given time to engage in System 2 thinking.<sup>14</sup> Recent behavioral economics 159 research has shown that a state of economic scarcity (i.e., living in poverty) induces people to focus most of their attention towards immediate needs,<sup>19,23,24</sup> and that 160 attention to these needs diminishes bandwidth.<sup>15,20</sup> Similarly, evidence suggests that 161 162 poverty affects time preferences by increasing future discounting (defined as the tendency to undervalue future outcomes).<sup>21</sup> In contrast, alleviating scarcity through 163 164 poverty alleviation has been shown to increase bandwidth and decrease discounting.<sup>16,22</sup> Whether poverty alleviation's effects on mental bandwidth and future 165 166 discounting translates into changes in future-oriented health behaviors remains an open

question. While this growing economics literature has focused on the effects of poverty on these psychological decision-making factors, our study is among the first to evaluate associations between these factors and a specific context like HIV within which poverty is a highly important determinant of outcomes.

171 This study has several limitations. First, this study was cross-sectional, which precludes 172 establishing causality for the identified associations. For example, living with HIV may 173 lead to reductions in mental bandwidth, but it's also possible that people with lower 174 mental bandwidth (because of poverty and other factors) may be more likely to acquire 175 HIV, or that lower mental bandwidth is co-correlated with other HIV risk factors (such as 176 substance use or housing insecurity). Regardless, the fact that PWH have lower mental 177 bandwidth indicates that they may especially benefit from economic interventions that increase bandwidth and potentially improve health behaviors like ART adherence.<sup>37</sup> 178 179 Second, as our findings come from a single center in a high-income country, they 180 should be considered preliminary and replicated in other contexts.

#### 181 Conclusion

In this cross-sectional study of low-income women living in Philadelphia, living with HIV was associated with lower mental bandwidth (on one of two measures) and risk aversion, whereas pregnancy was not associated with any psychological decisionmaking factors. Our findings suggest that PWH living in poverty may be especially constrained in their ability to make daily decisions to promote long-term health. We also found that these constraints are associated with important health outcomes like viral suppression and engagement in care. Consequently, interventions that increase mental

- 189 bandwidth and decrease future discounting, such as poverty alleviation, may lead to
- 190 important improvements in outcomes among PWH.

#### 191 Author Roles

- 192 Author contributions: Richterman had full access to all data in the study and takes
- responsibility for the integrity of the data and accuracy of the data analysis.
- 194 Concept and design: Richterman, Aitcheson, Momplaisir, Thirumurthy
- 195 Acquisition, analysis, interpretation of data: All authors
- 196 Drafting of the manuscript: Richterman
- 197 Critical revision of the manuscript. All authors
- 198 Statistical analysis: Richterman
- 199 **Competing interests**
- 200 The authors declare no conflicts of interest.

## 201 Funding

202 Penn Center for AIDS Research Center Developmental Pilot Award (P30AI045008) to

203 AR, and National Institute of Mental Health of the National Institutes of Health 204 (K23MH131464) to AR. **Table 1.** Associations (with 95% CIs) between mental bandwidth, risk aversion, and future discounting with HIV, viral suppression (among PWH), pregnancy, and antenatal care engagement (among pregnant participants). Effect measures are generated using multivariable linear regression models and are and standardized such that effect sizes are expressed in standard deviations of the outcomes. HIV and pregnancy models control for age, race, ethnicity, educational attainment, income, food security, pregnancy (HIV model), and HIV (pregnancy model). The viral suppression and antenatal care engagement models control for age and HIV or pregnancy. Sample sizes are as follows: HIV and Pregnancy model N=35; antenatal care model N=36.

	HIV <sup>1</sup> (N=86)	Viral Suppression (among PWH) <sup>2</sup> (N=35)	Pregnancy <sup>3</sup> (N=86)	Antenatal care engagement (among pregnant) <sup>4</sup> (N=36)
Mental				
Bandwidth				
(Psychomotor				
Vigilance				
Task)	-0.42 (-0.84 to -0.01)	1.12 (0.10 to 2.13)	0.09 (-0.31 to 0.49)	0.89 (-0.68 to 2.46)
Mental				
Bandwidth				
(Raven's				
Progressive				
Matrices)	-0.37 (-0.78 to 0.06)	-0.48 (-1.59 to 0.63)	0.07 (-0.34 to 0.47)	-0.32 (-1.79 to 1.15)
Future			-0.03 (-0.45 to	
Discounting	-0.07 (-0.51 to 0.37)	-0.11 (-1.10 to 0.89)	0.39)	-1.57 (-3.08 to -0.05)
	-		-0.62 (-1.48 to	
<b>Risk aversion</b>	-1.16 (-2.05 to -0.27)	0.08 (-1.72 to 1.88)	0.23)	-0.16 (-3.20 to 2.88)

212

<sup>1</sup> Models control for age, race, ethnicity, educational attainment, income, food security, and pregnancy.

<sup>2</sup> Models control for age and pregnancy.

<sup>3</sup>Models control for age, race, ethnicity, educational attainment, income, food security, and HIV status.

216 <sup>4</sup> Models control for age and HIV	status.
---	---------

# References

- Puttkammer N, Domerçant JW, Adler M, et al. ART attrition and risk factors among Option B+ patients in Haiti: A retrospective cohort study. *PloS one.* 2017;12(3):e0173123.
- 2. Institut Haïtien de l'Enfance IHEH, Icf. *Haiti EnquÎte Mortalitè, Morbiditè et Utilisation des Services 2016-2017 EMMUS-VI.* Pètion-Ville/Haïti: IHE/Haiti, ICF;2018.
- Knettel BA, Cichowitz C, Ngocho JS, et al. Retention in HIV Care During Pregnancy and the Postpartum Period in the Option B+ Era: Systematic Review and Meta-Analysis of Studies in Africa. *Journal of acquired immune deficiency syndromes (1999)*. 2018;77(5):427-438.
- 4. Nachega JB, Uthman OA, Anderson J, et al. Adherence to antiretroviral therapy during and after pregnancy in low-income, middle-income, and high-income countries: a systematic review and meta-analysis. *AIDS (London, England).* 2012;26(16):2039-2052.
- 5. Myer L, Phillips TK. Beyond "Option B+": Understanding Antiretroviral Therapy (ART) Adherence, Retention in Care and Engagement in ART Services Among Pregnant and Postpartum Women Initiating Therapy in Sub-Saharan Africa. *Journal of acquired immune deficiency syndromes (1999).* 2017;75 Suppl 2:S115-s122.
- 6. Adams JW, Brady KA, Michael YL, Yehia BR, Momplaisir FM. Postpartum Engagement in HIV Care: An Important Predictor of Long-term Retention in Care and Viral Suppression. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2015;61(12):1880-1887.
- 7. Patel M, Tedaldi E, Armon C, et al. HIV RNA Suppression during and after Pregnancy among Women in the HIV Outpatient Study, 1996 to 2015. *J Int Assoc Provid AIDS Care.* 2018;17:2325957417752259.
- 8. Chen JS, Pence BW, Rahangdale L, et al. Postpartum HIV care continuum outcomes in the southeastern USA. *AIDS (London, England).* 2019;33(4):637-644.
- 9. Momplaisir FM, Storm DS, Nkwihoreze H, Jayeola O, Jemmott JB. Improving postpartum retention in care for women living with HIV in the United States. *AIDS (London, England)*. 2018;32(2):133-142.
- 10. Philadelphia Department of Public Health. *HIV Surveillance Report 2019 HIV in Philadelphia.* 2020.
- 11. Centers for Disease Control and Prevention. *Behavioral and Clinical Characteristics of Persons with Diagnosed HIV Infection — Medical Monitoring Project, United States, 2019 Cycle (June 2019-May 2020). HIV Surveillance Special Report 28.* 2021.
- 12. Haushofer J, Fehr E. On the psychology of poverty. *Science (New York, NY).* 2014;344(6186):862.
- 13. Weiser SD, Palar K, Hatcher AM, Young S, Frongillo EA, Laraia B. Food Insecurity and Health: A Conceptual Framework. In: Ivers LC, ed. *Food Insecurity and Public Health*. Boca Raton, Fla, USA: CRC Press; 2015:23-41.
- 14. Schilbach F, Schofield H, Mullainathan S. The Psychological Lives of the Poor. *Am Econ Rev.* 2016;106(5):435-440.
- 15. Mani A, Mullainathan S, Shafir E, Zhao J. Poverty Impedes Cognitive Function. *Science (New York, NY).* 2013;341(6149):976.

- Ong Q, Theseira W, Ng IYH. Reducing debt improves psychological functioning and changes decision-making in the poor. *Proceedings of the National Academy of Sciences*. 2019;116(15):7244.
- 17. Shah AK, Shafir E, Mullainathan S. Scarcity Frames Value. *Psychological Science*. 2015;26(4):402-412.
- 18. Shah AK, Zhao J, Mullainathan S, Shafir E. Money in the mental lives of the poor. *Social Cognition*. 2018;36(1):4-19.
- 19. Shah AK, Mullainathan S, Shafir E. Some consequences of having too little. *Science (New York, NY).* 2012;338(6107):682-685.
- 20. Lichand G, Mani A. *Cognitive Droughts.* CSAE Working Paper Series. Centre for the Study of African Economies, University of Oxford;2020.
- 21. Bartoš V, Bauer M, Chytilová J, Levely I. Psychological Effects of Poverty on Time Preferences. *The Economic Journal.* 2021;131(638):2357-2382.
- 22. Ibrahim M, Chen Y, Pereira A, Musa T, Bathia D, Handa S. The Psychology of Poverty: A Replication Based on Zimbabwe's Harmonised Social Cash Transfer Programme. *Journal of African Economies*. 2023:ejad013.
- 23. Aarts H, Dijksterhuis A, De Vries P. On the psychology of drinking: being thirsty and perceptually ready. *Br J Psychol.* 2001;92(Pt 4):631-642.
- 24. Radel R, Clément-Guillotin C. Evidence of motivational influences in early visual perception: hunger modulates conscious access. *Psychol Sci.* 2012;23(3):232-234.
- 25. Coleman-Jensen A, Rabbitt MP, Gregory CA, Singh A. *Statistical Supplement to Household Food Security in the United States in 2020, AP-091.* U.S. Department of Agriculture, Economic Research Service;2020.
- 26. Dean EB, Schilbach F, Schofield H. 2. Poverty and Cognitive Function The Economics of Poverty Traps. In: Barrett CB, Carter M, Chavas J-P, Carter MR, eds. *The Economics of Poverty Traps*. University of Chicago Press; 2019:57-118.
- 27. Dinges DF, Pack F, Williams K, et al. Cumulative sleepiness, mood disturbance, and psychomotor vigilance performance decrements during a week of sleep restricted to 4-5 hours per night. *Sleep.* 1997;20(4):267-277.
- Schofield H. Behavioral Development Lab: Resources. <u>https://behavioraldevlab.org/resources/resource.html</u>. Published 2021. Accessed 11/4/2021.
- 29. Raven J. The Raven's progressive matrices: change and stability over culture and time. *Cogn Psychol.* 2000;41(1):1-48.
- 30. Kirby KN, Petry NM, Bickel WK. Heroin addicts have higher discount rates for delayed rewards than non-drug-using controls. *J Exp Psychol Gen.* 1999;128(1):78-87.
- 31. Lawyer SR, Schoepflin F, Green R, Jenks C. Discounting of hypothetical and potentially real outcomes in nicotine-dependent and nondependent samples. *Exp Clin Psychopharmacol.* 2011;19(4):263-274.
- 32. Kaplan BA, Amlung M, Reed DD, Jarmolowicz DP, Mckerchar TL, Lemley SM. Automating Scoring of Delay Discounting for the 21- and 27-Item Monetary Choice Questionnaires. *The Behavior Analyst.* 2016;39(2):293-304.

- 33. Chuang Y, Schechter L. Stability of experimental and survey measures of risk, time, and social preferences: A review and some new results. *Journal of Development Economics*. 2015;117:151-170.
- 34. Binswanger HP. Attitudes toward Risk: Experimental Measurement in Rural India. *American Journal of Agricultural Economics*. 1980;62(3):395-407.
- 35. Stanczyk AB. The Dynamics of U.S. Household Economic Circumstances Around a Birth. *Demography*. 2020;57(4):1271-1296.
- 36. Kahneman D, Frederick S. Representativeness Revisited: Attribute Substitution in Intuitive Judgment. In: Gilobich T, Griffin D, Kahneman D, eds. *Heuristics and Biases: The Psychology of Intuitive Judgment*. Cambridge, UK: Cambridge University Press; 2002:49-81.
- 37. Richterman A, Thirumurthy H. The effects of cash transfer programmes on HIV-related outcomes in 42 countries from 1996 to 2019. *Nature Human Behaviour.* 2022.