

Study Title: The Bambisana Study Protocol: A mixed methods pre- and post- test study assessing community and social media influence to increase influenza vaccination uptake among youth in Soweto, South Africa

Authors:

Janan J. Dietrich^{1,2*}, Catherine Hill^{3*}, Gugulethu Tshabalala^{1,4}, Tshepiso Msibi³, Stefanie Vermaak^{1,4}, Nellie Myburgh³, Sarah Malycha⁵, Izzy Goldstein⁵, Elliot Grainger⁵, Prima Alam⁵, Kimberley Gutu³, Kennedy Otwombe^{4,6}, Heidi J. Larson^{5,7}, Ziyaad Dangor³

Affiliations

1. African Social Sciences Unit of Research and Evaluation (ASSURE), Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
2. Health Systems Research Unit, South African Medical Research Council, Bellville, South Africa
3. South African Medical Research Council Vaccines and Infectious Diseases Analytics Research Unit, University of the Witwatersrand, Johannesburg, South Africa
4. Perinatal HIV Research Unit (PHRU), School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
5. Department of Infectious Disease Dynamics, London School of Hygiene & Tropical Medicine, London, United Kingdom
6. School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
7. University of Antwerp and KU Leuven, Belgium
8. School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

*Joint first authorship

Corresponding Author: Janan Dietrich, janan.dietrich@wits.ac.za

50 **Abstract**

51 **Background**

52 Seasonal influenza has an estimated global reach of 3 to 5 million infections with 290 000 to 650 000
53 influenza-related deaths yearly. Despite its efficacy in reducing morbidity and mortality, influenza vaccination
54 rates remain low globally and in South Africa. Youth between the ages of 18-34 years are not prioritised for
55 influenza vaccines although influenza surveillance in South Africa shows that individuals aged 19 to 44
56 present the highest asymptomatic episodes and the lowest medically attended illness. This creates an
57 opportunity to investigate if and how vaccine demand can be created in the absence of clear imperatives to
58 vaccinate. The study tests the effectiveness of tailored, context-specific education, community engagement,
59 including community and social media to increase influenza vaccination uptake. Tailored, context-specific
60 education, community engagement, reliable vaccine supply and free, localised access are all critical for
61 improving perceptions of, increasing confidence in, and motivating uptake of vaccination. This study explores
62 strategies to increase vaccine uptake amongst marginalised youth 18-34 years old in Soweto, South Africa,
63 where influenza vaccines are not universally accessible through the public health system for this age group.

64 **Methods**

65 The Bambisana Study uses an innovative approach - including community influencers and social media - to
66 increase uptake of influenza vaccines through designing and testing an integrated communications strategy
67 targeted at marginalised youth in Soweto, South Africa. The Bambisana study uses a mixed methods pre-test,
68 post-test intervention design to test the effects of the interventions.

69 **Conclusion and Significance**

70 Enhancing perceptions of, bolstering confidence in, and fostering uptake of vaccination relies heavily on the
71 efficacy of yearly influenza vaccination initiatives, personalized education tailored to specific contexts, active
72 community involvement, consistent vaccine availability, and easily accessible, cost-free distribution channels
73 at the local level.

74
75 **Keywords:** Influenza, Vaccines, Vaccine Confidence, Youth, South Africa, Social Media
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

101 Introduction

102 Globally seasonal influenza causes an estimated 3-5 million infections, and 290 000-650 000 deaths annually
103 [1, 2]. Estimates for influenza burden of disease are limited in Africa; however, hospitalisation and mortality
104 are higher in low- and middle-income countries (LMICs) than in high-income countries (HICs) [3, 4]. In South
105 Africa, an estimated 6 000-11 000 people die from influenza every year, with approximately half of these
106 deaths in the elderly, and people living with HIV [5]. Seasonal outbreaks of influenza pose challenges to the
107 healthcare system, leading to increased hospitalizations, outpatient visits, and strain on healthcare resources
108 [6]. Vulnerable populations, including the elderly, young children, pregnant women, and individuals with
109 underlying medical conditions are at highest risk of severe infection [7].

110
111 While non-pharmaceutical interventions can play a role in decreasing influenza infections, vaccination is
112 deemed pragmatic and endorsed by the World Health Organization (WHO) as the recommended method for
113 preventing seasonal influenza [1]. Overall vaccine efficacy to prevent influenza infection ranges from 40-
114 60%, with higher vaccine efficacy (82%) in reducing severe illness and hospitalization [8, 9]. Therefore, the
115 influenza vaccine is especially recommended for those who have chronic illnesses, those who are pregnant,
116 and the elderly [10-12]. Influenza vaccination campaigns typically prioritize these groups during the influenza
117 vaccination season [7, 13]. However, an investigation into the seasonal influenza burden in urban and rural
118 areas in South Africa showed that children younger than 12 years and those 19 and 44 years – have an elevated
119 risk of acquiring seasonal influenza [14]. Whilst severe disease is less likely, vaccinating younger adults has
120 the added indirect benefit of providing herd immunity to the family and communities [15, 16]. Furthermore,
121 vaccination offers the potential economic benefit related to the out-of-pocket expenses incurred by patients to
122 treat influenza related symptoms. An estimated annual cost of \$111.3 million is incurred by the government,
123 while out of pocket costs incurred by the patient were reported to be \$40.7 million [17]. Individuals are at risk
124 of financial loss due to absence at work and missed opportunity for those who are self-employed [18]. Despite
125 recommendations, seasonal influenza vaccine uptake, in the most vulnerable groups, is low in LMICs,
126 including South Africa [19]. Accessibility, lack of knowledge and risk perception have been cited as the most
127 significant barriers to vaccination in South Africa [20].

128
129 Despite the demonstrated effectiveness of vaccines in significantly reducing morbidity and mortality of
130 affected individuals, youth are less proactive in health-seeking behaviours due to low risk perception, are not
131 necessarily encouraged to vaccinate for influenza and other infectious diseases [9, 21-23]. For COVID-19,
132 youth were not initially prioritised for vaccination roll-out yet they might have been in contact with vulnerable
133 populations [24, 25]. Additionally, evidence suggests that low vaccination rates are in part due to low health-
134 seeking behaviours among youth. Health-seeking behaviours are the actions involved in achieving and
135 maintaining a healthy condition. [26, 27]. Various factors, both individual and related to healthcare providers,
136 can influence young people's healthcare-seeking behaviours [28].

137
138 Targeted health promotion interventions can increase health-seeking behaviour and uptake of vaccination [29].
139 Influenza surveillance shows that individuals aged 19 to 44 present the highest asymptomatic episodes and
140 the lowest medically attended illness [14, 30]. Studies undertaken in sub-Saharan Africa have shown that
141 leveraging community involvement in immunization programs positively contributes to vaccine confidence
142 and uptake [31, 32]. In the case of HIV, for example, awareness campaigns to lower the burden of HIV and
143 AIDS epidemics in Southern African countries have proven vital in reducing the rise of disease [33]. The use
144 of television series to promote positive sexual health and prevent HIV effectively engaged the youth by
145 challenging old models of education [34]). The use of social media campaigns have also been increasingly
146 used to promote behaviour change [35, 36], while sparking debate, especially when considering the abundance
147 of false and misinformation [37]. It is vital to consider the significance of social media to improve vaccine
148 uptake in LMICs.

149
150 Promoting health care through various platforms and to youth in LMICs may improve uptake of vaccination.
151 Therefore, as part of the Bambisana Project, which includes a diverse and multi-disciplinary team comprising
152 social and behavioural scientists, biostatisticians, population health surveillance specialists, health

communications researchers, community influence and engagement practitioners, and marketing and communication strategists, we aim to:

- (1) Evaluate the impact of influencer’s online, offline, and in collaboration, on influenza vaccine uptake, with a specific focus on young people, and assess the additional value added by social media engagement;
- (2) To design and implement an integrated communications strategy centred around community influencers and social media to increase the uptake of influenza vaccines, particularly among younger populations who are likely unemployed, partially or informally employed and living in Soweto, South Africa;
- (3) To explore motivations and barriers to health priorities, and to influenza vaccination, through the use of micro-influencers both online and offline in Soweto, South Africa.

The outcomes of the project will include increased influenza vaccination rates at the local healthcare clinics linked to campaign interventions, alongside an improved attitude towards the importance of vaccination in the target audience - this will be assessed using the Vaccine Confidence Index scale [38] and measured against the most recent Vaccine Confidence Index data from South Africa [39]. Additionally, the project will provide important health systems-strengthening recommendations which could potentially be implemented within current infrastructure.

Materials and Methods

Research Design

The primary research aim uses a prospective mixed methods study design including qualitative and quantitative data collection, to assess the effects of community and social media interventions to increase influenza vaccination uptake among youth in Soweto, South Africa. During the intervention phase, surveys will be conducted among clinic attendees accepting as well as declining influenza vaccination within the study clinics.

Study setting

The study is conducted in Soweto (South-Western Townships) which is a congregation of 29 townships within the Johannesburg Metropolis in South Africa [40]. Soweto is one of the poorest areas of Johannesburg with high rates of youth unemployment which is at 46.3% among the 15-34 year age group [41]

Sampling


The Bambisana Project leverages the Soweto Health Demographic Surveillance System (HDSS), led by Wits VIDA since 2017 and part of the Child Health and Mortality Prevention Surveillance programme (CHAMPS), provides access to longitudinal health and demographic data for specific enumerated areas of Soweto and the South of Johannesburg, since 2017[42]. The study population includes young adults 18-34 years from four HDSS economically marginalised and geographically distinct Soweto communities: namely Phiri/Senaone and Mapetla, Mofolo and Meadowlands, Thulani, and neighbouring Thembelihle. Adults (18 years and older), living in these clusters or accessing clinic services in the five study clinics (Senaone Clinic, Mofolo Community Health Centre, Meadowlands Zone 2 Clinic, Siphumlile Clinic, and Thembelihle Clinic) are deemed eligible for the study. The HDSS communities were included as below (Table 1):

Table 1: HDSS cluster summary data

Cluster #	Communities	Total Population	Youth 18-34	# households with youth	Total # of households
Cluster 1	Mofolo and Meadowlands	20986	5824	3478	5912
Cluster 2	Mapetla & Phiri/Senaone	23805	6801	4037	6626
Cluster 3	Thulani	23711	6848	4072	6262
Cluster 4	Thembalihle	19449	5648	3831	6586

197 Figure 1: Study Design

PROJECT TIMELINE: DATA POINTS & STUDY DESIGN



198

199

Community engagement

200

201

202

203

204

205

206

207

208

209

Formative qualitative data collection

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

- FGDs: 16 FGDs (8-12 participants per FGD)
- KIIs: 20 (10 male and 10 female participants)

226

227

Intervention

The intervention will be delivered through a public health messaging campaign in targeted clusters or ‘sub-places’ in Soweto, leveraging micro-influencers to provide information and motivate positive attitudes for vaccination without mandate in currently unvaccinated populations. The aim of the campaign is to create a behavioural change action: to go to your local clinic to get vaccinated. This has the secondary aim of highlighting the extent to which such interventions might be possible on wider issues of public health in the future. Secondary to this is the desire to investigate the value-add of social media targeting over wider forms of engagement, such as that which occurs in communities offline.

The campaign will be timed with points when the clinic is well supplied and prepared for such a public drive. The project will mitigate against the risk of potential vaccine shortages by making vaccine donations to partner clinics. The study will centre around the 2023 national influenza vaccination drive in 2023. Vaccines are generally provided in very short and sporadic supply to the local community clinics, so supplying vaccines is also part of the objective to make community-level impact as part of the project.

The intervention will be implemented after community engagement discussions have been conducted to obtain information which would be used to design the messaging and communication strategy. Surveys will be administered prior to, during, and following the communication interventions to establish baseline, midline, and endline measures for evaluating the intervention's impact. This will help assess the effectiveness of four distinct approaches aimed at boosting influenza vaccine uptake among youth across clinics in four diverse HDSS clusters.

1. **Standard of care:** Vaccine made available with standard government promotional activity without support from the project (i.e., based on standard government drive).
2. **Social media:** Social media campaign to highlight availability and drive to get vaccinated pushed to the target audience.
3. **Micro-influencers on and offline:** The social media campaign will be amplified through a coordinated effort by micro-influencers in their online and offline networks.
4. **Offline micro-influencers:** Non-social media focused; offline micro-influencer led engagement within communities.

The intervention components are detailed below:

1) Social media campaign to highlight availability and drive to get vaccinated pushed to the target audience (under 35s) in four distinct geographic areas

Utilizing Meta platforms Facebook and Instagram, along with other relevant social media channels identified through community engagement and marketing insights, we will disseminate information through static images, carousel posts, and video posts to promote the availability of, education around, and motivation to get vaccinated. Paid advertising on Meta will target people aged 18-34 in two of the four distinct geographic study clusters.

2) Micro-influencers on and offline: The social media campaign will be amplified through a coordinated effort by micro influencers in their online and offline networks

We will use community influencers - people with traditional influence such as community leaders, religious leaders, elders, and key business leaders - to amplify our social media campaign. They will engage their online and offline networks in coordination with our social media campaign to promote vaccine availability and encourage vaccination.

3) Offline micro-influencers: Non-social media focused; offline micro-influencer led engagement within communities

In these communities, our community influencers will focus only on offline engagement and will be specifically instructed to not do anything on social media. Using interpersonal conversations, and small-scale community events to share the importance of vaccines, they will encourage people to get vaccinated.

Quantitative Data Collection

Quantitative data collection will include five survey data points as outlined below.

All surveys will be developed using the REDCap software, a computer-assisted survey platform with built-in data encryption [44]. All surveys will be completed using dedicated study tablets which means participants would not be required to use their own internet data. Additionally, we will use a data-free survey completion option. The survey links would also be sent to participants who wish to complete the surveys on their own devices. There will be no data costs to participants for completing the assessments. Participants will be reimbursed ZAR 50 airtime towards their time.

A) **Household surveys** include two time points of data collection, a pre- and post-test design, as described below.

Sampling

Households were chosen through a simple random sampling method, and eligible participants were enrolled as follows: a minimum of 800 adult men and women aged 18 years and above, with 400 designated for the pre-test and 400 for the post-test. At least 100 households with index participants aged 18-34 years will be approached per community. Non-index participants, include other adults in the households who are aged 18-34, 35-49 and 50 years and older who will also be invited to complete surveys.

Pre-test and post-test surveys: Using a random sample of households having youth participants (18-34 years) generated from the HDSS database, adults will be approached in their households to complete a 15 minute online pre-test assessment survey to measure: socio-demographics (age, gender, food security, potential loss of income), knowledge of influenza vaccines and benefits, health beliefs and sources of health information (social media, television, radio, friends and family), and vaccine confidence[45]. The same households will be invited to participate in the post-test household survey. The post-test household survey will be conducted after the study campaign intervention.

B) **Clinic surveys** include data collection with vaccinators and those who declined vaccination as described below:

Vaccinated: All adults 18+ years, who received an influenza vaccination will be approached as such numbers may vary as this depends on who elects to vaccinate. However, a total of at least 400 (n=100 per community) youth (aged 18-34) participants who go to the intervention clinics will be purposively approached to complete a post vaccination assessment. The study team will work closely with healthcare workers at public health clinics in the study communities. Healthcare workers will refer patients after having received the influenza vaccine, on site at the clinics. These participants will be invited to complete a brief survey using the same measures as specified for the pre-test survey. The data will be covered as part of the project expenses. There will be no data costs to participants for completing the assessments. Participants will be reimbursed ZAR 50 airtime towards their time.

Additional measures will assess: motivations for influenza vaccines, where participants heard/learned about this or other drives for the influenza vaccination, if participants had seen any online messaging driving them to get the influenza vaccine or been exposed to specific community engagement activities, and whether it had influenced their decision to vaccinate, and online impressions, for example, video views rates (dependent on

319 final campaign strategy and media mix) and their impact on vaccinations in that region. Uptake of influenza
320 vaccines will be tracked in the four clusters across the study period.

321
322 **Clinic attendees who decline vaccination** will also be approached to complete surveys. These participants
323 will be 18+ years. Healthcare workers will refer patients who were offered vaccination as part of their clinic
324 attendance but who declined vaccination. Those participants who agree to participate will be invited to
325 complete a brief survey to understand reasons for declining vaccination. The survey measures will include: if
326 and where participants heard/learned about this drive for the influenza vaccination, if participants had seen
327 any online messaging driving them to get the influenza vaccine or been exposed to specific community
328 engagement activities, and whether it had influenced their decision to vaccinate.

329
330 As with all the surveys, the data costs will be covered as part of the project expenses. There will be no data
331 costs to participants for completing the assessments. Participants will be reimbursed ZAR 50 airtime towards
332 their time.

333 C) Qualitative evaluation

334
335 Additional FGDs, including 10-12 participants per FGD, will be conducted: 8 mid-way through the
336 intervention (during vaccination season) and 8 post-intervention. These FGDs will be mixed gender and will
337 be stratified by age group (18-24, 25-34), and community (two per cluster). Key informant interviews (KII)
338 (n=20) with key influencers including, religious leaders, traditional healers and youth leaders, as well as other
339 influencers that may identified through community engagement and communications activities.

340 Data Analysis

341 Qualitative data preparation and analysis plan

342 Qualitative data will be analysed using a Framework Analysis approach. Framework thematic analysis
343 provides a highly systematic method of categorizing and organizing data according to key themes, concepts
344 and emergent categories in grids or matrices [46]. The coding will follow a combined approach, i.e. both
345 deductive based on our hypothesized themes and inductive (data-driven) to allow for emerging themes. The
346 analysis will allow us to further refine the domains and will help identify sub-themes. Each transcript will be
347 coded in Dedoose, a qualitative data analysis software by at least two researchers and assessed for consistency
348 in coding of the first 5 transcripts [47]. The research team will convene to review the initial codes and apply
349 general categories that will make up the analytic framework. Any coding disagreements will be discussed until
350 agreement is reached. This analytical framework will then be applied to the remaining transcripts.

351 Statistical Analysis Plan

352 We will use descriptive statistics to describe study variables including: knowledge of influenza vaccines,
353 experiences of influenza vaccine, stigma and vaccine confidence. Reliability assessments using the Cronbach
354 alpha test will be evaluated for scale measures. Scales with poor reliability scores will be calibrated using
355 factor analysis. Data will be analysed overall and stratified by community cluster. Frequencies will be
356 determined for categorical measures whereas means (standard deviation) and medians (interquartile ranges)
357 will be determined for continuous measures. Pairwise comparisons of categorical measures between
358 communities will be evaluated by Chi-square or Fisher's exact test as appropriate; Similarly, pairwise
359 continuous measures will be analysed by the two-sample t-test, if satisfying normality distribution conditions
360 or the Wilcoxon Mann-Whitney test, if non-normal. Normality will be assessed by the Shapiro-Wilks method.
361 Drivers of knowledge, stigma and experiences of influenza vaccines will be determined by logistic regression
362 for binary outcomes and linear regression for continuous outcomes. Multivariable analysis will be done to
363 adjust for potential confounders. Model fit statistics will be assessed on all multivariable regression models to
364 ensure the validity of the findings. Data analyses will be conducted using SAS Enterprise Guide 7.15 [48].

365 Ethics statement

366 All research procedures were approved by the University of the Witwatersrand Human Research Ethics
367 Committee (Medical) (M230254) and the local Department of Health. Participants were reimbursed ZAR 50
368 airtime for the surveys and ZAR 150 for the FGDs and the KIIs.

Bias correction

While the study is designed to focus on four relatively distinct geographic areas, due to the nature of social media and health-seeking behaviours in the broader community of Soweto, it is possible that members from the intervention communities may interact with those not receiving the intervention. While potential interaction cannot be prevented entirely, research questions for all participants will include questions about exposure to intervention components and other potentially similar national or localised campaigns. In addition, a phased delivery of the interventions to ensure the offline intervention is not biased by the online intervention should help mitigate this.

It is also possible that those exposed to the intervention(s) may seek to be vaccinated in clinics that are not included in the study's geographically defined target-areas. An effort will be made to collaborate with the Department of Health to broadly understand influenza vaccinations during the campaign season, in broader Soweto and areas where residents typically seek healthcare. Vaccine and vaccination sentiments broadly in Soweto and South Africa will also be tracked during the course of the project for context and monitoring of potential risk.

Funding

This project was funded by The Vaccine Confidence Fund II, a philanthropic and charitable fund of Global Impact. Grant ID: VCFII-011. Wits VIDA is an extramural unit of the South African Medical Research Council. The work reported herein for Janan J Dietrich was also made possible through funding by the South African Medical Research Council through its Division of Research Capacity Development under the Early Investigators Programme through funding received from the South African National Treasury. The content and findings reported/illustrated are the sole deduction, view and responsibility of the authors and do not reflect the official position and sentiments of the Vaccine Confidence fund and the SAMRC.

Acknowledgements

We would like to extend our sincere gratitude to all study participants and beneficiaries, the Wits Vida study team, as well as the FreshNow Team for developing and delivering the content for the campaign.

Conflicts of Interest

The authors report no conflicts of interest.

References

1. World Health Organization. *Influenza (Seasonal)* 2023.
2. The National Institute for Communicable Diseases (NICD). 2023. *Increase in flu cases across the country*(5 June 2023).
3. Langer, J., et al., *High Clinical Burden of Influenza Disease in Adults Aged ≥ 65 Years: Can We Do Better? A Systematic Literature Review*. *Advances in Therapy*, 2023. **40**(4): p. 1601-1627.
4. Gharpure, R., et al., *Costs and cost-effectiveness of influenza illness and vaccination in low- and middle-income countries: A systematic review from 2012 to 2022*. *PLOS Medicine*, 2024. **21**(1): p. e1004333.
5. Motlogeloa, O., J.M. Fitchett, and N. Sweijd, *Defining the South African Acute Respiratory Infectious Disease Season*. *Int J Environ Res Public Health*, 2023. **20**(2).
6. Hagiwara, Y., et al., *Seasonal influenza, its complications and related healthcare resource utilization among people 60 years and older: A descriptive retrospective study in Japan*. *PLoS One*, 2022. **17**(10): p. e0272795.
7. McMorrow, M.L., et al., *Prioritization of risk groups for influenza vaccination in resource limited settings - A case study from South Africa*. *Vaccine*, 2019. **37**(1): p. 25-33.
8. Comber, L., et al., *Systematic review of the efficacy, effectiveness and safety of high-dose seasonal influenza vaccines for the prevention of laboratory-confirmed influenza in individuals ≥ 18 years of age*. *Rev Med Virol*, 2023. **33**(3): p. e2330.
9. Andre, F.E., et al., *Vaccination greatly reduces disease, disability, death and inequity worldwide*. *Bull World Health Organ*, 2008. **86**(2): p. 140-6.

- 424 10. Antonelli Incalzi, R., et al., *Influenza vaccination for elderly, vulnerable and high-risk subjects: a*
425 *narrative review and expert opinion*. Internal and Emergency Medicine, 2023.
- 426 11. The National Institute for Communicable Diseases (NICD). 2023. *Influenza: NICD recommendations*
427 *for the diagnosis, management, prevention and public health response 2023*.
- 428 12. Diop, D., *Influenza-Associated Morbidity and Mortality in Sub-Saharan Africa*. International Journal
429 of Vaccines & Vaccination, 2016. **2**(4).
- 430 13. Kaposy, C. and N. Bandrauk, *Prioritizing Vaccine Access for Vulnerable but Stigmatized Groups*.
431 Public Health Ethics, 2012. **5**(3): p. 283-295.
- 432 14. Cohen, C., et al., *Asymptomatic transmission and high community burden of seasonal influenza in an*
433 *urban and a rural community in South Africa, 2017–18 (PHIRST): a population cohort study*.
434 The Lancet Global Health, 2021. **9**(6): p. e863-e874.
- 435 15. Plans-Rubió, P., *The vaccination coverage required to establish herd immunity against influenza*
436 *viruses*. Prev Med, 2012. **55**(1): p. 72-7.
- 437 16. Glezen, W.P., *Herd protection against influenza*. Journal of Clinical Virology, 2006. **37**(4): p. 237-
438 243.
- 439 17. Tempia, S., et al., *Health and economic burden of influenza-associated illness in South Africa, 2013-*
440 *2015*. Influenza Other Respir Viruses, 2019. **13**(5): p. 484-495.
- 441 18. Blanchet Zumofen, M.H., J. Frimpter, and S.A. Hansen, *Impact of Influenza and Influenza-Like Illness*
442 *on Work Productivity Outcomes: A Systematic Literature Review*. Pharmacoeconomics, 2023. **41**(3):
443 p. 253-273.
- 444 19. Ortiz, J.R. and K.M. Neuzil, *Influenza Immunization in Low- and Middle-Income Countries:*
445 *Preparing for Next-Generation Influenza Vaccines*. The Journal of Infectious Diseases, 2019.
446 **219**(Supplement_1): p. S97-S106.
- 447 20. Alobwede, S.M., et al., *Influenza Vaccination Uptake and Hesitancy among Healthcare Workers in*
448 *Early 2021 at the Start of the COVID-19 Vaccine Rollout in Cape Town, South Africa*. Vaccines, 2022.
449 **10**(8): p. 1176.
- 450 21. Kim, A.W., et al., *Perceptions, risk and understandings of the COVID-19 pandemic in urban South*
451 *Africa*. S Afr J Psychiatr, 2021. **27**: p. 1580.
- 452 22. Quinn, S.C., et al., *Measuring vaccine hesitancy, confidence, trust and flu vaccine uptake: Results of*
453 *a national survey of White and African American adults*. Vaccine, 2019. **37**(9): p. 1168-1173.
- 454 23. Rodrigues, C.M.C. and S.A. Plotkin, *Impact of Vaccines; Health, Economic and Social Perspectives*.
455 Front Microbiol, 2020. **11**: p. 1526.
- 456 24. Muthoni, J., et al., *COVID-19 Vaccination Hesitancy among Youths in Soweto, South Africa*. Vaccines
457 (Basel), 2023. **11**(5).
- 458 25. Euser, S., et al., *Understanding COVID-19 vaccination willingness among youth: A survey study in*
459 *the Netherlands*. Vaccine, 2022. **40**(6): p. 833-836.
- 460 26. Otjombe, K., et al., *Health-seeking behaviours by gender among adolescents in Soweto, South Africa*.
461 Glob Health Action, 2015. **8**: p. 25670.
- 462 27. G, A.J.T., *Health Seeking Behaviours in South Africa: A Household Perspective using the General*
463 *Households Survey of 2007*, in *Statistics and population Studies*. 2010, University of the Western Cape
- 464 28. Latunji, O.O. and O.O. Akinyemi, *factors influencing health-seeking behaviour among civil servants*
465 *in ibadan, nigeria*. Ann Ib Postgrad Med, 2018. **16**(1): p. 52-60.
- 466 29. Hopkins, K.L., et al., *Community-Based Approaches to Increase COVID-19 Vaccine Uptake and*
467 *Demand: Lessons Learned from Four UNICEF-Supported Interventions*. Vaccines, 2023. **11**(7): p.
468 1180.
- 469 30. Hayward, A.C., et al., *Comparative community burden and severity of seasonal and pandemic*
470 *influenza: results of the Flu Watch cohort study*. The Lancet Respiratory Medicine, 2014. **2**(6): p. 445-
471 454.
- 472 31. Chantler, T., et al., *'We All Work Together to Vaccinate the Child': A Formative Evaluation of a*
473 *Community-Engagement Strategy Aimed at Closing the Immunization Gap in North-West Ethiopia*.
474 Int J Environ Res Public Health, 2018. **15**(4).
- 475 32. Jalloh, M.F., et al., *Association of community engagement with vaccination confidence and uptake: A*
476 *cross-sectional survey in Sierra Leone, 2019*. J Glob Health, 2022. **12**: p. 04006.

- 477 33. Johnson, L.F., et al., *The Effect of HIV Programs in South Africa on National HIV Incidence Trends*,
478 2000–2019. JAIDS Journal of Acquired Immune Deficiency Syndromes, 2022. **90**(2): p. 115-123.
- 479 34. Banerjee, A.V.a.L.F., Eliana and Orozco-Olvera, Victor H., T *The Entertaining Way to Behavioral*
480 *Change: Fighting HIV with Mtv*. NBER Working Paper No. w26096, 2019.
- 481 35. Neely, S., C. Eldredge, and R. Sanders, *Health Information Seeking Behaviors on Social Media During*
482 *the COVID-19 Pandemic Among American Social Networking Site Users: Survey Study*. J Med
483 Internet Res, 2021. **23**(6): p. e29802.
- 484 36. Ventola, C.L., *Social media and health care professionals: benefits, risks, and best practices*. Pt, 2014.
485 **39**(7): p. 491-520.
- 486 37. Jane, M., et al., *Social media for health promotion and weight management: a critical debate*. BMC
487 Public Health, 2018. **18**(1): p. 932.
- 488 38. Project, V.C. *Vaccine Confidence Index Data & Methodology*. 2022; Retrieved [17 January 2024],
489 from: <https://www.vaccineconfidence.org/vci/data-and-methodology/>.
- 490 39. The Centers for Disease Control and Prevention (CDC). 2022. *Wave 2 Vaccine Perceptions Survey in*
491 *South Africa*. 2022, Vaccine Confidence Project.
- 492 40. City of Johannesburg. *About Region D*. 2018. Retrieved [21 February 2024], from:
493 <https://joburg.org.za/about /regions/Pages/Region%20D%20-%20Greater%20Soweto/about-us.aspx>.
- 494 41. Statistics South Africa. *Youth still find it hard to secure jobs in South Africa*. 2021. Retrieved [11
495 December 2023], from: <https://www.statssa.gov.za/?p=14415>.
- 496 42. Cunningham, S.A., et al., *Health and Demographic Surveillance Systems Within the Child Health and*
497 *Mortality Prevention Surveillance Network*. Clinical Infectious Diseases, 2019. **69**(Supplement_4): p.
498 S274-S279.
- 499 43. Saunders, B., et al., *Saturation in qualitative research: exploring its conceptualization and*
500 *operationalization*. Qual Quant, 2018. **52**(4): p. 1893-1907.
- 501 44. *RedCap*. 2004.
- 502 45. de Figueiredo, A., et al., *Mapping global trends in vaccine confidence and investigating barriers to*
503 *vaccine uptake: a large-scale retrospective temporal modelling study*. Lancet, 2020. **396**(10255): p.
504 898-908.
- 505 46. Gale, N.K., et al., *Using the framework method for the analysis of qualitative data in multi-disciplinary*
506 *health research*. BMC Medical Research Methodology, 2013. **13**(1): p. 117.
- 507 47. 9.0.17, D.V., *cloud application for managing, analyzing, and presenting qualitative and mixed method*
508 *research data*, C.S.R. Consultants, Editor. 2021: Los Angeles.
- 509 48. Institutue, S., *SAS Enterprise Guide 7.15* 2022.
- 510
511
512