

Equal, equitable or exacerbating inequalities? Patterns and predictors of social prescribing referrals in 160,128 UK patients

Feifei Bu, PhD¹; Daniel Hayes, PhD¹; Alexandra Burton, PhD¹; Daisy Fancourt, PhD^{1*}

¹ Department of Behavioural Science and Health, Institute of Epidemiology & Health Care, University College London

* Correspondence to: Prof Daisy Fancourt, 1-19 Torrington Place, Department of Behavioural Science and Health, University College London WC1E 7HB, UK d.fancourt@ucl.ac.uk

Keywords: social prescribing, inequalities, mental health, wellbeing, social determinants

Abstract

Background: Social prescribing (SP) is growing rapidly globally as a way to tackle social determinants of health. However, whom it is reaching and how effectively it is being implemented remains unclear.

Aims: To gain a comprehensive picture of SP in the UK, from referral routes, reasons, to contacts with link workers and prescribed interventions.

Methods: This study undertook the first analyses of a large database of administrative data from over 160,000 individuals referred to SP across the UK. Data were analysed using descriptive analyses and regression modelling, including logistic regression for binary outcomes and negative binomial regression for count variables.

Results: Mental health was the most common referral reason and mental health interventions the most common interventions prescribed. Between 72% and 85% of SP referrals were from medical routes (primary or secondary health care). While these referrals demonstrate equality in reaching across socio-demographic groups, individuals from more deprived areas, younger adults, men, and ethnic minority groups were reached more equitably via non-medical routes (e.g. self-referral, school, charity). Despite 90% of referrals leading to contact with a link worker, only 38% resulted in any intervention being received. A shortage of provision of community activities - especially ones relevant to mental health, practical support, and social relationships - was evident. There was also substantial heterogeneity in how SP is being implemented across UK nations.

Conclusions: Mental health is the leading reason for SP referrals, demonstrating its relevance to psychiatrists. But there are inequalities in referrals. Non-medical referral routes could play an important role in addressing inequality in accessing social prescribing, therefore should be prioritised. Additionally, more financial and infrastructural resource and strategic planning are needed to address low intervention rates. Further investment into large-scale data platforms and staff training are needed to continue monitoring the development and distribution of social prescribing.

38 Introduction

39 Social determinants of health (SDOH), including income, education, employment, housing, childhood
40 experience, social support, have long been demonstrated to play an important role in determining health
41 outcomes.¹⁻³ SDOH are estimated to account for 47% of health outcomes compared to just the 16%
42 attributed to clinical care.¹ Social prescribing (SP) is proposed as a way to address SDOH and improve
43 health and wellbeing outcomes.⁴ Although SP has been advocated and practiced since 1980s, not until the
44 last decade has it been widely recognised and implemented across the UK.⁵ SP involves linking patients
45 with non-medical forms of support in their local communities, such as physical activity, social support,
46 mental health support, financial advice, housing support, and so forth.⁶ The predominant SP model
47 involves a healthcare professional referring patients on to a Link Worker, Community Connector,
48 Community Navigator, or Health Trainer, who then works with the patient to develop a personalised care
49 plan which connects them to community support and interventions.⁴

50
51 Evidence of mixed quality suggests that SP helps with a wide range of problems, including supporting
52 mental health, wellbeing and health behaviours.⁷⁻¹⁰ Preliminary evidence also suggests that SP either
53 reduces healthcare demand and costs or moves healthcare demand to primary care services rather than
54 more expensive tertiary care, and demonstrates a favourable return on investment.¹¹ Given its potential,
55 NHS England has committed to tripling the number Link Workers by 2036/37.¹² However, despite the
56 promises of SP, national evaluations remain scant. Instead, evidence on SP is often limited to small scale,
57 local evaluations and homogenous patient groups.¹³⁻¹⁶ Reviews into the use of SP suggest that referrals are
58 not equitable, with those who are under 16, male, and those from non-white backgrounds being less likely
59 to be offered SP.¹⁴ There are even concerns that SP is not effective for addressing social inequalities – it
60 cannot address upstream determinants of inequalities and could even be provided disproportionately
61 more to individuals who are less disadvantaged.¹⁷⁻¹⁹

62
63 To try and better understand the national picture of SP, studies have begun to explore patterns of SP in
64 electronic patient health records. For example, the SP Observatory has analysed administrative data on SP
65 referrals from over 1800 GP practices in England, showing a rapid increase in referrals since 2020,
66 especially during the COVID-19 pandemic.²⁰ Significant variation in rates of referrals across England has
67 been identified, but with no substantial variation in referral rates by gender, ethnicity or neighbourhood
68 deprivation.²⁰ Whilst findings such as this are of great importance, there are limitations. First, SP is not
69 limited to GP referrals, with referrals also made in secondary care,²¹ through voluntary and community
70 sector organisations (VCSOs), via statutory services (e.g. education, social care), self-referral and more.²²
71 Currently, the demographics of SP referrals via non-medical pathways are largely unknown. Moreover,
72 given that certain demographics, particularly people from ethnic minority backgrounds, are less likely to
73 report positive experiences accessing GP services²³ and young people report not wanting to access GP
74 support for their mental health,²⁴ it is important to establish if such groups are accessing SP via these
75 alternative pathways. Second, there is only very limited information available in SNOMED codes used by
76 GP practices, and these can include inconsistent coding, alongside poor recording of wider determinants of
77 health leading to conclusions that they cannot be used to assess equity of referrals.^{20,25} Indeed, little is
78 known about if SP referrals lead to any contact with link workers and actual interventions, why people are
79 referred to SP, what interventions are prescribed, and what aspects lead to a successful, engaged SP
80 referral. Third, national SP research has largely relied on data in England. How SP is implemented in other
81 devolved countries in the UK, is also unknown.

82
83 The growth in cloud-based referral-management platforms that allow multi-stakeholder data input about
84 SP pathways and integrate with primary care, secondary care and social care systems, is presenting new
85 opportunities for understanding how SP is working and who it is reaching.²⁶ However, data from these
86 platforms have not been properly capitalised on. Therefore, the present study was initiated to address the
87 knowledge gaps outlined above by using administrative data from one such SP platform: Access Elemental.

38 We aimed to explore demographic characteristics of people who were referred to SP, referral reasons
39 (including the importance of mental health within this), number of contacts with link workers, and
40 interventions being provided through SP pathways. We also aimed to explore whether these rates differed
41 across medical and non-medical referral pathways and whether there were any early indications of
42 differences in individual country SP models. Overall, these analyses aimed to provide a more
43 comprehensive understanding of SP roll-out across the UK, in particular for psychiatrists, with important
44 implications for future policy, service, and workforce planning.

35 Method

36 Data

37 Data were administrative records obtained from Access Elemental,²⁷ a digital SP platform used by health
38 and social care professionals, community development workers and other service providers to keep track
39 of SP activities and their impact from the point of referral. As a paid service, Elemental is the most widely
40 adopted SP platform in the UK to date. It serves a population of over 20 million people across England,
41 Scotland, Wales and Northern Ireland, and is used by over 37,740 health and care professionals and 4,405
42 social prescribers. To date, Elemental has recorded over 2.1 million contacts between patients and SP
43 teams, including over 438,000 referrals to community-based programmes, services, and interventions.
44 While Elemental does not comprise a random sample of sites or patients in the UK, it does have good
45 geographical coverage in England (45% of Integrated Care Boards), Wales (57% of Welsh University Health
46 Boards) and Northern Ireland (80% of Health and Social Care Trusts), alongside smaller representation in
47 Scotland (13% of Scottish Health and Social Care Partnerships). Further, due to the existence of multiple IT
48 systems in primary care in the UK, it is typical for research involving primary care to only involve a
49 proportion of the whole population.²⁸

10
11 Elemental collects detailed information tracking service users' SP journeys, including referral source,
12 referral reason, appointment with link worker, prescribed interventions, and so forth. They also collect
13 core demographic information, such as gender, age, ethnicity (optional). Elemental data are available since
14 January 2017, and this study used data up until November 2022. In this time, there were 201,037 SP cases
15 from 169,818 unique individuals. 94.3% of individuals provided valid postcodes (N=160,128), allowing for
16 data linkage with publicly available geographic information (see Figure S1). Figure S2 shows the
17 distribution of SP service users across local authorities in the UK. Figure S3 shows the number of SP service
18 users over time in the UK and across different countries. Most cases (88%) occurred between 2020-2022 in
19 the UK, after the formal launch of the scheme as part of the NHS England Long-Term Plan.²⁹ The research
20 project was approved by the UCL Research Ethics Committee (23909/001).

21 Measurement

22 Referral sources were used as binary medical vs non-medical referrals. Medical referrals were mainly from
23 GP practices, but also secondary care staff such as occupational therapists, physiotherapists, discharge
24 teams and hospital based social workers. Non-medical referrals could come from educational
25 establishments, local authorities, VSCOs, housing associations, self-referral, and so forth.

26
27 Referral reasons were coded in 600+ categories and we grouped them into seven domains: 1) mental
28 health and wellbeing, 2) physical health and wellbeing, 3) social relationships (e.g. loneliness and social
29 isolation), 4) lifestyle (e.g. exercise, diet, drinking, smoking), 5) employment, education and skills, 6)
30 practical support (e.g. housing, finance, legal), 7) other reasons (see Table S1 in the Supplement). The
31 grouping was done by two of the authors independently and any disagreement was discussed and resolved
32 by consensus. These domains were non-mutually exclusive, allowing people to have more than one referral
33 reasons.

34

35 Once people were referred to SP, they were connected to a SP link worker or a professional in a similar
36 role (see Figure S4). We were interested in how many contacts with a link worker were made related to
37 each SP case, referred as the number of contacts.

38
39 Contacts with link workers might or might not lead to individuals being connected with suitable community
40 resources, referred as interventions in the present study. The intervention variable concerns if any
41 prescription was issued by the link worker and what activities it entailed, without addressing service
42 attendance or engagement. To be consistent with referral reasons, interventions (400+ categories) were
43 also grouped into seven non-mutually exclusive domains (Table S1): 1) mental health and wellbeing, 2)
44 physical health and wellbeing, 3) social relationships, 4) lifestyle, 5) employment, education and skills, 6)
45 practical support, 7) other support.

46
47 Socio-demographic covariates included gender (female, male, other), age (0-17, 18-29, 30-39, 40-49, 50-59,
48 60-69, 70-79, 80+), urbanicity (urban, rural) and index of multiple deprivation (deciles: 1-most deprived to
49 10-least deprived). Both urbanicity and area of deprivation were derived via postcode linkage to urban
50 rural classification and index of multiple deprivation at Lower Layer Super Output Area (LSOA) in each
51 country. Although ethnicity was available, as an optional measure it had a high missing rate (>83%), which
52 was excluded from most analysis.

53

54 Statistical analysis

55 Descriptive analyses were undertaken to compare the socio-demographic profiles of people referred to SP
56 services across UK countries, the percentages of different referral reasons, types of interventions received,
57 and to what extent interventions matched to referral reasons. Regression analyses were conducted to
58 assess variables that were associated with SP services stratified by country, excluding Northern Ireland due
59 to a large proportion of missing data (82.9%) on referral route (Table S2). Referral route and interventions
60 received were analysed using logistic regression models. The number of contacts with link workers was
61 modelled using negative binomial regression to account for over-dispersion. All models controlled for
62 socio-demographic covariates. A small percentage of people had more than one SP case (<14%), so the
63 most recent record was used in regression analyses. Missing data were excluded in descriptive analyses
64 but imputed in regression models by using multiple imputation with chained equations, including all
65 covariates (m=30). All analyses were conducted in Stata V18.

56 Results

57 Demographic profiles

58 Among 160,128 unique individuals referred to SP, 141,011 people (88.1%) lived in England, 9,613 (6.0%) in
59 Wales, 2,374 (1.5%) in Scotland and 7,130 (4.5%) in Northern Ireland. This proportion was largely similar to
60 national statistics, with a slight overrepresentation of people from England (Table S3 in the Supplement).

61 There was an overrepresentation of females being referred to SP in all countries, especially in Northern
62 Ireland where 72.8% were females (Figure 1a, Table S4). Most service users (78.0%-87.2%) were aged 18 to
63 69 in each country (Figure 1b). However, Wales (5.2%) and Northern Ireland (11.5%) had a higher
64 percentage of people aged under 18. Scotland had a higher percentage of people in their 60s (32.7%). Only
65 5.6% of SP service users lived in rural areas in England, much lower than the rural population estimate of
66 17.1% (Figure 1c). However, this percentage was substantially higher in Wales (24.1% vs 20% rural
67 population), Scotland (27.2% vs 17% rural population) and Northern Ireland (35.5% vs 36% rural
68 population). SP service users were more likely to live in deprived areas (Figure 1d). In the general
69 population, roughly 10% of people living in each deprivation decile in each country. In England, 45.9% of
70 people referred to SP lived in the top three most deprived areas, 45.8% in Wales and 63.8% in Northern
71 Ireland. However, in Scotland, service users were more evenly distributed across deprivation deciles,
72 except for the top two least deprived deciles.

73

34 [Figure 1 here]

35 Referral routes

36 Most SP cases were referred from medical routes in England (85.3%), Wales (72.3%) and Scotland (84.8%).
37 The percentage was much lower in Northern Ireland (22.8%), possibly due to a high missing rate (82.9%,
38 see Table S2). Table S5 in the Supplement compares socio-demographic profiles of people referred from
39 medical and non-medical routes. Notably, SP service users referred via medical routes were more evenly
40 distributed across ethnic groups, and deprivation deciles; whereas those from non-medical routes were
41 more balanced across gender and urbanicity groups.

42 All socio-demographic factors were related to referral routes, but estimated associations differed across
43 countries (Figure 2 & Table S6). Key country differences were summarised in Table 1.

44 [Figure 2 here]
45 [Table 1 here]

37 Referral reasons

38 The most common referral reasons were related to mental health and wellbeing (33.5%), followed by
39 practical support (26.1%) and social relationship (22.5%); whereas employment or education/skills (10.3%)
40 and physical health (16.4%) related reasons were relatively less common (Figure 3a). Most SP cases (69.7%)
41 reported reasons from a single domain only. About a third reported reasons from more than one domain.
42 As shown in Figure 3b, 60.3% of cases referred for mental health reasons also had reasons from other
43 domains, 52.3% practical support, 63.4% social support and 50.7% physical health. Figure S5 shows
44 differences across countries. Physical health was among the most common reason in Wales, Scotland and
45 Northern Ireland. Lifestyle related reasons were also common in Scotland and Northern Ireland.

46 [Figure 3 here]

38 Contacts with link workers

39 After being referred, each case would be assigned to a link worker who typically initiated contacts.
40 Approximately 90% of SP cases had at least one contact with a link worker. Among cases without any
41 recorded contact with a link worker, 53.1% were active cases who might yet to be contacted. On average,
42 there were 4.2 contacts per case (SD=6.8, median=2). Among all the contacts, 11.5% were unsuccessful,
43 31.6% having reached clients with specific outcomes (e.g. appointment arranged, referral made,
44 signposted, declined service), and 56.9% unclear (e.g. contact made, email/message left). Most of contacts
45 (81.2%) were made via phone, video, email or letter, only 6.3% face-to-face, and another 12.6%
46 unspecified. The contacts typically lasted for 0-15 minutes (59.4%), but some were longer (Figure S6).
47 People referred from a medical route had a higher number of contacts across all countries (Figure S7 and
48 Table S7). Gender, age, urbanity and area deprivation were also associated with the number of contacts,
49 but some heterogeneities were founded across countries (Table 1).

21 Interventions

22 Among 201,037 cases, 38.3% (n=76,992) had at least one intervention prescribed. Intervention
23 prescription rates were 35.4% in England, 33.0% in Wales, but much higher in Scotland (51.9%) and
24 Northern Ireland (74.9%). Referral routes, number of contacts, referral reasons, age, urbanity were
25 associated with the odds of having an intervention prescribed (Figure S8 and Table S8). But there were
26 notably differences across countries (Table 1). For example, urban residents were more likely to have an
27 intervention in England, but less likely in Wales and Scotland than rural residents.

28
29 For SP cases with at least one intervention (n=76,971), there were 165,595 prescribed interventions (2.2
30 per case). Most of these interventions (91.7%) were recorded as free to access. Only 6.3% had out-of-
31 pocket costs and another 1.9% were subsidised (Table S9). England and Wales were similar in that over 95%
32 interventions were free. But only 50.3% interventions were free in Scotland and 78.7% in Northern Ireland
33 (Table S9).

34

35 Most prescribed interventions (82.9%) had information on service domains, with 62.9% of these
36 interventions falling under one service domain and 37.1% with more than one domain. As shown in Figure
37 4a, around 43.8% SP cases had an intervention related to mental health and wellbeing, followed by social
38 relationships (36.3%), lifestyle (28.7%), employment, education and skills (25.4%), practical support
39 (24.6%), physical health and wellbeing (19.0%) and other support (12.7%). Linking back to referral reasons,
40 Figure 4b shows the percentage of cases with an intervention prescribed by referral reasons in the UK. 69.3%
41 of cases who were referred due to mental health reasons had no intervention, 67.6% for physical health,
42 71.0% for social relationship, 65.9% for lifestyle, 70.5% for practical reasons, 72.8% for other reasons. The
43 percentage of cases with no intervention was much lower for those referred for employment related
44 reasons (37.1%). Among cases who did have an intervention, those who were referred for lifestyle related
45 reasons were more likely to have an intervention matched to these reasons (81.7%), followed by
46 employment (76.1%), mental health (70.3%), practical support (60.5%) and social relationships (60.1%)
47 reasons. Cases for physical health (32.7%) or other reasons (14.9%) were less likely to have a matched
48 intervention, but the patterns differed across countries (Figure S9). For instance, 35.9% of cases who were
49 referred due to lifestyle reasons had no intervention in Scotland, compared to 20.8% in Northern Ireland.
50 Moreover, only 18.6% of cases with social reasons had no intervention in Northern Ireland.

51

52

[Figure 4 here]

53 Discussion

54 This study is the first UK-wide study on SP referrals in the UK, with fine details on service users' socio-
55 demographic profiles, referral routes, reasons, contacts with link workers and actual interventions and
56 their costs. While relying on data from just one data management system, it provides importance evidence
57 to triangulate with previous research using GP records and smaller-scale evaluations by providing (i)
58 simultaneous data on both medical and non-medical referral sources, allowing the first direct comparison
59 of population characteristics, (ii) detail not just on whether a referral was made but the process by which
60 the referral pathway was carried out, and (iii) a large sample size with data over several years. Most
61 notably, it highlights that mental health is the leading cause of referrals across the UK. It also systematically
62 investigates the large-scale differences between medical and non-medical SP pathways across the UK and
63 provides early data on differences across the devolved countries in the UK. Understanding of the status-
64 quo of SP in the UK is crucial for identifying its future directions and supporting the roll-out of SP as a
65 public health intervention around the world.

56

57 As expected, the medical pathway is the dominant referral route in the UK. The seemingly exception of
58 Northern Ireland is likely to be driven by missing data. The medical pathway has its merits. Everyone in the
59 UK is entitled to free healthcare, and most people (if not all) are registered with a GP. This may explain why
60 SP service users were more evenly distributed across deprivation deciles and there was a higher
61 percentage of people from ethnic minority background among people from the medical route compared to
62 non-medical routes. This is consistent with previous findings of no ethnic or area deprivation difference in
63 SP when focusing on primary care data.²⁰ In this vein, the medical pathway prevails in promoting equality.
64 However, when it comes to actual service utilisation, there are notable socio-demographic differences. For
65 example, there is less use of primary healthcare services among young people and men,^{30,31} which may
66 limit their chances of being referred to SP through primary care. In addition to *equality*, we need to
67 consider *equity*. People from deprived areas generally have higher risks of health problems and more
68 complex needs, potentially requiring extra support and resources to achieve an equal outcome.^{32,33} Our
69 findings suggest that the non-medical pathway complements the medical pathway by reaching the
70 underrepresented in healthcare and those who are in greater need of support (e.g. those from more
71 deprived areas). The two pathways should work hand in hand to maximise the overall impact and
72 efficiency of SP.

32

33

34 Referral, regardless of route, is only an early step in the lifecycle of an SP case. A key question is whether a
35 referral leads to any intervention. Our analyses revealed that 9 in 10 referred SP cases had at least one
36 contact with a link worker, showing a successful initial transition. However, only 38.3% of cases were
37 subsequently referred to an intervention. What is unclear is whether this low level of intervention
38 engagement was voluntary (i.e. patients finding sessions of talking and a supportive relationship sufficient
39 for their needs) or due to other issues such as a Link Worker not being able to reach the client, referral
40 being declined, or a lack of appropriate resources or suitable interventions in the local community
41 (especially considering that over 90% of prescribed interventions are free of charge so depend on such
42 provision being available). However, several factors suggest that it is not entirely voluntary. First, referral
43 to an intervention is not even. People living in urban areas are more likely to be referred to an intervention
44 than rural residents, suggesting that a lack of provision may be an issue. Additionally, although mental
45 health, practical support and social relationship are the most common referral reasons, the percentage of
46 people having a matched intervention for each of these reasons is only around 20%, compared to 47.9%
47 for employment – provision of free employment support which has long been a priority in the UK,
48 suggesting that when free interventions are readily available, uptake is higher.³⁴ Therefore, there is urgent
49 work that is needed either building further community infrastructure in domains outside employment to
50 enable free services to be available, or providing further ringfenced funding for link workers to utilise in
51 under-resourced intervention domains. Additionally, more behavioural research is recommended to
52 identify what the barriers to accessing interventions are and how they can be addressed.

33

34 Another finding that stands out is the heterogeneity of SP across countries in the UK. Socio-demographic
35 profiles of SP service users across countries. For example, in Wales and Northern Ireland, there are higher
36 percentages of children and young people under 18, as well as people from the most deprived decile being
37 referred to SP. This could reflect different approaches being adopted with respect to equality and equity in
38 different countries, which is in line with our finding of differential associations of socio-demographic
39 factors with referral routes, contacts with link workers and prescribed interventions across countries.
40 Differences in service user compositions could also, to some extent, explain the country differences in the
41 distribution of referral reasons. For example, over 50% of SP cases are referred due to practical reasons in
42 Wales, and lifestyle is the most common referral reason in Northern Ireland (one of the least common
43 reasons in England and Wales). Intervention rates are notably higher in Scotland and Northern Ireland,
44 which could be related to reasons of referral and the availability of required services. Notably, England and
45 Wales have a much lower percentage of interventions that are costed or subsidised than Scotland and
46 Northern Ireland, providing further evidence that relying on free services due to a lack of funding could be
47 a major obstacle to an efficient and effective service.

18

19 However, with regards to limitations, the country differences should be interpreted with caution as we are
20 unable to assess the representativeness of the data. This is particularly concerning for Wales, Scotland and
21 Northern Ireland where the sample sizes are small. Although Access Elemental is the largest digital
22 platform for SP across the UK, we cannot rule out the possibilities that the country differences might be
23 driven by different clienteles of the Access Elemental in these countries compared to England. Indeed,
24 despite being used across a diverse range of places in the UK, Elemental has a low coverage in Scotland
25 (18%) and is not used in some areas e.g. Greater Glasgow and Clyde in Scotland, which have high levels of
26 deprivation, accounting for the much lower representation we had in our dataset of people from the
27 lowest two deciles of deprivation in Scotland. Given that all primary care data in the UK is non-
28 representative given the multiple different data platforms in use both amongst GPs and for more specialist
29 purposes like SP, this challenge is not easily solved, but further work cross-validating the findings
30 presented here is encouraged if we are to fully understand cross-country differences in SP. In these
31 analyses, although ethnicity is recorded in the data, the fact that it is 'optional' for data input means it is
32 missing for most people, so we were unable to examine any SP differences related to ethnicity.

33 Additionally, there is limited information on people's medical conditions which may influence the SP
34 services that they receive. Continued efforts are needed to improve data quality and accessibility for a
35 better understanding of SP implementation and its impacts. Our study has the strength of using data from
36 the most widely adopted SP platform in the UK to date, serving a large and diverse population with
37 detailed information about diverse referral routes. However, due to its nature, we were unable to compare
38 our sample to those who have never been considered for SP. Development of further large-scale rigorous
39 evaluations of SP as a service are needed.

40
41 Overall, this analysis shows that SP needs to be understood more by psychiatrists given the central role of
42 mental health. It has previously been argued that SP is aligned with the principles of modern psychiatric
43 care, emphasising what matters to the individual, prioritising belonging and community inclusion,
44 supporting emotion processing and regulation, and facilitating behavioural adaptations.³⁵ SP is also timely,
45 given existing challenges in psychiatry such as medication adherence, availability of traditional psychiatric
46 services, and shifts in thinking about the concept of recovery.³⁵ Psychiatrists are encouraged to consider
47 the suitability of social prescribing as an additional clinical service they can refer or signpost their patients
48 to alongside traditional psychiatric services. This can be carried out through existing Link Worker models
49 using Link Workers available within primary care. However, novel programmes embedding SP into
50 psychiatric services such as in mental health service waiting lists in the UK and as part of shared decision
51 making in psychiatric hospitals in Australia to support discharge plans are underway, and further such work
52 is also encouraged.³⁶ Nonetheless, although the medical pathway in SP is dominant across the UK, it has a
53 number of challenges associated, including that it does not manage to reach certain demographic groups
54 such as young people, men, and people from more deprived areas as effectively. Consequently, there is a
55 clear need for non-medical referral routes to be prioritised in the policy and delivery planning and funding
56 of future SP in the UK. Psychiatrists also have a role to play in supporting the development of non-clinical
57 referral pathways for mental health, such as peer support SP pathways,³⁷ to ensure that they are
58 developed in a way that is safe, feasible, and appropriate to the mental health needs of those who will be
59 using them.

50
51 Although SP is an individual-focused intervention that does not address the structural factors that may
52 have given rise to those inequalities in the first place, results from non-medical referrals suggest SP could
53 provide additional support to individuals experiencing inequalities.^{38,39} Additionally, despite a high number
54 of SP referrals taking place, only a third are leading to an intervention after link worker consultations, with
55 rates of interventions being particularly low on medical referral pathways and certain types of
56 interventions (such as those for physical health) being very limited. Challenges in funding for community
57 resources may be key here, with most financial resource for SP currently going to the pathway and
58 affiliated staff rather than community resources. Additional financial resource for certain intervention
59 domains for which there is not suitable free provision could help to readdress this balance. Third, given our
60 findings of heterogeneity in how SP is being delivered, for whom, across England and devolved nations,
61 specific strategies for the development of SP that meet the differing health and social needs in the
62 different countries are recommended moving forwards. Finally, in order to be able to continue monitoring
63 the development and distribution of SP in the UK, further investment into large-scale data platforms such
64 as Access Elemental and staff training in good quality data capture is vital.

65
66 **Declaration of interests**
67 We declare no competing interests.

68 **Funding**
69 This work was supported by the Prudence Trust INSPYRE grant [PT-0040] and the MRC [MR/Y01068X/1].
70 DF was supported by a British Academy Mid-Career Fellowship [MCFSS22\220006]. The funders had no
71 role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.
72

32 Acknowledgments

33 We thank Access Elemental and all data holders for granting us the data access.

34 Author contributors

35 DF and FB conceptualised the study and acquired the data. FB conducted the data analysis and designed
36 the data visualisation. All authors (FB, DH, AB and DF) contributed to the interpretation of results and
37 writing of the article, and had final responsibility for the decision to submit for publication. DF and FB
38 directly accessed and verified underlying data.

39 Data sharing

40 Due to the sensitive nature of the data, the research data can not be shared publicly. Data access can be
41 requested from the Access Elemental subject to ethical restrictions.

42

33 Reference

34

- 35 1 Hood CM, Gennuso KP, Swain GR, Catlin BB. County Health Rankings: Relationships Between Determinant
36 Factors and Health Outcomes. *Am J Prev Med* 2016; **50**: 129–135.
- 37 2 Braveman P, Egerter S, Williams DR. The Social Determinants of Health: Coming of Age.
38 <https://doi.org/10.1146/annurev-publhealth-031210-101218> 2011; **32**: 381–398.
- 39 3 Benach J, Friel S, Houweling T, Labonte R, Muntaner C, Schrecker T *et al*. A conceptual framework for action
40 on the social determinants of health. 2010.
- 41 4 Drinkwater C, Wildman J, Moffatt S. Social prescribing. *BMJ* 2019; **364**. doi:10.1136/BMJ.L1285.
- 42 5 Morse DF, Sandhu S, Mulligan K, Tierney S, Polley M, Giurca BC *et al*. Global developments in social
43 prescribing. *BMJ Glob Health* 2022; **7**: e008524.
- 44 6 Muhl C, Mulligan K, Bayoumi I, Ashcroft R, Godfrey C. Establishing internationally accepted conceptual and
45 operational definitions of social prescribing through expert consensus: a Delphi study. *BMJ Open* 2023; **13**:
46 e070184.
- 47 7 Bickerdike L, Booth A, Wilson PM, Farley K, Wright K. Social prescribing: less rhetoric and more reality. A
48 systematic review of the evidence. *BMJ Open* 2017; **7**: e013384.
- 49 8 Chatterjee HJ, Camic PM, Lockyer B, Thomson LJM. Non-clinical community interventions: a systematised
50 review of social prescribing schemes. <https://doi.org/10.1080/1753301520171334002> 2017; **10**: 97–123.
- 51 9 Howarth M, Brettle A, Hardman M, Maden M. What is the evidence for the impact of gardens and gardening
52 on health and well-being: a scoping review and evidence-based logic model to guide healthcare strategy
53 decision making on the use of gardening approaches as a social prescription. *BMJ Open* 2020; **10**: e036923.
- 54 10 Nguyen PY, Astell-Burt T, Rahimi-Ardabili H, Feng X. Effect of nature prescriptions on cardiometabolic and
55 mental health, and physical activity: a systematic review. *Lancet Planet Health* 2023; **7**: e313–e328.
- 56 11 Kimberlee R, Bertotti M, Dayson C, Asthana S, Polley M, Burns L *et al*. The economic impact of social
57 prescribing. London, 2022[https://socialprescribingacademy.org.uk/media/carfrp2e/evidence-review-](https://socialprescribingacademy.org.uk/media/carfrp2e/evidence-review-economic-impact.pdf)
58 [economic-impact.pdf](https://socialprescribingacademy.org.uk/media/carfrp2e/evidence-review-economic-impact.pdf) (accessed 18 Jul2023).
- 59 12 NHS England. NHS Long Term Workforce Plan. 2023[https://www.england.nhs.uk/long-read/accessible-nhs-](https://www.england.nhs.uk/long-read/accessible-nhs-long-term-workforce-plan/#:~:text=This%20Plan%20sets%20out%20modelling,over%20the%20next%20two%20years.)
60 [long-term-workforce-](https://www.england.nhs.uk/long-read/accessible-nhs-long-term-workforce-plan/#:~:text=This%20Plan%20sets%20out%20modelling,over%20the%20next%20two%20years.)
61 [plan/#:~:text=This%20Plan%20sets%20out%20modelling,over%20the%20next%20two%20years.](https://www.england.nhs.uk/long-read/accessible-nhs-long-term-workforce-plan/#:~:text=This%20Plan%20sets%20out%20modelling,over%20the%20next%20two%20years.) (accessed 18
62 Jul2023).

- 23 13 Elliott M, Davies M, Davies J, Wallace C. Exploring how and why social prescribing evaluations work: a realist
24 review. *BMJ Open* 2022; **12**: e057009.
- 25 14 Cartwright L, Burns L, Akinyemi O, Carder-Gilbert H, Tierney S, Elston J *et al.* Who is and isn't being referred to
26 social prescribing? London, 2022.
- 27 15 Mercer SW, Fitzpatrick B, Grant L, Chng NR, McConnachie A, Bakhshi A *et al.* Effectiveness of Community-
28 Links Practitioners in Areas of High Socioeconomic Deprivation. *The Annals of Family Medicine* 2019; **17**: 518-
29 525.
- 30 16 Kiely B, Croke A, O'Shea M, Boland F, O'Shea E, Connolly D *et al.* Effect of social prescribing link workers on
31 health outcomes and costs for adults in primary care and community settings: a systematic review. *BMJ Open*
32 2022; **12**: e062951.
- 33 17 Moscrop A. Social prescribing is no remedy for health inequalities. *BMJ* 2023; **381**. doi:10.1136/BMJ.P715.
- 34 18 NALW. URGENT: Act Now to Address Alarming Disparities in Social Prescribing Access!
35 2023.<https://www.nalw.org.uk/urgent-act-now-to-address-alarming-disparities-in-social-prescribing-access/>
36 (accessed 4 Mar2024).
- 37 19 Calderón-Larrañaga S, Milner Y, Clinch M, Greenhalgh T, Finer S. Tensions and opportunities in social
38 prescribing. Developing a framework to facilitate its implementation and evaluation in primary care: a realist
39 review. *BJGP Open* 2021; **5**: 1-13.
- 40 20 Jani A, Liyanage H, Okusi C, Sherlock J, De Lusignan S. Social prescribing observatory: A learning health system
41 approach for using data to improve practice. 2020.
- 42 21 Family Action. Social Prescribing in Secondary Care Pilot Service Evaluation Report. 2018[www.family-
43 action.org.uk/what-we-do](http://www.family-action.org.uk/what-we-do) (accessed 18 Jul2023).
- 44 22 Kimberlee R, Bertotti M, Dayson C, Elston J, Polley M, Burns L *et al.* (Sustainable) funding models for social
45 prescribing. 2022.
- 46 23 Fisher R, Fraser C. Who gets in? What does the 2020 GP patient survey tell us about access to general practice?
47 <https://www.health.org.uk/news-and-comment/charts-and-infographics/who-gets-in> (accessed 18 Jul2023).
- 48 24 Young Minds, The Children's Society. First port of call: The role of GPs in early support for young people's
49 mental health. London, 2021.
- 50 25 Khan K, Al-Izzi R, Montasem A, Gordon C, Brown H, Goldthorpe J. The feasibility of identifying health
51 inequalities in social prescribing referrals and declines using primary care patient records. *NIHR Open*
52 *Research* 2023 3:1 2023; **3**: 1.
- 53 26 Elemental Software. Social Prescribing and the Digital Landscape . 2018.
- 54 27 Access Elemental social prescribing software. [https://www.theaccessgroup.com/en-gb/our-
55 brands/elemental/](https://www.theaccessgroup.com/en-gb/our-brands/elemental/) (accessed 8 Aug2023).
- 56 28 Wolf A, Dedman D, Campbell J, Booth H, Lunn D, Chapman J *et al.* Data resource profile: Clinical Practice
57 Research Datalink (CPRD) Aurum. *Int J Epidemiol* 2019; **48**: 1740.
- 58 29 NHS England. The NHS Long Term Plan. 2019www.longtermplan.nhs.uk (accessed 7 Feb2024).
- 59 30 Mukhtar TK, Bankhead C, Stevens S, Perera R, Holt TA, Salisbury C *et al.* Factors associated with consultation
60 rates in general practice in England, 2013-2014: a cross-sectional study. *The British Journal of General*
61 *Practice* 2018; **68**: e370.
- 62 31 Wang Y, Hunt K, Nazareth I, Freemantle N, Petersen I. Do men consult less than women? An analysis of
63 routinely collected UK general practice data. *BMJ Open* 2013; **3**: e003320.

- 54 32 Skapinakis P, Lewis G, Araya R, Jones K, Williams G. Mental health inequalities in Wales, UK: Multi-level
55 investigation of the effect of area deprivation. *The British Journal of Psychiatry* 2005; **186**: 417–422.
- 56 33 Pearce JR, Richardson EA, Mitchell RJ, Shortt NK. Environmental justice and health: the implications of the
57 socio-spatial distribution of multiple environmental deprivation for health inequalities in the United Kingdom.
58 *Transactions of the Institute of British Geographers* 2010; **35**: 522–539.
- 59 34 Finn D. Jobcentres and the delivery of employment services and benefits. In: Millar J, Sainsbury R (eds).
70 *Understanding social security*. Policy Press, 2018, pp 217–234.
- 71 35 Zisman-Ilani Y, Hayes D, Fancourt D. Promoting Social Prescribing in Psychiatry—Using Shared Decision-Making
72 and Peer Support. *JAMA Psychiatry* 2023; **80**: 759.
- 73 36 Zisman-Ilani Y, Roe D, Elwyn G, Kupermintz H, Patya N, Peleg I *et al.* Shared Decision Making for Psychiatric
74 Rehabilitation Services Before Discharge from Psychiatric Hospitals. *Health Commun* 2019; **34**: 631–637.
- 75 37 Zisman-Ilani Y, Byrne L. Shared Decision Making and Peer Support: New Directions for Research and Practice.
76 <https://doi.org/10.1176/appi.ps20220407> 2022; **74**: 427–428.
- 77 38 Gibson K, Pollard TM, Moffatt S. Social prescribing and classed inequality: A journey of upward health
78 mobility? *Soc Sci Med* 2021; **280**: 114037.
- 79 39 Moscrop A. Social prescribing is no remedy for health inequalities. *BMJ* 2023; **381**. doi:10.1136/BMJ.P715.


30

Tables and Figures


Table 1. Summary of key differences across countries from regression analyses

	England	Wales	Scotland
Referral routes (Figure 2, Table S6)	<ul style="list-style-type: none"> • Males were less likely to be referred via medical routes. • People from older age groups (40+) had higher odds of being referred via medical routes (vs. aged 30-39). • Urban residents had 3.24 times higher odds of being referred via medical routes. • People from less deprived areas were more likely to be from medical routes. 	<ul style="list-style-type: none"> • Males were less likely to be referred via medical routes. • Children and young adults (0-29) were more likely to be referred via medical routes (vs. aged 30-39). • Urban residents had 1.63 times higher odds of being from medical routes. • People from less deprived areas were less likely to be from medical routes. 	<ul style="list-style-type: none"> • No difference by gender. • Children under 18 and older adults in their 60s were less likely to be referred via medical routes (vs. aged 30-39). • No difference by urbanicity. • People from less deprived areas were less likely to be from medical routes.
Contacts with link workers (Figure S7, Table S7)	<ul style="list-style-type: none"> • Males had fewer contacts. • Urban residents had fewer contacts. • Older adults aged 60+ had fewer contacts. • People from less deprived areas had more contacts 	<ul style="list-style-type: none"> • Males had fewer contacts. • Urban residents had fewer contacts. • Older adults aged 80+ had fewer contacts. • People from less deprived areas had more contacts, but no difference between the most and least deprived deciles 	<ul style="list-style-type: none"> • Males had more contacts. • No difference by urbanicity. • Older adults in their 60s and those aged 80+ had more contacts. • People from less deprived areas had more contacts, but no difference between the most and least deprived deciles
Interventions (Figure S8, Table S8)	<ul style="list-style-type: none"> • Those from medical routes were less likely to have an intervention. • People with employment related reasons had 2.52 times higher odds of having an intervention; whereas those with lifestyle or practical reasons had lower odds. • Male were less likely to have an intervention. • Children under 18 were less likely to have an intervention. People aged 40-69 were more likely to have an intervention. • Urban residents had 1.27 times higher odds of having an intervention. • People from less deprived areas were less likely to have an intervention. 	<ul style="list-style-type: none"> • Those from medical routes were more likely to have an intervention. • People with employment and practical reasons had higher odds of having an intervention, lower odds for lifestyle reasons. • No difference by gender. • Children under 18 were more likely to have an intervention. No difference between those aged 40-69 and 30-39. • Urban residents were less likely to have an intervention. • People from less deprived areas were more likely to have an intervention, except for the top three least deprived deciles. 	<ul style="list-style-type: none"> • Those from medical routes were less likely to have an intervention. • People with lifestyle reasons had 2.03 times higher odds of having an intervention, but no difference by employment or practical reasons. • No difference by gender. • Children under 18 were more likely to have an intervention. No difference between those aged 40-69 and 30-39. • Urban residents were less likely to have an intervention. • Limited difference by area deprivation.


(a) Gender



(b) Age groups



(c) Urbanicity



(d) Index of multiple deprivation deciles




Figure 1 Socio-demographic characteristics of unique individuals by country




Figure 2 Odds ratios and 95% confidence intervals from the logistic regression model on being referred via medical routes by country




Figure 3 Referral reasons (a) percentage of cases with referral reasons from each domain (b) percentages of cases with reasons from one or more different domain for each domain




Figure 4 Intervention domains (a) among cases with an intervention, percentage of cases belong to each domain (b) percentages of cases receiving an intervention matched to referral reason in each referral reason domain