1	Title
2	Sedation practices in patients intubated in the emergency department compared to the intensive
3	care unit.
4	
5	Short title
6	Sedation practices in the ED versus ICU
7	
8	Jariya Sereeyotin <sup>1,2</sup> , Christopher Yarnell <sup>2,3</sup> , Sangeeta Mehta <sup>2*</sup>
9	
10	<sup>1</sup> Department of Anesthesiology, Division of Critical Care Medicine, King Chulalongkorn
11	Memorial Hospital and Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand
12	<sup>2</sup> Department of Medicine, Sinai Health; Interdepartmental Division of Critical Care Medicine,
13	University of Toronto, Toronto, ON, Canada
14	<sup>3</sup> Department of Critical Care Medicine and Research Institute, Scarborough Health Network,
15	Toronto, ON, Canada
16	
17	*Corresponding author
18	E-mail: geeta.mehta@utoronto.ca (SM)
19	
20	Author contributions
21	JS- Conceptualization, Data curation, Formal analysis, Methodology, Investigation, Writing -
22	original draft, Writing – review & editing
23	CY- Formal analysis, Methodology, Writing – review & editing
24	SM- Conceptualization, Formal analysis, Methodology, Supervisor, Writing - review & editing

### 25 Abstract

26 Purpose

27 This study aimed to compare sedation management during and after intubation in the emergency

- 28 department (ED) versus the intensive care unit (ICU).
- 29 Methods

30 This was a single-center retrospective cohort study of adults intubated in the ED or in the ICU

and received mechanical ventilation between January 2018 and February 2022. We collected

32 data from the electronic medical record. The primary outcome was duration from intubation to

- 33 first documentation of light sedation, defined as a Sedation Agitation Scale score (SAS) of 3-4.
- 34 Results
- The study included 264 patients, with 95 (36%) intubated in the ED and 169 (64%) in the ICU.
- 36 Regarding anesthetic agents used for intubation, ketamine was the most commonly used drug in
- the ED and was used more frequently than in the ICU (61% vs 40%, p=0.001). Propofol was the
- 38 predominant sedative used in the ICU, with a higher prevalence compared to the ED (50% vs

39 33%, p=0.01). Additionally, benzodiazepines and fentanyl were more frequently used in the ICU

40 (39% vs 6%, p<0.001 and 68% vs 9.5%, p<0.001, respectively). Within 24 hours after

41 intubation, 68% (65/95) ED patients and 82% (138/169) patients intubated in ICU achieved light

42 sedation, with median durations of 13.5 hours and 10.5 hours. Patient location in the ED at

43 intubation was associated with decreased probability of achieving light sedation at 24 hours

44 (adjusted odds ratio 0.64, p=0.04).

45 Conclusion

46 Critically ill patients intubated in the ED are at risk of deeper sedation and a longer time to

47 achieve light sedation compared to patients intubated in the ICU.

## 48 Introduction

Effective sedation management is crucial for facilitating intubation, ensuring comfort, and promoting patient-ventilator synchrony in mechanically ventilated adults. However, this management poses challenges in both the emergency department (ED) and the intensive care unit (ICU) due to limited physiological reserves and the potential for serious complications in critically ill patients.[1, 2] Moreover, inappropriate or excessive sedation has been associated with adverse outcomes in critically ill patients.

55 Several studies have reported that deep sedation in mechanically ventilated patients 56 during the first 48 hours after ICU admission is associated with a higher risk of death and 57 delirium, as well as delayed time to extubation.[3-7] Despite the 2018 Pain, Agitation/Sedation, 58 Delirium, Immobility and Sleep Disruption (PADIS) guidelines[8] recommending light sedation 59 in critically ill mechanically ventilated adults, deep sedation is commonly used for intubated 60 patients in the ED. In a recent multicenter, prospective cohort study of 324 patients receiving 61 mechanical ventilation in the ED, 52.8% of intubated patients were deeply sedated [defined as Sedation Agitation Scale (SAS) score of 1 or 2], and deep sedation continued throughout the first 62 63 48 hours of ICU admission.[9] Therefore, an early emphasis on sedation minimization in 64 critically ill patients may be beneficial.

Many critically ill patients are initially intubated and managed in the ED, yet there are
few studies evaluating sedation practices for mechanically ventilated patients in the ED.[9, 10]
The overall purpose of our study was to compare sedation management during and after
intubation in patients intubated in the ED versus those intubated in the ICU. Given that our study
period included the COVID-19 pandemic, during which rapid sequence intubation (RSI) was
recommended for all patients with suspected or confirmed COVID-19 infection[11, 12], we also

71	explored the change in sedation management prior to and during the COVID-19 pandemic.
72	Moreover, we aimed to identify patient factors and other potentially modifiable factors
73	associated with deep sedation to improve sedation management in critically ill patients.
74	

## 75 Materials and Methods

76 This retrospective cohort study was conducted at a tertiary care university-affiliated 77 hospital and approved by the Sinai Health Research Ethics Board (REB#23-0002-C) on February 78 15<sup>th</sup>, 2023. Due to the retrospective research design and the fact that no identifying information 79 would be collected, the need for informed consent was waived. We identified eligible 80 consecutive patients from an ICU research screening database and electronic medical record 81 (EMR) prior to (January 1<sup>st</sup>, 2018-January 31<sup>st</sup>, 2020) and during the COVID-19 pandemic 82 (February 1st, 2020-February 28th, 2022). We accessed individual patient information between 83 February 15<sup>th</sup>, 2023 and August 31<sup>st</sup>, 2023; however, the identifying information was not 84 recorded during data collection. 85 The study included adult patients aged 18 years or older who were intubated either in the 86 ED or within the first 24 hours of ICU admission and received mechanical ventilation. Exclusion 87 criteria were: need for deep sedation or neuromuscular blocking agents after intubation (e.g., 88 therapeutic hypothermia after cardiac arrest, moderate to severe acute respiratory distress 89 syndrome, or status epilepticus); death within 48 hours of intubation; transferred to another 90 department after intubation; and not expected to achieve light sedation within 48 hours (e.g., 91 intracranial hemorrhage, brain stem infarction, or encephalopathy). 92 The primary outcome was the time from intubation to the first documentation of light 93 sedation, defined as SAS 3-4. Secondary outcomes included SAS at 6, 12, 24 and 48 hours after

94 intubation: intubation technique (RSI and non-RSI: RSI was defined as the administration of 95 sedative agents followed by rapid onset neuromuscular blocking agents without positive pressure 96 ventilation unless it was necessary); drug types and dosage for intubation and the postintubation 97 period; potential factors associated with deep sedation after intubation; and incidence of hemodynamic instability (defined as hypotension SBP < 90 mmHg or at least 20% decrease of 98 99 initial MAP for > 30 minutes; new requirement for or increase in dose of vasopressors; fluid 100 bolus > 15 ml/kg to maintain target blood pressure; or new onset of arrhythmia) within 30 101 minutes from the start of intubation.

102

#### 103 Statistical analysis

104 Demographic variables were presented by descriptive analysis, using frequency (percentage) for 105 categorical data. For continuous data, Kolmogorov-Smirnov was used for normality testing and 106 data expressed as mean (standard deviation, SD) or median (interguartile range, IOR) as 107 appropriate. The primary outcome was analyzed using a Cox proportional hazards model that 108 adjusted for relevant covariates. Covariates included age, Acute Physiologic Assessment and 109 Chronic Health Evaluation (APACHE) II score, obesity (body mass index  $\geq 30 \text{ kg/m}^2$ ), renal 110 insufficiency (defined as  $GFR < 30 \text{ mL/min}/1.73 \text{ m}^2$ ), end-stage liver disease, baseline Glasgow Coma Scale (GCS) and potential confounders for achieving light sedation including 111 112 administration of benzodiazepines and neuromuscular blocking agents after intubation. From the 113 primary analysis we reported the odds ratio of achieving light sedation according to patient 114 location at the time of intubation (ED versus ICU).

- 115 For the secondary analysis, we used logistic regression models to identify potential
- 116 predictors for the following binary outcomes: RSI versus non-RSI, deep sedation versus light

sedation, presence of hemodynamic instability versus none.

118 All analyses were performed using SPSS software version 28 and a two-sided P value <

119 0.05 was considered statistically significant.

120

## 121 **Results**

122 Between January 2018 and February 2022, a total of 314 patients were eligible for the inclusion 123 criteria, of whom 50 patients were excluded due to the reasons as shown in Fig 1. Ultimately, 124 264 patients were included in the primary analysis, with 95 (36%) intubated in the ED and 169 (64%) intubated in the ICU. The mean (SD) age was 60 (18) years and 153 (58%) were male. 125 126 The two groups were similar at baseline prior to intubation, other than a higher proportion of 127 patients with GCS  $\leq 8$  in the ED group (68% vs 25%), and higher mean age in the ICU group (63 128 vs 56 years). The main reason for intubation in the ED was neurological dysfunction while 129 respiratory failure was the most common reason for intubation in the ICU. Patient characteristics 130 are shown in Table 1. 131 132 Fig 1. Flow of participants in a study of the sedation practices in patients intubated in the

**133** emergency department compared to the intensive care unit. ED = Emergency Department,

134 ICU = Intensive Care Unit.

135

136 Table 1. Patient characteristics prior to endotracheal intubation

Variables	Emergency	Intensive Care	P value
	Department N=95	Unit N=169	
Age, years, mean (SD)	56 (21)	63 (15)	0.004
Male sex, n (%)	55 (58)	98 (58)	0.99
BMI (kg/m <sup>2</sup> ), median (IQR)	26 (21-28)	26 (23-31)	0.30
Race, n (%)*			
White	68 (72)	110 (65)	0.34
Black	9 (9)	10 (6)	
Other person of color	18 (19)	49 (29)	
APACHE II, median (IQR)	22 (17-27)	22 (17-27)	0.80
Baseline GCS before intubation			
$GCS \leq 8, n (\%)$	61 (68)	36 (25)	< 0.001
GCS > 8, n (%)	29 (32)	110 (75)	
COVID infection, n (%)			
Positive	6 (6)	6 (4)	0.30
Negative	89 (94)	163 (96)	
Underlying disease, n (%)			
Hypertension	34 (36)	72 (43)	0.28
Diabetes mellitus	26 (27)	40 (24)	0.51
COPD	9 (9)	20 (12)	0.56
Renal insufficiency	12 (13)	28 (17)	0.39
Hepatic dysfunction	10 (11)	27 (16)	0.22
End-stage liver disease	0 (0)	3 (2)	0.19

Reason for intubation, n (%)			
Airway obstruction	8 (8)	11 (7)	0.56
Respiratory failure	25 (26)	107 (63)	<0.001
Cardiogenic shock	4 (4)	4 (2)	0.40
Neurological dysfunction	46 (48)	22 (13)	< 0.001
Sepsis	3 (3)	24 (14)	0.004
Cardiac arrest	7 (7)	4 (2)	0.05
Other	2 (2)	0 (0)	0.06

137 \* Race was visually identified and recorded in the ICU research database by the research

139 BMI = Body Mass Index; APACHE II = Acute Physiologic Assessment and Chronic Health

140 Evaluation II score; GCS = Glasgow Coma Scale; COPD = Chronic Obstructive Pulmonary

141 Disease; SD = standard deviation; IQR = interquartile range

142

### 143 **Primary outcome and sedation agitation scale**

144 Within the first 24 hours after intubation, 68.4% (65/95) ED patients and 81.7% (138/169) ICU

145 patients achieved light sedation, with median durations of 13.5 hours and 10.5 hours,

146 respectively. Patient location in the ED at the time of intubation was associated with decreased

probability of achieving light sedation at 24 hours, with adjusted odds ratio of 0.64 (p=0.04; 95%

148 CI, 0.42 to 0.97) (S1 Table in S1 Appendix). Cumulative hazard curves assessing time from

intubation to the first documentation of achieving light sedation are shown in Fig 2.

<sup>138</sup> coordinator.

#### 151 Fig 2. Cumulative hazard of the time to achieve light sedation at 24 hours, by patients'

#### 152 location of intubation

- 153 Within the first 24 hours after intubation, the median time to achieve light sedation was 13.5
- hours for patients intubated in the ED and 10.5 hours for those intubated in the ICU. The mean
- odds ratio for achieving light sedation at 24 hours was 0.69 for the ED (p=0.01; 95% confidence
- 156 interval [CI], 0.51-0.92).

157

- 158 Deep sedation (SAS 1 or 2) was more frequently observed in the ED group at all time points,
- especially at 12 and 48 hours (60.2% vs. 46.3%, p =0.03; and 26.5% vs. 13.0%, p=0.02,
- 160 respectively) (Fig 3). More detail for the number of patients who were deeply sedated and lightly

sedated at each time point is reported in S2 Table in S1 Appendix.

162

#### 163 Fig 3. Sedation Agitation Scale at each time point in ED and ICU patients.

164 The Sedation Agitation Scale (SAS) at each time point is shown for patients intubated in the ED

and the ICU. Deep sedation (SAS 1 or 2) was more frequently observed in the ED group, and a

higher prevalence of light sedation (SAS 3 or 4) was found in the ICU group at all time points,

- with statistically significant differences at 12 and 48 hours (60.2% vs. 46.3%, p=0.03; and 26.5%
- 168 vs. 13.0%, p=0.02, respectively for deep sedation; 39.8% vs. 53.7%, p =0.03 and 73.5% vs.
- 169 87.0%, p=0.02, respectively for light sedation).

170

#### 171 Intubation technique, drug types and drug dosages

- 172 RSI was significantly more frequent in the ED compared to the ICU (82.1% vs. 30.2%, p
- 173 <0.001). During the COVID-19 pandemic, the incidence rate of RSI was significantly increasing

174	from 75% to 92.3% (p=0.03) in the ED and from 17.3% to 50.8% in the ICU	J (p<0.001).
-----	---	--------------

- 175 Intubation technique prior to and during the COVID-19 pandemic in the ED and ICU appear in
- 176 S3 Table and S1 Fig in S1 Appendix.
- 177
- 178 Regarding anesthetic agents used for intubation (Table 2), ketamine was the most commonly
- used drug in the ED and was used more frequently than in the ICU (61% vs 40% patients,
- 180 p=0.001). Propofol was the predominant sedative used in the ICU, with a higher prevalence
- 181 compared to the ED (50% vs 33%, p=0.01). Additionally, benzodiazepines, opiates and topical
- 182 xylocaine were more frequently used in the ICU (39% vs 6%, p 0.001; 68% vs 10%, p<0.001;
- 183 and 15% vs 3%, p=0.003; respectively).
- 184

#### 185 Table 2. Sedative, analgesic, and paralytic medications used for intubation and post-

186 intubation in the ED versus ICU

Intubation period	ED	ICU	P value
	N=95	N=169	
Induction drug administration			
Xylocaine topical	3 (3.2)	25 (14.8)	0.003
Fentanyl	9 (9.5)	115 (68)	< 0.001
- Dosage, mcg/kg	1.1 (0.6-1.4)	1.0 (0.6-1.4)	0.50
Ketamine	58 (61.1)	68 (40.2)	0.001
- Dosage, mg/kg	1.2 (0.8-1.4)	0.9 (0.5-1.2)	< 0.001
Propofol	31 (32.6)	85 (50.3)	0.01
- Dosage, mg/kg	1.2 (0.9-1.7)	0.7 (0.4-1.0)	< 0.001

Midazolam	6 (6.3)	66 (39.1)	< 0.001
- Dosage, mg/kg	0.07 (0.03-0.1)	0.02 (0.02-0.03)	0.02
Rocuronium	44 (46.3)	64 (37.9)	0.18
- Dosage, mg/kg	1.2 (1.0-1.4)	0.8 (0.7-1.1)	<0.001
Succinylcholine	36 (37.9)	0	< 0.001
- Dosage, mg/kg	1.4 (1.2-1.9)	0	< 0.001
Post-intubation drug administration	I	I	I
Opioids	24 (25.3)	116 (68.6)	< 0.001
Opioid dosage (fentanyl equivalent <sup>a</sup> )			
- Bolus, mcg/kg	0.9 (0.3-1.3)	0.5 (0.3-1.2)	0.37
- Infusion, mcg/kg/hr	0.8 (0.6-1.5)	0.9 (0.6-1.3)	0.76
Ketamine	16 (16.8)	8 (4.7)	0.001
Ketamine dosage			
- Bolus, mg/kg	0.2 (0.1-0.4)	0.7 (0.5-1.2)	0.09
- Infusion, mg/kg/hr	0.4 (0.3-0.5)	0.06 (0.05-0.15)	0.07
Propofol	48 (50.5)	110 (65.1)	0.02
Propofol dosage			
- Bolus, mg/kg	0.6 (0.3-1.1)	0.7 (0.4-0.8)	0.78
- Infusion, mcg/kg/min	30 (20-50)	30 (20-40)	0.046
Benzodiazepines	32 (33.7)	14 (8.3)	< 0.001
Benzodiazepine dosage (midazolam			
equivalents <sup>b</sup> )			
- Bolus, mg/kg	0.04 (0.03-0.09)	0.04 (0.02-0.05)	0.37

- Infusion, mg/kg/hr	0.05 (0.03-0.08)	0.04 (0.03-0.05)	0.44
Intermittent NMBA administration	4 (4.2)	9 (5.3)	0.69
Method of opioid administration, peri-			
intubation			
- Bolus	12 (12.6)	43 (25.4)	
- Infusion	11 (11.6)	30 (17.8)	< 0.001
- Both	3 (3.2)	77 (45.6)	
- None	69 (72.6%)	19 (11.2)	
Method of sedative administration, peri-			
intubation			
- Bolus	12 (12.6)	35 (20.7)	
- Infusion	3 (3.2)	9 (5.3)	0.15
- Both	78 (82.1)	117 (69.2)	
- None	2 (2.1)	8 (4.7)	

187 Data are shown as N (%) or median (IQR). ED = Emergency Department, ICU = Intensive Care

188 Unit, NMBA = neuromuscular blocking agent, mg/kg = milligram per kilogram, mcg/kg =

189 microgram per kilogram, mg/kg/hr = milligram per kilogram per hour, mcg/kg/min = microgram

190 per kilogram per minute, mcg/kg/hr = microgram per kilogram per hour

191 The most common benzodiazepine used in ED and ICU was midazolam.

<sup>a</sup> Fentanyl equivalent parenteral dose; fentanyl 0.1 mg = morphine 10 mg = hydromorphone 1.5

193 mg (Barr et al. Clinical Practice Guidelines for the Management of Pain, Agitation, and Delirium

in Adult Patients in the Intensive Care Unit. Critical Care Medicine. 2013:41(1); 263-306)

<sup>b</sup> Midazolam equivalent parenteral dose; midazolam 1.5 mg = diazepam 5 mg (Reuben Straye.
www.maimonidesem.org/benzodiazepines)

197

198 In terms of drug dosages, higher doses of sedatives and neuromuscular blocking agents were

used for intubation in the ED, while no significant difference in opioid dosage was found. We

200 identified 3 patients intubated in the ED who received neuromuscular blocking agents alone

201 without documented sedatives: two were diagnosed with drug intoxication and had baseline GCS

202 3 with no improvement after therapeutic naloxone administration, and the other had baseline

203 GCS 5 and was intubated due to COVID-19 infection and septic shock.

204

After intubation, opioids were less commonly used (25.3% vs. 68.6%, p <0.001), while ketamine</li>
and benzodiazepines were more frequently used (16.8% vs 4.7%, p=0.001; and 33.7% vs. 8.3%,
p <0.001, respectively) in the ED compared to the ICU. The bolus and infusion dosages were</li>
similar in ED and ICU except for a higher dose of propofol infusion in the ED [30 (20-50) vs 30
(20-40) mcg/kg/min, p=0.046].

210

211 Compared to prior to COVID-19 pandemic, rocuronium was used for intubation at higher doses

both in the ED and the ICU during the COVID-19 pandemic [0.9 (0.6-1.0) vs 1.3 (1.2-1.6)

213 mg/kg, p<0.001 for ED; and 0.7 (0.6-0.8) vs 1.1 (0.7-1.4) mg/kg, p<0.001 for ICU]. After

intubation, higher prevalence of using propofol and benzodiazepines as well as higher doses of

215 propofol infusion were observed in the ICU while higher prevalence of ketamine was observed

216 in the ED (S4.1 and S4.2 Tables in S1 Appendix).

Following intubation in the ICU, SAS scores were documented in the patient record every 4
hours. In the ED, GCS was recorded before transfer to the ICU, whereas no sedation assessment
tools were documented.

221

#### 222 Other outcomes

223 A multivariable logistic regression analysis was performed to identify the effects of APACHE II 224 score, baseline GCS, renal insufficiency, end-stage liver disease, use of midazolam for induction 225 and sedative administration methods (bolus, infusion, or both) on the likelihood that patients 226 would be deeply sedated 24 hours after intubation. In adjusted analysis, patients with renal 227 insufficiency (GFR  $< 30 \text{ mL/min}/1.73\text{m}^2$ ) were 2.78 times (p=0.03) more likely to be deeply 228 sedated at 24 hours than those without renal insufficiency. In addition, patients with baseline 229  $GCS \le 8$  were 2.17 times (p=0.03) more likely to be deeply sedated than those with baseline 230 GCS > 8 (S5 Table in S1 Appendix). Neurologic dysfunction as a reason for intubation and 231 being intubated during the COVID-19 pandemic were two independent factors strongly 232 associated with greater use of the RSI technique (OR 2.9, p<0.001; and OR 3.6, p<0.001, 233 respectively) (S6 Table in S1 Appendix).

234

Regarding outcomes after intubation, the incidence of cardiovascular instability requiring
vasopressor treatment was higher in the ICU group compared to the ED group (68.6% vs. 45.3%,
p<0.001) (Table 3). Increasing age, a diagnosis of sepsis and fentanyl use for induction were</li>
found to be associated with the risk of cardiovascular instability, while sedative agents used for
induction, including propofol, ketamine, and benzodiazepine, were not (S7 Table in S1
Appendix). Days of mechanical ventilation, hospital length of stay and 28-day mortality were

higher in the ICU group compared with the ED group (4 (2-9) vs 2 (1-3) days, p<0.001; 14 (8-</li>
47) vs 8 (3-16) days, p<0.001 and 37.7% vs. 12.3%, p<0.001, respectively). However, patients</li>
who were deeply sedated at any time point had longer days of mechanical ventilation, longer
hospital length of stay and fewer ventilator-free days at 28 days than the lightly sedated group
(Fig 4). In addition, deeply sedated patients were less likely to be discharged home and were
more commonly transferred to a rehabilitation facility than the lightly sedated group. More
details are provided in S8 Table in S1 Appendix

249 Table 3. Cardiovascular instability and ICU outcomes between the ED and ICU group

Outcomes	ED	ICU	P value
	N=95	N=169	
Cardiovascular instability, n (%)	52 (54.7)	130 (76.9)	< 0.001
Vasopressor requirement	43 (45.3)	116 (68.6)	< 0.001
Requirement for fluid bolus	36 (37.9)	66 (39.1)	0.85
Duration of mechanical ventilation, day, median (IQR)	2 (1-3)	4 (2-9)	< 0.001
Ventilator-free days at 28 days, median (IQR)	26 (24-27)	17 (0-25)	<0.001
Hospital length of stay, day, median (IQR)	8 (3-16)	14 (8-47)	< 0.001
28-day mortality, n/total collected data (%)	10/81 (12.3)	61/162 (37.7)	< 0.001
Discharge disposition, n (%)			
Home	52 (54.7)	30 (17.8)	
Death	10 (10.5)	77 (45.6)	< 0.001
Rehabilitation or long-term care institution	21 (22.1)	23 (13.6)	
Acute care hospital	12 (12.6)	39 (23.1)	

250	Data are shown as N (%) or median (IQR). ED = Emergency Department, ICU = Intensive Care
251	Unit.
252	Cardiovascular instability was defined as hypotension SBP < 90 mmHg or at least 20% decrease
253	of initial MAP for > 30 minutes; new requirement for or increase in dose of vasopressors; or
254	fluid bolus > 15 ml/kg to maintain target blood pressure; new onset of arrhythmia within 30
255	minutes from the start of intubation.
256	
257	Fig 4. Duration of mechanical ventilation, ventilator-free days at 28 days, and hospital
258	length of stay between deeply sedated and lightly sedated groups.
259	Deeply sedated is defined as SAS 1-2; Lightly sedated is defined as SAS $\geq$ 3., * statistical
260	significance
261	A. Duration of mechanical ventilation was longer in deeply sedated group compared to lightly
262	sedated group at any time point with statistical significance at 12, 24 and 48 hours after
263	intubation., B. Patients who were lightly sedated at 6, 24 and 48 hours after intubation had more
264	ventilator-free days at 28 days than those who were deeply sedated., C. At 6, 12 and 24 hours
265	after intubation, deeply sedated patients had significantly longer hospital length of stay than
266	lightly sedated patients.
267	
l 268	Discussion

In this single-center retrospective cohort study, patients who were intubated and had initiation of
invasive ventilation in the ED had a longer time to achieve light sedation than patients intubated
in the ICU. Moreover, deep sedation was more frequently observed in the ED group during the
first 48 hours of mechanical ventilation.

273

274	Our study emphasizes the impact of ED sedation practice for mechanically ventilated patients on
275	ICU outcomes, including fewer ventilator-, ICU- and hospital-free days.[9, 10] Our results are
276	consistent with the ED-SED study, which evaluated ED sedation practices in mechanically
277	ventilated patients.[9] They found that deep sedation was frequently started in the ED and
278	continued throughout the first 48 hours of ICU admission. More than 70% of deeply sedated
279	patients in their ICU received sedation management starting from the ED. Likewise, this study
280	also found a higher frequency of deep sedation in patients intubated in the ED at 6, 12, 24 and
281	until 48 hours after intubation compared to those intubated in the ICU.
282	
283	There is limited high-quality evidence to clarify the optimal dose of sedatives for intubation in
284	critically ill adults.[13-15] Higher doses of sedatives and neuromuscular blocking agents were
285	used for intubation in the ED despite the finding that the most common indication for intubation
286	in the ED was neurologic dysfunction (48%). Concerningly, 3 patients with low baseline GCS
287	(3-5) were intubated in the ED with the administration of neuromuscular blocking agents alone
288	without sedation, highlighting an opportunity for improvement.
289	
290	In a study of intubation practice in 3,659 critically ill patients across 29 countries, the main
291	reasons for intubation were respiratory failure (52%) and neurological dysfunction (31%),
292	similar to our study (50% respiratory failure, 26% neurological dysfunction). In their study the
293	most common drugs used for induction were propofol (41.5%), midazolam (36.4%), etomidate
294	(17.8%) and ketamine (14.2%). Propofol was significantly associated with cardiovascular

instability compared with etomidate (64% vs 50% respectively, absolute difference 14%; 95% CI

1.4-27.0%, p=0.02) when administered to patients with hemodynamic instability.[15] In our
study, ketamine was the most commonly used drug in the ED (61.1%), while propofol and
ketamine were the two most predominant sedatives used in the ICU (50.3% and 40.2%,
respectively). The use of specific sedatives for induction was not found to be a risk factor for
deep sedation at 24 hours nor for cardiovascular instability.

301

302 Regarding postintubation sedation, 33.7% of patients in the ED received benzodiazepines,

303 compared with 8.3% in the ICU. In addition, only 25.3% of patients in the ED group received

304 opioids for postintubation analgesia. These sedation practices were inconsistent with the 2018

305 PADIS Guidelines[8], which recommend an assessment-driven, protocol-based, stepwise

approach for pain and sedation management in critically ill adults, using analgesia first.

307 Moreover, benzodiazepines are not recommended for sedation.[16, 17] Another area for

improvement is that sedation assessment tools should be used in the ED to better target lightsedation.

310

Our logistic regression analysis identified that patients with renal insufficiency and patients with
low baseline GCS (≤8) were more likely to have deep sedation following intubation,[18-20]
suggesting that a cautious approach to sedative administration may be wise in these groups.
Furthermore, it was not surprising that the COVID-19 pandemic and neurologic dysfunction as
the reason for intubation were independent factors strongly associated with performing RSI
technique. The most likely reason for patients with neurological dysfunction was concern for
increased intracranial pressure during laryngoscopy.

318

319 Regarding outcomes after intubation, increasing age, sepsis diagnosis and pretreatment with 320 fentanyl were associated with cardiovascular instability in our study, which is consistent with 321 prior studies.[21-23] This could explain the higher incidence of cardiovascular instability in the 322 ICU group. Although this study has not shown an association between deep sedation and 323 mortality outcome as previous studies did, [5, 6, 10, 24] we found that fewer patients were 324 discharged home and more were discharged to a rehabilitation hospital in the early deep sedation 325 group. Moreover, fewer ventilator-free days and longer hospital length of stay were observed in 326 this group.

327

328 To our knowledge, this is the first study comparing sedation practices for intubation and post-329 intubation between the ED and the ICU. Strengths of this study include detailed data on drug 330 types and dosages, evaluation during and prior to the COVID-19 pandemic, complete patient 331 follow-up, and identification of predictors of deep sedation. This study has limitations. First, it 332 was a single-center study, so these data may not represent broader clinical practice. However, the 333 prevalence of deep sedation in the ED in our study was similar to previous studies. Second, the 334 duration from intubation to the first documentation of light sedation was analyzed for the 335 primary outcome, which mainly depended on the frequency of nurse assessments and might not 336 be the actual time for achieving light sedation. For this reason, we also collected the sedation 337 score at four specific time points. The data was in the same direction: a higher frequency of deep 338 sedation at every time point and a longer time required to achieve light sedation were found in 339 the ED group. Third, unmeasured confounders between the location of intubation and the use of 340 post-intubation sedation could partially explain our findings. Finally, due to the retrospective

- study design, the association between deep sedation and longer-term outcomes (e.g., 90-day
- 342 mortality, cognitive outcomes, etc.) could not be evaluated.

343

# 344 Conclusion

- 345 Patients intubated in the ED were more deeply sedated and took longer to achieve light sedation
- than patients intubated in the ICU.

# 348 Acknowledgements

- 349 We thank Sumesh Shah, research coordinator and Stanley Oei, respiratory therapist for sharing
- 350 their screening database.

351

### 353 **References**

354 1. Rinderknecht AS, Dyas JR, Kerrey BT, Geis GL, Ho MH, Mittiga MR. Studying the 355 Safety and Performance of Rapid Sequence Intubation: Data Collection Method Matters. Acad 356 Emerg Med. 2017;24:411-21. 357 2. Tayal VS, Riggs RW, Marx JA, Tomaszewski CA, Schneider RE. Rapid-sequence 358 intubation at an emergency medicine residency: success rate and adverse events during a two-359 year period. Acad Emerg Med. 1999;6:31-7. 360 Shehabi Y, Bellomo R, Reade MC, Bailey M, Bass F, Howe B, et al. Early intensive care 3. 361 sedation predicts long-term mortality in ventilated critically ill patients. Am J Respir Crit Care 362 Med. 2012;186:724-31. Shehabi Y, Bellomo R, Kadiman S, Ti LK, Howe B, Reade MC, et al. Sedation Intensity 363 4. 364 in the First 48 Hours of Mechanical Ventilation and 180-Day Mortality: A Multinational 365 Prospective Longitudinal Cohort Study. Crit Care Med. 2018;46:850-9. 366 5. Tanaka LM, Azevedo LC, Park M, Schettino G, Nassar AP, Réa-Neto A, et al. Early 367 sedation and clinical outcomes of mechanically ventilated patients: a prospective multicenter 368 cohort study. Crit Care. 2014;18:R156. 369 Balzer F, Weiß B, Kumpf O, Treskatsch S, Spies C, Wernecke KD, et al. Early deep 6. 370 sedation is associated with decreased in-hospital and two-year follow-up survival. Crit Care. 371 2015;19:197. 372 7. Stephens RJ, Dettmer MR, Roberts BW, Fowler SA, Fuller BM. Practice patterns and

372 7. Stephens RJ, Dettmer MR, Roberts BW, Fowler SA, Fuller BM. Practice patterns and
373 outcomes associated with early sedation depth in mechanically ventilated patients: a systematic
374 review protocol. BMJ Open. 2017;7:e016437.

375	8. Devlin JW, Skrobik Y, Gélinas C, Needham DM, Slooter AJC, Pandharipande PP, et al.
376	Clinical Practice Guidelines for the Prevention and Management of Pain, Agitation/Sedation,
377	Delirium, Immobility, and Sleep Disruption in Adult Patients in the ICU. Crit Care Med.
378	2018;46:e825-e73.
379	9. Fuller BM, Roberts BW, Mohr NM, Knight WAt, Adeoye O, Pappal RD, et al. The ED-
380	SED Study: A Multicenter, Prospective Cohort Study of Practice Patterns and Clinical Outcomes
381	Associated With Emergency Department SEDation for Mechanically Ventilated Patients. Crit
382	Care Med. 2019;47:1539-48.
383	10. Stephens RJ, Ablordeppey E, Drewry AM, Palmer C, Wessman BT, Mohr NM, et al.
384	Analgosedation Practices and the Impact of Sedation Depth on Clinical Outcomes Among
385	Patients Requiring Mechanical Ventilation in the ED: A Cohort Study. Chest. 2017;152:963-71.
386	11. Cook TM, El-Boghdadly K, McGuire B, McNarry AF, Patel A, Higgs A. Consensus
387	guidelines for managing the airway in patients with COVID-19: Guidelines from the Difficult
388	Airway Society, the Association of Anaesthetists the Intensive Care Society, the Faculty of
389	Intensive Care Medicine and the Royal College of Anaesthetists. Anaesthesia. 2020;75:785-99.
390	12. Orser BA. Recommendations for Endotracheal Intubation of COVID-19 Patients. Anesth
391	Analg. 2020;130:1109-10.
392	13. Higgs A, McGrath BA, Goddard C, Rangasami J, Suntharalingam G, Gale R, Cook TM.
393	Guidelines for the management of tracheal intubation in critically ill adults. Br J Anaesth.

394 2018;120:323-52.

395 14. Tarwade P, Smischney NJ. Endotracheal intubation sedation in the intensive care unit.
396 World J Crit Care Med. 2022;11:33-9.

397	15. Russotto V, Myatra SN, Laffey JG, Tassistro E, Antolini L, Bauer P, et al. Intubation
398	Practices and Adverse Peri-intubation Events in Critically Ill Patients From 29 Countries. JAMA
399	2021;325:1164-72.
400	16. Pandharipande PP, Pun BT, Herr DL, Maze M, Girard TD, Miller RR, et al. Effect of
401	Sedation With Dexmedetomidine vs Lorazepam on Acute Brain Dysfunction in Mechanically
402	Ventilated PatientsThe MENDS Randomized Controlled Trial. JAMA. 2007;298:2644-53.
403	17. Riker RR, Shehabi Y, Bokesch PM, Ceraso D, Wisemandle W, Koura F, et al.
404	Dexmedetomidine vs Midazolam for Sedation of Critically Ill Patients: A Randomized Trial.
405	JAMA. 2009;301:489-99.
406	18. Bauer TM, Ritz R, Haberthür C, Ha HR, Hunkeler W, Sleight AJ, et al. Prolonged
407	sedation due to accumulation of conjugated metabolites of midazolam. Lancet. 1995;346:145-7.
408	19. Motayagheni N, Phan S, Eshraghi C, Nozari A, Atala A. A Review of Anesthetic Effects
409	on Renal Function: Potential Organ Protection. Am J Nephrol. 2017;46:380-9.
410	20. Domi R, Huti G, Sula H, Baftiu N, Kaci M, Bodeci A, Pesha A. From Pre-Existing Renal
411	Failure to Perioperative Renal Protection: The Anesthesiologist's Dilemmas. Anesth Pain Med.
412	2016;6:e32386.
413	21. Smischney NJ, Kashyap R, Khanna AK, Brauer E, Morrow LE, Seisa MO, et al. Risk
414	factors for and prediction of post-intubation hypotension in critically ill adults: A multicenter
415	prospective cohort study. PLoS One. 2020;15:e0233852.

416 22. Perbet S, De Jong A, Delmas J, Futier E, Pereira B, Jaber S, Constantin JM. Incidence of

417 and risk factors for severe cardiovascular collapse after endotracheal intubation in the ICU: a

418 multicenter observational study. Crit Care. 2015;19:257.

- 419 23. Takahashi J, Goto T, Okamoto H, Hagiwara Y, Watase H, Shiga T, Hasegawa K.
- 420 Association of fentanyl use in rapid sequence intubation with post-intubation hypotension. Am J
- 421 Emerg Med. 2018;36:2044-9.
- 422 24. Stephens RJ, Dettmer MR, Roberts BW, Ablordeppey E, Fowler SA, Kollef MH, Fuller
- 423 BM. Practice Patterns and Outcomes Associated With Early Sedation Depth in Mechanically
- 424 Ventilated Patients: A Systematic Review and Meta-Analysis. Crit Care Med. 2018;46:471-9.

425

426

# 427 Supporting information

428 S1 Appendix Table E1-E8 and Fig E1

- 430
- 431
- 432
- 433



# Figure 1



Time to achieve light sedation (hours)

Figure 2

# % patients 6-hour 12-hour 24-hour 48-hour

# Figure 3







# Figure 4