

# **Applying Deep Learning in Heart Failure: Hospital Readmission is Not Like Other Health Quality Metrics**

## **Short Title: Predicting Readmission in Heart Failure**

**Authors:** Hailey M. Shepherd, MD<sup>1</sup>, Jeffrey T. Heaton, PhD<sup>2</sup>, Theodore Marghitu, BA<sup>3</sup>, Yun Z. Bai, MD<sup>1</sup>, Melanie P. Subramanian, MD, MPHS<sup>1</sup>, Sophia H. Roberts, MD<sup>1</sup>, Martha M.O. McGilvray, MD<sup>1</sup>, Amit A. Pawale, MD<sup>1</sup>, Gregory A. Ewald, MD<sup>4</sup>, Brian P. Cupps, PhD<sup>1</sup>, Michael K. Pasque, MD<sup>1</sup>, Randi E. Foraker, PhD<sup>5</sup>

<sup>1</sup>Division of Cardiothoracic Surgery, Department of Surgery, School of Medicine, Washington University in St. Louis, St. Louis, Missouri

<sup>2</sup>Sever Institute, McKelvey School of Engineering, Washington University in St. Louis, St. Louis, Missouri

<sup>3</sup>School of Medicine, Washington University in St. Louis, St. Louis, Missouri

<sup>4</sup>Division of Cardiovascular Medicine, Department of Medicine, School of Medicine, Washington University in St. Louis, St. Louis, Missouri

<sup>5</sup>Institute for Informatics, Division of General Medical Sciences, School of Medicine, Washington University in St. Louis, St. Louis, Missouri

### **Corresponding Author:**

Hailey M. Shepherd, MD  
Washington University School of Medicine  
Campus Box 8109, 660 South Euclid Avenue  
St. Louis, MO 63110  
Phone: (314) 482-0458  
Fax: (314) 367-8459  
Email: [hailey.shepherd@wustl.edu](mailto:hailey.shepherd@wustl.edu)

**Manuscript Word Count: 2,349**

## **Abstract**

### **Background**

Early identification of heart failure patients at increased risk for near-term adverse outcomes would assist clinicians in efficient resource allocation and improved care. Deep learning can improve identification of these patients.

### **Methods**

This retrospective study examined adult heart failure patients admitted to a tertiary care institution between January 2009 and December 2018. A deep learning model was constructed with a dense input layer, three long short-term memory (LSTM) layers, and a dense hidden layer to cohesively extract features from time-series and non-time-series EHR data. Primary outcomes were all-cause hospital readmission or death within 30 days after hospital discharge.

### **Results**

Among a final subset of 49,675 heart failure patients, we identified 171,563 hospital admissions described by 330 million EHR data points. There were 22,111 (13%) admissions followed by adverse 30-day outcomes, including 19,122 readmissions (87%) and mortality in 3,330 patients (15%). Our final deep learning model achieved an area under the receiver-operator characteristic curve (AUC) of 0.613 and precision-recall (PR) AUC of 0.38.

### **Conclusions**

This EHR-based deep learning model developed from a decade of heart failure care achieved marginal clinical accuracy in predicting very early hospital readmission or death despite previous accurate prediction of 1-year mortality in this large study cohort. These findings suggest that

factors unavailable in standard EHR data play pivotal roles in influencing early hospital readmission.

**Abbreviations:**

AUC = Area under the receiver operating characteristic curve

BMI = Body mass index

ECMO = Extracorporeal membrane oxygenation

EHR = Electronic health record

ICD = International Classification of Diseases

LOS = Length of stay

LSTM = Long short-term memory

PR AUC = Precision-recall area under the curve

ReLU = Rectified linear activation unit function

## **Clinical Perspective**

**What is new?** We developed an EHR-based deep learning model trained by 330 million data points from one of the largest cohorts of heart failure patients to date. Despite this model's highly accurate prediction of long-term outcomes, such as mortality and disease progression, our findings suggest that EHR data alone offers limited predictive power for predicting the short-term outcomes of 30-day hospital readmission or death.

**What are the clinical implications?** Our study supports the notion that hospital readmission, in contrast to other health outcomes, is uniquely driven by additional factors beyond traditional EHR variables. Once identified, incorporation of these determinants into future deep learning models could allow for accurate heart failure risk-stratification at hospital discharge to facilitate more efficient allocation of limited resources to the most vulnerable patients.

## **Introduction**

Heart failure remains a significant cause of morbidity and mortality in the United States with an annual mortality rate ranging from 22-30% (1-5). Heart failure patients represent the most rapidly growing and resource-intensive subset among patients with cardiovascular disease (6,7). Additionally, patients with heart failure have among the highest rates of unplanned 30-day hospital readmissions, ranging from 20-56% (5-7), many of which may be preventable (5-11). To this end, there has been increased emphasis on identifying heart failure patients who are at increased risk for adverse outcomes early after hospital discharge.

Deep learning models employ automated learning techniques to identify outcome-predictive data patterns in large datasets (12,13). Prior investigations using machine learning

models to predict hospital readmission in heart failure patients have reported inconsistent results (14-31). This may be attributed to difficulty in obtaining consistent, high-quality granular data for this heterogeneous patient group, or simply due to a lack of predictive signal in standard clinical metrics. Many other potential predictors, such as medication adherence and socioeconomic status, are not explicitly captured in EHR data but have been identified as critical determinants of care metrics including hospital readmission (32).

Although prior studies have not identified any single EHR variable capable of accurately predicting heart failure readmission with the precision required to drive clinical decision-making, predictive associations between groups of EHR variables and heart failure outcomes have been reported (17,24,32,33). As an alternative strategy to traditional regression analysis models, this investigation employs the pattern-recognition capabilities of deep learning to exploit the outcome-predictive capabilities associated with often complex and non-intuitive groupings of variables found in the medical records of heart failure patients.

EHR data platforms present patient-specific clinical variables in a uniquely structured format that may simplify and accelerate direct integration into machine learning models, facilitating application of their advanced analytic capabilities in risk prediction. These attributes give EHR-based deep learning models considerable potential for direct clinical utility in the identification of heart failure patients at high-risk for readmission or mortality. The accurate identification of these high-risk patients may improve distribution of available resources to the most vulnerable heart failure patients.

## **Methods**

## Data Source

The human studies *Institutional Review Board* at Washington University School of Medicine approved this study. All machine learning clinical variables analyzed in this study were obtained from the EHR of patients at Barnes-Jewish Hospital at Washington University Medical Center in St. Louis, Missouri.

## Cohort and Study Design

The study population consisted of adult patients between the ages of 18 to 90 years old with any heart failure International Classification of Diseases (ICD) codes admitted between January 2009 and December 2018. Admissions within 30 days before the conclusion of data collection or a length of stay (LOS) under 48 hours were excluded. Admissions with no discharge date or ending in death were excluded as *index* hospitalizations, although not as subsequent *readmissions*. Finally, at least 5 unstructured records (e.g., diagnosis, treatment, procedures, etc.), 1 vital sign record, and 1 laboratory value were required to include a recorded medical encounter as a hospitalization. These criteria helped exclude brief admissions scheduled for diagnostic purposes, such as cardiac catheterization and transesophageal echocardiography, which are often ordered on an outpatient basis but performed in the hospital due to their invasive nature.

The unit of analysis used to train our machine learning model was hospital admission, in contrast to other studies that have performed analysis at a patient (29) or unspecified (27) level. In patients with serial hospitalizations, each admission may qualify as a unique index hospitalization as well as a potential readmission if within 30 days after a prior discharge. By analyzing each instance as a unique opportunity for learning, data utilized for model training can be maximized. Moreover, this approach more accurately represents clinical practice and

enhances prediction applicability, since heart failure clinicians are routinely tasked with evaluating the risk of readmission at the end of each hospitalization.

Our primary positive outcomes were all-cause readmission lasting  $\geq 48$  hours or death within 30 days after discharge following an index hospitalization. These were classified as “positive events”. The 30-day window after hospital discharge has been identified as a key health policy metric for health care quality improvement (34,35). Readmissions within this time frame often reflect poor patient risk evaluation and inadequate recognition of post-discharge needs (36). Ventricular assist device placement, extracorporeal membrane oxygenation (ECMO) support, or cardiac transplantation are not considered separately since all are captured as hospitalization events. Model inputs included EHR data prior to index hospitalization discharge date, including all EHR data obtained during prior and current hospitalizations and in the outpatient setting.

### Feature Extraction and Deep Learning Network Design

The deep learning model in **Figure 1** is comprised of three LSTM layers to extract features from time-series EHR data, including vital signs, medication, laboratory, assessment, procedural, and diagnostic coding variables (37). Additionally, a dense layer extracts non-time-series patient attributes, such as age, gender, race; and a dense hidden layer which extracts additional features from the output features generated by the three LSTM input layers and the dense input layer. The intermediate features are extracted and pooled to a final hidden dense layer containing 16 neurons and a bias neuron. The model employs the rectified linear activation unit function (ReLU) for all dense hidden layers, and sigmoid activation for the final output neuron.



Most patients had available data for a subset of the 70 laboratory values evaluated (**Table 1**). These values are encoded by a 70-length vector which initially starts as the mean data set value for each laboratory test. As new results are obtained over time, the previous value in the vector is replaced and remains in place until new testing is performed. Therefore, the vector contains the most recent test results or the mean value for the dataset if no results are available for a specific patient. Vectors are entered as input sequences for the LSTM layers in a sliding window fashion with the oldest vectors sliding off once they reach the end of the sequence. Vital signs are similarly encoded using a 7-length vector whose values are updated as more recent vital signs are recorded in the EHR.

The coding variables for basic procedures, surgical procedures, diagnostics, assessments, and medications were transformed into 100-length sequences of 32-dimensional vectors using Python's Genism Word2Vec (38). These sequences were then entered into an LSTM layer. Since the data were high-cardinality and sparse, no missing data management was required. The encoding process only considered present values and ignored those that were not present in the time series.

The data were randomly divided into three sets: training (80%), validation (10%), and testing (10%). The model's performance was evaluated using AUC, PR AUC, log loss, and Brier score. Feature importance was determined using the permutation feature importance method, where each feature in the trained neural network was permuted and the change in AUC analyzed. Additionally, multiple features were permuted together to analyze grouped feature importance.

## **Results**

### Heart Failure Dataset Baseline Characteristics

Our initial patient cohort comprised 52,806 heart failure patients. After the previously described exclusions, our final study set consisted of 49,675 patients, of whom, 20% (9,975) experienced positive events during the 10-year study period. Of those who experienced positive events, 84% (8,419) experienced 30-day readmissions and 28% (2,751) suffered 30-day all-cause mortality. The baseline demographic characteristics and vital signs of our training and internal validations cohorts are listed in **Table 2**.

### Heart Failure Dataset 30-day Adverse Events

Of 171,563 study hospitalizations, 22,111 (13%) met our primary positive outcome (**Table 3**). Of these positive events, 19,122 (11%) involved 30-day all-cause readmission and 3,330 (2%) involved 30-day all-cause mortality. Some patients experienced both mortality and re-admission within 30 days (**Table 3**).

### Heart Failure Dataset Hospital LOS Data

The overall average LOS for the index hospitalizations was 122 hours. The average LOS of positive versus negative observations was 148 hours versus 133 hours, respectively.

### Heart Failure Dataset Variable Importance

**Table 1** shows the importance of the 70 variables evaluated for individual impact on 30-day all-cause readmission and mortality. Patient height, C-reactive protein level, and unstructured row count (which included variables such as diagnosis, treatment, procedures, etc.) demonstrated the highest importance at a univariate level. **Figure 2** demonstrates the relative predictive importance of vital sign and laboratory test variables depicted on the x and y axes,

respectively. Variables are listed in descending order of univariate importance. Red hues represent those with highest impact on model prediction (darker red hues representing factors with the most important impact) and blue hues highlighting low impact variables (dark blue hues representing the least important impact). Brain natriuretic peptide levels and serum sodium levels exhibited the highest impact among the clinical variables evaluated (vertical gradient), and there was no apparent association between clinical variable importance and vital sign values (horizontal gradient).

### Model Performance

The performance of the model for predicting 30-day readmission or death in heart failure patients was measured using the ROC curve with a test set ROC AUC of 0.613 (95% CI: 0.59-0.64) (**Figure 3**). **Table 4** shows additional performance metrics, including PR AUC of 0.38 (CI: 0.33-0.41) shown in **Figure 4**. The model's predictive performance remained stable across a range of hospitalization durations, including hospitalizations with a brief LOS (**Table 1**). The log loss was 0.57 with a Brier Score of 0.19

### **Discussion**

This deep learning model predicting 30-day all-cause hospital readmission or death was developed using a decade of EHR data from a tertiary care hospital and included 330 million data points from 171,563 hospital admissions in 49,675 heart failure patients. The model employed time-sensitive LSTM networks to analyze subtle fluctuations in clinical EHR variables over time and utilized densely connected networks for fixed, non-time-dependent demographics. Notably, we previously demonstrated that a nearly identical model successfully generated

accurate predictions of long-term adverse outcomes in the same cohort of heart failure patients, achieving an AUC of 0.91 (39). Our current findings indicate that EHR data alone is insufficient to generate similar predictive accuracy of 30-day hospital readmission or death in heart failure patients. This study underscores the unique nature of hospital readmission in comparison to other health quality metrics and the need to identify additional factors beyond traditional EHR variables that influence preventable hospital readmissions in this patient population.

Our study supports a growing body of literature heightening awareness of socioeconomic factors (such as income, education, place of residence, disability, ethnicity, race) as pivotal determinants contributing to hospital readmission in heart failure patients (40,41). While it is unlikely that targeting individual socioeconomic factors alone will be enough to reduce readmissions (41), the predictive model we seek likely requires cohesive integration of a cumulative burden of socioeconomic determinants in combination with key clinical variables that can be extracted from the EHR. This task is one for which machine learning is uniquely suited, given its unique proficiency at identifying non-intuitive, multiparametric outcome predictive patterns, and its seemingly boundless capability for comprehensive data analysis and integration (14,15,17,19,22-24). Additionally, incorporating time-based metrics through recurrent neural network units allows for the detection of even subtle progression in heart failure status, which can uncover time-dependent relationships between variables to enhance prediction and accelerate learning. These results suggest only modest capability of time-sensitive neural networks for predicting post-discharge adverse outcomes within the framework of each individual patient's unique heart failure course.

Prior research utilizing machine learning to predict hospital readmission in heart failure patients has yielded varying degrees of accuracy, with predictive AUC values ranging from 0.62

to 0.96 (14,15,17,19,22-24,26-29). Our final deep learning model was developed in one of the largest cohorts of heart failure patients to date and lends support to previous model results at the lower end of that spectrum of predictive accuracy. Although our model AUC does not attain the high level of accuracy achieved by Lv et al in a smaller cohort of 13,600 patients (17), our results are comparable to those reported by larger models and may accurately represent current potential of large scale implementation of patient-specific outcome prediction using an embedded EHR model.

By excluding hospitalizations with LOS  $\leq$  48 hours as index admissions or readmissions, our model attempts to exclude routine follow-up admissions, such as those related to elective, regularly scheduled endomyocardial biopsy and other surveillance investigations, many of which are unique to this complex patient subpopulation. This type of readmission may have contributed to overestimation of readmission for actual heart failure decompensation in previously published investigations (14,24,28). Instead, our model focuses upon longer hospitalizations, such as those for severe heart failure decompensation.

Several limitations must be considered when interpreting the findings of this study. Its retrospective design and the use of data from a single institution may restrict the generalizability of the model. Additionally, retrograde leakage of data that may falsely improve predictive accuracy is always a possibility (42-44). Our definition of a positive event as all-cause readmission or mortality may thereby include a small proportion of patients for whom 30-day readmission or death was not related to heart failure. Although not ideal, this primary outcome definition is the most effective and clinically applicable in our large population of medically complex heart failure patients. The inclusion of multiple hospitalizations from each patient in predictive model development may potentially skew results to the subpopulation with heightened

disease severity and worse overall health status. However, multiple readmissions are common in heart failure patients and their impact should be represented in predictive models. Clinical models assist but should not substitute clinical judgement. Additional validation is essential to verify the applicability of this model in other heart failure patient populations.

Our results suggest that EHR data alone offers limited predictive power for certain care quality metrics, such as early hospital readmission or death, and support the notion that other factors not routinely recorded in hospital medical records may play a pivotal role in determining these outcomes. To this end, further investigation is necessary to determine which of these non-standard metrics are most important in this patient population. Once identified, incorporation of these variables into future deep learning models could allow for accurate heart failure risk-stratification at hospital discharge to facilitate more efficient allocation of limited resources to the most vulnerable patients.

## References

1. Husaini BA, Mensah GA, Sawyer D et al. Race, sex, and age differences in heart failure-related hospitalizations in a southern state: implications for prevention. *Circ Heart Fail* 2011;4:161-9.
2. Johansen H, Strauss B, Arnold JM, Moe G, Liu P. On the rise: The current and projected future burden of congestive heart failure hospitalization in Canada. *Can J Cardiol* 2003;19:430-5.
3. McCullough PA, Philbin EF, Spertus JA et al. Confirmation of a heart failure epidemic: findings from the Resource Utilization Among Congestive Heart Failure (REACH) study. *J Am Coll Cardiol* 2002;39:60-9.
4. Healthcare Cost and Utilization Project. Agency for Healthcare Research and Quality, 2019.
5. Chen J, Normand SL, Wang Y, Krumholz HM. National and regional trends in heart failure hospitalization and mortality rates for Medicare beneficiaries, 1998-2008. *JAMA* 2011;306:1669-78.
6. Heidenreich PA, Albert NM, Allen LA et al. Forecasting the impact of heart failure in the United States: a policy statement from the American Heart Association. *Circ Heart Fail* 2013;6:606-19.
7. Virani SS, Alonso A, Aparicio HJ et al. Heart Disease and Stroke Statistics-2021 Update: A Report From the American Heart Association. *Circulation* 2021;143:e254-e743.
8. Yancy CW, Jessup M, Bozkurt B et al. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2013;128:1810-52.
9. Krumholz HM, Merrill AR, Schone EM et al. Patterns of hospital performance in acute myocardial infarction and heart failure 30-day mortality and readmission. *Circ Cardiovasc Qual Outcomes* 2009;2:407-13.

10. Khan MS, Sreenivasan J, Lateef N et al. Trends in 30- and 90-Day Readmission Rates for Heart Failure. *Circ Heart Fail* 2021;14:e008335.
11. Unplanned Hospital Visits - National. 7/27/22 ed: The Centers for Medicare & Medicaid Services, 2022.
12. Choi RY, Coyner AS, Kalpathy-Cramer J, Chiang MF, Campbell JP. Introduction to Machine Learning, Neural Networks, and Deep Learning. *Transl Vis Sci Technol* 2020;9:14.
13. Deo RC. Machine Learning in Medicine. *Circulation* 2015;132:1920-30.
14. Sharma V, Kulkarni V, McAlister F et al. Predicting 30-Day Readmissions in Patients With Heart Failure Using Administrative Data: A Machine Learning Approach. *J Card Fail* 2022;28:710-722.
15. Wang Z, Chen X, Tan X et al. Using Deep Learning to Identify High-Risk Patients with Heart Failure with Reduced Ejection Fraction. *J Health Econ Outcomes Res* 2021;8:6-13.
16. Sarijaloo F, Park J, Zhong X, Wokhlu A. Predicting 90 day acute heart failure readmission and death using machine learning-supported decision analysis. *Clin Cardiol* 2021;44:230-237.
17. Lv H, Yang X, Wang B et al. Machine Learning-Driven Models to Predict Prognostic Outcomes in Patients Hospitalized With Heart Failure Using Electronic Health Records: Retrospective Study. *J Med Internet Res* 2021;23:e24996.
18. Najafi-Vosough R, Faradmal J, Hosseini SK, Moghimbeigi A, Mahjub H. Predicting Hospital Readmission in Heart Failure Patients in Iran: A Comparison of Various Machine Learning Methods. *Healthc Inform Res* 2021;27:307-314.
19. Shameer K, Johnson KW, Yahi A et al. Predictive Modeling of Hospital Readmission Rates Using Electronic Medical Record-Wide Machine Learning: A Case-Study Using Mount Sinai Heart Failure Cohort. *Pac Symp Biocomput* 2017;22:276-287.
20. Ashfaq A, Sant'Anna A, Lingman M, Nowaczyk S. Readmission prediction using deep learning on electronic health records. *J Biomed Inform* 2019;97:103256.



21. Park J, Zhong X, Babaie Sarijaloo F, Wokhlu A. Tailored risk assessment of 90-day acute heart failure readmission or all-cause death to heart failure with preserved versus reduced ejection fraction. *Clin Cardiol* 2022;45:370-378.
22. Beecy AN, Gummalla M, Sholle E et al. Utilizing electronic health data and machine learning for the prediction of 30-day unplanned readmission or all-cause mortality in heart failure. *Cardiovasc Digit Health J* 2020;1:71-79.
23. Jiang W, Siddiqui S, Barnes S et al. Readmission Risk Trajectories for Patients With Heart Failure Using a Dynamic Prediction Approach: Retrospective Study. *JMIR Med Inform* 2019;7:e14756.
24. Golas SB, Shibahara T, Agboola S et al. A machine learning model to predict the risk of 30-day readmissions in patients with heart failure: a retrospective analysis of electronic medical records data. *BMC Med Inform Decis Mak* 2018;18:44.
25. Mahajan SM, Ghani R. Using Ensemble Machine Learning Methods for Predicting Risk of Readmission for Heart Failure. *Stud Health Technol Inform* 2019:243-247.
26. Mortazavi BJ, Downing NS, Bucholz EM et al. Analysis of Machine Learning Techniques for Heart Failure Readmissions. *Circ Cardiovasc Qual Outcomes* 2016;9:629-640.
27. Awan SE, Bennamoun M, Sohel F, Sanfilippo FM, Chow BJ, Dwivedi G. Feature selection and transformation by machine learning reduce variable numbers and improve prediction for heart failure readmission or death. *PLoS One* 2019;14:e0218760.
28. Allam A, Nagy M, Thoma G, Krauthammer M. Neural networks versus Logistic regression for 30 days all-cause readmission prediction. *Sci Rep* 2019;9:9277.
29. Frizzell JD, Liang L, Schulte PJ et al. Prediction of 30-Day All-Cause Readmissions in Patients Hospitalized for Heart Failure: Comparison of Machine Learning and Other Statistical Approaches. *JAMA Cardiol* 2017;2:204-209.

30. Turgeman L, May JH. A mixed-ensemble model for hospital readmission. *Artif Intell Med* 2016;72:72-82.
31. Xiao C, Ma T, Dieng AB, Blei DM, Wang F. Readmission prediction via deep contextual embedding of clinical concepts. *PLoS One* 2018;13:e0195024.
32. Segar MW, Hall JL, Jhund PS et al. Machine Learning-Based Models Incorporating Social Determinants of Health vs Traditional Models for Predicting In-Hospital Mortality in Patients With Heart Failure. *JAMA Cardiol* 2022;7:844-854.
33. Dharmarajan K, Hsieh AF, Lin Z et al. Diagnoses and timing of 30-day readmissions after hospitalization for heart failure, acute myocardial infarction, or pneumonia. *JAMA* 2013;309:355-63.
34. Vader JM, LaRue SJ, Stevens SR et al. Timing and Causes of Readmission After Acute Heart Failure Hospitalization-Insights From the Heart Failure Network Trials. *J Card Fail* 2016;22:875-883.
35. Leppin AL, Gionfriddo MR, Kessler M et al. Preventing 30-day hospital readmissions: a systematic review and meta-analysis of randomized trials. *JAMA Intern Med* 2014;174:1095-107.
36. McIlvennan CK, Eapen ZJ, Allen LA. Hospital readmissions reduction program. *Circulation* 2015;131:1796-803.
37. Mpanya D, Celik T, Klug E, Ntsinjana H. Machine learning and statistical methods for predicting mortality in heart failure. *Heart Fail Rev* 2021;26:545-552.
38. Choi E, Xiao C, Stewart W, Sun J. MiME: Multilevel medical embedding of electronic health records for predictive healthcare. *arXiv* 2018;1810.09593v1 [cx.LG].

39. McGilvray MMO, Heaton J, Guo A et al. Electronic Health Record-Based Deep Learning Prediction of Death or Severe Decompensation in Heart Failure Patients. *JACC Heart Fail* 2022;10:637-647.
40. Murray F, Allen M, Clark CM, Daly CJ, Jacobs DM. Socio-demographic and -economic factors associated with 30-day readmission for conditions targeted by the hospital readmissions reduction program: a population-based study. *BMC Public Health* 2021;21:1922.
41. Sterling MR, Ringel JB, Pinheiro LC et al. Social Determinants of Health and 30-Day Readmissions Among Adults Hospitalized for Heart Failure in the REGARDS Study. *Circ Heart Fail* 2022;15:e008409.
42. Hochreiter S, Schmidhuber J. Long short-term memory. *Neural Comput* 1997;9:1735-80.
43. Ma M, Liu C, Wei R, Liang B, Dai J. Predicting machine's performance record using the stacked long short-term memory (LSTM) neural networks. *J Appl Clin Med Phys* 2022;23:e13558.
44. Yu Y, Si X, Hu C, Zhang J. A Review of Recurrent Neural Networks: LSTM Cells and Network Architectures. *Neural Comput* 2019;31:1235-1270.

## Figure 1

A deep learning model was constructed with a dense input layer, three long short-term memory (LSTM) layers, and a dense hidden layer to cohesively extract features from time-series and non-time-series EHR data.

## Figure 2

Heat map demonstrating the relative predictive importance of vital sign and laboratory test variables depicted on the x and y axes, respectively. Variables are listed in descending order of univariate importance, with red hues representing those with highest impact on model prediction and blue hues highlighting low impact variables.

### **Figure 3**

The area under the receiver-operating characteristic curve (AUC) (or C-statistic) of 0.61 with a 95% CI of 0.59-0.64 demonstrates marginal clinical accuracy in short-term prediction of adverse 30-day outcomes in this large study cohort.

#### **Figure 4**

The precision recall area under the curve (PR AUC) of 0.37 with a 95% CI of 0.33-0.41 further supports the notion that standard EHR data lacks crucial factors, such as socioeconomic determinants, that lay pivotal roles in influencing early readmission and mortality in this cohort.

**Table 1**

Univariate importance of the 70 clinical variables evaluated for individual impact on 30-day all-cause readmission and mortality.

Variable	AUC Without Variable	Importance
<b>Vital Signs</b>		
Height	0.593	1.000
Weight	0.594	0.996
Heart rate	0.598	0.987
Diastolic blood pressure	0.598	0.987
Oxygen flow rate	0.599	0.984
Systolic blood pressure	0.603	0.974
Oxygen saturation	0.603	0.974
<b>Laboratory Values</b>		
C-reactive protein	0.503	1.000
Sodium	0.510	0.987
Lactate dehydrogenase	0.513	0.980
Brain natriuretic peptide	0.517	0.973
Hematocrit	0.521	0.966
Fasting glucose	0.521	0.965
Monocytes	0.524	0.958
Albumin	0.525	0.956
Fraction of inspired oxygen	0.526	0.954
Lactic acid	0.529	0.949
Uric acid	0.529	0.948
Plasma potassium	0.530	0.946



Ammonia	0.531	0.945
Cannabinoids	0.532	0.943
Hemoglobin	0.533	0.940
Amylase	0.534	0.939
Total hemoglobin	0.535	0.936
Venous partial pressure of oxygen	0.537	0.932
Triglycerides	0.541	0.925
Hemoglobin A1c	0.541	0.925
Chloride	0.541	0.925
Potassium	0.541	0.924
Carbon dioxide	0.545	0.916
Carboxyhemoglobin	0.546	0.915
Calculated oxygen saturation	0.546	0.914
Venous oxygen content	0.546	0.914
Oxygen gradient	0.547	0.912
Ketones	0.550	0.907
Alanine aminotransferase	0.550	0.906
Basophils	0.552	0.903
Base excess	0.552	0.901
Methemoglobin	0.553	0.901
Blood urea nitrogen	0.553	0.899
Troponin I	0.555	0.897
Benzodiazepines	0.556	0.894
Pulmonary artery oxygen	0.558	0.891
Partial thromboplastin time	0.558	0.891
Magnesium	0.558	0.890
Alkaline phosphatase	0.559	0.888

Glucose	0.560	0.887
Partial pressure of carbon dioxide	0.561	0.885
Pulmonary artery oxyhemoglobin	0.563	0.880
International normalized ratio	0.563	0.879
Cocaine	0.564	0.878
Creatinine	0.569	0.867
Direct bilirubin	0.572	0.862
High-density lipoprotein	0.574	0.858
Prothrombin time	0.575	0.856
Phosphorous	0.575	0.855
D dimer	0.584	0.839
Aspartate aminotransferase	0.586	0.833
Fraction of inspired oxygen	0.595	0.816
Plasma protein	0.595	0.815
Anion gap	0.596	0.813
Calcium	0.598	0.809
pH	0.599	0.807
Eosinophils	0.599	0.807
Fibrinogen	0.600	0.806
Urine specific gravity	0.600	0.806
Lymphocytes	0.600	0.806
Indirect Coombs	0.600	0.806
Lipase	0.600	0.805
Bicarbonate	0.602	0.802
Creatine kinase	0.602	0.802
Platelets	0.602	0.802
Bilirubin	0.603	0.800

Partial pressure of oxygen	0.603	0.800
Thyroid stimulating hormone	0.603	0.799
Neutrophils	0.603	0.799
Cholesterol	0.604	0.798
Oxygen saturation	0.604	0.798
<b>Static Variables</b>		
Unstructured row count	0.540	1.000
Laboratory test row count	0.567	0.940
Length of stay (LOS) hours	0.591	0.889
Vital sign row count	0.597	0.875
Gender	0.602	0.865
Race	0.604	0.859
Age at admission	0.606	0.857

**Table 2**

Baseline characteristics in training and internal validation cohorts.

	<b>Training Cohort</b>	<b>Internal Validation Cohort</b>
Number of participants	19,884	2,486
Age at admission, years	63 (SD:15)	63 (SD:15)
Gender		
Female, %	9,052 (46)	1,095 (44)
Male, %	10,832 (55)	1,390 (46)
Unknown, %	0 (0)	1 (0)
Ethnicity		
White, %	9,036 (45)	1,093 (44)
Black, %	9,313 (47)	1,202 (48)
Asian, %	87 (0)	7 (0)
Other, %	1,448 (7)	184 (7)
LOS, hours	268 (SD:718)	290 (SD:832)
Systolic blood pressure, mmHg	125 (SD:15)	125 (SD:14)
Diastolic blood pressure, mmHg	70 (SD:9)	71 (SD:9)
Heart rate, bpm	82 (SD:8)	82 (SD:8)
O2 saturation, %	97 (SD:1)	97 (SD:1)

**Table 3**

Comparison of positive and negative observations.

	<b>Negative Observation</b>	<b>Positive Observation</b>
Number of participants	49,675	9,975
Age at admission, years	62 (SD:13)	62 (SD:15)
Gender		
Female, %	22,956 (46)	4,488 (45)
Male, %	26,717 (54)	5,487 (55)
Unknown, %	2 (0)	0 (0)
Ethnicity		
White, %	19,373 (39)	3,591 (36)
Black, %	27,321 (55)	5,386 (54)
Asian, %	496 (1)	1 (0)
Other, %	2,485 (5)	997 (10)
LOS, hours	98 (SD:453)	128 (SD:439)
Systolic blood pressure, mmHg	127 (SD:20)	125 (SD:21)
Diastolic blood pressure, mmHg	71 (SD:13)	71 (SD:13)
Heart rate, bpm	81 (SD:13)	82 (SD:13)
O2 saturation, %	97 (SD:2)	97 (SD:2)

**Table 4**

Predictive model performance results.

<b>Measure</b>	<b>Result</b>
ROC AUC	0.613
Log Loss	0.570
PR AUC	0.380
Brier Score	0.190



Electronic Health Records  
171,563 hospitalizations  
52,806 heart failure patients  
350 million datapoints

### Time Series Data

(Procedural, medication, assessment, diagnostic coding variables, vital signs, laboratory tests)

### Patient Data

(Age, gender, race, length of stay)

#### A1: Input Layer

Unstructured Data  
32 vector x 100  
sequences

#### LSTM1

Output: 256  
vector

#### B1: Input Layer

Laboratory Data  
70 vector x 100  
sequences

#### LSTM2

Output: 256  
vector

#### C1: Input Layer

Vital Signs  
7 vector x 100  
sequences

#### LSTM3

Output: 256  
vector

#### D1: Input Layer


Static Values  
7 vector

#### Dense 1

Static Values  
Output 16

### Dense 2: Model Output


Input: 256+256+256+16  
Output: 1



	SYSTOLIC	WDRGT	PULSE	DIASTOLIC	HEIGHT	O2 SAT	O2 FLOW
Brain natriuretic peptide	0.967	0.960	0.965	0.972	0.958	0.958	1.000
Sodium	0.863	0.863	0.844	0.843	0.833	0.858	0.895
Venous partial pressure of oxygen	0.862	0.860	0.862	0.841	0.836	0.868	0.836
Lactate dehydrogenase	0.830	0.818	0.804	0.801	0.816	0.831	0.820
Hematocrit	0.767	0.761	0.766	0.781	0.798	0.741	0.787
Albumin	0.782	0.797	0.777	0.777	0.788	0.772	0.783
C-reactive protein	0.767	0.770	0.754	0.781	0.781	0.738	0.768
Chloride	0.765	0.778	0.780	0.780	0.774	0.758	0.767
Fraction of inspired oxygen	0.733	0.747	0.738	0.727	0.748	0.752	0.748
Hemoglobin	0.756	0.747	0.787	0.751	0.746	0.727	0.739
Carnitine	0.707	0.680	0.701	0.722	0.725	0.695	0.719
Calculated oxygen saturation	0.728	0.710	0.745	0.738	0.745	0.740	0.713
Total hemoglobin	0.735	0.722	0.705	0.735	0.732	0.728	0.713
Amylase	0.680	0.686	0.684	0.691	0.694	0.702	0.701
International normalized ratio	0.691	0.726	0.689	0.698	0.705	0.694	0.695
Potassium	0.694	0.700	0.709	0.709	0.699	0.681	0.689
Benzodiazepines	0.673	0.664	0.682	0.682	0.697	0.677	0.687
Oxygen gradient	0.680	0.678	0.683	0.683	0.685	0.690	0.681
Venous oxygen content	0.664	0.666	0.679	0.666	0.678	0.678	0.678
Monocytes	0.681	0.687	0.682	0.689	0.705	0.671	0.678
Uric acid	0.674	0.699	0.701	0.670	0.692	0.684	0.678
Plasma creatinum	0.668	0.676	0.686	0.680	0.693	0.698	0.676
Lactic acid	0.669	0.664	0.680	0.683	0.677	0.670	0.667
Methemoglobin	0.664	0.648	0.660	0.665	0.647	0.657	0.664
Carboxyhemoglobin	0.664	0.679	0.673	0.679	0.674	0.675	0.660
Partial thromboplastin time	0.665	0.659	0.655	0.668	0.664	0.659	0.659
Eosinophils	0.668	0.675	0.662	0.664	0.661	0.659	0.651
Pulmonary artery oxyhemoglobin	0.636	0.656	0.666	0.655	0.655	0.659	0.649
Ammonia	0.664	0.660	0.663	0.663	0.667	0.654	0.636
Partial pressure of carbon dioxide	0.643	0.621	0.645	0.622	0.632	0.630	0.636
Glucose	0.623	0.608	0.629	0.625	0.616	0.614	0.636
Carbon dioxide	0.639	0.630	0.647	0.631	0.632	0.639	0.635
Triglycerides	0.628	0.635	0.646	0.642	0.628	0.641	0.635
Prothrombin time	0.641	0.644	0.615	0.625	0.625	0.648	0.635
Direct bilirubin	0.624	0.627	0.625	0.617	0.627	0.627	0.634
Troponin I	0.635	0.633	0.642	0.629	0.648	0.635	0.633
Fasting glucose	0.628	0.617	0.623	0.630	0.633	0.635	0.631
Hemoglobin A1c	0.631	0.631	0.626	0.624	0.648	0.635	0.629
Ketones	0.634	0.643	0.635	0.643	0.647	0.611	0.628
Bone mass	0.665	0.662	0.666	0.649	0.648	0.634	0.625
Blood urea nitrogen	0.605	0.606	0.625	0.600	0.603	0.603	0.618
Alkaline phosphatase	0.620	0.611	0.634	0.631	0.648	0.625	0.617
Pulmonary artery oxygen	0.591	0.597	0.586	0.600	0.603	0.593	0.606
Alanine aminotransferase	0.587	0.601	0.584	0.589	0.592	0.590	0.601
Cocaine	0.605	0.599	0.605	0.601	0.592	0.605	0.601
High-density lipoprotein	0.575	0.582	0.583	0.598	0.570	0.582	0.587
Magnesium	0.598	0.584	0.598	0.591	0.601	0.598	0.587
Aspartate aminotransferase	0.547	0.576	0.578	0.583	0.598	0.561	0.567
Phosphorus	0.563	0.562	0.561	0.565	0.568	0.545	0.566
Creatinine	0.543	0.535	0.538	0.544	0.540	0.529	0.544
Plasma protein	0.513	0.528	0.524	0.528	0.524	0.520	0.525
D dimer	0.504	0.511	0.512	0.514	0.514	0.511	0.514
Fraction of inspired oxygen	0.509	0.506	0.501	0.514	0.509	0.509	0.511
pH	0.502	0.505	0.505	0.507	0.505	0.507	0.505
Calcium	0.507	0.507	0.507	0.508	0.508	0.506	0.506
Thyroid stimulating hormone	0.501	0.503	0.503	0.508	0.503	0.503	0.504
Induct. Coombs	0.502	0.502	0.505	0.508	0.503	0.503	0.504
Cholesterol	0.502	0.503	0.502	0.503	0.503	0.503	0.504
Creatine kinase	0.501	0.503	0.503	0.504	0.504	0.503	0.503
Urine specific gravity	0.500	0.501	0.501	0.504	0.502	0.503	0.503
Bicarbonate	0.501	0.503	0.503	0.508	0.502	0.503	0.503
Platelets	0.501	0.502	0.503	0.503	0.503	0.503	0.503
Lipase	0.501	0.502	0.502	0.504	0.502	0.503	0.502
Anion gap	0.502	0.504	0.504	0.503	0.506	0.496	0.502
Partial pressure of oxygen	0.502	0.504	0.504	0.505	0.502	0.503	0.502
Bilirubin	0.501	0.501	0.503	0.504	0.503	0.500	0.502
Eosinophils	0.498	0.503	0.500	0.502	0.499	0.499	0.502
Neutrophils	0.500	0.501	0.502	0.504	0.500	0.500	0.502
Fibrinogen	0.500	0.503	0.501	0.505	0.502	0.503	0.500
Lymphocytes	0.500	0.500	0.501	0.503	0.502	0.500	0.500



# Receiver Operating Characteristic Curve



Precision Recall Curve

