

Post-stroke Epilepsy and its Association with Stroke Location – Systematic review and Meta-analysis

Sihyeong Park^{1, 2*}, Yasar T. Esengul¹, Khaled Gharaibeh¹, Sidra Saleem¹, Ajaz Sheikh¹

¹University of Toledo, Department of Neurology

² Mayo Clinic, Department of Neurology, Division of Epilepsy

*Correspondence to: Sihyeong Park

Sihyeong.park90@gmail.com

200 1st St SW, Rochester, MN 55901

Abstract

Objective

There have been several meta-analytic studies that investigated potential risk factors for post-stroke epilepsy (PSE)^{1,2}. Cortical location is known to be associated with increased risk of PSE. However, relationship between the involvement of specific lobe with stroke and PSE has not been evaluated in a meta-analysis due to unavailability of relevant data. To the best of our knowledge, this is the first meta-analysis on this subject matter.

Methods

The PubMed and Embase databases were comprehensively assessed and searched for studies that evaluated the association between the involvement of different lobes with PSE. Studies that evaluated post-stroke epilepsy or recurrent post-stroke seizure were included, and acute symptomatic seizure was excluded. Following search strategy was used, “post-stroke” OR “stroke” OR “intracerebral hemorrhage” AND “epilepsy” OR “seizure” AND “lobe” OR “territory” OR “location”. Joanna Briggs Institute appraisal tool was utilized to perform quality assessment of the included studies. A random effects model was used to estimate pooled odds ratio of the selected outcomes with 95 percent confidence intervals (CIs), and the study heterogeneity was analyzed.

Results

Nine studies³⁻¹¹ were identified with 1328 stroke patients. 9/9 studies showed association of frontal lobe involvement with higher likelihood of PSE (OR=2.10, 95% CI=1.37-3.21, p-value=0.0007). All studies showed association of parietal stroke with increased likelihood of PSE (OR=2.85, 95% CI=1.93-4.22, p-value<0.00001). Previous temporal lobe stroke was associated with increased risk of PSE (OR=3.55, 95% CI=1.95-6.45, p-value<0.00001). In

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

contrast, association between occipital lobe involvement and PSE was not statistically significant (OR=1.37, 95% CI=0.95-1.96, p-value<0.09), and this remained true when subgroup analysis was performed.

Discussion

Our meta-analysis shows that PSE is associated with frontal, parietal, and temporal involvement of previous stroke. There was no statistically significant association between occipital lobe involvement and PSE. Further high quality studies are needed to investigate if stroke in one lobe/territory can be more epileptogenic than those in others.

Keywords post-stroke epilepsy, stroke, epilepsy, meta-analysis

Key Points

- Cerebrovascular disease is the most common etiology of newly diagnosed epilepsy
- Prior stroke involving the frontal, temporal, or parietal lobe is associated with increased risk of post-stroke epilepsy
- Occipital lobe stroke is not associated with increased risk of post-stroke epilepsy, regardless of the age group.

1. Introduction

An epidemiological study that included all age group reported that cerebrovascular disease was the most common etiology in patients with defined cause of newly diagnosed epilepsy¹². With the ageing population, the importance of new-onset epilepsy in the elderly has become more significant. A number of previous literature reported cerebrovascular disease as the most common etiology of the late-onset epilepsy. A large epidemiological study reported stroke incidence of 13.7 million and stroke prevalence of 80.1 million in 2016¹³.

There have been a few different definitions of post-stroke epilepsy (PSE) over the years¹⁴. Previous literature has steadily defined late-onset seizures occurring after two weeks or more from stroke, which increases the risk of PSE. The incidence of PSE was reported to be 7% in a meta-analysis that included 102008 stroke patients¹⁵. There have been a few meta-analytic studies that reviewed the risk factors for PSE^{1,2}. Cortical location is known to be associated with increased risk of PSE. However, relationship between the involvement of specific lobe with stroke and PSE has not been evaluated in a meta-analysis due to unavailability of relevant data¹, although there has been some investigations that demonstrated association of parietal lobe infarction with the increased risk of PSE⁸. To the best of our knowledge, this is the first meta-analysis on this subject matter.

2. Methods

2.1. Search strategy and selection criteria

The PubMed and Embase databases were comprehensively assessed and searched from their inception to July 2023. Studies that evaluated the association between the involvement of different lobes with PSE were identified. Following search strategy was used, “post-stroke” OR “stroke” OR “intracerebral hemorrhage” AND “epilepsy” OR “seizure” AND “lobe” OR “territory” OR “location”. The full search strategy is outlined in the Supplementary Material (Supp 1). Results were reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Two authors (S.P. and Y.T.E.) independently performed the search, screened titles and abstracts, and assessed eligibility. For included studies, S.P. collected the data from each study. Following data were extracted: first author name, year of publication, study design, total sample size, number of patients with a history of infarct with involvement of a given lobe and infarct sparing that lobe, and number of patients with PSE in each group with lobar involvement. The number of the patients were directly obtained or derived from the available information in the manuscripts of the included studies. With a study that specified concurrent involvement of different lobes (Kopyta 2021), e.g. frontal only vs frontal, temporal, and parietal, the cases with frontal lobe involvement were added, and the cases with no involvement of frontal lobe were added to evaluate for the association of frontal lobe infarcts and poststroke epilepsy. Same strategy was utilized to derive the number of the cases with infarcts affecting the other lobes.

Inclusion criteria were as follows: (1) studies that evaluated post-stroke epilepsy or recurrent post-stroke seizure and (2) original articles. Following studies were excluded: (1) studies that investigated acute symptomatic seizure, (2) studies that did not clarify the chronicity of the

seizure onset after stroke, (3) studies with insufficient information regarding anatomic location of the stroke, (4) irrelevant studies, (5) animal or in vitro studies, (6) case reports, conference abstracts, or reviews, and (7) studies in language other than English.

2.2 Quality assessment

Joanna Briggs Institute (JBI) appraisal tool was utilized to perform quality assessment of the included studies.

2.3. Statistical analysis

Given the variability in study design and the patient population in the studies, random effects model was assumed and Mantel-Haenszel method was utilized to estimate the pooled odds ratio of the selected outcomes with 95 percent confidence intervals (CIs). Study heterogeneity was analyzed with I^2 statistic. For the I^2 statistic, 25% to 50%, 50% to 75%, and 75% were considered low, moderate, and high heterogeneity, respectively. Data analysis was conducted using Review Manager 5.3 and R 4.1.1.

3. Results

3.1. Study selection

Our search revealed 1952 studies through PubMed, and 2664 studies were identified through EMBASE. Of those studies, 852 articles were duplicated and removed from the screening process. After excluding the articles as outlined in Methods, 9 studies were included in the qualitative and quantitative analysis (Figure 1).

3.2. Study characteristics

Table 1 illustrates the characteristics of these 9 studies³⁻¹¹, which included a total of 1328 patients with history of stroke. Four of the studies were performed with pediatric population, and the other five included adult population. All the studies were retrospective and observational. The mean age was 56 years, and males represented 57% of total patients. Among 1328 included patients, 255 had PSE (19%). Quality assessment of the included studies with JBI appraisal tool revealed average score of 9.6, with range of 7-11 (Table 2).

3.3. Location of stroke and risk of PSE

All nine studies described the association of frontal lobe stroke and PSE. Prior infarcts with frontal lobe involvement was associated with increased risk of PSE (OR=2.10, 95% CI=1.37-3.21, p-value=0.0007) (Figure 2). I^2 value was 35%. Subgroup analysis was performed based on the age group. Analysis of the adult subgroup continued to show statistically significant associated between frontal lobe infarct and PSE (OR=2.11, 95% CI=1.43-3.10, p-value=0.0002; I^2 =16%). On contrary, there was no significant association of such in the pediatric group, with higher heterogeneity (OR=2.04, 95% CI=0.51-8.15, p-value=0.31; I^2 = 61%).

All included studies showed association of parietal stroke with increased likelihood of PSE

(OR=2.85, 95% CI=1.93-4.22, p-value<0.00001). I^2 value was 24%. Subgroup analysis was performed based on the age group. Risk of PSE was higher in both adult and pediatric subgroups (OR=2.50, 95% CI=1.68-3.70, p-value<0.00001; I^2 =15% and OR=4.15, 95% CI=1.71-10.08, p-value<0.002; I^2 =26%, respectively) (Figure 3).

Previous temporal lobe stroke was associated with increased risk of PSE (OR=3.55, 95% CI=1.95-6.45, p-value<0.00001). I^2 value was 65%. Risk of PSE in the adult subgroup remained high (OR=2.87, 95% CI=1.67-4.95, p-value=0.0001; I^2 =50%), and in the pediatric subgroup (OR=5.51, 95% CI=1.19-25.45, p-value=0.03; I^2 =75%), although there was a significant heterogeneity in the latter (Figure 4).

In contrast, association between occipital lobe involvement and PSE was not statistically significant (OR=1.37, 95% CI=0.95-1.96, p-value=0.09). I^2 value was 0%. Neither the adult group (OR=1.44, 95% CI=0.98-2.09, p-value=0.06; I^2 =0), nor the pediatric group (OR=0.77, 95% CI=0.22-2.76, p-value=0.69; I^2 =0) showed statistical significance in subgroup analysis (Figure 5).

3.3. Risk of bias in studies

Funnel plots of the studies did not reveal significant asymmetry (Figure 6), although the number of the studies was not sufficient to demonstrate a true asymmetry if one existed.

4. Discussion

To our knowledge, this is the first meta-analysis to investigate the association between the anatomic location of the ischemic infarct and the risk of PSE. Previous frontal lobe infarct was associated with PSE in adults, but the association was not significant in pediatric population. Parietal lobe infarct and temporal lobe infarct were associated with PSE. There was no such association was observed in patients with occipital lobe infarcts.

There has been growing body of literature regarding the pathophysiology of poststroke epilepsy. The role of astrocytic proliferation and glutamate-releasing channel in excitotoxic complications, such as seizure, after stroke has been postulated in an animal study¹⁶. Another in vitro study with primary hippocampal culture demonstrated that the neurons survived from stroke-like injury showed spontaneous, recurrent epileptiform discharges¹⁷. Genetic polymorphism associated with PSE has been reported in previous human studies^{18,19}.

Interestingly, we found no association between previous occipital lobe infarcts and PSE. It can be postulated that the occipital lobe is resistant to epileptogenesis. Another explanation is that the occipital lobe has fewer or less susceptible ictal networks with other cortical or subcortical regions compared to other lobes. Such postulation is supported by studies that described lack of correlation between occipital lobe involvement of glioma and epilepsy²⁰. In addition, it is possible that the rate of reported seizures in patients with occipital lobe infarcts was falsely low due to the relative subtlety of their semiology compared to seizures originating from other cortical areas.

Limitations exist in our study. There was only one prospective study and the rest were retrospective in nature. Also, the prevalence of PSE in the pooled population was lower than that of a previous study²¹. In addition, not all of the studies confirmed seizure onset zones, which may be discordant with the locations of the previous infarcts, although diagnosis of epilepsy does not require this information.

Our analysis was limited by how variables were reported by the included studies. Specifically, only one study⁸ reported the median proportion of each lobe involved by infarction and median infarction volume in each lobe. Although other studies reported other variables, these were not reported separately for stroke involving each lobe, which is required to perform meta-regression analysis.

Analysis showed low to moderate heterogeneity. In general, higher degree of heterogeneity was observed with studies that included pediatric population. Studies that classified the pre-existing infarcts into vascular territories, rather than lobes, were not included, and this may have contributed to the bias and heterogeneity.

Current literature recommends against initiating antiseizure medications in patients who suffered stroke, partly due to the low incidence of PSE. More risk factors in stroke patients for developing PSE are being identified and investigated, including location of the stroke. Further high-quality prospective studies and basic research are needed to deepen the understanding on the mechanism of epileptogenesis of different stroke locations and guide therapeutic approaches for these patients.

5. Conclusion

Our meta-analysis shows that PSE is associated with frontal, parietal, and temporal involvement of previous stroke. However, there was no statistically significant association between occipital lobe involvement and PSE. Further high quality studies are needed to investigate if stroke in certain lobe/territory can be more epileptogenic than those in others.

Consent for Publication

Not applicable.

Declaration of Competing Interest

Authors have no competing interests to declare.

Author Contributions

S.P. designed the study protocol. S.P. and Y.T.E. performed the systematic review. S.P. carried out the statistical analyses. S.P. drafted the manuscript. All authors critically revised the manuscript.

Declaration of Generative AI and AI-assisted technologies in the writing process

There is nothing to disclose.

Funding

None

Reference

1. Zhang C, Wang X, Wang Y, et al. Risk factors for post-stroke seizures: a systematic review and meta-analysis. *Epilepsy Res.* Dec 2014;108(10):1806-16. doi:10.1016/j.eplepsyres.2014.09.030
2. Gasparini S, Ascoli M, Brigo F, et al. Younger age at stroke onset but not thrombolytic treatment predicts poststroke epilepsy: An updated meta-analysis. *Epilepsy Behav.* Mar 2020;104(Pt B):106540. doi:10.1016/j.yebeh.2019.106540
3. Wojcik E, Gráf R, László AM, Gyebnar G, Seri I. Long-term neurodevelopmental outcome of neonates born at term with perinatal haemorrhagic stroke: A population-based study. *Dev Med Child Neurol.* Aug 2022;64(8):971-978. doi:10.1111/dmcn.15149
4. Ouerdiene A, Messelmani M, Derbali H, et al. Post-stroke seizures: risk factors and management after ischemic stroke. Article in Press. *Acta Neurologica Belgica.* 2021;doi:10.1007/s13760-021-01742-x
5. López-Espejo MA, Chávez MH, Huete I. Short-term outcomes after a neonatal arterial ischemic stroke. *Childs Nerv Syst.* Apr 2021;37(4):1249-1254. doi:10.1007/s00381-020-04931-w
6. Kopyta I, Sarecka-Hujar B, Racziewicz D, Gruszczyńska K, Machnikowska-Sokołowska M. Assessment of Post-Stroke Consequences in Pediatric Ischemic Stroke in the Context of Neuroimaging Results-Experience from a Single Medical Center. *Children (Basel).* Apr 8 2021;8(4)doi:10.3390/children8040292
7. Yamada S, Nakagawa I, Tamura K, et al. Investigation of poststroke epilepsy (INPOSE) study: a multicenter prospective study for prediction of poststroke epilepsy. Article. *Journal of Neurology.* 2020;267(11):3274-3281. doi:10.1007/s00415-020-09982-2
8. Takase KI. Cardiogenic cerebral infarction in the parietal lobe predicts the development of post-stroke epilepsy. *Seizure.* Aug 2020;80:196-200. doi:10.1016/j.seizure.2020.06.018
9. Jia Q, Jiang F, Ma D, Li J, Wang F, Wang Z. Association Between IL-6 and Seizure Recurrence in Patients with the First Post-Ischemic Stroke Seizure. *Neuropsychiatr Dis Treat.* 2020;16:1955-1963. doi:10.2147/ndt.S257870
10. Laugesaar R, Vaher U, Lõo S, et al. Epilepsy after perinatal stroke with different vascular subtypes. *Epilepsia Open.* Jun 2018;3(2):193-202. doi:10.1002/epi4.12104
11. Keller L, Hobohm C, Zeynalova S, Classen J, Baum P. Does treatment with t-PA increase the risk of developing epilepsy after stroke? Article. *Journal of Neurology.* 2015;262(10):2364-2372. doi:10.1007/s00415-015-7850-0
12. Hauser W, Annegers JF, Kurland LT. Incidence of epilepsy and unprovoked seizures in Rochester, Minnesota: 1935-1984. *Epilepsia.* May 1993 1993;34(3):453-68. doi:10.1111/j.1528-1157.1993.tb02586.x
13. Collaborators GBDS. Global, regional, and national burden of stroke, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* May 2019;18(5):439-458. doi:10.1016/S1474-4422(19)30034-1
14. Sarecka-Hujar B, Kopyta I. Poststroke epilepsy: current perspectives on diagnosis and treatment. *Neuropsychiatr Dis Treat.* 2019;15:95-103. doi:10.2147/NDT.S169579

15. Zou S, Wu X, Zhu B, Yu J, Yang B, Shi J. The pooled incidence of post-stroke seizure in 102 008 patients. *Top Stroke Rehabil.* Dec 2015;22(6):460-7. doi:10.1179/1074935715Z.000000000062
16. Yang J, Vitery MDC, Chen J, Osei-Owusu J, Chu J, Qiu Z. Glutamate-Releasing SWELL1 Channel in Astrocytes Modulates Synaptic Transmission and Promotes Brain Damage in Stroke. *Neuron.* May 22 2019;102(4):813-827 e6. doi:10.1016/j.neuron.2019.03.029
17. Sun DA, Sombati S, DeLorenzo RJ. Glutamate injury-induced epileptogenesis in hippocampal neurons: an in vitro model of stroke-induced "epilepsy". *Stroke.* Oct 2001;32(10):2344-50. doi:10.1161/hs1001.097242
18. Yang H, Song Z, Yang GP, et al. The ALDH2 rs671 polymorphism affects post-stroke epilepsy susceptibility and plasma 4-HNE levels. *PLoS One.* 2014;9(10):e109634. doi:10.1371/journal.pone.0109634
19. Zhang B, Chen M, Yang H, Wu T, Song C, Guo R. Evidence for involvement of the CD40/CD40L system in post-stroke epilepsy. *Neurosci Lett.* May 1 2014;567:6-10. doi:10.1016/j.neulet.2014.03.003
20. Zhang J, Yao L, Peng S, Fang Y, Tang R, Liu J. Correlation between glioma location and preoperative seizures: a systematic review and meta-analysis. *Neurosurg Rev.* Sep 2019;42(3):603-618. doi:10.1007/s10143-018-1014-5
21. Graham NS, Crichton S, Koutroumanidis M, Wolfe CD, Rudd AG. Incidence and associations of poststroke epilepsy: the prospective South London Stroke Register. *Stroke.* Mar 2013;44(3):605-11. doi:10.1161/STROKEAHA.111.000220

Study	Year	Country	Study design	Population	Stroke type	Age	Total number	Male(%)	PSE prevalence (%)
Vojcek	2022	Hungary	Retrospective	Pediatric	Hemorrhagic	Median 60 mo (IQR: 35-88)	55	34 (62)	9 (16)
Ouerdiene	2021	Tunisia	Retrospective	Adult	Ischemic	Mean 55.1 yrs	52	39 (75)	31 (60)
Lopez	2021	Chile	Retrospective	Pediatric	Ischemic	Median 2 d (IQR 1-5.6)	33	21 (64)	9 (27)
Kopyta	2021	Poland	Retrospective	Pediatric	Ischemic	Mean 8.4 ± 5.5 yrs	75	43 (57)	11 (15)
Yamada	2020	Japan	Prospective	Adult	Ischemic/Hemorrhagic	Mean 68.2 ± 11.1 yrs	436	257 (58.9)	26 (6)
Takase	2020	Japan	Retrospective	Adult	Ischemic	Median case 79.0 (IQR 71 – 86); control 78.5 (IQR 74 – 82)	93	47 (51)	43 (46)
Jia	2020	China	Retrospective	Adult	Ischemic	Mean 62.93 ± 8.82 yrs	209	117 (55.98)	105 (50)
Laugesaar	2018	Estonia	Retrospective	Pediatric	Ischemic/Hemorrhagic	Median 50 mo (1 mo - 18.4yrs)	73	39 (53)	21 (29)
Keller	2015	Germany	Retrospective	Adult	Ischemic	Mean 66.3 ± 12.9 yrs	302	161 (53.3)	42(13.9)

Table 1. Characteristics of the included studies. d, days; IQR, interquartile range; mo, months; yrs, years

ID	Study	JB I (from 11)
1	Vojcek (2022)	10
2	Ouerdiene (2021)	8
3	Lopez (2021)	7
4	Kopyta (2021)	8
5	Yamada (2020)	10
6	Takase (2020)	11
7	Jia (2020)	11
8	Laugesaar (2018)	11
9	Keller (2015)	11

Table 2. Quality assessment of the studies. *JB I: Joanna Briggs Institute appraisal tool score*




Figure 1. A flow diagram illustrating the study selection process for the review.




Figure 2. Forest plot illustrating meta-analysis on frontal lobe stroke and PSE.




Figure 3. Forest plot illustrating meta-analysis on parietal lobe stroke and PSE.




Figure 4. Forest plot illustrating meta-analysis on temporal lobe stroke and PSE.




Figure 5. Forest plot illustrating meta-analysis on occipital lobe stroke and PSE.




Figure 6. Funnel plots of the studies. (a) frontal lobe, (b) parietal lobe, (c) temporal lobe, and (d) occipital lobe infarcts.