

1 **Effect of parental autoimmune diseases on type 1 diabetes in offspring can be partially explained by** 2 **HLA and non-HLA polymorphisms: a nationwide registry and biobank-based study in 7.2M Finns**

3 Feiyi Wang^{†1,2}, Aoxing Liu^{†1,3,4,5}, Zhiyu Yang¹, Pekka Vartiainen¹, Sakari Jukarainen¹, Satu Koskela⁶,
4 Richard A. Oram⁷, Lowri Allen⁸, Jarmo Ritari⁶, Jukka Partanen⁶, FinnGen, Markus Perola⁹, Tiinamaija
5 Tuomi^{*1,10,11,12}, Andrea Ganna^{*1,3,4,5}

- 6 1. Institute for Molecular Medicine Finland (FIMM), Helsinki Institute of Life Science (HiLIFE), University of
7 Helsinki, Helsinki, Finland
- 8 2. Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics,
9 University of Edinburgh, Edinburgh, Scotland, UK
- 10 3. Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA, USA
- 11 4. Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT, Cambridge, MA, USA
- 12 5. Analytic and Translational Genetics Unit, ATGU, Massachusetts General Hospital, Boston, MA, USA
- 13 6. Finnish Red Cross Blood Service, Helsinki, Finland
- 14 7. University of Exeter, Exeter, UK
- 15 8. Cardiff University and University Hospital of Wales, Cardiff, UK
- 16 9. Finnish Institute for Health and Welfare (THL), Helsinki, Finland
- 17 10. Abdominal Center, Endocrinology, Helsinki University Hospital, Helsinki, Finland
- 18 11. Folkhälsan Research Center, Helsinki, Finland
- 19 12. Lund University Diabetes Center, Malmo, Sweden

20 † Joint first authorship

21 * Joint senior authorship

22 Corresponding author: Andrea Ganna (andrea.ganna@helsinki.fi) and Tiinamaija Tuomi (tiinamaija.tuomi@hus.fi)

23

24 **Abstract**

25 Type 1 diabetes (T1D) and other autoimmune diseases (AIDs) co-occur in families. We studied the
26 aggregation of 50 parental AIDs with T1D in offspring and the contribution of a shared genetic background,
27 which was partitioned into HLA and non-HLA variation. Leveraging nationwide registers of 7.2M Finns,
28 including 58,284 family trios, we observed that 15 parental AIDs, such as coeliac disease and rheumatoid
29 arthritis, were associated with an increased risk of T1D in offspring. The identified epidemiological
30 associations were then tested by comprehensive genetic analyses performed on 470K Finns genotyped in
31 the FinnGen study (12,563 trios). The within-family genetic transmission analysis further demonstrated that
32 the aggregation of parental AIDs with offspring T1D could be partially explained by HLA and non-HLA
33 polymorphisms in a disease-dependent manner. For example, the associations with offspring T1D for
34 coeliac disease and psoriasis were mainly driven by HLA while autoimmune hypothyroidism and
35 rheumatoid arthritis also had non-HLA contributors. We, therefore, proposed a novel parental polygenic
36 score (PGS), integrating variations in both HLA and non-HLA genes, to understand the cumulative risk
37 pattern of T1D in offspring. This raises an intriguing possibility of considering parental PGS, in conjunction
38 with clinical diagnoses, to inform individuals about T1D risk in their offspring.


39

40 Type 1 diabetes (T1D) is an autoimmune disease (AID) characterized by a severe insulin deficiency
41 resulting from the destruction of insulin-producing pancreatic beta cells. While the pathogenic mechanism
42 remains largely unknown, T1D usually appears in genetically susceptible individuals^{1,2} and triggered by
43 environmental exposures^{3,4} - mostly before 20 years of age, with incidence increasing from birth and
44 peaking at age 10-14 during puberty⁵⁻⁷. Notably, in many populations such as Europeans, Asians, and Latins,
45 genetic polymorphisms in human leukocyte antigen (HLA) genes were reported to account for up to half
46 of the T1D heritability, for which the strongest effects were attributable to two HLA class II haplotypes,
47 HLA-DR3-DQ2 and DR4-DQ8⁸⁻¹⁴. In addition to the well-documented large-effect HLA contribution, non-
48 HLA susceptibility also plays a role². Large-scale genome-wide association studies (GWAS) have
49 identified dozens of independent non-HLA signals across the genome¹⁵⁻¹⁸, that together with HLA
50 polymorphisms can be used to construct polygenic scores (PGS) to evaluate genetic predisposition of T1D
51 prior to the disease onset¹⁹⁻²¹ as well as better differentiate T1D from other types of diabetes²²⁻²⁴.

52 Interestingly, T1D tends to co-occur with some other AIDs, both in the same individuals and within families
53 (**Supplementary notes 1**). Previous studies mainly explored the co-occurrence from two angles. A popular
54 and straightforward one was to adopt a population-based design and epidemiological analysis for the risk
55 of T1D among people whose relatives were diagnosed with some AIDs. We have provided a detailed review
56 of these studies in **Table S1**. To obtain conclusive findings, a sufficient number of trios or families is
57 needed²⁵⁻²⁷. Another set of studies attempted to leverage genetic information for a better understanding of
58 T1D-AID aggregation among genetically related individuals²⁸⁻³⁰. In one such study, among the one-third of
59 children with T1D who were found to have a relative affected by an AID, having HLA-DR3-DQ2 haplotype
60 was associated with celiac disease (CD) in their extended family²⁹. Whereas this type of studies mainly
61 collected family history from questionnaires and focused only on HLA haplotype analysis due to data
62 limitation, other studies systematically researched the disease associations of common variants in either
63 HLA or non-HLA genes. Instead of individual-level data, many studies utilized publicly accessible
64 summary statistics generated from population-based analyses. For example, a recent study captured
65 intercorrelation of AIDs, such as T1D, rheumatoid arthritis (RA), and autoimmune hypothyroidism
66 (HYPO), in two large HLA hotspots (locus 961 and 964)³¹. Further, GWAS revealed shared genetic
67 susceptibility loci beyond the HLA regions for T1D and five other AIDs, including CD, RA, multiple
68 sclerosis (MS), Crohn's diseases, and systemic lupus erythematosus³².

69 To date, there has been a lack of comprehensive investigation that combines the two angles and
70 encompasses a wider range of AID spectrum than CD, RA, HYPO and MS. Meanwhile, it remains an
71 enigma why individuals with similar genetic backgrounds can develop such heterogeneous diseases ranging
72 from potentially lethal insulin deficiency to mild hypothyroidism - even among members of the same family
73 who also tend to share the same environment. One way to approach this question is to combine evidence
74 from population-based multi-generational cohorts and large-scale biobanks with high-resolution diagnostic
75 data, preferably of the same population to avoid any bias or confounding caused by differences in across-
76 population socio-cultural contexts and clinical practices. In this study, we aim to comprehensively evaluate
77 the genetic determinants of familial aggregation of T1D and other AIDs using the nationwide multi-
78 generational health registers of the whole Finnish population in FinRegistry³³, as well as the genomic and
79 family trio data available in Finnish biobanks through the FinnGen Study³⁴. Considering the world's highest
80 incidence of T1D in Finland³⁵, these detailed and structured data resources provide us a unique opportunity
81 to quantify the associations between parental AIDs and offspring T1D, and more importantly, to answer
82 three fundamental questions: 1) Which parental diagnoses of AIDs are associated with T1D in offspring?
83 2) To what extent is the shared genetic background between AIDs and T1D driven by genetic components,
84 separately for HLA and non-HLA variants? And, 3) Can AID-related genetic information in parents
85 complement disease family history be used to estimate the risk of having a child developing T1D?

86



87

88 **Figure 1:** An overview of the study design and study populations. Panel A, the study population of
 89 FinRegistry (7.2 million individuals) represents every Finn alive in 2010 (5.3 million individuals) and their
 90 first-degree relatives. To maximize the coverage of diagnoses of AIDs for parents and T1D for children,
 91 we included only family trios with both parents born before 1976 and children between 1960-1999 (the
 92 follow-up time in 2019 was at least 45 years for parents and 20 years for children). The solid lines denote
 93 the birth year range of the study population and the dashed lines the years of follow-up. Among the 2.4
 94 million family trios, 14,571 had a child ever diagnosed with T1D (T1D trios), and for each of the T1D-trios,
 95 we matched three control trios based on sex, birth year, birthplace, and the number of siblings of the child,
 96 as well as birth years of both parents. In total, we included 14,571 T1D-trios and their matched 43,713
 97 control trios. Panel B, the study population of FinnGen includes 473.7K Finns enrolled in a nationwide
 98 network of Finnish biobanks, of which 12,563 family trios can be accurately constructed with available
 99 genomic information. FinnGen includes 3,370 T1D cases, 385,786 controls, as well as 1,129 T1D-trios and
 100 11,434 control trios. Panel C illustrates a matched case-control study conducted in the 58,284 FinRegistry
 101 family trios to examine the association between parents' AID and the T1D status of offspring. The different
 102 symbols denote father, mother, and offspring, and colors the disease status (red, offspring with T1D; green,
 103 parents with AIDs; grey, individuals unaffected by T1D or other AIDs). Panel D depicts the examination
 104 of shared genetic components between T1D and other AIDs using population-based analyses (left panel
 105 with blue background) or trio-based family analyses (right panel with yellow background). A haplotype
 106 and PGS-based analysis of HLA variants conducted in 473.7K FinnGen participants (left); a genetic
 107 correlation analysis of the non-HLA variants utilized GWAS summary statistic data (middle); and a
 108 polygenic transmission disequilibrium test (pTDT) examined in FinnGen participants whether the AID-
 109 associated common variants as a whole were over-transmitted from AID-unaffected parents to their T1D-
 110 affected offspring (right). Panel E, the average AID PGS of the parents and its predictive performance of

111 T1D in offspring among 12,563 genotyped family trios in FinnGen. AID, autoimmune disease; T1D, type
112 1 diabetes; HLA, human leukocyte antigen; PGS, polygenic score; GWAS, genome-wide association study.

113
114

115 **Results**

116

117 **Study populations and disease diagnoses**

118

119 We leveraged nationwide socio-demographic and health registers of the entire Finnish population collected
120 in the FinRegistry study³³ (N = 7.2 million) and a subset of 473,681 genotyped Finns from the FinnGen
121 study³⁴ (**Figure 1A** and **1B**). To avoid potentially biased estimates caused by incomplete medical records,
122 we excluded individuals who had migrated in or out of Finland by 2019 (572,640 were excluded for
123 FinRegistry and 9,196 for FinnGen). To have comparatively complete coverage of AID diagnoses for
124 parents and T1D diagnoses for children, we considered parents born before 1976 and children born between
125 1960 and 1999, corresponding to at least 20 years of follow-up for children by the end of 2019 (**Figure**
126 **S2.2**). We defined T1D cases as those who had their first diagnosis recorded before the age of 40. In total,
127 we included 14,571 T1D cases in FinRegistry (0.61%) and 3,668 in FinnGen (0.83%). To examine the
128 association between AIDs in parents and T1D in offspring, as well as the extent to which genetic factors
129 could contribute to the identified association, we constructed 2.4 million family trios in FinRegistry based
130 on Finnish multi-generation registers and 12,563 genotype-inferred trios in FinnGen (**Methods**). Of these
131 trios, 14,571 in FinRegistry and 1,129 in FinnGen had children with T1D.

132 To have broad coverage of AIDs affecting different tissues or organ systems, we defined 50 AIDs or
133 autoimmune-related disorders based on Finnish versions of the International Classification of Diseases
134 (ICD) codes 8-10 (**Supplementary notes 2.1**). After excluding 13 AIDs with less than 50 parental cases in
135 FinRegistry and 11 AIDs that were highly correlated with other AIDs (**Table S2.1**), we included 26 AIDs
136 for further analyses: T1D, adrenocortical insufficiency (ADDISON), autoimmune hyperthyroidism
137 (HYPER), autoimmune hypothyroidism (HYPO), autoimmune haemolytic anaemias (AIHA), allergic
138 purpura, vitamin B12 deficiency anemia (B12A), idiopathic thrombocytopenic purpura (ITP), sarcoidosis,
139 primary biliary cholangitis (PBC), celiac disease (CD), inflammatory bowel disease (IBD), IgA
140 nephropathy, ankylosing spondylitis (AS), mixed connective tissue disease (MCTD), rheumatoid arthritis
141 (RA), Sjögren's syndrome (SjS), systemic sclerosis, Wegener granulomatosis (WG), systemic lupus
142 erythematosus (SLE), Guillain-Barré syndrome (GBS), multiple sclerosis (MS), myasthenia gravis (MG),
143 alopecia areata (AA), psoriasis, and vitiligo. The number of parental cases ranged from 55 for AIHA
144 (corresponding to a prevalence of 0.05%) to 11,485 for HYPO (prevalence = 9.85%) (**Table S2.1** and
145 **S3.1.1**).

146 We designed four sets of complementary analyses (**Figure 1**). We first examined the association between
147 parental AID and offspring T1D risk in FinRegistry. For AIDs showing significant epidemiological
148 associations, we explored the HLA-related associations with haplotype and PGS-based analyses in FinnGen,
149 and the genome-wide non-HLA associations with a summary statistic-based genetic correlation analysis. A
150 polygenic transmission disequilibrium test (pTDT) was then used to examine the within-family
151 transmission of AID genetic signals from AID-unaffected parents to their T1D-affected and unaffected
152 offspring. Lastly, we proposed a novel parental PGS method, integrating both HLA and non-HLA
153 contributors, to assess the risk of developing T1D in offspring.

154

155

156


157

158

159 **Epidemiological associations between parental AIDs and T1D in offspring**

160 We first wanted to examine the associations between parental AIDs and T1D in children using the large
161 number of family trios available in the FinRegistry study. For each of the 14,571 trios with T1D-affected
162 children, we employed a 1:3 matched case-control design considering the information of both the child (sex,
163 birth year, birthplace, and number of siblings) and parents (birth year) (**Methods** and **Figure 1C**). For other
164 available socio-demographic factors that were not used in matching, we saw limited differences between
165 T1D trios and their matched controls (**Table S3.1.2**). Overall, children with T1D (42.2% [95% CI 41.4%,
166 43.0%]) were more likely to have AID-affected parent(s) compared to those without T1D (31.9% [31.5%,
167 32.3%]).

168 Of 26 parental AIDs examined in FinRegistry, fifteen were associated with increased risk of T1D in
169 offspring at a nominal P-value <0.05, and nine remained statistically significant after Bonferroni correction
170 for multiple-testing ($0.05/26 = 0.002$) (**Figure 2** and **Table S3.2.1.1**). The strongest association was seen
171 for T1D (odds ratio (OR) [95% CI], 6.77 [5.44, 8.42], $P=3.8 \times 10^{-66}$), followed by CD (2.14 [1.90, 2.42],
172 $P=2.5 \times 10^{-35}$), B12A (1.76 [1.54, 2.02], $P=1.4 \times 10^{-16}$), MCTD (1.54 [1.24, 1.92], $P=8.5 \times 10^{-5}$), HYPO (1.53
173 [1.46, 1.60], $P=4.8 \times 10^{-73}$), RA (1.48 [1.37, 1.60], $P=1.8 \times 10^{-23}$), sarcoidosis (1.45 [1.28, 1.65], $P=1.1 \times 10^{-8}$),
174 HYPERS (1.42 [1.20, 1.68], $P=4.3 \times 10^{-5}$), and SjS (1.33 [1.12, 1.59], $P=1.6 \times 10^{-3}$). In terms of sex, we
175 didn't observe significant differences either for parents or for offspring (**Figure S3.2.2**). However, as
176 previously reported in the literature³⁶⁻³⁸, fathers with T1D had a higher risk of having children with T1D
177 than mothers with T1D (7.49 [5.68, 9.86] vs. 5.71 [4.01, 8.13], $P_{\text{difference}}=0.24$), but this was not significant.
178 We also noted that, a paternal history of T1D could significantly impact the onset age of T1D among boys,
179 resulting in T1D being diagnosed 1.85 [0.29, 3.41] years earlier compared to having a T1D-unaffected
180 father ($P=0.02$) (**Supplementary notes 3.2.3**). A sensitivity analysis restricted to early-onset T1D (before
181 age 20) yielded stronger associations for parental early-onset T1D and parental SLE from the main analysis
182 (before age 40). For example, the OR increased from 6.77 [5.44, 8.42] to 10.90 [7.90, 15.04] for parental
183 early-onset T1D ($P_{\text{difference}}=0.02$) and from 1.29 [0.96, 1.73] to 1.77 [1.24, 2.51] for parental SLE
184 ($P_{\text{difference}}=0.18$) (**Table S3.2.1.2**).
185



186
 187 **Figure 2:** Epidemiological associations between parental AIDs and T1D in offspring ordered by P-value.
 188 Upper panel: Association (OR, and 95% CI) between parental AIDs and T1D in offspring using a matched
 189 case-control design in Finnish nationwide registry data (FinRegistry). Lower panel: number of T1D cases
 190 with parents having a given AID diagnosis. The dark green diamonds and bars indicate AIDs that are
 191 significantly associated with T1D after multiple testing corrections. AID, autoimmune disease; T1D, type
 192 1 diabetes; OR, odds ratio; CI, confidence interval.

193
 194

195 Shared genetic components from HLA and non-HLA variants at a population level

196 Parents and offspring share many environmental exposures and genetic components. We next aimed to
 197 quantify the extent to which the identified epidemiological associations are attributable to a shared genetic
 198 background. Given that T1D is known to be impacted by both HLA and non-HLA variations^{2,8-18}, we
 199 analyzed HLA and non-HLA regions separately at a population level (**Methods**). We first examined the
 200 association of the main T1D predisposing / protective HLA haplotypes with other AIDs to understand to
 201 what extent a single haplotype can drive the identified epidemiological associations. Next, we combined
 202 multiple HLA imputed alleles to create multi-allelic scores for HLA for each AIDs and examine the
 203 association with T1D. The analyses were conducted on 439,817 individuals from FinnGen (**Figure 1B**,
 204 **Table S3.1.1**). For non-HLA variations, we estimated genetic correlation between T1D and other AIDs
 205 using the latest GWAS summary statistics.

206

207 *Shared effect of HLA haplotypes for T1D and other AIDs*

208 Given that for T1D, the strongest genetic associations are for Class II haplotypes, in particular HLA-DR3
209 and DR4 haplotypes^{8-14,39}, we constructed 19 *DRB1-DQA1-DQB1* haplotypes based on 64 imputed alleles
210 of these genes in FinnGen⁴⁰ (**Methods**).

211 We examined 19 haplotypes (frequencies > 0.5%, **Table S3.3.1.1**), of which six were strongly associated
212 with T1D ($P < 1.0 \times 10^{-50}$), with *DRB1*04:01-DQA1*03:01-DQB1*03:02* (OR=7.54 [7.00, 8.13], $P < 1.0$
213 $\times 10^{-602}$) conferring the strongest susceptibility and *DRB1*15:01-DQA1*01:02-DQB1*06:02* (0.06 [0.04,
214 0.07], $P = 2.0 \times 10^{-106}$) the strongest protection (**Table S3.3.1.2**). We then tested the association of these six
215 haplotypes with the 24 other included AIDs using logistic regression and considering the age, sex, and first
216 ten principal components (PCs) as covariates (**Figure 3A**). Many of these T1D-associated haplotypes were
217 also associated with other AIDs ($P < 1.0 \times 10^{-4}$) (**Table S3.3.1.3**). For example, the T1D risk haplotype
218 *DRB1*03:01-DQA1*05:01-DQB1*02:01* increased the risk of both CD (15.83 [14.56, 17.20], $P < 1.0 \times 10^{-$
219 ⁹¹⁸) and autoimmune hyperthyroidism (2.42 [2.23, 2.63], $P = 6.3 \times 10^{-100}$), and *DRB1*04:01-DQA1*03:01-*
220 *DQB1*03:02* increased the risk of RA (1.93 [1.84, 2.02], $P = 5.1 \times 10^{-174}$). Of note, opposite effects were
221 seen for some AIDs: the strongest haplotype protecting against T1D (*DRB1*15:01-DQA1*01:02-*
222 *DQB1*06:02*) was the lead risk haplotype for MS⁴¹ (2.85 [2.60, 3.12], $P = 7.9 \times 10^{-112}$) and the strongest T1D
223 susceptibility haplotype (*DRB1*04:01-DQA1*03:01-DQB1*03:02*) protected against IBD (0.85 [0.80,
224 0.90], $P = 4.7 \times 10^{-8}$).

225

226 *Multi-allelic scores (PGS) of HLA alleles for AIDs and T1D*

227 Having observed that multiple HLA haplotypes could contribute to the susceptibility to T1D and other
228 AIDs independently, we constructed polygenic HLA scores (HLA PGSs) to summarize the overall multi-
229 allelic HLA effects for each AIDs. Considering the widespread variation in HLA allele frequencies across
230 populations⁴², we opted to construct HLA PGSs within the FinnGen samples 187 imputed Class I and Class
231 II alleles using weighted ridge regression (**Methods**). We were able to construct reliable HLA PGSs
232 ($P \leq 0.05/26$ and partial correlation $|\rho| \geq 2\%$ for predicting the susceptibility of the disease itself) in 23,336
233 individuals for 14 AIDs (including T1D). Of these, HLA PGSs for 13 AIDs exhibited significant
234 associations with T1D susceptibility after multiple-test correction ($0.05/14 = 0.003$), with exceptions for
235 psoriasis ($P = 3.6 \times 10^{-3}$) and AS ($P = 6.1 \times 10^{-2}$) (**Figure 3B** and **Table S3.3.2.1**). Overall, the effects of AID
236 HLA PGS on individuals' own T1D risk were highly correlated with the effects of parental AIDs on
237 offspring T1D obtained from our FinRegistry epidemiological analyses (ρ from a Spearman's rank
238 correlation = 0.63, $P = 6.1 \times 10^{-4}$). Among all AIDs other than T1D (OR=5.33 [4.79, 5.93], $P = 3.74 \times 10^{-207}$),
239 strongest associations were seen for ADDISON (OR=2.30 [2.13, 2.49], $P = 4.2 \times 10^{-100}$), RA (OR=2.10 [1.95,
240 2.27], $P = 1.8 \times 10^{-78}$), and CD (OR=2.07 [1.94, 2.22], $P = 2.4 \times 10^{-96}$). Consistent with the HLA haplotype
241 analysis, we observed negative associations with T1D for HLA PGS of IBD (OR=0.64 [0.60, 0.69],
242 $P = 1.2 \times 10^{-30}$) and MS (OR=0.81 [0.75, 0.88], $P = 2.4 \times 10^{-7}$). These associations were further replicated with
243 a distinct analytic approach - leave-one-group-out (LOGO) (**Supplementary notes 2.3.2** and **Table**
244 **S3.3.2.2**).


245

246 *Genome-wide non-HLA correlation between T1D and AIDs*

247 In addition to the major HLA contributors, non-HLA variants identified in the GWAS studies are known
248 to impact AID susceptibility as well. We, therefore, explored the shared correlation between T1D and other
249 AIDs using a summary statistics-based method - LD score regression (LDSC)^{43,44}, excluding the HLA
250 regions (**Methods**). Significant positive genetic correlations (r_g) were observed between T1D and ten AIDs
251 ($P < 0.05/25$ after Bonferroni correction) (**Figure 3C**), including HYPO, CD, RA, B12A, sarcoidosis,
252 HYPER, vitiligo, PBC, MG, and SLE. The first six AIDs were also among the parental AIDs that exerted
253 significant effects on offspring T1D in our FinRegistry epidemiological analyses (ρ from a Spearman's
254 rank correlation=0.30, $P=0.15$). The highest r_g were seen for B12A ($r_g=0.48$ [0.34, 0.63], $P=6.4 \times 10^{-11}$),
255 followed by HYPO (0.43 [0.34, 0.51], $P=2.9 \times 10^{-23}$) and RA (0.43 [0.25, 0.61], $P=3.9 \times 10^{-6}$) (**Table**
256 **S3.3.3.1**). Contrary to the negative HLA association, we did not observe a significant non-HLA genetic
257 correlation between T1D and MS or between T1D and IBD. A sensitivity analysis excluding all
258 chromosome 6 variants yielded similar estimates, suggesting that the observed non-HLA results were not
259 driven by variants in highly linkage disequilibrium with HLA alleles (**Table S3.3.3.2**).

260 In summary, both the HLA and non-HLA analyses recapitulated the multi-generational epidemiological
261 effects observed in nationwide registers. While the results conferred that the risk of T1D was positively
262 associated with the risk of many other AIDs regarding both HLA and non-HLA genetic variants (e.g.,
263 HYPO, CD, RA), the opposite held for MS and IBD concerning the HLA susceptibility whereas no genetic
264 correlation was observed in non-HLA regions.

265



266

267 **Figure 3:** Shared genetic background between T1D and other AIDs, stratified by HLA and non-HLA
 268 variations at a population level. **Panel A**, HLA haplotypes strongly associated with T1D and their
 269 associations with other AIDs in FinnGen (3,668 T1D cases and 436, 149 T1D controls). The strength of the
 270 association is shown as a heatmap with blue depicting a susceptible haplotype associated with the AID in
 271 question. The haplotypes are sorted by the P-value for T1D as shown on the left. **Panel B**, upper panel:
 272 association (OR, and 95% CI) between HLA PGSs for different AIDs and T1D in FinnGen. The dark blue
 273 squares indicate that HLA PGS for a given disease has a significant association with T1D after multiple
 274 testing corrections. "x" denotes that an HLA PGS could not be robustly constructed for that AID (**Methods**).
 275 Bottom panel: blue bars represent the number of individuals with the given disease. **Panel C**, upper panel:
 276 non-HLA based genetic correlations (r_g and 95% CI) between AIDs and T1D using GWAS summary

277 statistics from European populations. Bottom panel: the number of cases for the given disease. The dark
278 orange squares or bars indicate that the AIDs have a significant r_g with T1D after multiple testing
279 corrections. T1D, type 1 diabetes; AID, autoimmune disease; HLA, human leukocyte antigen; OR, odds
280 ratio; CI, confidence interval; PGS, polygenic score; r_g , genetic correlation; GWAS, genome-wide
281 association study.

282
283
284


285 **Transmitted and non-transmitted genetic liability within families**

286
287
288

287 *Polygenic transmission disequilibrium for HLA and non-HLA genetic factors*

289 Following the population-based epidemiological evidence for clustering of other AIDs and T1D in the trios,
290 and the genetic evidence for a shared genetic origin of the AIDs and T1D at a population level, we wanted
291 to examine the transmission of the HLA and non-HLA within families using a polygenic transmission
292 disequilibrium test (pTDT)⁴⁵. The pTDT examines whether the AID risk variants are observed to be over-
293 transmitted (or under-transmitted) to the offspring with T1D (compared to the expected transmission rate)
294 and thus it is immune to many of the potential confounders arising from population studies on unrelated
295 individuals. We considered 12,563 trios (genotyped in FinnGen), of which 1,159 had offspring with T1D
296 (9.2%). The prevalence of T1D in these trios was higher than expected in the general population because
297 of the inclusion of several studies have specifically targeted individuals with T1D. We analyzed 10 AIDs
298 with reliable PGSs ($\rho \geq 2\%$) for both HLA and non-HLA genes and additional six diseases with only reliable
299 PGSs for the HLA regions (**Methods; Supplementary notes 3.3.2 and Table S3.4.1**). For offspring with
300 T1D, both HLA and non-HLA PGSs for T1D deviated significantly from their mid-parent value (1.23 [1.16,
301 1.30], $P=2.6 \times 10^{-168}$ and 0.69 [0.62, 0.75], $P=9.6 \times 10^{-74}$) while no such deviation was observed in unaffected
302 siblings (-0.06 [-0.13, 0.01], $P=0.08$ and 0.00 [-0.06, 0.05], $P=0.88$) (**Figure 4 and Table S3.4.2**). Overall,
303 two major patterns were seen across the analyzed AIDs. The first group encompassed T1D, HYPO, RA,
304 and SLE, for which significant over-transmission in offspring with T1D was seen for both HLA and non-
305 HLA PGSs. The second group exhibited significant transmission only for HLA but not non-HLA PGSs:
306 while psoriasis exhibited significant over-transmission in offspring with T1D compared to unaffected
307 siblings, IBD and MS had a significant under-transmission. Taking IBD as an example, among 2,040
308 genotyped trios, 756 offspring with T1D presented under-transmitted HLA PGS (-0.38 [-0.45, -0.31],
309 $P=3.1 \times 10^{-22}$) while their 1,269 unaffected siblings had comparable PGS to their parents (-0.01 [-0.08, 0.06],
310 $P=0.17$). No diseases showed significant pTDT only regarding non-HLA PGS without a significant
311 association for HLA-PGS.

312



313
314
315
316
317
318
319
320
321
322
323
324

Figure 4: Polygenic transmission disequilibrium tests (pTDT) assess whether AID PGSs transmitted to offspring significantly deviated from the parental PGS in T1D-affected offspring and their unaffected siblings. Deviations from parental PGS (mean and 95% CI) of 1 SD change in offspring PGS are shown separately for HLA and non-HLA PGSs, with the P value obtained from a two-sided t-test. Red color denotes offspring with T1D and gray unaffected siblings. AID, autoimmune disease; PGS, polygenic score; T1D, type 1 diabetes; CI, confidence interval; SD, standard deviation; HLA, human leukocyte antigen.

Offspring T1D risk based on parental AIDs clinical history and genetic information

325
326
327
328
329
330
331
332
333
334
335
336

The PGS for T1D has shown potential to predict disease susceptibility before its onset¹⁹⁻²¹. We asked whether T1D risk in offspring can be assessed using a couple's genetic information before they plan to have a child, and how this compares to estimating the risk based on family history of T1D and other AIDs, as is currently being done in clinical practice. To address this question, we adopted a novel strategy to construct a full PGS (Full-PGS) by considering the contribution ratio of HLA PGS relative to non-HLA PGS (**Methods; Supplementary notes 3.5**). We tested eight AIDs (T1D, HYPO, RA, SLE, CD, psoriasis, IBD, and MS) with at least one reliable PGS in HLA regions or non-HLA regions for T1D prediction ($|p| \geq 1\%$) (**Table S3.5.1**) using five-fold cross validation. By modelling the HLA and non-HLA variations separately and using imputed HLA alleles (187 alleles in 10 classical HLA genes rather than common genotyped SNPs), the Full-PGSs we proposed outperformed the standard SNP-based PGS methods such as PRS-CS⁴⁶ at an individual level (**Table S3.6.1; Table S3.6.2**) and when predicting T1D in offspring using AID-specific parental PGS among the 12,563 FinnGen genotyped trios (**Table S3.6.3; Figure S3.6.4**).

337
338
339
340


Having established that a parental Full-PGS for T1D was strongly associated with T1D risk in offspring ($AUC_{mean} = 0.817$), we tested whether the parental Full-PGSs for all the eight AIDs could add any additional information to the parental Full-PGS for T1D. However, limited difference was seen by integrating the PGS of eight AIDs ($AUC_{mean}: 0.817$ vs 0.820 , $P_{difference} = 0.77$). This suggested that, instead of a direct effect, the

341 impact of parental AID PGS on offspring T1D was most likely to be mediated through the genetic
342 correlation between T1D and AID, and therefore provided limited additional information to the offspring
343 T1D prediction when parental T1D PGS was already included and sufficiently accurate.

344 We thus focused on parental Full-PGS for T1D and estimated the cumulative risk of T1D in offspring using
345 a Cox proportional hazards model adjusted for the first 10 PCs and birthyear of the child among genotyped
346 trios (**Methods**). When stratified PGS into four groups by percentiles (0-50th, 50-90th, 90-99th, 99-100th),
347 we observed distinct trajectories in terms of T1D cumulative incidence rates (0.24%, 0.74%, 2.29%, 6.96%
348 by age 20) (**Figure 5A**). That is, children whose parents were in the top percentile of T1D Full-PGS had a
349 29-fold higher risk of developing T1D than children whose parents were within the bottom 50% of T1D
350 Full-PGS. When further stratified by the sex of children (**Figure 5B** and **5C**), the adjusted survival curves
351 suggested that sons overall had a higher cumulative incidence than daughters across all PGS groups. For
352 example, in the top T1D parental Full-PGS group, sons (11.48%) could have a 2-fold higher cumulative
353 risk than daughters (5.29%) by age 20. Overall, the cumulative incidence curves of T1D started to level off
354 around the age of 14 to 16.

355 We then wanted to understand how informative the Full-PGS was compared to parental AID clinical
356 history. In general, the offspring of T1D couples had a higher cumulative incidence of T1D risk (**Figure**
357 **5D**) than T1D-unaffected couples and such differences were larger for boys (**Figure 5E**) than girls (**Figure**
358 **5F**). Having either father or mother with T1D (prevalence_{father} = 0.22%, prevalence_{mother} = 0.15%) resulted
359 in a lower cumulative risk than having both parents in the top percentile of Full-PGS for T1D (1.49% and
360 4.16% vs 6.95% by age 20 for children). Having both parents affected by T1D (a very rare event with only
361 a prevalence of 8.5 per million) resulted in a higher cumulative risk (11.30%) than having parents in the
362 top percentile of Full-PGS for T1D (6.95%). Among couples only affected by AIDs other than T1D,
363 offspring also had a higher cumulative incidence of T1D than the general population (**Fig. 5 G, H, and I**)
364 (0.96 % vs 0.48%), although the cumulative incidence would be smaller than that for the offspring from
365 couples with physician-diagnosed T1D or couples in the top decile of the Full-PGS for T1D.

366



367
368
369
370
371
372
373
374
375
376
377
378
379
380

Figure 5: Adjusted survival curves from Cox proportional hazards models for cumulative incidence of T1D in offspring by age 20, stratified by parental full T1D PGS percentiles comprising both HLA and non-HLA variants (**Panels A-C**; 8,872 FinnGen genotyped trios), parents' T1D status (**Panels D-F**; 3,048,812 FinRegistry trios), parents' other AIDs (**Panels G-I**; 3,037,723 FinRegistry trios after removing trios having parent(s) diagnosed with T1D), including T1D, autoimmune hypothyroidism, rheumatoid arthritis, systemic lupus erythematosus, coeliac disease, psoriasis, inflammatory bowel disease, and multiple sclerosis; The panels show pooled data for both sexes (left), daughters (middle), and sons (right). T1D, type 1 diabetes; AID, autoimmune disease; HLA, human leukocyte antigen; PGS: polygenic score.

381 Discussion

382 In this study, we comprehensively explored the genetic determinants of the familial aggregation of T1D
383 and other AIDs with data from high-dimensional nationwide registers and rich genetic information of the
384 Finnish population. Our long follow-up (≥ 20 years) in offspring and the early age of T1D onset (median
385 age = 12.7) allows us to maximize the number of T1D cases identified in the study. With systematically
386 designed epidemiological and genetic analyses, we seek to answer three key but underexplored questions:
387 which parental AIDs are related to T1D in offspring, to what extent the identified parental AID-offspring
388 T1D association is attributable to genetic polymorphisms of HLA and non-HLA genes, and last, for any
389 couples that are planning to have children, how well we could use their genetic information to evaluate the
390 T1D risk of their offspring.

391 The nationwide registers of 7.2 Finns collected in the FinRegistry study cover longitudinal health and
392 sociodemographic information of 58,284 family trios which provides us a unique opportunity to perform a
393 comprehensive exploration of 50 autoimmune diseases for their impacts on offspring T1D, on a scale and
394 detailed level that could be difficult to achieve in questionnaire- or survey-based studies. This rich familial
395 and health information enabled us to perform an efficient matched case-control design to better control for
396 unmeasurable confounding factors than the standard unmatched design used in previous large population-
397 based studies²⁵⁻²⁷. In total, we detected ten parental AID-offspring T1D associations, including T1D, HYPO,
398 CD, RA, HYPHER, and B12A, that have previously been linked to T1D risk in offspring in large population-
399 based studies (**Table S4**), and more importantly, six novel associations encompassing AIHA, MCTD, MG,
400 AA, psoriasis, and vitiligo.

401 Following the epidemiological analyses, we leveraged the genetic information of 470K genotyped Finns
402 enrolled in the FinnGen study to better understand the genetic causes of the observed family aggregations.
403 Given that T1D is known to be impacted by both HLA and non-HLA variations^{2,8-18}, we designed a set of
404 genetic analyses in a hypothesis-free manner to separate the contributions from HLA and non-HLA regions
405 for each analyzed AID. These genetic analyses were conducted both at a population level, studying the
406 shared effects of HLA and non-HLA variations among unrelated individuals, and, within families using
407 pTDT in 12,563 FinnGen family trios, which to the best of our knowledge, is the largest family-based
408 genetic analysis performed for T1D and AIDs. Similar to the epidemiological family-based study that is
409 widely accepted to yield less biased estimates than population-based studies, pTDT can condition out the
410 impact of shared familial factors through within-family comparisons, allowing, for the first time, to examine
411 under- or over-transmission of genetic risk factors for AIDs in individuals with T1D and their unaffected
412 siblings.

413 Generally, two major patterns were seen for the transmission of AID-associated variants to offspring with
414 T1D, with one group encompassing T1D, HYPO, RA, and SLE showing significant over-transmission for
415 both HLA and non-HLA variants, and another group comprising IBD, MS, CD, and psoriasis that exerted
416 significant deviated transmission only for HLA. We noted that even among the disease group presenting
417 over-transmission for both HLA and non-HLA, more significant transmission was seen regarding HLA
418 compared to non-HLA. This is consistent with the previous studies which reported strong HLA associations
419 for many AIDs^{2,47,48}.

420 Previous works over decades have shown evidence for variation in genes in both HLA and non-HLA
421 regions being associated with and between AIDs, especially after the advances in the development of
422 GWAS and multi-omics data in recent years^{47,49-51}. Specifically, genes modulating adaptive immunity
423 through T cell activation and signaling (e.g., *PTPN22*^{52,53}, *TAGAP*⁵⁴⁻⁵⁷, *CD226*⁵⁸⁻⁶⁰), B cell activation (e.g.,
424 *KIF5A/PIP4K2C*⁶¹⁻⁶³, *IL10*⁶⁴⁻⁶⁶), T helper cells (e.g., *STAT1-STAT4*⁶⁷⁻⁷², *PRKCQ*⁷³⁻⁷⁵), and regulatory T
425 cells^{57,76-79}, are involved. Variation has also been shown in genes impacting innate immunity, which may
426 play an important role in viral or microbial infections through the pathogen recognition receptor pathways,

427 transcription factors, apoptosis, autophagy, and immune-complex clearance. A cross-disease genomic
428 analysis of nine AIDs, such as T1D, RA, SLE, suggested that the top prioritized genes of each analyzed
429 AID converged on the same common pathways relevant to T cell activation and signaling, although distinct
430 genes were prioritized across diseases⁸⁰. Many of these works indicated that T1D shared biological
431 pathways with other AIDs, especially RA, CD, and SLE, which to some extent explained why they were
432 tightly associated with T1D in our analyses.

433 The interplay of HLA and non-HLA variants on disease susceptibility turned out to be complex and disease-
434 dependent. Especially, our HLA haplotype analysis showed that multiple T1D lead haplotypes exerted
435 strong effects on other AIDs, but largely in a disease-dependent manner. The HLA effects regarding other
436 AIDs and T1D might also be in opposite directions as illustrated by some HLA haplotypes. For example,
437 being the most protective haplotype of T1D, *DRB1*15:01-DQAI*01:02-DQB1*06:02* meanwhile strongly
438 increased the risk of MS. This opposite pleiotropic HLA was consistently observed by multi-allelic HLA
439 PGS. However, in non-HLA regions, we observed no link between T1D and MS from either genetic
440 correlation or pTDT. IBD had a similar pattern in these analyses. These findings, taken together, can point
441 to potential explanations for the discrepancies observed between the epidemiological evidence of familial
442 aggregation and the genetic results. In particular, our results suggest that HLA risk factors for IBD and MS
443 are protective of T1D, while their non-HLA risk factors are not associated with T1D. The overall lack of
444 familial aggregation suggested that this HLA protective effect could be counterbalanced by other non-
445 captured genetic effects or by environmental effects shared within the family.

446 Most AIDs are more prevalent in women than in men and disease progression and severity could also differ
447 between the two sexes^{37,81}. T1D is an exception in that a slight majority of at least pediatric cases are boys
448 and twice as many fathers than mothers of the patients with T1D also have T1D^{37,38}. In this study, we also
449 observed that among individuals with T1D, men had a higher risk of having offspring with T1D than women,
450 while significant sex-specific associations affecting transmission were overall not observed among parental
451 AIDs. Further, when at least one parent had T1D, especially if it was the father, the male offspring would
452 have a higher risk of developing T1D than the female offspring. To date, the precise cause of a higher rate
453 of T1D transmission from fathers than mothers has not been identified, but potential hypotheses largely
454 revolve around in utero exposure to different aspects of maternal T1D and or its consequences e.g. exposure
455 to maternal hyperglycemia, exogenous insulin, maternal islet antibodies, and maternal enterovirus
456 antibodies³⁸. It has also been proposed that a protective effect of maternal T1D could be mediated by an
457 *E. coli* dominant maternal microbiome, which fosters the development of neonatal T cells in the infant⁸²⁻⁸⁵.

458 Previous studies also showed that people with AIDs, are less likely to have children⁸⁶. This is likely to
459 reflect multiple factors, including a higher rate of miscarriage amongst women with AIDs including T1D,
460 amongst whom the rate of miscarriage is 15-30% higher than in the background population⁸⁷⁻⁹⁰. In some
461 cases, individuals with AIDs may choose not have children due to a fear of transmitting disease to offspring.
462 Being able to estimate early T1D risk trajectories for a couple's offspring, utilizing their PGS and AID
463 disease status, will equip them with more accurate information to make such decisions around family
464 planning, and provide reassurance to those whose offspring have a low predicted risk of T1D. For
465 individuals with offspring predicted to be at high risk of disease, recent advances in T1D research may
466 mean that targeted screening and/or the option of early preventative therapies may be a realistic option.

467 So far, we are not aware of any previous study using parental PGS in estimating an AID. We introduced a
468 new approach to calculate PGSs for diseases that have a strong HLA signal, by training different models
469 for HLA and non-HLA regions and combining them based on their expected contribution to the disease.
470 We showed that parental PGS for T1D is strongly predictive of T1D risk before age of 20 and that parental
471 PGS for other AIDs does not help in better predicting T1D. We showed that a T1D PGS that represents the
472 genetic risk factors from parents is positively associated with developing T1D in offspring, especially sons.
473 The cumulative risk trajectories start to plateau around the age of 14-16, especially for children with high

474 parental PGS (top 10%), which is consistent with the incidence peak at 10-14 years reported in previous
475 studies⁵⁻⁷. The effect of high parental PGS on T1D risk in children was in many instances higher than a
476 diagnosis of T1D in their mother or father. For example, having a mother with T1D resulted in a cumulative
477 T1D risk of 1.49% by age 20, while the top 10% of parents with the highest PGS had a cumulative risk of
478 2.88%. Considering that only 0.15% of mothers with children had T1D in Finland, the PGS provides
479 additional value on top of family history. These results indicate the intriguing possibility of considering
480 parental PGS, in conjunction with clinical diagnoses, to inform parents about T1D risk in their offspring.

481 This study also has several limitations. One limitation was that AID diagnoses for some patients might not
482 be well captured in the healthcare register. For example, the specific ICD-9 code for T1D was not
483 implemented until 1987 although the Care Register for Finnish Health Care began as early as in 1969.
484 Another limitation was that rather than being nationally representative, the amalgamated Finnish biobanks
485 represent a collection of biobanks with diverse methods of data collection. Some of the diseases might have
486 a higher prevalence in the Finnish biobanks compared to the nationally representative FinRegistry.
487 Although these aspects might limit the generalizability of the results, we noted that our findings in the
488 Finnish biobanks were consistent with those we observed from FinRegistry and matched well to previous
489 studies^{26,29,31,80,81}. Thirdly, rather than extending the analysis to siblings or other relatives, in this study, we
490 focused on parents and offspring. We note the HLA and non-HLA association patterns with T1D across
491 AIDs would be similar to what we observed here although the effect sizes would differ. Finally, while many
492 risk factors (e.g., socioeconomic factors, genetic nurture) might have an impact on the development of T1D,
493 our study mainly focused on family history of diseases and genetic factors by controlling these factors with
494 delicate analytical designs.

495 In conclusion, our results, while confirming the existence of general familial aggregation of AIDs and T1D,
496 highlight a substantial heterogeneity in the impact of different AIDs on T1D. This heterogeneity is partially
497 explained by different genetic effects within and outside the HLA regions and demonstrated two different
498 transmission patterns regarding shared genetic liabilities in HLA and non-HLA. Overall, genetic effects
499 inside and outside HLA regions are consistent with observational analyses, but in the case of IBD and MS
500 we revealed unexpected divergence between the genetic effects and epidemiological observations.
501 Investigating the mechanisms behind these findings may provide valuable insights into the origins of T1D
502 and the etiology of the familial aggregations of AIDs. Moreover, the relative contribution of HLA and non-
503 HLA genetic risk can be leveraged to create powerful PGSs that can complement a family history of AIDs
504 as a tool to inform parents about expected T1D risk in their offspring.

505

506 **References**

- 1 Kavoura, Fotini K., and John PA Ioannidis. "CTLA-4 gene polymorphisms and susceptibility to type 1 diabetes mellitus: a HuGE Review and meta-analysis." *American journal of epidemiology* 162.1 (2005): 3-16.
- 2 Pociot, F., and M. F. McDermott. "Genetics of type 1 diabetes mellitus." *Genes & Immunity* 3.5 (2002): 235-249.
- 3 Åkerblom, Hans K., et al. "Environmental factors in the etiology of type 1 diabetes." *American journal of medical genetics* 115.1 (2002): 18-29.
- 4 Knip, Mikael, et al. "Environmental triggers and determinants of type 1 diabetes." *Diabetes* 54.suppl_2 (2005): S125-S136.
- 5 Maahs, David M., et al. "Epidemiology of type 1 diabetes." *Endocrinology and Metabolism Clinics* 39.3 (2010): 481-497.
- 6 Gregory, Gabriel A., et al. "Global incidence, prevalence, and mortality of type 1 diabetes in 2021 with projection to 2040: a modelling study." *The Lancet Diabetes & Endocrinology* 10.10 (2022): 741-760.

- 7 Hormazábal-Aguayo, Ignacio, et al. "Incidence of type 1 diabetes mellitus in children and adolescents under 20 years of age across 55 countries from 2000 to 2022: A systematic review with meta-analysis." *Diabetes/Metabolism Research and Reviews* (2023): e3749.
- 8 Noble, Janelle A., et al. "The role of HLA class II genes in insulin-dependent diabetes mellitus: molecular analysis of 180 Caucasian, multiplex families." *American journal of human genetics* 59.5 (1996): 1134.
- 9 Thomson G, Robinson WP, Kuhner MK et al. Genetic heterogeneity, modes of inheritance, and risk estimates for a joint study of Caucasians with insulin-dependent diabetes mellitus *Am J Hum Genet* 1988 43: 799–816
- 10 Lambert, A. Paul, et al. "Absolute risk of childhood-onset type 1 diabetes defined by human leukocyte antigen class II genotype: a population-based study in the United Kingdom." *The Journal of Clinical Endocrinology & Metabolism* 89.8 (2004): 4037-4043.
- 11 Luo, Shuoming, et al. "HLA genetic discrepancy between latent autoimmune diabetes in adults and type 1 diabetes: LADA China Study No. 6." *The Journal of Clinical Endocrinology & Metabolism* 101.4 (2016): 1693-1700.
- 12 Ikegami, Hiroshi, et al. "Genetics of type 1 diabetes: similarities and differences between Asian and Caucasian populations." *Annals of the New York Academy of Sciences* 1079.1 (2006): 51-59.
- 13 Kawabata, Yumiko, et al. "Asian-specific HLA haplotypes reveal heterogeneity of the contribution of HLA-DR and-DQ haplotypes to susceptibility to type 1 diabetes." *Diabetes* 51.2 (2002): 545-551.
- 14 Rojas-Villarraga, Adriana, Diana Botello-Corzo, and Juan-Manuel Anaya. "HLA-Class II in Latin American patients with type 1 diabetes." *Autoimmunity reviews* 9.10 (2010): 666-673.
- 15 Todd, John A., et al. "Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes." *Nature genetics* 39.7 (2007): 857-864.
- 16 Barrett, Jeffrey C., et al. "Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes." *Nature genetics* 41.6 (2009): 703-707.
- 17 Evangelou, Marina, et al. "A method for gene-based pathway analysis using genomewide association study summary statistics reveals nine new type 1 diabetes associations." *Genetic epidemiology* 38.8 (2014): 661-670.
- 18 Robertson, Catherine C., et al. "Fine-mapping, trans-ancestral and genomic analyses identify causal variants, cells, genes and drug targets for type 1 diabetes." *Nature genetics* 53.7 (2021): 962-971.
- 19 Winkler, Christiane, et al. "Feature ranking of type 1 diabetes susceptibility genes improves prediction of type 1 diabetes." *Diabetologia* 57 (2014): 2521-2529.
- 20 Sharp, Seth A., et al. "Development and standardization of an improved type 1 diabetes genetic risk score for use in newborn screening and incident diagnosis." *Diabetes care* 42.2 (2019): 200-207.
- 21 Padilla-Martínez, Felipe, et al. "Systematic review of polygenic risk scores for type 1 and type 2 diabetes." *International journal of molecular sciences* 21.5 (2020): 1703.
- 22 Patel, Kashyap A., et al. "Type 1 diabetes genetic risk score: a novel tool to discriminate monogenic and type 1 diabetes." *Diabetes* 65.7 (2016): 2094-2099.
- 23 Yaghootkar, H., et al. "Type 1 diabetes genetic risk score discriminates between monogenic and Type 1 diabetes in children diagnosed at the age of < 5 years in the Iranian population." *Diabetic Medicine* 36.12 (2019): 1694-1702.
- 24 Billings, Liana K., et al. "Utility of polygenic scores for differentiating diabetes diagnosis among patients with atypical phenotypes of diabetes." *The Journal of Clinical Endocrinology & Metabolism* 109.1 (2024): 107-113.
- 25 Nielsen, Nete M., et al. "Type 1 diabetes and multiple sclerosis: a Danish population-based cohort study." *Archives of neurology* 63.7 (2006): 1001-1004.
- 26 Hemminki, K., et al. "Familial association between type 1 diabetes and other autoimmune and related diseases." *Diabetologia* 52.9 (2009): 1820-1828.
- 27 Yen, Fu-Shun, et al. "Maternal autoimmune disease associated with a higher risk of offspring with type 1 diabetes: A nationwide mother-child cohort study in Taiwan." *Diabetes & Metabolism* (2023): 101443.
- 28 Vyse, Timothy J., and John A. Todd. "Genetic analysis of autoimmune disease." *Cell* 85.3 (1996): 311-318.
- 29 Parkkola, Anna, et al. "Extended family history of autoimmune diseases and phenotype and genotype of children with newly diagnosed type 1 diabetes." *European Journal of Endocrinology* 169.2 (2013): 171-178.
- 30 Wägner, Ana M., et al. "Predictors of associated autoimmune diseases in families with type 1 diabetes: results from the Type 1 Diabetes Genetics Consortium." *Diabetes/metabolism research and reviews* 27.5 (2011): 493-498.

- 31 Werme, Josefin, et al. "An integrated framework for local genetic correlation analysis." *Nature genetics* 54.3 (2022): 274-282.
- 32 Lettre, Guillaume, and John D. Rioux. "Autoimmune diseases: insights from genome-wide association studies." *Human molecular genetics* 17.R2 (2008): R116-R121.
- 33 Viippola, Essi, et al. "Data Resource Profile: Nationwide registry data for high-throughput epidemiology and machine learning (FinRegistry)." *International Journal of Epidemiology*(2023): dyad091.
- 34 Kurki, Mitja I., et al. "FinnGen provides genetic insights from a well-phenotyped isolated population." *Nature* 613.7944 (2023): 508-518.
- 35 Patterson, Christopher C., et al. "Incidence trends for childhood type 1 diabetes in Europe during 1989–2003 and predicted new cases 2005–20: a multicentre prospective registration study." *The Lancet* 373.9680 (2009): 2027-2033.
- 36 Parkkola, Anna, et al. "Extended family history of type 1 diabetes and phenotype and genotype of newly diagnosed children." *Diabetes care* 36.2 (2013): 348-354.
- 37 Harjutsalo, Valma, Antti Reunanen, and Jaakko Tuomilehto. "Differential transmission of type 1 diabetes from diabetic fathers and mothers to their offspring." *Diabetes* 55.5 (2006): 1517-1524.
- 38 Allen, Lowri A., et al. "Maternal type 1 diabetes and relative protection against offspring transmission." *The Lancet Diabetes & Endocrinology* (2023).
- 39 Erlich, Henry, et al. "HLA DR-DQ haplotypes and genotypes and type 1 diabetes risk: analysis of the type 1 diabetes genetics consortium families." *Diabetes* 57.4 (2008): 1084-1092.
- 40 Ritari, Jarmo, et al. "Increasing accuracy of HLA imputation by a population-specific reference panel in a FinnGen biobank cohort." *NAR genomics and bioinformatics* 2.2 (2020): lqaa030.
- 41 Ramagopalan, Sreeram V., and George C. Ebers. "Multiple sclerosis: major histocompatibility complexity and antigen presentation." *Genome medicine* 1.11 (2009): 1-5.
- 42 Arrieta-Bolaños, Esteban, Diana Iraíz Hernández-Zaragoza, and Rodrigo Barquera. "An HLA map of the world: A comparison of HLA frequencies in 200 worldwide populations reveals diverse patterns for class I and class II." *Frontiers in Genetics* 14 (2023): 866407.
- 43 Bulik-Sullivan, Brendan K., et al. "LD Score regression distinguishes confounding from polygenicity in genome-wide association studies." *Nature genetics* 47.3 (2015): 291-295.
- 44 Bulik-Sullivan, Brendan, et al. "An atlas of genetic correlations across human diseases and traits." *Nature genetics* 47.11 (2015): 1236-1241.
- 45 Weiner, Daniel J., et al. "Polygenic transmission disequilibrium confirms that common and rare variation act additively to create risk for autism spectrum disorders." *Nature genetics* 49.7 (2017): 978-985.
- 46 Ge, Tian, et al. "Polygenic prediction via Bayesian regression and continuous shrinkage priors." *Nature communications* 10.1 (2019): 1776.
- 47 Anaya, Juan-Manuel, et al. "Autoimmunity: from bench to bedside [Internet]." (2013).
- 48 Ritari, J., Koskela, S., Hyvärinen, K. and Partanen, J., 2022. HLA-disease association and pleiotropy landscape in over 235,000 Finns. *Human Immunology*, 83(5), pp.391-398.
- 49 Serrano, Norma C., Paula Millan, and María-Carolina Páez. "Non-HLA associations with autoimmune diseases." *Autoimmunity reviews* 5.3 (2006): 209-214.
- 50 Simmonds, M. J., and S. C. L. Gough. "The HLA region and autoimmune disease: associations and mechanisms of action." *Current genomics* 8.7 (2007): 453-465.
- 51 Shiina, Takashi, et al. "The HLA genomic loci map: expression, interaction, diversity and disease." *Journal of human genetics* 54.1 (2009): 15-39.
- 52 Bottini, Nunzio, et al. "A functional variant of lymphoid tyrosine phosphatase is associated with type I diabetes." *Nature genetics* 36.4 (2004): 337-338.
- 53 Begovich, Ann B., et al. "A missense single-nucleotide polymorphism in a gene encoding a protein tyrosine phosphatase (PTPN22) is associated with rheumatoid arthritis." *The American Journal of Human Genetics* 75.2 (2004): 330-337.
- 54 Hunt, Karen A., et al. "Newly identified genetic risk variants for celiac disease related to the immune response." *Nature genetics* 40.4 (2008): 395-402.
- 55 Smyth, Deborah J., et al. "Shared and distinct genetic variants in type 1 diabetes and celiac disease." *New England Journal of Medicine* 359.26 (2008): 2767-2777.
- 56 Raychaudhuri, Soumya, et al. "Genetic variants at CD28, PRDM1 and CD2/CD58 are associated with rheumatoid arthritis risk." *Nature genetics* 41.12 (2009): 1313-1318.

- 57 Todd, John A., et al. "Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes." *Nature genetics* 39.7 (2007): 857-864.
- 58 Hafler, Jason P., et al. "CD226 Gly307Ser association with multiple autoimmune diseases." *Genes & Immunity* 10.1 (2009): 5-10.
- 59 Löfgren, Sara E., et al. "A 3'-untranslated region variant is associated with impaired expression of CD226 in T and natural killer T cells and is associated with susceptibility to systemic lupus erythematosus." *Arthritis & Rheumatism* 62.11 (2010): 3404-3414.
- 60 Qiu, Zhi-Xin, et al. "CD226 Gly307Ser association with multiple autoimmune diseases: a meta-analysis." *Human immunology* 74.2 (2013): 249-255.
- 61 Raychaudhuri, Soumya, et al. "Common variants at CD40 and other loci confer risk of rheumatoid arthritis." *Nature genetics* 40.10 (2008): 1216-1223.
- 62 Fung, E. Y. M. G., et al. "Analysis of 17 autoimmune disease-associated variants in type 1 diabetes identifies 6q23/TNFAIP3 as a susceptibility locus." *Genes & Immunity* 10.2 (2009): 188-191.
- 63 Cohen, Jose E., Svetlana Boitsova, and Eyal Itshayek. "Cerebral venous sinus thrombosis." *Isr Med Assoc J* 11.11 (2009): 685-688.
- 64 Franke, Andre, et al. "Sequence variants in IL10, ARPC2 and multiple other loci contribute to ulcerative colitis susceptibility." *Nature genetics* 40.11 (2008): 1319-1323.
- 65 Barrett, Jeffrey C., et al. "Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes." *Nature genetics* 41.6 (2009): 703-707.
- 66 Gateva, Vesela, et al. "A large-scale replication study identifies TNIP1, PRDM1, JAZF1, UHRF1BP1 and IL10 as risk loci for systemic lupus erythematosus." *Nature genetics* 41.11 (2009): 1228-1233.
- 67 International Consortium for Systemic Lupus Erythematosus Genetics (SLEGEM), et al. "Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PXX, KIAA1542 and other loci." *Nature genetics* 40.2 (2008): 204-210.
- 68 Martinez, A., et al. "Association of the STAT4 gene with increased susceptibility for some immune-mediated diseases." *Arthritis & Rheumatism* 58.9 (2008): 2598-2602.
- 69 Tong, Guanghui, et al. "Association between polymorphism in STAT4 gene and risk of rheumatoid arthritis: a meta-analysis." *Human immunology* 74.5 (2013): 586-592.
- 70 Liang, Ya-ling, et al. "Association of STAT4 rs7574865 polymorphism with autoimmune diseases: a meta-analysis." *Molecular biology reports* 39 (2012): 8873-8882.
- 71 Barrett, Jeffrey C., et al. "Genome-wide association defines more than 30 distinct susceptibility loci for Crohn's disease." *Nature genetics* 40.8 (2008): 955-962.
- 72 Jakkula, Eveliina, et al. "Genome-wide association study in a high-risk isolate for multiple sclerosis reveals associated variants in STAT3 gene." *The American Journal of Human Genetics* 86.2 (2010): 285-291.
- 73 Cooper, Jason D., et al. "Meta-analysis of genome-wide association study data identifies additional type 1 diabetes risk loci." *Nature genetics* 40.12 (2008): 1399-1401.
- 74 Raychaudhuri, Soumya, et al. "Genetic variants at CD28, PRDM1 and CD2/CD58 are associated with rheumatoid arthritis risk." *Nature genetics* 41.12 (2009): 1313-1318.
- 75 Hinks, Anne, et al. "Overlap of disease susceptibility loci for rheumatoid arthritis and juvenile idiopathic arthritis." *Annals of the rheumatic diseases* 69.6 (2010): 1049-1053.
- 76 Adamovic, Svetlana, et al. "Association study of IL2/IL21 and FcγRIIa: significant association with the IL2/IL21 region in Scandinavian coeliac disease families." *Genes & Immunity* 9.4 (2008): 364-367.
- 77 Maiti, Amit K., et al. "Confirmation of an association between rs6822844 at the IL2–IL21 region and multiple autoimmune diseases: evidence of a general susceptibility locus." *Arthritis & Rheumatism: Official Journal of the American College of Rheumatology* 62.2 (2010): 323-329.
- 78 Daha, Nina A., et al. "Confirmation of STAT4, IL2/IL21, and CTLA4 polymorphisms in rheumatoid arthritis." *Arthritis & Rheumatism* 60.5 (2009): 1255-1260.
- 79 Hinks, A., et al. "Association of the AFF3 gene and IL2/IL21 gene region with juvenile idiopathic arthritis." *Genes & Immunity* 11.2 (2010): 194-198.
- 80 Demela, Pietro, Nicola Pirastu, and Blagoje Soskic. "Cross-disorder genetic analysis of immune diseases reveals distinct gene associations that converge on common pathways." *Nature Communications* 14.1 (2023): 2743.
- 81 Conrad, Nathalie, et al. "Incidence, prevalence, and co-occurrence of autoimmune disorders over time and by age, sex, and socioeconomic status: a population-based cohort study of 22 million individuals in the UK." *The Lancet* 401.10391 (2023): 1878-1890.

- 82 Warram, James H., et al. "Differences in risk of insulin-dependent diabetes in offspring of diabetic mothers and diabetic fathers." *New England Journal of Medicine* 311.3 (1984): 149-152.
- 83 Yang, Siyoung, et al. "Regulatory T cells generated early in life play a distinct role in maintaining self-tolerance." *Science* 348.6234 (2015): 589-594.
- 84 Roth-Schulze, Alexandra J., et al. "Type 1 diabetes in pregnancy is associated with distinct changes in the composition and function of the gut microbiome." *Microbiome* 9 (2021): 1-21.
- 85 Gao, Yuan, et al. "Maternal gut microbiota during pregnancy and the composition of immune cells in infancy." *Frontiers in Immunology* 13 (2022): 986340.
- 86 Liu, Aoxing, et al. "Evidence from Finland and Sweden on the relationship between early-life diseases and lifetime childlessness in men and women." *Nature Human Behaviour* (2023): 1-12.
- 87 Miodovnik, Menachem, et al. "Spontaneous abortion among insulin-dependent diabetic women." *American journal of obstetrics and gynecology* 150.4 (1984): 372-376.
- 88 Dorman, Janice S., et al. "Temporal trends in spontaneous abortion associated with type 1 diabetes." *Diabetes research and clinical practice* 43.1 (1999): 41-47.
- 89 Lorenzen, T., et al. "A population-based survey of frequencies of self-reported spontaneous and induced abortions in Danish women with Type 1 diabetes mellitus." *Diabetic medicine* 16.6 (1999): 472-476.
- 90 Penney, Gillian C., et al. "Outcomes of pregnancies in women with type 1 diabetes in Scotland: a national population-based study." *BJOG: an International Journal of Obstetrics & Gynaecology* 110.3 (2003): 315-318.

507
508
509
510

Acknowledgments

511 We thank the entire FinRegistry and FinnGen team for making the data available for this study. We also acknowledge
512 Richard Oram (Institute of Biomedical and Clinical Science, University of Exeter Medical School, Exeter, UK) for
513 his discussion on this study, Samuel Jones (Institute for Molecular Medicine Finland (FIMM), Finland) for discussion
514 on HLA PGS analysis, Bradley Jermy (FIMM, Finland) for discussion on genetic correlation analysis, Mary P Reeve
515 (FIMM, Finland) for discussion on disease definitions, Om Dwivedi (FIMM, Finland) for discussion on T1D PGS,
516 and Alessio Gerussi (University of Milano-Bicocca, Italy) for discussion on results from a clinical standpoint. The
517 We want to acknowledge the participants and investigators of FinnGen study. The FinnGen project is funded by two
518 grants from Business Finland (HUS 4685/31/2016 and UH 4386/31/2016) and the following industry partners: AbbVie
519 Inc., AstraZeneca UK Ltd, Biogen MA Inc., Bristol Myers Squibb (and Celgene Corporation & Celgene International
520 II Sàrl), Genentech Inc., Merck Sharp & Dohme LCC, Pfizer Inc., GlaxoSmithKline Intellectual Property
521 Development Ltd., Sanofi US Services Inc., Maze Therapeutics Inc., Janssen Biotech Inc, Novartis Pharma AG, and
522 Boehringer Ingelheim International GmbH. Following biobanks are acknowledged for delivering biobank samples to
523 FinnGen: Auria Biobank (www.auria.fi/biopankki), THL Biobank (www.thl.fi/biobank), Helsinki Biobank
524 (www.helsinginbiopankki.fi), Biobank Borealis of Northern Finland (<https://www.ppshep.fi/Tutkimus-japetus/Biopankki/Pages/Biobank-Borealis-briefly-in-English.aspx>), Finnish Clinical Biobank Tampere
525 (www.tays.fi/en-US/Research_and_development/Finnish_Clinical_Biobank_Tampere), Biobank of Eastern Finland
526 (www.ita-suomenbiopankki.fi/en), Central Finland Biobank (www.ksshp.fi/fi-FI/Potilaille/Biopankki), Finnish Red
527 Cross Blood Service Biobank (www.veripalvelu.fi/verenluovutus/biopankkitoiminta), Terveystalo Biobank
528 (www.terveystalo.com/fi/Yritystietoa/Terveystalo-Biopankki/Biopankki/) and Arctic Biobank
529 (<https://www oulu.fi/en/university/faculties-and-units/faculty-medicine/northern-finland-birth-cohorts-and-arctic-biobank>). All Finnish Biobanks are members of BBMRI.fi infrastructure (www.bbmri.fi). Finnish Biobank
530 Cooperative -FINBB (<https://finbb.fi/>) is the coordinator of BBMRI-ERIC operations in Finland. The Finnish biobank
531 data can be accessed through the Fingenious® services (<https://site.fingenious.fi/en/>) managed by FINBB.
532
533

534
535

Author contributions

536 FW, AL, TT, and AG designed the study and wrote the manuscript, with comments from all authors. FW processed
537 the registry data, conducted the statistical analyses, and generated all figures and tables; AL and FW preprocessed the
538 FinRegistry pedigree data and the FinRegistry team preprocessed all other registry data of FinRegistry; the FinnGen
539 data team preprocessed the FinnGen data. TT, PV, and the FinnGen clinical team defined disease endpoints; TT

540 oversaw the project and interpreted the findings from a clinical standpoint. AL, ZY, SJ, and AG advised the registry-
541 based epidemiological analyses; AL, ZY, and AG advised genetic analyses; AL, SK, JR, JP, and TT advised HLA
542 haplotype analyses. AG and TT supervised the study. MP is the principal investigator of the FinRegistry project. All
543 authors discussed the results, revised the manuscript, and had final responsibility for the decision to submit for
544 publication.

545 546 **Funding**

547 FW was funded by the University of Helsinki and the University of Edinburgh joint PhD program in Human
548 Genomics. AG was supported by the European Research Council (ERC) under the European Union's Horizon 2020
549 research and innovation programme [grant number 945733], starting grant AI-Prevent. AG was supported by the
550 Academy of Finland (grant no. 323116). This project also received funding from the European Union's Horizon 2020
551 Research and Innovation Programme under grant agreement no. 101016775. TT was supported by grants from
552 Folkhälsan Research Foundation, The Academy of Finland (336822, 312072 and 336826), and the University of
553 Helsinki.

554 555 **Competing interests**

556 AG is the founder of Real World Genetics.

557 RO holds a UK Medical Research Council Institutional Confidence in Concept Grant to develop a 10 SNP biochip
558 T1D genetic test in collaboration with Randox.

559 560 561 **Ethics statement**

562 FinRegistry is a collaboration project of the Finnish Institute for Health and Welfare (THL) and the Data Science
563 Genetic Epidemiology research group at the Institute for Molecular Medicine Finland (FIMM), University of Helsinki.
564 The FinRegistry project has received approvals for data access from THL (THL/1776/6.02.00/2019 and subsequent
565 amendments), Digital and Population Data Services Agency (VRK/5722/2019–2), Finnish Center for Pension
566 (ETK/SUTI 22003) and Statistics Finland (TK-53–1451-19). The FinRegistry project has received IRB approval from
567 THL (Kokous 7/2019).

568 The use of the biobank donor samples is in accordance with the biobank consent and meets the requirements of the
569 Finnish Biobank Act 688/2012. Alternatively, separate research cohorts, collected prior the Finnish Biobank Act came
570 into effect (in September 2013) and start of FinnGen (August 2017), were collected based on study-specific consents
571 and later transferred to the Finnish biobanks after approval by Fimea (Finnish Medicines Agency), the National
572 Supervisory Authority for Welfare and Health. Recruitment protocols followed the biobank protocols approved by
573 Fimea. The Coordinating Ethics Committee of the Hospital District of Helsinki and Uusimaa (HUS) statement number
574 for the FinnGen study is Nr HUS/990/2017.

575 The FinnGen study is approved by Finnish Institute for Health and Welfare (permit numbers: THL/2031/6.02.00/2017,
576 THL/1101/5.05.00/2017, THL/341/6.02.00/2018, THL/2222/6.02.00/2018, THL/283/6.02.00/2019,
577 THL/1721/5.05.00/2019 and THL/1524/5.05.00/2020), Digital and population data service agency (permit numbers:
578 VRK43431/2017-3, VRK/6909/2018-3, VRK/4415/2019-3), the Social Insurance Institution (permit numbers: KELA
579 58/522/2017, KELA 131/522/2018, KELA 70/522/2019, KELA 98/522/2019, KELA 134/522/2019, KELA
580 138/522/2019, KELA 2/522/2020, KELA 16/522/2020), Findata permit numbers THL/2364/14.02/2020,
581 THL/4055/14.06.00/2020, THL/3433/14.06.00/2020, THL/4432/14.06/2020, THL/5189/14.06/2020,
582 THL/5894/14.06.00/2020, THL/6619/14.06.00/2020, THL/209/14.06.00/2021, THL/688/14.06.00/2021,
583 THL/1284/14.06.00/2021, THL/1965/14.06.00/2021, THL/5546/14.02.00/2020, THL/2658/14.06.00/2021,
584 THL/4235/14.06.00/2021, Statistics Finland (permit numbers: TK-53-1041-17 and TK/143/07.03.00/2020 (earlier

585 TK-53-90-20) TK/1735/07.03.00/2021, TK/3112/07.03.00/2021) and Finnish Registry for Kidney Diseases
586 permission/extract from the meeting minutes on 4th July 2019.

587 The Biobank Access Decisions for FinnGen samples and data utilized in FinnGen Data Freeze 11 include: THL
588 Biobank BB2017_55, BB2017_111, BB2018_19, BB_2018_34, BB_2018_67, BB2018_71, BB2019_7, BB2019_8,
589 BB2019_26, BB2020_1, BB2021_65, Finnish Red Cross Blood Service Biobank 7.12.2017 (000-2018), Helsinki
590 Biobank HUS/359/2017, HUS/248/2020, HUS/430/2021 §28, §29, HUS/150/2022 §12, §13, §14, §15, §16, §17, §18,
591 §23, §58 and §59, Auria Biobank AB17-5154 and amendment #1 (August 17 2020) and amendments BB_2021-0140,
592 BB_2021-0156 (August 26 2021, Feb 2 2022), BB_2021-0169, BB_2021-0179, BB_2021-0161, AB20-5926 and
593 amendment #1 (April 23 2020) and it's modification (Sep 22 2021), BB_2022-0262, BB_2022-0256, Biobank
594 Borealis of Northern Finland 2017_1013, 2021_5010, 2021_5018, 2021_5015, 2021_5015 Amendment, 2021_5023,
595 2021_5023 Amendment, 2021_5017, 2022_6001, 2022_6006 Amendment, BB22-0067, 2022_0262, Biobank of
596 Eastern Finland 1186/2018 and amendment 22§/2020, 53§/2021, 13§/2022, 14§/2022, 15§/2022, 27§/2022, 28§/2022,
597 29§/2022, 33§/2022, 35§/2022, 36§/2022, 37§/2022, 39§/2022, 7§/2023, Finnish Clinical Biobank Tampere MH0004
598 and amendments (21.02.2020 & 06.10.2020), 8§/2021, 9§/2021, §9/2022, §10/2022, §12/2022, 13§/2022, §20/2022,
599 §21/2022, §22/2022, §23/2022, 28§/2022, 29§/2022, 30§/2022, 31§/2022, 32§/2022, 38§/2022, 40§/2022, 42§/2022,
600 1§/2023, Central Finland Biobank 1-2017, BB_2021-0161, BB_2021-0169, BB_2021-0179, BB_2021-0170,
601 BB_2022-0256, and Terveystalo Biobank STB 2018001 and amendment 25th Aug 2020, Finnish Hematological
602 Registry and Clinical Biobank decision 18th June 2021, Arctic biobank P0844: ARC_2021_1001.

603
604
605
606

607 **Methods**

608 **Data sources**

609 Most of our analyses were conducted by leveraging two large datasets in Finland – FinRegistry³³ and
610 FinnGen³⁴ (**Table S3.1.1**).

611 FinRegistry is a national registry database in Finland combining disease registries with comprehensive
612 records on individuals' demographics, socioeconomic status, death, drug purchases, prescriptions, and
613 administrative information. It includes all the residents in Finland who were alive on 1st January 2010, and
614 their first-degree relatives (in total approximately 7.2 million individuals). 572,640 individuals were
615 excluded as they migrated in or out of Finland by 2019, remaining 3,412,326 individuals with both father
616 and mother information.

617 FinnGen (R11) is a national research project using samples and data from Finnish biobanks, comprising
618 genome-wide genotype data and healthcare registry data for approximately 0.47 million Finnish biobank
619 donors. After excluding 9,196 individuals who had migrated in or out of Finland by 2019, removing 10,588
620 individuals without imputed HLA data and 8,627 individuals who have other missing values in the cohort,
621 we included 3,668 T1D cases and 441,602 controls without T1D included in FinnGen and observed lifetime
622 prevalence and number of cases in FinnGen for each selected AID. The study population features slightly
623 more women (N=251,294, 56.4%).

624 By using the datasets from the same country, we alleviate concerns about potential bias caused by
625 differences in genetic background, healthcare system, or socio-cultural context.

626

627 **Finnish population-based case-control study**

628 The epidemiology association between AIDs in parents and T1D in their children was investigated by a
629 case-control study within FinRegistry. To include most of the individuals and their parents in this analysis,
630 we selected all individuals born in Finland between 1960 and 1999 who had both father's and mother's
631 information available. The follow-up period started either from the date they were born or from the year
632 the codes for our selected diseases were included in national patient register and ended on 31 December
633 2019, which is the latest follow-up date in FinRegistry. We then restricted our analysis to offspring whose
634 father was born between 1917 and 1976 and whose mother was born between 1922 and 1976. Finally, we
635 removed those individuals who died during the follow-up period.

636 T1D cases were individuals diagnosed with T1D before the age of 40 years. For each trio with T1D-affected
637 children, three controls without T1D diagnosis were matched by following variables: sex, birth year (five-
638 year bin, 8 levels), and birthplace for the child (19 levels), birth year of father (five-year bin, 12 levels),
639 birth year of mother (five-year bin, 11 levels), and family size (represented by the number of siblings: 0, 1,
640 2, 3, >=4). The differences in socio-demographic characteristics between cases and controls, including
641 socio-economic status, marital status, number of offspring, occupation, education, and mother tongue, were
642 further examined to ensure that the two groups were comparable. Cohen's D^{91} (≥ 0.2) for ordinal
643 characteristics such as number of siblings and number of children and Chi-squared test ($P < 0.05$) for
644 categorical variables such as education and marital status were used to test whether the differences are
645 minor.

646 Exposures of interest were all the parental endpoints related to AIDs mainly defined by Finnish versions 8-
647 10 of International Classification of Diseases (ICD) (**Supplementary notes 2.1**). AIDs with fewer than 50
648 cases among parents in the final study population in the case-control study were filtered out from the list.
649 We then excluded AIDs with an unclear definition or AID codes that were subtypes of other codes. The
650 final AID list was used across all analyses in this study.

651 A conditional logistic regression, adjusted by birth year, birth year of father, birth year of mother, and the
652 exact number of siblings, was applied to estimate the effect of having a given AID in parents on the risk of
653 developing T1D in offspring. Bonferroni multiple testing correction was applied, and only the diseases
654 reached significant associations ($P \leq 0.05/26$) were included in the following analyses. To examine whether
655 the identified associations are shared between father and mother, we used conditional logistic regression
656 stratified by maternal AIDs status and paternal AIDs status. We also investigated whether parental AID
657 status can influence the onset age of disease in offspring by regressing the age at onset of T1D in offspring
658 on the AID status in their parents, with the same covariates described above adjusted. Statsmodels v.0.13.5⁹²
659 and SciPy v.1.7.3⁹³ were used for the analyses in this section in the context of Python v.3.7.11.

660

661 **Shared genetic components at a population level**

662 To explore, to what extent, the familial aggregations between AIDs and T1D identified in the population-
663 based case-control study are contributed by genetic factors, we conducted three analyses in both HLA
664 regions and non-HLA regions at individual level. Given the strong associations between HLA and AIDs⁴,
665 we first performed an HLA haplotype analysis and an HLA-based PGS analysis using imputed HLA alleles.
666 Then, we performed a genetic correlation analysis using variants in non-HLA regions.

667 *HLA haplotype analysis*

668 The HLA alleles were imputed by a population-specific reference panel using FinnGen data at the
669 individual level³⁷. We considered all the 187 unique available alleles on 10 classical genes in class I and II
670 regions for analysis, including 27 alleles for *HLA-A*, 40 for *HLA-B*, 23 for *HLA-C*, 24 for *HLA-DPB1*, 15

671 for *HLA-DQA1*, 16 for *HLA-DQB1*, 33 for *HLA-DRB1*, and 3 each for *HLA-DRB3*, *HLA-DRB4*, and *DRB5*
672 separately. According to previous studies^{8-10,36}, several HLA haplotypes in Class II have been significantly
673 associated with an increased risk of developing T1D and other AIDs. To understand whether HLA genetic
674 liability to T1D is associated with other AIDs, we conducted a haplotype analysis (**Figure S2.4.1**) and only
675 considered possible haplotypes from gene combination *HLA-DRB1-DQA1-DQB1*. We first removed
676 individuals with any ambiguous genotypes in the given haplotype. We define an ambiguous genotype as a
677 genotype with imputed allele dosage in [0.4, 0.6] or [1.4,1.6]. Next, we extracted the potential haplotypes
678 using the expectation–maximization algorithm and removed those haplotypes whose frequency is less than
679 0.005. We identified the most prevalent HLA haplotypes that were either positively or negatively associated
680 with T1D using a haplotype score test ($P < 1.0 \times 10^{-10}$), a statistical test to calculate the contribution of a
681 specific haplotype on a certain phenotype when the linkage phase is unknown⁹⁵. Multiple testing corrections
682 were applied for haplotype selection. We then only selected the three haplotypes with the highest P-value
683 and the three with the lowest. Afterward, a multivariate logistic regression was built to estimate the
684 associations between the remaining haplotypes and AID status, adjusting for age, sex and the first ten
685 principal components (PCs). Haplotype analysis was done in Python v.3.7.13 using statsmodels v.0.13.2⁹²
686 and in R v.4.2.2 using gap v.1.3-1⁹⁶.

687

688 *HLA multi-allelic score*

689 To tackle the complex linkage disequilibrium (LD) structure amongst HLA alleles, we introduced a PGS
690 method to construct PGS for T1D and other AIDs specific to HLA regions using FinnGen R11. To handle
691 the downstream analyses in later Panels, we considered all individuals of genotyped family trios as our
692 target population and the rest as a training set for weight calculation.

693 In the training set, to account for the complex LD and high polymorphism among HLA alleles, we applied
694 to calculate the weight for each HLA allele using weighted ridge classifiers with optimal regularization
695 strength, which can address this problem by imposing a penalty on the size of the coefficients. For
696 covariates, we adjusted for age, sex, and the first ten PCs. For the genotyped trios, we calculated PGS by
697 summing up all the allele dosage multiplied by its weight from the corresponding training set. After PGS
698 normalization, we first measured the partial correlation between each AID and its PGS. We dropped all the
699 AIDs whose partial correlation with PGS for the AIDs themselves was lower than 2%, to filter out diseases
700 with low sample sizes or with weak HLA signals. Next, we conducted logistic linear regression to examine
701 the association between PGS and T1D. The Bonferroni correction was applied to account for false positives
702 related to multiple comparisons. Weighted ridge regression and logistic regression were applied using
703 sklearn v.1.0.2 and statsmodels v.0.13.2⁹² in Python v.3.7.13 and partial correlation were calculated using
704 ppcor v1.1⁹⁷ in R v.4.3.2.

705

706 *non-HLA genetic correlation*

707 To evaluate the genetic overlap between T1D and other AIDs due to shared non-HLA effects, we utilized
708 largeset GWAS summary statistics from European ancestry individuals (**Table S2.3.1**) and estimated
709 genetic correlation in non-HLA regions using linkage disequilibrium score regressions (LDSC), which
710 capitalized the patterns of LD among common genetic variants of the whole genome except for the HLA
711 regions^{43,44}. To separate the contribution of HLA from non-HLA genetic signals, we excluded single-
712 nucleotide polymorphisms (SNPs) in the extended HLA regions on chromosome 6 (25–34 Mb). In a
713 sensitivity analysis, we further examined whether the genetic correlations changed significantly while
714 excluding all the SNPs on chromosome 6. We followed the suggested protocol for LDSC analyses from

715 <https://github.com/bulik/ldsc> and used the package LDSC v.1.0.1⁴³ in Python v.3.7.11. We considered
716 HapMap3⁹⁸ SNPs and constructed LD structures with European ancestry samples in 1000 Genomes
717 project⁹⁹.

718 719 **Analyses of inter-generation cross-trait transmission via polygenic transmission disequilibrium test**

720 To understand the transmission of parental AIDs to T1D in offspring, we further estimated the contribution
721 of parental transmitted and non-transmitted genetic liability to T1D in offspring in both non-HLA regions
722 and HLA regions using data in FinnGen. While offspring are expected to receive, on average, half of their
723 parents' risk alleles for a disease, some will over or under-inherit alleles associated with the disease, which
724 will impact their disease risk. To understand how much of the shared genetic liability is on average
725 transmitted to the next generation, we applied a polygenic transmission disequilibrium test (pTDT) to assess
726 whether the mean of the offspring PGS distribution is consistent with its parentally derived expected value⁴².

727
728 To construct non-HLA PGS, we used the same GWAS summary statistics used in genetic correlation
729 analysis and adopted PRS-CS⁴⁶ (**Table S2.3.2**). We included 12,563 genotyped trios where both parents
730 and at least one offspring were directly genotyped. We considered HLA PGSs and non-HLA PGSs that
731 were robustly associated with the corresponding AID ($P < 0.05/19$ and $|\rho| \geq 2\%$) (**Table S2.3.2**). This
732 resulted in ten diseases with both robust HLA PGS and non-HLA PGS: T1D, HYPO, CD, RA, SLE,
733 sarcoidosis, psoriasis, IBD, MS, and AS, as well as additional six diseases for HLA PGS (B12A, HYPER,
734 PBC, MCTD, SjS, and ADDISON) (**Supplementary notes 3.3.2 and Table S3.3.4**).

735 For each remaining parental AD, we first removed genotyped families with affected parents and then
736 subtracted mid-parent PGS for that AID from offspring PGS for the same disease. We normalized the
737 obtained difference by dividing it by the standard deviation of the mid-parent distribution. For each child,
738 we grouped his or her family trios according to whether this child was affected, so that we could compare
739 how the mean PGS of children deviate from their mid-parent PGS in the affected sibling group and
740 unaffected sibling group separately, as well as the difference between the two groups. We conducted pTDT
741 using SciPy v.1.7.3⁹³ in Python v.3.7.13.

742 743 744 **Predicting T1D risk in offspring using parental PGS**

745 To assess how well parental PGSs could be linked to T1D in offspring, we designed several analyses to test
746 the associations and predicting abilities. Considering the prediction performance of PGS for parental AIDs,
747 we only focused on those AIDs that showed significance from previous PGS analyses in HLA regions and
748 non-HLA regions. We introduced a new approach, Full-PGS, combining separately generated HLA and
749 non-HLA PGSs.

750 751 *Full-PGS construction*

752 Theoretically, with pTDT, four patterns could be defined for parental AID - offspring T1D transmissions:
753 1) both HLA PGS and non-HLA PGS over-/under-transmitted from unaffected parents to affected children;
754 2) only HLA PGS over-/under-transmitted; 3) only non-HLA PGS over-/under-transmitted; 4) neither was
755 transmitted.
756

757 We wondered whether we could improve PGS prediction across diseases by better considering genetic
 758 architecture of a target disease. For our target disease, T1D, we constructed HLA PGS and non-HLA PGS
 759 separately, and further considered the contribution ratio of HLA PGS relative to non-HLA PGS in a full
 760 PGS context.

761
 762 We proposed an equation to quantify the exact contribution of HLA PGS and non-HLA PGS ($ratio_{new}$)
 763 for diseases following pattern 1 ($effect_{HLA} \neq 0; effect_{non-HLA} \neq 0$), by taking HLA vs non-HLA
 764 contributions for both T1D and another AID into account.

$$765 \quad ratio_{new} = \frac{2 \times w_{HLA}}{2 \times w_{HLA} + 1 \times w_{non-HLA}} : \frac{1 \times w_{non-HLA}}{2 \times w_{HLA} + 1 \times w_{non-HLA}}$$

767
 768 Where the contribution ratio between HLA PGS and non-HLA PGS for T1D is 2:1, w_{HLA} and $w_{non-HLA}$
 769 are the weight of HLA PGS and the weight of non-HLA PGS for the AID when predicting the AID itself,
 770 $ratio_{new}$ denotes the contribution ratio between HLA PGS and non-HLA PGS when predicting T1D.

771
 772 For the diseases following pattern 2 ($effect_{HLA} \neq 0; effect_{non-HLA} = 0$), since only HLA PGS is
 773 assumed to be over-/under-transmitted, we only used HLA PGS for these diseases to predict T1D.

$$774 \quad ratio_{new} = 1:0$$

775
 776 For the diseases following pattern 3 ($effect_{HLA} = 0; effect_{non-HLA} \neq 0$), similarly, we only considered
 777 non-HLA PGS for these diseases to predict T1D.

$$778 \quad ratio_{new} = 0:1$$

779
 780 For the diseases following pattern 4 ($effect_{HLA} \neq 0; effect_{non-HLA} \neq 0$), neither HLA PGS nor non-
 781 HLA PGS was used for T1D prediction.

$$782 \quad ratio_{new} = 0:0$$

783
 784 In a scenario of cross-disease prediction, e.g., using PGS for RA to predict T1D, we conducted the analyses
 785 with the following steps: 1) assigned RA to one of the four groups according to its pTDT results, 2)
 786 quantified the contribution ratio between HLA PGS and non-HLA PGS using $\rho_{HLA \text{ PGS}}$, $\rho_{non-HLA \text{ PGS}}$ and the
 787 proposed equation, and 3) built a Full-PGS using HLA PGS, non-HLA PGS and the ratio of their
 788 contributions.

789
 790 We proposed that this equation could be extended to broader conditions for cross-disease prediction to fully
 791 utilize PGS information of a correlated disease to boost the prediction power of a target disease.

$$792 \quad ratio_{new} = \begin{cases} 0:0, & effect_1 = 0; effect_2 = 0 \\ 1:0, & effect_1 \neq 0; effect_2 = 0 \\ 0:1, & effect_1 = 0; effect_2 \neq 0 \\ \frac{w_{T1} \times w_{E1}}{w_{T1} \times w_{E1} + w_{T2} \times w_{E2}} : \frac{w_{T2} \times w_{E2}}{w_{T1} \times w_{E1} + w_{T2} \times w_{E2}}, & effect_1 \neq 0; effect_2 \neq 0 \end{cases}$$

793
 794 where w_{T1} and w_{T2} are the weights of two separate components of the target disease under a presumed
 795 genetic architecture (e.g., partition into HLA variants vs non-HLA variants, or common variants vs rare
 796 variants), w_{E1} and w_{E2} are the weights of two separate components of the correlated disease, $ratio_{new}$ is

800 the contribution ratio between the two genetic components for the correlated disease when predicting the
801 target disease.
802

803 *Impact of parental AID information and cumulative risk of T1D in children*

804 For each parental AID, we constructed a mid-parent Full-PGS using HLA PGS and non-HLA PGS
805 considering their ratio for cross-disease prediction and examined the association with T1D in offspring. For
806 comparisons, we ran additional models including only using parental diagnosis, only mid-parent PGS
807 calculated by a traditional PGS approach that takes genome-wide variants altogether for calculation, e.g.,
808 PRS-CS⁴⁶, or both HLA PGS and non-HLA PGS from our previous analyses, assumed to be at least as good
809 as Full-PGS. We used AUC as a primary metric to evaluate model performances. To understand how much
810 additive value parental PGS for other AIDs can be added to a T1D prediction model, we then compare a
811 model with all mid-parent PGSs and one with only mid-parent T1D PGS. We evaluated the two models by
812 comparing the AUCs among 12,563 genotyped trios in FinnGen. To avoid overfitting, the analyses were
813 done using five-fold cross validation.

814 We then conducted a Cox proportional hazards model to assess the T1D cumulative incidence risk in
815 offspring before the age of 20 years, adjusting for birthyear and the first 10 PCs of offspring. To maximize
816 the statistical power, we considered all the genotyped trios with children born between 1960-2010. Among
817 the 8,827 trios we analyzed, 1,035 had a child with T1D (11.7%). We then grouped the trios by dividing
818 the parental Full-PGS percentile into 0-50th, 50-90th, 90-99th, and 99-100th and estimated cumulative risk
819 within each group given the mean of children's birthyear and PCs. We also stratified the model by the sex
820 of offspring. To make the result generalizable, we calibrated the FinnGen results based on the FinRegistry
821 nationwide T1D prevalence among all children born between 1960-2010 (mean birthyear=1984;
822 N=3,048,812; prevalence=0.6%).

823 For the analyses in FinRegistry, we used the same cohort as we used for FinnGen calibration. We considered
824 four groups: 1) parents without T1D, 2) maternal T1D only, 3) paternal T1D only, and 4) both parents with
825 T1D among the whole study population. For the analyses excluding children with parental T1D, we
826 considered four groups: 1) parents without AID, 2) only the mother had AID(s), 3) only the father had
827 AID(s), and 4) both parents had AID(s). The parental AIDs in this analysis covered all the significant
828 associations identified from the previous analyses. All the analyses in this session were done using
829 statsmodels v.0.13.2⁹² and sklearn v.1.0.2 in Python v.3.7.13 and survival v.3.5.7 in R v.4.3.2.

830

831 **Code availability**

832 Code for the complete analyses is available at https://github.com/dsgelab/parentalAIDs_T1D.

833

834 **Reference**

- 91 Cohen, Jacob. Statistical power analysis for the behavioral sciences. Academic press, 2013.
- 92 Seabold, Skipper, and Josef Perktold. "Statsmodels: econometric and statistical modeling with python." *SciPy* 7 (2010): 1.
- 93 Virtanen, Pauli, et al. "SciPy 1.0: fundamental algorithms for scientific computing in Python." *Nature methods* 17.3 (2020): 261-272.

- 94 Douek, I. F., et al. "Three generations of autoimmune diabetes: an extended family study." *Diabetologia* 46 (2003): 1313-1318.
- 95 Schaid, Daniel J., et al. "Score tests for association between traits and haplotypes when linkage phase is ambiguous." *The American Journal of Human Genetics* 70.2 (2002): 425-434.
- 96 Zhao, Jing Hua. "gap: Genetic analysis package." *Journal of Statistical Software* 23 (2008): 1-18.
- 97 Kim, Seongho. "ppcor: an R package for a fast calculation to semi-partial correlation coefficients." *Communications for statistical applications and methods* 22.6 (2015): 665.
- 98 International HapMap 3 Consortium. "Integrating common and rare genetic variation in diverse human populations." *Nature* 467.7311 (2010): 52.
- 99 1000 Genomes Project Consortium. "A global reference for human genetic variation." *Nature* 526.7571 (2015): 68.

835