

Pace of Aging in older adults matters for healthspan and lifespan

Balachandran A¹, Pei H^{1,2}, Beard J^{1,2,3}, Caspi A^{4,5}, Cohen A^{1,6}, Domingue BW⁷, Eckstein Indik C¹, Ferrucci L⁸, Furuya A^{1,2}, Kothari M¹, Moffitt TE^{5,6}, Ryan C¹, Skirbekk V¹⁰, Zhang Y^{1,11}, Belsky DW^{1,2,3*}

*Correspondence to daniel.belsky@columbia.edu

1. Robert N Butler Columbia Aging Center, Columbia University Mailman School of Public Health, New York, NY, USA
2. Department of Epidemiology, Columbia University Mailman School of Public Health, New York, NY, USA
3. Department of Health Policy & Management, Columbia University Mailman School of Public Health, New York, NY, USA
4. Department of Psychology & Neuroscience, Duke University, Durham, NC, USA
5. Social, Genetic, and Developmental Psychiatry Unit, Institute of Psychiatry, Kings College, University of London, London, UK
6. Department of Environmental Health Sciences, Columbia University Mailman School of Public Health, New York, NY, USA
7. Graduate School of Education, Stanford University, Palo Alto, CA, USA
8. National Institute on Aging, Bethesda, MD, USA
9. ICAP, Columbia University Mailman School of Public Health, New York, NY, USA
10. Norwegian Institute for Public Health, Oslo, Norway
11. Department of Sociomedical Sciences, Columbia University Mailman School of Public Health, New York, NY, USA

Funding: This research was supported by National Institutes of Health grant R01AG061378, Russel Sage Foundation BioSS Grant 1810-08987, and the Robert N Butler Columbia Aging Center. AF is supported by T32AI114398. DWB is a fellow of the CIFAR CBD Network.

ABSTRACT

As societies age, policy makers need tools to understand how demographic aging will affect population health and to develop programs to increase healthspan. The current metrics used for policy analysis do not distinguish differences caused by early-life factors, such as prenatal care and nutrition, from those caused by ongoing changes in people's bodies due to aging. Here we introduce an adapted Pace of Aging method designed to quantify differences between individuals and populations in the speed of aging-related health declines. The adapted Pace of Aging method, implemented in data from N=13,626 older adults in the US Health and Retirement Study, integrates longitudinal data on blood biomarkers, physical measurements, and functional tests. It reveals stark differences in rates of aging between population subgroups and demonstrates strong and consistent prospective associations with incident morbidity, disability, and mortality. Pace of Aging can advance the population science of healthy longevity.

INTRODUCTION

Population aging is the central demographic phenomenon of the century and is unprecedented in human history^{1,2}. Current trends are driven primarily by gains in life expectancy for older adults³⁻⁵. In some places and population subgroups, these gains in life expectancy have been matched by gains in healthy years of life ("healthspan")^{6,7}. However, in other cases, healthspan and lifespan are diverging, with potential to disrupt healthcare systems and economies^{3,8,9}. The current toolkit available to demographers and public health planners allows differentiation of populations with more and less successful aging only in terms of completed lifespans or healthspans¹⁰⁻¹². These metrics are effective in summarizing differences between populations accumulated across the full life course. However, they do not distinguish deficits in health established early in life from the ongoing changes in people's bodies that are the essence of aging¹³ and the targets of public health interventions to promote healthy longevity¹⁴. As a consequence, the metrics in our existing toolkit may not be adequately sensitive to the effects of such interventions. Measurements are needed that can distinguish ongoing aging-related changes in organs, tissues, and capacities from differences in health that are legacies of early life in order to provide optimally-sensitive metrics for population surveillance and intervention evaluation. We previously developed the Pace of Aging method to quantify individual differences in aging trajectories with the goal of informing design and evaluation of clinical interventions^{15,16}. Here we adapt this methodology with the goal of informing evaluations programs and policies for aging societies.

Our original Pace of Aging method was developed from analysis of health changes from young adulthood to midlife in the Dunedin Study 1972-73 birth cohort¹⁵. To be most useful for comparative biodemographic analysis used by planners to evaluate efforts to promote healthy longevity, the Pace of Aging method needs to be adapted to a different context: samples of individuals representing a wide range of birth cohorts for whom follow-up begins later in the life course. In addition, whereas the Dunedin Study collected extensive biochemical and

physical examination data from participants, the studies used by planners typically have access to much sparser measurement panels.

Here, we introduce an adapted method for calculation of Pace of Aging in a sample composed of a wide range of birth cohorts with follow-up in midlife and older age and a sparse panel of biomarkers. Our adapted method is designed to generalize across the Gateway to Global Aging family of harmonized cohort studies¹⁷, which comprises cohorts across Europe, Asia, and Central and South America that are routinely cited in planning and policy analysis. We implement the method and test proof of concept using data from the US Health and Retirement Study (HRS). As the other cohorts extend their biomarker follow-up, the method introduced here can be applied to conduct cross-national comparative analysis.

We compiled data from dried-blood spot, physical exam, and functional test protocols conducted by the HRS during 2006-2016 (six assessment waves). We identified nine parameters measured at all six waves that met criteria for inclusion in the Pace of Aging analysis: C-reactive protein (CRP), Cystatin-C, glycated hemoglobin (HbA1C), diastolic blood pressure, waist circumference, lung capacity (peak flow), tandem balance, grip strength, and gait speed. A total of 13,626 individuals provided data on at least six of these nine biomarkers across at least two of the follow-up assessments. We modeled longitudinal change in these biomarkers to estimate person-specific slopes for each of them. Then we combined slope information across biomarkers to compute each participants' Pace of Aging. Our validation analyses tested associations of Pace of Aging with measures of morbidity, disability, and survival through 2021. Finally, we evaluated socioeconomic and demographic disparities in the Pace of Aging among US older adults.

METHODS

Sample

The Health and Retirement Study (HRS) is a nationally representative longitudinal survey of US residents ≥ 50 years of age and their spouses. The HRS has been fielded every two years since 1992. Participants are asked about four broad areas: income and wealth; health, cognition, and use of healthcare services; work and retirement; and family connections. A new cohort of 51–56-year-olds and their spouses is enrolled every six years to maintain representativeness of the U.S. population over 50 years of age. Response rates over all waves of the HRS range from 81–91%. As of the most recent data release, HRS included data collected from 42,515 individuals in 26,600 households. We linked HRS data curated by RAND Corporation¹⁸ with dried-blood-spot biomarker data collected during 2006–2016¹⁹.

Measures

Pace of Aging. We measured Pace of Aging from blood biomarker, physical assessment, and functional test performance data collected during home visits to participants during 2006–2016. Biomarkers were included in Pace of Aging analysis based on prior knowledge of monotonic change across the age range included in the HRS. Three blood biomarkers (HbA1c, C-reactive protein, cystatin-C), three physical assessments (diastolic blood pressure, peak-flow lung-function testing, waist circumference), and three functional tests (gait speed, balance, grip strength) were included in analysis. Participants received home visits at 4-year intervals (2006, 2010, and 2014 or 2008, 2012 and 2016). We measured Pace of Aging using data from participants with at least 2 repeated measures of at least six of the nine biomarkers representing all three types of data (N=13,626). Baseline observations were recorded in 2006 for 39% of the sample, in 2008 for 34%, in 2010 for 14% and in 2012 for 13%. Three repeated measures were available for 55% of the sample. Characteristics of each of these groups of

participants are reported in **Supplemental Table 1**. Biomarker measurements are described in detail in **Supplemental Table 2**. Biomarker summary statistics are reported in **Supplemental Table 3**.

We modeled Pace of Aging in 4 steps. First, we standardized biomarkers to a common distribution by centering values on the sex-specific mean for participants aged <65 years of age and dividing the centered values by the sex-specific standard deviation of that group (Gait speed was measured in participants aged 65 and older; we standardized values based on distributions for participants aged 65-75). Age-group specific means and standard deviations for men and women included in the analysis are reporting in **Supplemental Table 3**. For biomarkers that decline with aging (diastolic blood pressure, peak flow, balance, and grip strength), standardized values were reversed so that higher values corresponded to poorer organ-system function. Gait speed was measured as the natural log of time to complete the walk test and so did not need to be reversed.

Second, we modeled change over time in each biomarker using linear mixed-effects regression models with participant-specific random intercepts and slopes. Models included baseline year of biomarker measurement, follow-up time, chronological age at baseline (modeled as a 3rd degree B-spline) and an interaction between age at baseline and follow-up time as covariates. Interactions between baseline-age terms and time were included to allow for variation in slopes of change depending on participants' ages at baseline. Models were fitted separately for men and women.

Third, for each biomarker, we measured pace of change for each participant by combining fixed and random slope components.

Finally, we computed each participant's Pace of Aging as the average pace of change across the nine biomarkers. We scaled Pace of Aging based on the sex-specific average of participants under age 65. For the resulting measure, a value of 1 represents average biomarker change per chronological year in HRS participants aged <65. A value of 1.5 would indicate a Pace of Aging

50% faster relative to this average. A value of 0.5 would indicate a Pace of Aging 50% slower relative to the average.

Mortality, morbidity, disability and cognitive impairment. We evaluated criterion validity of the Pace of Aging by testing associations with four components of healthy lifespan: survival, incidence of chronic disease, incidence of disability and decline in cognition. Survival was measured by HRS through the 2020 follow-up assessment. We analyzed survival from the baseline biomarker measurement. Chronic disease incidence was measured from participant reports of physician diagnosed chronic disease. Disability incidence was measured from participant reports about Activities of Daily Living (ADL) limitations and Instrumental Activity of Daily Living (IADL) limitations. Cognitive decline and incident cognitive impairment were measured according to the Langa-Weir approach²⁰ from cognitive-test and interview data. Survival, chronic disease, disability, and cognition measures are described in detail in **Supplemental Table 4**. All outcomes other than mortality were measured at the 2020 measurement wave or the last wave a participant contributed data subsequent to the end of follow-up for pace of aging measurement.

Analysis

We used regression analysis to test Pace of Aging associations with healthspan phenotypes. We fit Cox proportional hazard models to estimate hazard ratios (HRs) and 95% CIs for mortality. We fit Poisson regression models to estimate incidence rate ratios (IRRs) and 95% confidence intervals (CIs) for incident chronic disease and limitations to activities of daily living (ADLs) and instrumental activities of daily living (IADLs). We use multinomial logit models to estimate odds-ratios (OR) for incident cognitive impairment and dementia. We use linear regression models to estimate change in cognitive-function scores between baseline and follow-up and to test differences in Pace of Aging between population sub-groups. All models were adjusted for sex, race/ethnicity, age, age-squared, and a set of terms encoding baseline measurement year and whether the participant contributed two or three repeated measures of biomarker

data. Models of healthspan outcomes other than mortality included an additional covariate for follow-up time.

RESULTS

We analyzed data from US Health and Retirement Study (HRS) participants aged 40 or older at the time of their first biomarker measurement who contributed at least two repeated measures of six or more biomarkers over 2006-2016 (N=13,626; 41% male, mean age at baseline=65, SD=10). This analysis sample was slightly younger and better educated in comparison to the overall HRS sample (**Supplemental Table 1**). Baseline observations were recorded in 2006 for 39% of the sample, in 2008 for 34%, in 2010 for 14% and in 2012 for 13%. Three repeated measures were available for 55% of the sample. Summary statistics for biomarker measurements at baseline are reported in **Supplemental Table 3**.

HRS participants showed signs of correlated decline in multiple indicators of system integrity over 4-8 years of follow-up. Of the nine biomarkers included in analysis, eight showed the expected pattern of change: Gait speed, grip strength, balance, diastolic blood pressure, and peak-flow declined; cystatin-C, HbA1c, and waist circumference increased. For CRP, change was in the expected positive direction for men, but declined slightly for women. Slopes of aging-related decline were steepest for the functional test biomarkers. Aging-related biomarker changes were greater for men as compared to women, except for peak flow, for which change was greater for women. On average, there were no changes in HbA1C and waist circumference for women. Biomarker slopes of change are reported in **Supplemental Table 5** and are plotted in **Figure 1**.

Participants who were older at baseline showed steeper slopes of change in most biomarkers. Biomarker-slope correlations with baseline chronological age were positive ($r=0.01-0.32$), except waist circumference and grip strength. Net of chronological age, biomarker slopes of change showed positive correlation with one another, except for waist circumference (mean Pearson's $r=-0.07$ for women and -0.09 for men; **Supplemental Figure 2**).

In the next step, we composited slopes of change across biomarkers to quantify the overall rate of decline in system integrity.

Pace of Aging. We computed each participant's Pace of Aging as the average pace of change across the nine biomarkers. We scaled Pace of Aging based on the sex-specific average value for participants under age 65. Resulting values can be interpreted as years of biological change per calendar year relative to the reference group. Pace of Aging values were approximately normally distributed and indicated faster aging in men as compared to women and older as compared to younger participants (Pace of Aging mean=1.33 (SD=0.68); correlation with chronological age at baseline $r=0.61$; male-female difference Cohen's $d=0.21$, 95% CI [0.19-0.24]; **Figure 2**). Comparison of Pace of Aging between participants identifying as White ($n=9,301$), Black ($n=2,269$), Hispanic ($n=1,674$), and Other ($n=377$) race/ethnicity showed that, compared to White-identifying participants, Black- and Hispanic identifying participants had faster Pace of Aging (for Black, Cohen's $d=0.18$, 95% CI [0.16-0.21]; for Hispanic, Cohen's $d=0.06$, 95% CI [0.03-0.10]; **Supplemental Table 7**). Subsequent analyses included participants' sex, age, and race/ethnicity as covariates. Differences in Pace of Aging across demographic groups are illustrated in **Figure 3**.

Participants with faster Pace of Aging were at increased risk of cognitive impairment, incident chronic disease, disability, and mortality. We conducted criterion validity analysis by testing if participants with faster Pace of Aging more often developed cognitive impairment, new chronic disease or disability over follow up, and if they were more likely to die. We measured chronic disease and disability from participant reports of physician-diagnosed conditions and limitations to activities of daily living (ADLs) and instrumental activities of daily living (IADLs) at baseline and in 2020 ($n=8,623$). Those with faster Pace of Aging reported more new diagnoses of chronic diseases and more new ADLs and IADLs (chronic diseases IRR=1.08 95% CI [1.06-1.10]; ADLs IRR=1.58 [1.49-1.65]; IADLs 1.59 [1.40-1.66]; all p -values<0.001). We conducted analysis of mortality using follow-up data accumulated through 2019. We included data on all deaths occurring subsequent to a participants' second biomarker data collection (i.e. after the

minimum follow-up required to compute Pace of Aging). Analysis included N=13,626 participants who contributed mean follow-up time of 10 years (SD=2) over which 3,124 deaths were recorded. Participants with faster Pace of Aging were at increased risk of mortality (HR=1.73 [1.66-1.81], $p < 0.001$; **Figure 4**). Effect-sizes for all healthspan metrics are shown in **Figure 5** and **Supplemental Table 6**. We observed some differences in effect-sizes for associations of Pace of Aging with measures of healthspan and lifespan between demographic groups of participants. Effect-sizes were generally somewhat larger for white participants, men, and those under age 65 as compared with non-white participants, women, and older participants. Results are reported in **Supplemental Table 8**, **Supplemental Figure 4** and **Supplemental Figure 5**.

Pace of Aging associations with cognitive impairment, morbidity, disability and mortality were independent of smoking, obesity, and education. We investigated whether Pace of Aging associations with cognitive impairment, morbidity, disability, and mortality were accounted for by baseline socio-economic and health behavior risk factors. Participants with lower levels of education, who were overweight or obese, and who were current or former smokers tended to have a faster Pace of Aging (Cohen's d range 0.08 – 0.38; **Supplemental Table 7**). We therefore repeated analyses of cognitive function and impairment, morbidity, disability and mortality including these risk factors as covariates. Adjusted results were similar to unadjusted results (**Supplemental Table 8**).

DISCUSSION

The Pace of Aging method remains an understudied approach to quantification of biological aging. The method was first proposed in the Dunedin Longitudinal Study¹⁶, which has followed a single-year birth cohort over five decades²¹. The Dunedin analysis included individuals all born in the same year and followed-up at the same ages from young adulthood to midlife. Here, we show that Pace of Aging can be modeled in the very different setting of a mixed-age cohort of

older adults followed-up at ages ranging from the sixth through ninth decades of life. In addition, we report three key findings that advance knowledge of Pace of Aging.

First, Pace of Aging accelerates at more advanced ages. HRS participants who were older at their baseline biomarker assessment showed more rapid change across subsequent follow-ups as compared to those who were younger. This observation is consistent with biodemographic data showing that mortality risk accelerates at older ages²², with biomarker analysis in the Baltimore Longitudinal Study on Aging²³ and the Cardiovascular Health Study²⁴, and with analysis of the Pace of Aging epigenetic clock, DunedinPACE^{16,25}, and advances the hypothesis that the pace of aging accelerates later in life.

Second, Pace of Aging is faster in sociodemographic groups characterized by shorter lifespan. Men tended to experience faster Pace of Aging as compared with women. Those with less education tended to experience faster Pace of Aging as compared to those with more education, consistent with observations of a socioeconomic gradient in the pace of aging from the Dunedin Cohort and a Swiss cohort^{27,28}. In addition, we observed a faster Pace of Aging in Black- and Hispanic-identifying as compared to White-identifying Americans. While race/ethnic disparities in healthy aging outcomes are well established²⁶, our observation of faster Pace of Aging in Black as compared to White Americans indicates healthy-aging disparities continue to accumulate into later life. This finding suggests that intervention/prevention in this older-aged group has the potential to reduce aging-related health disparities.

Third, midlife and older adults with faster Pace of Aging were at increased risk of incident chronic disease, disability, and mortality. In the Dunedin Study, where we first introduced Pace of Aging, participants are still middle aged; it is not yet possible to test associations with aging-related health problems and mortality. In HRS, older adults with faster Pace of Aging more often developed new chronic diseases and disabilities and were at increased risk of death. Moreover, these associations were independent of smoking, obesity, and educational attainment. These findings contribute evidence that the adapted Pace of Aging method introduced here captures differences in aging processes that are important to healthspan and

lifespan. The variation observed across social and demographic groups suggests potential to modify pace of aging through changes in the organization of aging societies and in the environment and behavior of older adults.

These findings have implications for future research. Pace of Aging summarizes changes occurring across multiple systems in the body to provide a dynamic measure of a person's healthy aging trajectory. The availability of a Pace of Aging measure within the HRS can inform research in the fields of Sociology and Economics to understand how social transitions such as retirement, spousal loss, and caregiving responsibilities affect trajectories of healthy aging. In medicine and gerontology, Pace of Aging measures can help reveal how health shocks such as new diagnoses, accidents, or elective surgeries modify aging trajectories. In life-course epidemiology, Pace of Aging measures can help differentiate effects of early-life exposures on aging-related health decline from health deficits established earlier in life. In health equity research, Pace of Aging measures can be used to quantify disparities across population subgroups in health aging trajectories and illuminate how economic and health shocks contribute to inequalities²⁹. These lines of research, in turn, can deliver new knowledge critical to the development of policies and programs to promote healthy longevity, consistent with the priorities of World Health Organization¹⁴ and other public health actors³⁰⁻³².

We acknowledge limitations. The HRS measurement battery available to measure Pace of Aging is more limited as compared with the Dunedin Study. Some parameters are measured with lower precision instruments (e.g. peak flow meters as compared to spirometry for assessment of lung function). In other cases, measurements available the in Dunedin Study were not collected by HRS (e.g. periodontal health, cardiorespiratory fitness). In addition, the older ages of HRS participants limits utility of some biomarkers that are available. For example, medical management of systolic blood pressure and cholesterol levels in older adults along with non-linearities in the patterning of these markers with aging in late life limit their utility to Pace of Aging analysis. Nevertheless, HRS data do provide for measurement of changes in pulmonary, vascular, immune, renal, endocrine, metabolic, and musculoskeletal systems and therefore

make possible Pace of Aging analysis in a large, national sample of older adults with significant race/ethnic diversity.

HRS biomarker data have been collected at up to only three timepoints for each participant. As a result, we were unable to model non-linear changes in biomarkers. Consequently, we must infer the acceleration in Pace of Aging at older chronological ages from comparisons of younger and older participants. As additional waves of biomarker data are collected, it will be possible to model non-linear slopes of change and more rigorously test the hypothesis that aging accelerates towards the end of life.

Because HRS enrolls some participants in their 70s and 80s, the sample we analyzed may over-represent those with slower Pace of Aging. If, as data reported here suggest, faster Pace of Aging increases risk of death, the oldest segments of the sample will tend to include those with slower Pace of Aging. Importantly, despite this limitation, we still observe evidence of faster Pace of Aging in older as compared to younger participants. Moreover, faster Pace of Aging was associated with incident morbidity, disability, cognitive impairment, and mortality, in both older and younger segments of the sample.

Beyond the potential bias toward slower agers, our analysis sample was limited to participants contributing at least two repeated observations of biomarker data, requiring follow-up over at least four years. The subset of HRS participants meeting this requirement tended to be better educated, less diverse, and younger than the overall sample. This may also contribute to the observed distribution of the Pace of Aging being somewhat slower than in the general population HRS is designed to represent.

Consistent with observations in the original Pace of Aging analyses^{15,33}, we found that patterns of decline were correlated across biological systems although, as in the original analyses, these correlations were modest. Other investigators have interpreted similarly modest cross-system correlations as evidence of stochastic variation in aging^{34,35}. Whether Pace of Aging reflects an underlying set of biological processes causing correlated change across systems or instead

summarizes changes across systems experiencing stochastic aging trajectories is an important question for further research. Critically, either interpretation supports application of Pace of Aging within population health science to quantify trajectories of healthy aging.

To summarize, we developed a measure of Pace of Aging in HRS. It is predictive of morbidity, disability, cognitive impairment, and mortality and reflects known social gradients in healthy aging. It provides a new tool for researchers seeking to understand how features of societal organization, built and social environments, and individual behavior contribute to healthy aging trajectories in populations around the world.

References

1. Lutz W, Sanderson W, Scherbov S. The coming acceleration of population ageing. *Nature*. 2008;(451):716-716. doi:10.1038/nature06516
2. Balachandran A, de Beer J, James KS, van Wissen L, Janssen F. Comparison of Population Aging in Europe and Asia Using a Time-Consistent and Comparative Aging Measure. *Journal of Aging and Health*. Published online 2020:089826431882418-089826431882418. doi:10.1177/0898264318824180
3. Crimmins EM. Lifespan and healthspan: past, present, and promise. *The Gerontologist*. 2015;55(6):901-911.
4. Partridge L, Deelen J, Slagboom PE. Facing up to the global challenges of ageing. *Nature*. 2018;561(7721):45-56.
5. Beard JR, Si Y, Liu Z, Chenoweth L, Hanewald K. Intrinsic capacity: validation of a new WHO concept for healthy aging in a longitudinal Chinese study. *The Journals of Gerontology: Series A*. 2022;77(1):94-100.
6. Crimmins EM, Saito Y. Trends in healthy life expectancy in the United States, 1970-1990: Gender, racial, and educational differences. *Social Science and Medicine*. 2001;52(11):1629-1641. doi:10.1016/S0277-9536(00)00273-2
7. Ailshire JA, Beltran-Sanchez H, Crimmins EM. Social Characteristics and Health Status of Exceptionally Long-Lived Americans in the Health and Retirement Study. *Journal of the American Geriatrics Society*. 2011;59(12):2241-2248. doi:10.1111/j.1532-5415.2011.03723.x
8. Seaman R, Höhn A, Lindahl-Jacobsen R, Martikainen P, Van Raalte A, Christensen K. Rethinking morbidity compression. *European journal of epidemiology*. 2020;35(5):381-388.
9. Payne CF. Expansion, compression, neither, both? Divergent patterns in healthy, disability-free, and morbidity-free life expectancy across US birth cohorts, 1998–2016. *Demography*. 2022;59(3):949-973.
10. Balachandran A. Population ageing in Europe and Asia : Beyond traditional perspectives University of Groningen Population ageing in Europe and Asia : Beyond traditional perspectives. 2020;(October).
11. Skirbekk V, Dieleman JL, Stonawski M, Fejkiel K, Tyrovolas S, Chang AY. The health-adjusted dependency ratio as a new global measure of the burden of ageing: a population-based study. *The Lancet Healthy Longevity*. 2022;3(5):e332-e338.
12. Skirbekk VF, Staudinger UM, Cohen JE. How to measure population aging? The answer is less than obvious: A review. *Gerontology*. 2019;65(2):136-144. doi:10.1159/000494025
13. Ferrucci Luigi, Levine Morgan E., Kuo Pei-Lun, Simonsick Eleanor M. Time and the Metrics of Aging. *Circulation Research*. 2018;123(7):740-744. doi:10.1161/CIRCRESAHA.118.312816
14. Beard JR, Officer A, de Carvalho IA, et al. The World report on ageing and health: a policy framework for healthy ageing. *The Lancet*. 2016;387(10033):2145-2154. doi:10.1016/S0140-6736(15)00516-4
15. Belsky DW, Caspi A, Houts R, et al. Quantification of biological aging in young adults. *Proceedings of the National Academy of Sciences*. 2015;112(30):E4104-E4110.

16. Belsky DW, Caspi A, Arseneault L, et al. Quantification of the pace of biological aging in humans through a blood test, the DunedinPoAm DNA methylation algorithm. *Elife*. 2020;9:e54870-e54870.
17. Lee J, Phillips D, Wilkens J, Team G to GAD. Gateway to Global Aging Data: Resources for cross-national comparisons of family, social environment, and healthy aging. *The Journals of Gerontology: Series B*. 2021;76(Supplement_1):S5-S16.
18. Bugliari D, Carroll J, Hayden O, et al. RAND HRS Longitudinal File 2020 (V1) Documentation. *Aging*. Published online 2023.
19. Kim JK, Faul J, Weir DR, Crimmins EM. Dried blood spot based biomarkers in the Health and Retirement Study: 2006 to 2016. *American Journal of Human Biology*. Published online 2023:e23997.
20. Langa KM, Kabeto M, Weir D. Report on race and cognitive impairment using HRS in 2010 Alzheimer's disease facts and figures. Retrieved July. 2010;12:2010.
21. Poulton R, Guiney H, Ramrakha S, Moffitt TE. The Dunedin study after half a century: reflections on the past, and course for the future. *Journal of the Royal Society of New Zealand*. 2023;53(4):446-465.
22. Finch CE, Crimmins EM. Constant molecular aging rates vs. the exponential acceleration of mortality. *PNAS*. 2016;113(5):1121-1123. doi:10.1073/pnas.1524017113
23. Kuo PL, Schrack JA, Levine ME, et al. Longitudinal phenotypic aging metrics in the Baltimore Longitudinal Study of Aging. *Nature aging*. 2022;2(7):635-643.
24. Newman AB, Sanders JL, Kizer JR, et al. Trajectories of function and biomarkers with age: the CHS All Stars Study. *Int J Epidemiol*. 2016;45(4):1135-1145. doi:10.1093/ije/dyw092
25. Belsky DW, Caspi A, Corcoran DL, et al. DunedinPACE, a DNA methylation biomarker of the pace of aging. *Elife*. 2022;11:e73420-e73420.
26. Newman AB, Odden MC, Cauley JA. Epidemiology of aging. In: *Handbook of Epidemiology*. Springer; 2023:1-37.
27. Belsky DW, Caspi A, Cohen HJ, et al. Impact of early personal-history characteristics on the Pace of Aging: implications for clinical trials of therapies to slow aging and extend healthspan. *Aging Cell*. 2017;16(4):644-651. doi:10.1111/accel.12591
28. Schrepft S, Belsky DW, Draganski B, et al. Associations Between Life-Course Socioeconomic Conditions and the Pace of Aging. *J Gerontol A Biol Sci Med Sci*. 2022;77(11):2257-2264. doi:10.1093/gerona/glab383
29. Schmitz LL, Duque V. In utero exposure to the Great Depression is reflected in late-life epigenetic aging signatures. *Proceedings of the National Academy of Sciences*. 2022;119(46):e2208530119-e2208530119.
30. Ye P, Jin Y, Er Y, et al. A Scoping Review of National Policies for Healthy Ageing in Mainland China from 2016 to 2020. *The Lancet Regional Health - Western Pacific*. 2021;12:100168. doi:10.1016/j.lanwpc.2021.100168
31. Healthy Aging - PAHO/WHO | Pan American Health Organization. Accessed March 11, 2024. <https://www.paho.org/en/healthy-aging>
32. Innovation and Leadership in Healthy Aging. Accessed March 11, 2024. <https://www.aarpinternational.org/resources/healthy-aging>

33. Elliott ML, Caspi A, Houts RM, et al. Disparities in the pace of biological aging among midlife adults of the same chronological age have implications for future frailty risk and policy. *Nat Aging*. 2021;1(3):295-308. doi:10.1038/s43587-021-00044-4
34. Sanders JL, Ding V, Arnold AM, et al. Do Changes in Circulating Biomarkers Track With Each Other and With Functional Changes in Older Adults? *J Gerontol A Biol Sci Med Sci*. 2014;69A(2):174-181. doi:10.1093/gerona/glt088
35. Cohen AA, Ferrucci L, Fülöp T, et al. A complex systems approach to aging biology. *Nat Aging*. 2022;2(7):580-591. doi:10.1038/s43587-022-00252-6

Figure 1. Pace of Aging and biomarker slopes of change correlations with chronological age at baseline in women and men. Figure shows changes in biomarker values across baseline and four- and eight-year follow-up assessments among participants in the US Health and Retirement Study (HRS, N=13,626). The HRS collected biomarker data from participants beginning either in 2006 or 2008. The figure combines data from these groups of participants; e.g., the baseline (time-0) data include observations made in 2006 of one group and in 2008 of the other group. Y-axis values are z-scores (M=0, SD=1) based on sex-specific distributions of values in participants aged <65. For gait speed, which was measured only in participants aged 65 and older, distributions were formed from participants aged 65-75. Biomarkers known to decline with aging were reverse coded so that higher values on the Y-axis corresponds to “older” levels of the biomarkers. Therefore, positive values for slopes of change indicate aging related change in the direction expected.





Figure 2. Pace of Aging in the US Health and Retirement Study. Figure shows data from men and women in the US Health and Retirement Study (HRS, N=13,626). Panel A shows the association of chronological age (X-axis) with Pace of Aging (Y-axis). The histograms above the plot show distributions of age at baseline. The histograms to the right of the plot show the histograms of Pace of Aging. Data for women are graphed in pink. Data for men are graphed in blue. Panel B shows model-predicted trajectories of system integrity scores for women (pink) and men (blue). Predicted trajectories were computed based on model-predicted intercepts and slopes for all biomarkers included in the analysis. Intercepts and slopes were averaged across biomarkers to compute values.

Panel A



Panel B





Figure 3. Differences in Pace of Aging among US Older Adults by Race and Ethnicity. Figure shows data from non-Hispanic White (N=9,301), Black (N=2,269), and Hispanic (N=1,674) - identifying older adults in the US Health and Retirement Study. Panel A shows the differences in distribution of Pace of Aging between White, Black, and Hispanic participants. Densities reflect distributions of Pace of Aging after adjustment for chronological age. White lines show group means. Panel B shows effect-size estimates for the differences in Pace of Aging relative to White participants. The figure illustrates overall faster pace of aging in Hispanic- and Black-identifying older adults as compared with White-identifying older adults.

Panel A



Panel B




FIGURE 4. Association of Pace of Aging with mortality. The figure plots Kaplan-Meier curves for four groups of US Health and Retirement Study participants defined by Pace of Aging (slowest quartile graphed in dark blue, fastest quartile graphed in dark red, middle quartiles graphed in lighter shades). Mortality follow-up was conducted from the time of the second biomarker measurement through the 2020 data collection wave. Over follow-up there were 1568 deaths in the slowest Pace of Aging quartile, 1990 deaths in the quartile containing the 25th-50th percentiles, 1824 deaths in the quartile containing the 50th-75th percentiles, and 1946 deaths in the fastest quartile. Numbers at risk at baseline, four, eight, and twelve years of follow-up are shown below the plot.




Figure 5. Effect sizes for associations of Pace of Aging with mortality and incident chronic disease, ADLs, IADLs and cognitive impairment. The figure shows effect-sizes for associations of Pace of Aging with mortality (N=13,621) and with incident chronic disease (n=11,619), ADLs and IADLs (n=11,611), and cognitive impairment (n=10,666) among participants followed-up during 2016-2020. Incident chronic disease, ADLs, IADLs, and cognitive impairment were determined by comparing measurements taken at participants' biomarker baseline wave with measurements taken in 2020 or the last measurement wave they participated in. Effect-sizes for mortality are reported as hazard ratios (HRs) estimated from Cox regressions. Effect-sizes for incidence of chronic disease, ADLs, and IADLs are reported as incidence rate ratios (IRRs) from Poisson regressions. Effect-sizes for incidence of cognitive impairment are reported as odds ratios (ORs) estimated from multinomial logistic regressions in which outcomes were normal (reference), cognitively impaired but not demented (CIND), and demented. Effect-sizes are reported for a one-standard-deviation difference in Pace of Aging. All regression models included covariate adjustment for sex, race, age, age-squared, a set of terms encoding baseline measurement year for Pace of Aging, follow-up time to outcome assessment, and whether the participant was aged 65 or older at baseline. Figure

