

1 **DISRUPTIONS IN LIVELIHOODS, HEALTHCARE ACCESS AND HEALTH** 2 **OUTCOMES IN WEST AFRICA DURING EMERGING INFECTIOUS DISEASE** 3 **OUTBREAKS**

4 Rashid Ansumana^{1*}, Alfred S. Bockarie², Monya Konneh¹, Najmul Haider³, Eniola Cadmus⁹,
5 Alberta Korvah⁵, Joyce Fayiah⁵, Yolaine Kate Waka-Metzger⁵, Angella M. George¹ Gassimu
6 Mallah⁹ Joseph M. Lamin¹, Osman Dar⁸, Susannah Mayhew⁷, Doris Bah¹⁰, Pepe Bilivigoi⁹,
7 Nfaly Maggasouba⁴, Simeon Cadmus,⁶ Robyn Alders⁸.

8 9 **Affiliations**

- 10 1. School of Public Health, College of Medical Sciences, Njala University, Bo Campus
- 11 2. Institute of Environmental Management and Quality Control, School of Environmental
12 Sciences, Njala University, Njala Campus
- 13 3. Keele University, United Kingdom
- 14 4. University of Gamel Abdel Nasser of Conakry, Conakry Guinea
- 15 5. National Public Health Institute of Liberia, Monrovia, Liberia
- 16 6. University of Ibadan, Ibadan, Nigeria
- 17 7. London School of Hygiene and Tropical Medicine, UK
- 18 8. Australia National University
- 19 9. Ministry of Health, Guinea
- 20 10. Ministry of Health, Sierra Leone

21
22 ***Correspondence:** Rashid Ansumana(PhD), School of Public Health, College of Medical
23 Sciences, Njala University. Email: r.ansumana@njala.edu.sl

24 25 **ABSTRACT**

26 **Background:** Outbreaks of infectious diseases like Ebola virus disease, Lassa fever, and
27 COVID-19 have severely strained infrastructural systems and social services across West Africa.
28 We investigated the disruptions caused by emerging outbreaks on access to healthcare, health
29 outcomes, and livelihoods in West Africa.

30
31 **Method:** A mixed-methods approach was utilized, conducting extensive studies in Nigeria,
32 Sierra Leone, Guinea, and Liberia through structured questionnaires, in-depth interviews with
33 key informants, and focus group discussions. Using a device-to-cloud system guided by GIS for
34 randomized sampling across the four nations, this technique allowed us to comprehensively
35 analyze implications across imposed lifestyle changes on health and wellbeing due to disrupted
36 healthcare.

37 38 **Results**

39
40 Our findings indicate drastic shifts in food consumption patterns and healthcare access. In
41 Guinea, self-reported “Once Daily” meals astonishingly surged from 148 to 775 individuals
42 ($p < 0.001$), with analogous substantial increases observed in Liberia and Sierra Leone. Nigeria
43 exhibited a varied response, with notable rises both in “Once Daily” and “Twice Daily” meal
44 frequencies ($p < 0.001$), reflecting broad dietary adaptations out of necessity. Additionally, there
45 was a significant decrease in consumption of traditional protein sources like bushmeat, beef, and
46 mutton, mainly because of disrupted supply chains and heightened concerns over food insecurity.

47 Conversely, fish consumption slightly fell possibly due to its perceived safety or accessibility
48 amidst the outbreak.

49
50 Healthcare services faced severe disruptions, particularly acute in Sierra Leone and Liberia
51 compared to Guinea and Nigeria. The interruption of services drastically impacted everything
52 from immunization rates to mental health, with a rise in reported anxiety and depression
53 alongside public dissatisfaction towards the healthcare disruptions.

54
55 **Conclusion:** This study demonstrates the dramatic effects of infectious disease outbreaks on
56 health access and diets in West Africa. The research calls for integrating health initiatives with
57 social protection to strengthen the resilience of societies to meet all types of health challenges. It
58 is imperative districts establish robust health systems and social security mechanisms to counter
59 future public health crises.

60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92

93

94 **BACKGROUND**

95 The intricate relationship between health and socio-economic well-being is deeply rooted in the
96 fabric of human society. At the confluence of these critical dimensions lies the healthcare
97 system, a fundamental pillar that ensures a population's health and vitality and acts as a sentinel
98 for broader socio-economic stability. West Africa, with its unique socio-cultural, economic, and
99 political landscapes, has unfortunately been a recurrent epicentre for infectious disease
100 outbreaks, notably the Ebola Virus Disease (EVD) in 2014-2016(Ansumana et al., 2014;
101 Ansumana et al., 2017), Lassa fever (Ogbu, Ajuluchukwu and Uneke, 2007; Tambo, Adetunde
102 and Olalubi, 2018; Kofman, Choi and Rollin, 2019) and, more recently, the COVID-19
103 pandemic(Ansumana, Sankoh and Zumla, 2020; Sumner, Hoy and Ortiz-Juarez, 2020). These
104 outbreaks, while primarily health crises have invariably cast ripple effects, perturbing healthcare
105 access, undermining livelihoods, and precipitating adverse health outcomes.

106

107 The importance of consistent and robust healthcare access cannot be overstated. In the context of
108 West Africa, healthcare access is not only pivotal for managing endemic diseases such as
109 malaria, tuberculosis, and HIV/AIDS but also for providing essential maternal and child health
110 services(Smith and Whittaker, 2014). Outbreaks tend to exacerbate challenges faced by
111 healthcare systems in resource-poor countries.

112 The economic backbone of many West African nations is agriculture, which employs a
113 significant portion of the population. During outbreaks, lockdowns, travel restrictions, and
114 market closures disrupt agricultural activities and supply chains. This has far-reaching
115 consequences, including job loss, decreased income, and food insecurity, especially in rural
116 areas(Egeru et al., 2020; Kipchumba et al., 2023).

117 For instance, during the EVD outbreak, many healthcare facilities in affected countries either
118 closed or significantly reduced their operations due to fear of transmission, lack of protective
119 equipment, or workforce attrition(Parpia *et al.*, 2016). This curtailment reduced access to
120 essential health services, increasing morbidity and mortality from non-Ebola causes(J W T
121 Elston *et al.*, 2017).

122

123 Furthermore, the socio-economic ramifications of outbreaks in West Africa are profound.
124 Livelihoods, particularly in informal sectors, are often the first to be affected. As movement
125 restrictions and lockdowns are implemented to curb the spread of disease, daily wage earners,
126 small-scale traders, and artisans face immense challenges in sustaining their income(Vinck *et al.*,
127 2019). This economic disruption, in turn, affects purchasing power and exacerbates food
128 insecurity, further compromising vulnerable populations' nutritional and health status(WFP,
129 2020).

130

131 Animal health is linked to human health and livelihoods, particularly in regions like West Africa.
132 The concept of 'One Health' recognises that the health of people, animals, and the environment is
133 interconnected(Conrad, Meek and Dumit, 2013). Outbreaks often have zoonotic origins, where
134 infectious diseases are transmitted between animals and humans. For instance, the EVD outbreak
135 is believed to have originated from bats and potentially transmitted to humans via the
136 consumption of bushmeat(Olival and Hayman, 2014). Similarly, other pathogens with potential
137 for outbreaks, such as Lassa and Marburg infectious disease and monkeypox, have animal
138 reservoirs(Mylne *et al.*, 2015). Disruptions caused by outbreaks can impact veterinary care and

139 animal vaccination programs, elevating the risk of disease transmission between animals and
140 humans.

141 Additionally, many of West Africa's livelihoods depend on livestock and agriculture. Disease
142 outbreaks among livestock can thus compound the socioeconomic challenges posed by human
143 health crises (Rich and Wanyoike, 2010). Recognizing and addressing the intersection of animal
144 and human health is essential for a comprehensive understanding of outbreak impacts and for
145 devising effective intervention strategies.

146 Moreover, the broader health outcomes associated with these disruptions are multifaceted.
147 Beyond the immediate morbidity and mortality from the outbreak, there is a cascade of
148 secondary health implications. Reduced immunization coverage, for example, predisposes
149 populations to vaccine-preventable diseases (Cooper *et al.*, 2023). Additionally, mental health
150 issues, often overlooked, emerge as significant concerns, with people grappling with anxiety,
151 depression, and post-traumatic stress exacerbated by stigma and societal disruption (Shultz *et al.*,
152 2016).

153 This paper aims to elucidate the interconnections and implications of disruptions in healthcare
154 access, livelihoods, and health outcomes in West Africa during outbreaks. Through a
155 comprehensive review of the literature and primary data, we will explore the multi-dimensional
156 impacts of these disruptions and reflect upon strategies to mitigate their consequences.

157

158 **METHODS**

159

160 **Study Design**

161 This study adopted a mixed-methods approach, blending quantitative and qualitative data
162 collection techniques, to delve into the disruptions in healthcare access, livelihoods, and health
163 outcomes in West Africa amid outbreaks. Both prospective and retrospective data gathering were
164 undertaken, offering a holistic depiction of the healthcare landscape during the outbreak
165 episodes.

166 **Setting and Population**

167 The research spanned four pivotal West African countries: Nigeria, Sierra Leone, Guinea, and
168 Liberia. These countries were deliberately chosen considering their distinct experiences with
169 outbreaks, enabling a multifaceted perspective on healthcare disruptions.

170 **Data Collection Tools and Strategy**

171 A structured survey was curated and disseminated via the Open Data Kit (ODK) platform for the
172 quantitative dimension. Responses from these surveys were securely stored and managed on the
173 ONA server at <https://ona.io/waohecostudy>.

174 Parallely, the qualitative dimension encompassed in-depth interviews and focus group
175 discussions designed to extract richer insights into individual and community experiences,
176 perceptions, and recommendations surrounding healthcare disruptions.

177 Random sampling was employed to ensure representativeness across diverse population subsets.
178 The survey was directed using Geographic Information Systems (GIS), where geolocations of
179 potential participants were randomly chosen. This GIS-directed approach enabled a spatially
180 representative sample, ensuring that diverse geographical areas and their associated populations
181 were adequately represented.

182 **Sample Size Determination**

183 Using the OpenEpi tool, each country's sample sizes were independently deduced, anchored on
184 their respective SARS-CoV-2 seroprevalence rates. These computations were designed to

185 achieve a 95% confidence interval (CI), factoring in elements like design effect, projected
186 response rates, and possible contingencies such as non-responses or partial data.

187 **Quality Assurance**

188 To maintain the highest data quality and ensure methodological rigor, daily dashboard
189 monitoring was instituted to track and review incoming data in real time.

190 Surveyors documented their activities with photographs taken on-site, serving as a validation
191 mechanism. Regular supervisory site visits were conducted to oversee the data collection
192 process. Peer monitoring was employed, where teams cross-checked and validated each other's
193 work, ensuring consistency and accuracy.

194 **Data Analysis**

195 Quantitative datasets were subjected to rigorous statistical analysis, with descriptive statistics
196 elucidating the overall trends and inferential statistics employed for hypothesis testing and
197 inference derivation.

198
199 Concurrently, qualitative data, sourced from interviews and discussions was transcribed
200 verbatim. Thematic content analysis was then performed to discern emergent patterns and
201 themes contextualized within the overarching healthcare disruption narrative.

203 **Ethical Considerations**

204
205 Before their involvement, all participants rendered informed consent. Utmost confidentiality was
206 observed throughout, with data being anonymised or pseudonymised. Ethical clearance was
207 sought and obtained from relevant ethical review boards across the participating nations.

209 **RESULTS**

210 **Outbreaks and Meal Frequency**

211 We assessed the impact of an outbreak (SARS-CoV-2) on meal frequency patterns in Guinea,
212 Liberia, Sierra Leone, and Nigeria. We categorized meal frequencies as 'Once Daily,' 'Twice
213 Daily,' and 'Thrice Daily,' conducting statistical analyses to determine the significance of the
214 changes observed (Table 1).

215
216 In Guinea, there was a statistically significant increase in individuals consuming 'Once Daily'
217 meals, from 148 before the outbreak to 775 during the outbreak ($p < 0.001$). Liberia displayed a
218 similar trend in the increase of 'Once Daily' meal frequency, from 2,178 to 4,345 occurrences (p
219 < 0.001). Sierra Leone followed this regional pattern, with 'Once Daily' meal frequency rising
220 from 1,806 to 2,872 ($p < 0.001$). A significant reduction was also noted in 'Twice Daily' meals,
221 from 3,374 to 2,620 ($p < 0.001$).

222
223
224 Nigeria, while experiencing an increase in 'Once Daily' meal frequency from 11 to 98 ($p <$
225 0.001), showed an increase in 'Twice Daily' meals from 244 to 634 ($p < 0.001$).

226
227 The observed increase in 'Once Daily' meal frequencies across all countries examined suggests a
228 region-wide shift in dietary patterns due to the outbreak's impact, echoing the findings of global
229 health crises in the literature (Hendriks *et al.*, 2022; Koyratty *et al.*, 2022, 2022; Marti and
230 Puertas, 2022; Lewis *et al.*, 2023). The significant reduction in 'Thrice Daily' meals is consistent

231 with patterns observed in food-scarce situations elsewhere in the world(González-Monroy *et al.*,
232 2021).

233
234 The decline in 'Twice Daily' meals in Liberia, Sierra Leone and Guinea, contrary to the Nigerian
235 data, indicates varied national responses to the outbreak. The reduction may reflect a more severe
236 disruption of food systems compared to Nigeria, where an increase in 'Twice Daily' meals could
237 suggest a different form of dietary adaptation, perhaps due to varying economic resilience or
238 policy responses(Belton *et al.*, 2021; Schreiber *et al.*, 2022).

239
240 Statistically significant p-values affirm that these dietary changes are associated with the
241 outbreak conditions, supporting the theory that health crises exert considerable influence on food
242 security and nutritional behaviors (Beyene, 2023). These changes underscore the necessity for
243 robust food security interventions during health emergencies to prevent long-term adverse health
244 outcomes.

245
246

247 **Outbreaks and Household Protein Source Consumption in West Africa**

248 Our investigation into household protein source consumption across Sierra Leone, Liberia,
249 Nigeria, and Guinea provides critical insights into the dietary impacts of the recent outbreak.
250 Consistent with disruptions observed in food systems during public health emergencies, there
251 was a notable decline in the consumption of various protein sources during the outbreak(Béné,
252 2020; FAO, 2021).

253
254 Fig 1: Change in Protein Source Consumption During SARS-CoV-2

255
256 Before the outbreak, our data showed that common protein sources included bushmeat, beef, and
257 mutton(Fig.1). In the subsequent period, the intake of these proteins significantly decreased, with
258 the most pronounced reduction noted for bushmeat. This aligns with previous studies that have
259 highlighted bushmeat as a potential zoonotic risk, particularly in West African contexts (Karesh
260 *et al.*, 2005; Wolfe *et al.*, 2005). The decline in consumption of beef and mutton may be
261 attributed to disrupted supply chains and heightened food security concerns, phenomena which
262 are well-documented during health crises(World Bank, 2020).

263
264 Interestingly, the data suggest a more moderate decrease in the consumption of fish, which might
265 indicate a strategic pivot towards protein sources perceived to be safer or more readily available
266 during the outbreak(Headey and Martin, 2016). The differential changes across the various
267 protein sources suggest a complex interplay of factors influencing household dietary choices
268 during the outbreak(Barrett, 2010).

269
270 The observed shifts in consumption patterns underscored the profound effect of health crises on
271 food security and dietary habits, emphasizing the need for robust food systems that can adapt to
272 such challenges and protect vulnerable populations in the West African region(FAO, 2018).

273
274 Further, our qualitative study from West Africa suggests a complex interplay between
275 community skepticism towards health interventions and the practical aspects of food safety and
276 handling, with implications for the spread of foodborne diseases. Direct quotes from the study

277 encapsulate the concerns: In Sierra Leone, it was noted that "Community beliefs about disease
278 transmission from wild animals could lead to changes in hunting practices and dietary habits,
279 affecting traditional food sources." The study further records that lockdowns brought significant
280 challenges: "During lockdowns, community members find it difficult to feed their animals due to
281 restrictions on movement, potentially impacting livestock health and food availability."
282 Similarly, the situation in Nigeria mirrored these difficulties with animals also subjected to
283 movement restrictions by their owners. The repercussions in Liberia included shifts in social
284 behaviors: "Altered social norms and practices during outbreaks, such as reduced communal
285 gatherings, can impact food distribution and consumption patterns within communities." In
286 addition, the spread of misinformation about the origins of diseases from certain foods has
287 resulted in altered dietary choices: "Misinformation or beliefs about disease origins from certain
288 foods could lead to changes in dietary habits, affecting demand for specific food items." These
289 insights from the field reveal the breadth and depth of the pandemic's impact on traditional ways
290 of life and the potential risks that arise from altered food-related practices.

291
292 **Healthcare Service Interruptions in West African Countries**
293 During the SARS-CoV-2 pandemic, healthcare service interruptions were observed across
294 several countries. We focused our analysis on Sierra Leone, Liberia, Guinea, and Nigeria to
295 understand the extent of these interruptions.
296 In Sierra Leone, Liberia, and Guinea, the patterns of healthcare service interruptions were
297 notably similar (Fig 2). The three most frequently interrupted services in these countries were:
298 immunization, malaria treatment and routine bed net distribution.

299
300 Despite exhibiting a similar trend with immunization being the most interrupted service, Nigeria
301 displayed distinct differences in the ranking of other services compared to Sierra Leone, Liberia,
302 and Guinea.
303 The chart below provides a visual representation of the healthcare service interruptions in these
304 countries:
305

306 Fig. 2: Healthcare Services Disruption Reported by Country

307
308 By quantifying the interruptions, our analysis reveals the significant impact of the SARS-CoV-2
309 pandemic on essential healthcare services in West Africa. The disruption of crucial services,
310 such as immunisation, underscores the cascading effects of global health crises on regional
311 healthcare systems.
312 Liberia and Sierra Leone had more disruptions than Guinea and Nigeria (Fig 3).

313
314
315
316 Fig. 3: Choropleth Map of Healthcare Service Disruptions in West Africa

317
318 Additional in-depth analyses of one country, Sierra Leone, which had a high level of disruptions
319 revealed further the extent of service disruptions by outbreaks such as SARS-CoV-2.

320
321 **Disruptions in HIV Health Services**

322 Our detailed examination of the repercussions of the SARS-CoV-2 pandemic on specific HIV
323 services between 2019 and 2020 underscores significant variations in service provision (Fig 4).

324
325 Fig 4: Disruptions in HIV Services

326
327 The care and services provided to HIV-exposed infants were noticeably impacted: There was a
328 pronounced decline observed in the "HIV-exposed infant follow-up visit" service, indicating
329 potential barriers to continuous care during the early phase of the pandemic. The "HIV-exposed
330 infant first visit" also faced a substantial reduction, suggesting delays or challenges in initial care
331 access for this vulnerable group.

332 For adult HIV treatment, while some adult services highlighted resilience, others indicated
333 vulnerabilities: "ART Adult defaulted" for both males and females increased, emphasizing
334 potential challenges in maintaining treatment adherence during the pandemic. Interestingly,
335 "ART Adult given ARV new" for both genders witnessed growth. This could be interpreted as an
336 increase in new HIV diagnoses or improved access to treatment during this challenging period.

337 Child HIV Treatment services demonstrated mixed outcomes: The "ART Child defaulter"
338 service showcased stability, with only a slight decline, indicating the relative robustness of
339 treatment adherence mechanisms for children.

340 On the contrary, "Child HIV positive Cotrimoxazole prophylaxis start" experienced a decrease,
341 suggesting potential disruptions in preventive care for HIV-positive children.

342 In treatment transfers on treatment changes, the "ART Adult switched to 2nd line" service, a
343 marker for treatment change, showed a significant drop. This could point to potential barriers to
344 accessing alternative treatments when required.

345 Moreover, while the SARS-CoV-2 pandemic introduced challenges across the board in HIV
346 services during 2020, the degree and nature of these disruptions varied widely. Specific services,
347 especially those related to infants and treatment changes, faced pronounced challenges. These
348 findings emphasize the importance of bolstering resilience in healthcare systems, especially for
349 vulnerable demographics, in the face of global crises.

350
351 **Disruptions in TB Services**

352 We observed a decline in both the number of HIV-positive TB patients receiving antiretroviral
353 therapy (ART) and those receiving co-Trimoxazole preventive therapy from 2019 to 2020. This
354 trend is likely indicative of the disruptions caused by the COVID-19 pandemic, which affected
355 healthcare services globally, leading to challenges in accessing treatment for TB patients
356 (Fig.5)

357
358 Fig. 5: Trends in TB Treatment (2019-2021)

359
360 There was a slight increase in the number of patients receiving both forms of treatment from
361 2020 to 2021. This suggests a recovery or adaptation of TB services, possibly through the
362 implementation of new strategies to deliver care amidst pandemic constraints.

363
364 **Inpatient Service Disruption**

365

366 There was a dive in hospital admissions between 2019 and 2020 when SARS-CoV-2 started in
367 Sierra Leone and a resurgence thereafter (Fig 6). The dive could be attributed to initial shocks
368 from the outbreak.

369

370 Fig.6: Hospital Admissions and Year-Over-Year Changes in Sierra Leone

371 There is likely skepticism at the onset of outbreaks. A focus group discussion in Liberia noted
372 thus: "Whenever they come those who are in front of it, when they tell us, we always have the
373 belief that these people, that money they looking for again so that the notion we always carry.
374 Most of the time we can say that the government naa bring the sickness to put it on the people so
375 we can't listen to any advice." This reflects skepticism towards health interventions and a belief
376 that disease outbreaks might be financially motivated or artificially induced, which can
377 significantly affect healthcare seeking behavior and access.

378 There was a subsequent gradual, substantial rise in healthcare demand during the COVID-19
379 period, further emphasizing the pandemic's profound effect on hospital admissions.

380 There was a mix of increases and decreases in inpatient admissions across different categories
381 from 2018 to 2022. The years 2020 and 2021, which align with the height of the COVID-19
382 pandemic, showed pronounced changes in admissions, with some categories experiencing
383 significant disruptions (Fig. 7).

384

385 Fig.7: Hospital Admissions Trends

386

387 For the Intensive Care Admissions, a noticeable increase was observed from 2019 to 2021,
388 possibly due to the severity of COVID-19 cases. However, there was a decline in 2022.

389 There was a significant increase in the number of children with pneumonia admissions in 2021
390 and a slight decrease in 2022. The symptoms of pneumonia and COVID-19 can overlap,
391 potentially leading to more admissions.

392 The number of admissions for malaria in individuals aged 15 and above peaked in 2019, declined
393 in 2020 and 2021, and slightly rebounded in 2022. In the case of children with malaria, a decline
394 from 2019 to 2020 was observed, followed by a steady increase in the subsequent years.

395 The shifts in inpatient admissions could be attributed to multiple factors, including the direct
396 and indirect effects of the COVID-19 pandemic, changes in healthcare-seeking behaviors,
397 hospital resource allocations, and other external events. The increase in intensive care and
398 pneumonia-related admissions during the pandemic underscores the strain on healthcare
399 systems and the potential overlap in symptoms between COVID-19 and other illnesses.
400 Fluctuations in malaria admissions might reflect changes in healthcare access, seasonal
401 patterns, or broader public health interventions.

402

403 **Community Perceptions of Healthcare Access and Utilization**

404 The study in West Africa revealed a multifaceted understanding of disease outbreaks, availability
405 of healthcare infrastructure and health practices among community members. Respondents
406 highlighted key insights:

407 In the case of healthcare access and utilization: challenges in accessing quality healthcare were
408 prominent in all four countries, with many relying on pharmacies or traditional healers due to
409 distant or inadequate health facilities. There was a notable reliance on traditional medicine for
410 ailments during outbreaks.

411 "Most times, we go to the baba (traditional healer) because the hospital people don't treat us
412 well...." (Nigeria Qualitative Study Transcription Notes).

413 In Liberia, lights were shed on the decision to seek care and the gender equal opportunity for
414 community awareness campaigns for care. "In time pass it was just men but right after the Ebola
415 outbreak there are people train in the community to carry on awareness. In case of anything they
416 suspect they create awareness and to advise for the person to go to the facility even though it can
417 be difficult for some of them to go to the facility." (Liberian In-depth Interview). This reflects a
418 change in who takes charge of health safety in the community, shifting from a male-dominated
419 decision-making process to a more inclusive approach post-Ebola.

420 Despite the potential gender equal awareness campaigns, at the onset of outbreaks, there is a
421 tendency to boycott the healthcare facilities. "For us when they talk it, we can't believe it...They
422 will be explaining it to us we be going different way but when it attacks us naa then we will run
423 to the health center." (Liberian In-depth Interview). This quote suggests a common initial denial
424 of disease outbreaks among community members, followed by a reactive approach to seeking
425 healthcare once the disease has already affected them. Similar observations were noted in
426 Guinea, Sierra Leone and Nigeria.

427 **DISCUSSIONS**

428 This study has investigated the profound impacts of infectious disease outbreaks on livelihoods,
429 healthcare access, and health outcomes in West Africa, focusing on emerging diseases like
430 COVID-19. Our findings reveal the intricate link between health and socio-economic stability,
431 demonstrating how such outbreaks strain healthcare systems and disrupt socio-economic
432 structures (WHO, 2016).

433 The recurring outbreaks in West Africa challenge the region's fragile healthcare infrastructure.
434 Health facilities often face closures or reduced operations due to transmission fears, lack of
435 protective equipment, and workforce attrition, leading to decreased healthcare services (Kieny et
436 al., 2014). This exacerbates the burden of endemic diseases and hampers essential health services
437 like maternal and child care, increasing morbidity and mortality from non-outbreak-related
438 causes (Elston *et al.*, 2017).

439 Moreover, the socio-economic impact of these outbreaks extends beyond immediate health
440 concerns. Livelihoods are severely affected, with communities facing job losses, economic

441 instability, and increased poverty. This disruption in economic activities complicates recovery
442 and resilience(UNDP, 2020).

443 Our study underscores the need for robust healthcare systems and comprehensive preparedness
444 and response strategies that address both health and socio-economic aspects. Strengthening
445 healthcare infrastructure, ensuring an adequate supply of protective equipment, and maintaining
446 essential health services during outbreaks are crucial(WHO, 2015). Additionally, socio-economic
447 support for affected communities and measures to sustain livelihoods are vital in mitigating the
448 overall impact of such crises(Maxmen, 2020).

449 450 **CONCLUSIONS**

451 In conclusion, our research highlights the extensive disruptions caused by infectious disease
452 outbreaks in West Africa, affecting healthcare access, health outcomes, and socio-economic
453 contexts. The recurrent nature of these outbreaks necessitates an urgent and holistic approach to
454 healthcare system strengthening and preparedness(Kieny and Dovlo, 2015).

455 Our findings call for integrated strategies that address both the health and socio-economic
456 dimensions of outbreak response. Ensuring robust healthcare systems, alongside socio-economic
457 support mechanisms, is essential for the resilience of West African countries(Kieny and Dovlo,
458 2015; Gebremeskel *et al.*, 2021).

459
460 Future research should focus on developing and evaluating interventions that effectively bridge
461 the gap between health and socio-economic needs during outbreaks. The lessons learned from
462 West Africa are crucial for global strategies in managing infectious disease outbreaks,
463 emphasizing a multidimensional approach that considers both health and livelihoods(UNDP,
464 2020).

465
466 The resilience of West Africa against future health crises depends on learning from past
467 experiences and preparing proactively for multifaceted challenges. This study contributes to that
468 understanding and advocates for continued efforts to strengthen the region against the challenges
469 posed by infectious disease outbreaks(Maxmen, 2020).

470 471 472 **References**

473
474 Ansumana, R. *et al.* (2014) 'Ebola in Sierra Leone: A call for action', *The Lancet*, p. 303. doi:
475 10.1016/S0140-6736(14)61119-3.

476 Ansumana, R., Sankoh, O. and Zumla, A. (2020) 'Effects of disruption from COVID-19 on
477 antimalarial strategies', *Nature Medicine*. doi: 10.1038/s41591-020-1047-5.

- 478 Barrett, C. B. (2010) ‘Measuring Food Insecurity’, *Science*, 327(5967), pp. 825–828. doi:
479 10.1126/science.1182768.
- 480 Belton, B. *et al.* (2021) ‘COVID-19 impacts and adaptations in Asia and Africa’s aquatic food
481 value chains’, *Marine Policy*, 129, p. 104523. doi: 10.1016/j.marpol.2021.104523.
- 482 Béné, C. (2020) ‘Resilience of local food systems and links to food security - A review of some
483 important concepts in the context of COVID-19 and other shocks.’, *Food security*, 12(4), pp.
484 805–822. doi: 10.1007/s12571-020-01076-1.
- 485 Beyene, S. D. (2023) ‘The impact of food insecurity on health outcomes: empirical evidence
486 from sub-Saharan African countries’, *BMC Public Health*, 23(1), p. 338. doi: 10.1186/s12889-
487 023-15244-3.
- 488 Conrad, P. A., Meek, L. A. and Dumit, J. (2013) ‘Operationalizing a One Health approach to
489 global health challenges.’, *Comparative immunology, microbiology and infectious diseases*,
490 36(3), pp. 211–6. doi: 10.1016/j.cimid.2013.03.006.
- 491 Cooper, S. *et al.* (2023) ‘Vaccination coverage in rural Burkina Faso under the effects of
492 COVID-19: evidence from a panel study in eight districts.’, *BMC health services research*,
493 23(1), p. 1016. doi: 10.1186/s12913-023-10029-1.
- 494 Egeru, A., Dejene, S. W. and Siya, A. (2020) ‘Short report on implications of Covid-19 and
495 emerging zoonotic infectious diseases for pastoralists and Africa.’, *Pastoralism* □: *research,*
496 *policy and practice*, 10(1), p. 12. doi: 10.1186/s13570-020-00173-2.
- 497 Elston, J W T *et al.* (2017) ‘The health impact of the 2014-15 Ebola outbreak.’, *Public health*,
498 143, pp. 60–70. doi: 10.1016/j.puhe.2016.10.020.
- 499 Elston, J. W.T. *et al.* (2017) ‘The health impact of the 2014–15 Ebola outbreak’, *Public Health*.
500 doi: 10.1016/j.puhe.2016.10.020.
- 501 FAO (2018) *Nutrition and food systems. A report by the High Level Panel of Experts on Food*
502 *Security and Nutrition of the Committee on World Food Security*. ROME. Available at:
503 <https://www.fao.org/documents/card/en?details=I7846E>.
- 504 FAO (2021) *The State of Food and Agriculture 2021: Making agrifood systems more resilient to*
505 *shocks and stresses*. ROME: FAO. Available at:
506 <https://www.fao.org/documents/card/en?details=cb4476en>.
- 507 Gebremeskel, A. T. *et al.* (2021) ‘Building resilient health systems in Africa beyond the COVID-
508 19 pandemic response’, *BMJ Global Health*, 6(6), p. e006108. doi: 10.1136/bmjgh-2021-
509 006108.
- 510 González-Monroy, C. *et al.* (2021) ‘Eating Behaviour Changes during the COVID-19 Pandemic:
511 A Systematic Review of Longitudinal Studies’, *International Journal of Environmental*
512 *Research and Public Health*, 18(21), p. 11130. doi: 10.3390/ijerph182111130.
- 513 Headey, D. D. and Martin, W. J. (2016) ‘The Impact of Food Prices on Poverty and Food
514 Security’, *Annual Review of Resource Economics*, 8(1), pp. 329–351. doi: 10.1146/annurev-
515 resource-100815-095303.

- 516 Hendriks, S. L. *et al.* (2022) ‘Global environmental climate change, covid-19, and conflict
517 threaten food security and nutrition.’, *BMJ (Clinical research ed.)*, 378, p. e071534. doi:
518 10.1136/bmj-2022-071534.
- 519 Kieny, M. P. and Dovlo, D. (2015) ‘Beyond Ebola: a new agenda for resilient health systems’,
520 *The Lancet*, 385(9963), pp. 91–92. doi: 10.1016/S0140-6736(14)62479-X.
- 521 Kipchumba Kipruto, H. *et al.* (2023) ‘Estimating the economic impact of COVID-19 disruption
522 on access to sexual and reproductive health and rights in Eastern and Southern Africa.’,
523 *Frontiers in public health*, 11, p. 1144150. doi: 10.3389/fpubh.2023.1144150.
- 524 Kofman, A., Choi, M. J. and Rollin, P. E. (2019) ‘Lassa fever in travelers from West Africa,
525 1969–2016’, *Emerging Infectious Diseases*. doi: 10.3201/eid2502.180836.
- 526 Koyratty, N. *et al.* (2022) ‘Food Insecurity and COVID-19 Food-Related Perceptions, Practices,
527 and Problems: A 3-State Descriptive Study.’, *Disaster medicine and public health preparedness*,
528 17, p. e288. doi: 10.1017/dmp.2022.250.
- 529 Lewis, M. *et al.* (2023) ‘Healthy Food Prices Increased More Than the Prices of Unhealthy
530 Options during the COVID-19 Pandemic and Concurrent Challenges to the Food System.’,
531 *International journal of environmental research and public health*, 20(4). doi:
532 10.3390/ijerph20043146.
- 533 Marti, L. and Puertas, R. (2022) ‘Link between structural risk factors for adverse impacts of
534 COVID-19 and food insecurity in developed and developing countries.’, *Environment*,
535 *development and sustainability*, pp. 1–20. doi: 10.1007/s10668-022-02749-x.
- 536 Maxmen, A. (2020) ‘Ebola prepared these countries for coronavirus — but now even they are
537 floundering’, *Nature*, 583(7818), pp. 667–668. doi: 10.1038/d41586-020-02173-z.
- 538 Mylne, A. Q. N. *et al.* (2015) ‘Mapping the zoonotic niche of Lassa fever in Africa.’,
539 *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 109(8), pp. 483–92. doi:
540 10.1093/trstmh/trv047.
- 541 Ogbu, O., Ajuluchukwu, E. and Uneke, C. J. (2007) ‘Lassa fever in West African sub-region: an
542 overview.’, *Journal of vector borne diseases*, 44, pp. 1–11.
- 543 Olival, K. J. and Hayman, D. T. S. (2014) ‘Filoviruses in bats: current knowledge and future
544 directions.’, *Viruses*, 6(4), pp. 1759–88. doi: 10.3390/v6041759.
- 545 Parpia, A. S. *et al.* (2016) ‘Effects of Response to 2014-2015 Ebola Outbreak on Deaths from
546 Malaria, HIV/AIDS, and Tuberculosis, West Africa.’, *Emerging infectious diseases*, 22(3), pp.
547 433–41. doi: 10.3201/eid2203.150977.
- 548 R., A. *et al.* (2017) ‘Impact of infectious disease epidemics on tuberculosis diagnostic,
549 management, and prevention services: experiences and lessons from the 2014-2015 Ebola virus
550 disease outbreak in West Africa’, *International Journal of Infectious Diseases*.
- 551 Rich, K. M. and Wanyoike, F. (2010) ‘An assessment of the regional and national socio-
552 economic impacts of the 2007 Rift Valley fever outbreak in Kenya.’, *The American journal of*
553 *tropical medicine and hygiene*, 83(2 Suppl), pp. 52–7. doi: 10.4269/ajtmh.2010.09-0291.

- 554 Schreiber, K. *et al.* (2022) ‘Diverse adaptation strategies helped local food producers cope with
555 initial challenges of the Covid-19 pandemic: Lessons from Québec, Canada’, *Journal of Rural*
556 *Studies*, 90, pp. 124–133. doi: 10.1016/j.jrurstud.2022.02.002.
- 557 Shultz, J. M. *et al.* (2016) ‘The Role of Fear-Related Behaviors in the 2013-2016 West Africa
558 Ebola Virus Disease Outbreak.’, *Current psychiatry reports*, 18(11), p. 104. doi:
559 10.1007/s11920-016-0741-y.
- 560 Smith, C. and Whittaker, M. (2014) ‘Beyond mobile populations: a critical review of the
561 literature on malaria and population mobility and suggestions for future directions.’, *Malaria*
562 *journal*, 13, p. 307. doi: 10.1186/1475-2875-13-307.
- 563 Sumner, A., Hoy, C. and Ortiz-Juarez, E. (2020) ‘Estimates of the impact of COVID-19 on
564 global poverty’, *Unuwidder*.
- 565 Tambo, E., Adetunde, O. T. and Olalubi, O. A. (2018) ‘Re-emerging Lassa fever outbreaks in
566 Nigeria: Re-enforcing “One Health” community surveillance and emergency response practice’,
567 *Infectious Diseases of Poverty*. doi: 10.1186/s40249-018-0421-8.
- 568 UNDP (2020) *Human Development Report. The Next Frontier: Human Development and the*
569 *Anthropocene*. New York.
- 570 Vinck, P. *et al.* (2019) ‘Institutional trust and misinformation in the response to the 2018-19
571 Ebola outbreak in North Kivu, DR Congo: a population-based survey.’, *The Lancet. Infectious*
572 *diseases*, 19(5), pp. 529–536. doi: 10.1016/S1473-3099(19)30063-5.
- 573 WFP (2020) *COVID-19 will double number of people facing food crises unless swift action is*
574 *taken*. Available at: [https://www.wfp.org/news/covid-19-will-double-number-people-facing-](https://www.wfp.org/news/covid-19-will-double-number-people-facing-food-crises-unless-swift-action-taken)
575 [food-crises-unless-swift-action-taken](https://www.wfp.org/news/covid-19-will-double-number-people-facing-food-crises-unless-swift-action-taken) (Accessed: 10 August 2023).
- 576 WHO (2015) *Health worker Ebola infections in Guinea, Liberia and Sierra Leone*. Available at:
577 http://www.who.int/hrh/documents/21may2015_web_final.pdf (Accessed: 5 October 2016).
- 578 WHO (2016) ‘After Ebola in West Africa — Unpredictable Risks, Preventable Epidemics’, *New*
579 *England Journal of Medicine*, 375(6), pp. 587–596. doi: 10.1056/NEJMSr1513109.
- 580 World Bank (2020) *Responding to the Emerging Food Security Crisis*. Washington DC.
581 Available at: [https://documents.worldbank.org/en/publication/documents-](https://documents.worldbank.org/en/publication/documents-reports/documentdetail/775981606955884100/responding-to-the-emerging-food-security-crisis)
582 [reports/documentdetail/775981606955884100/responding-to-the-emerging-food-security-crisis](https://documents.worldbank.org/en/publication/documents-reports/documentdetail/775981606955884100/responding-to-the-emerging-food-security-crisis).

583

584 **CONFLICT OF INTEREST**

585 All authors declare that there is not conflict of interest

586

587 **AUTHOR CONTRIBUTIONS**

588

589 RA, ASB, SC, JG, NM, SM, NH, OD, GM were involved in study conception. RA, ASB, JML,
590 MK were involved in drafting the manuscript. All authors reviewed and revised the manuscript.

591 JML and ASB developed data collection tool. RA, ASB, MK, JML, JG, SC, EC, NM, supervised
592 data collection. RA, ASB, analyzed data.
593

594 **FUNDING**

595 Funding for this study was provided by the Canadian, International Development Research
596 Center under the Collaborative One Health Research Initiative on Epidemics (COHRIE).

597 **ACKNOWLEDGMENTS**

598 We acknowledge all data collectors in Sierra Leone, Guinea, Liberia and Nigeria.

599 **DATA AVAILABILITY STATEMENT**

600 Datasets generated for this study can be found at <https://ona.io/waohecostudy/197244> and
601 <https://sl.dhis2.org/>. Transcripts of qualitative data are available on request.
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627


Table 1: Meal Frequency Table


Country	Meal Frequency	Before Outbreak	During Outbreak	P-Value
Guinea	Once Daily	148	775	0.00

	Thrice Daily	3687	3078	0.00
	Twice Daily	1721	1689	0.72
Liberia	Once Daily	2178	4345	0.00
	Thrice Daily	310	70	0.00
	Twice Daily	3222	1280	0.00
Nigeria	Once Daily	11	98	0.00
	Thrice Daily	1212	755	0.00
	Twice Daily	244	634	0.00
Sierra Leone	Once Daily	1806	2872	0.00
	Thrice Daily	523	203	0.00
	Twice Daily	3374	2620	0.00

628

629





a: Pharmaceutical services	k: Interrupted Pentavalent vaccination
b: Interrupted Antenatal Care	l: interrupted Reproductive Health Services
c: Interrupted Family Planning	m: Interrupted Routine Bednet Distribution
d: Interrupted HIV & AIDS Services	n: Interrupted Services for NCDs
e: Interrupted Immunization	o: Interrupted Severe Acute Malnutrition Services
f: Interrupted Intermittent Preventive Treatment in Pregnancy (IPTp)	p: Interrupted Surgery
g: interrupted Malaria Treatment	q: Interrupted TB Services
h: interrupted Nutrition Preventive	r: Interrupted Vitamin A supplementation for 12_59_months
i: Interrupted Other	s: Interrupted BCG among Under 1-year Children
j: Interrupted Outpatient Services	


Fig. 2: Healthcare Services Disruption Reported by Country

Rate of Healthcare Service Disruptions by Country




Top Disruptions in HIV Services from 2019 to 2020


Service




Trends in TB Treatment (2019-2021)



Hospital Admissions and Year-Over-Year Change in Sierra Leone



Hospital Admissions Trends (2018-2022)



- Intensive Care Admissions
- Pneumonia Admissions (Children)
- Malaria Admissions (15+)
- Malaria Admissions (Children)
- Admission - Child with Diarrhoea
- Separation - Surgical
- Admission - Antenatal Client with Malaria
- Separation - TB
- Admission - Ebola Survivor
- Separation - Psychiatry

medRxiv preprint doi: <https://doi.org/10.1101/2024.04.27.24101010>; this version posted April 29, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.