Anti-tetanus vaccination is associated with reduced occurrence and slower progression of Parkinson's disease

Ariel Israel^{1,2}*, Eli Magen³, Eugene Merzon^{1,4}, Eytan Ruppin⁵, Shlomo Vinker^{1,6}, Nir Giladi⁷

5 **Affiliations:**

	¹ Leumit Research Institute, Leumit Health Services; Tel-Aviv, Israel
	² Department of Epidemiology and Preventive Medicine, School of Public Health, Faculty of Medical & Health Sciences, Tel Aviv University; Tel-Aviv, Israel
10	³ Medicine A Department, Assuta Ashdod University Medical Center, Ben Gurion University of the Negev; Beer Sheva, Israel
	⁴ Adelson School of Medicine, Ariel University; Ariel, Israel.
	⁵ Cancer Data Science Laboratory, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, USA
	⁶ Department of Family Medicine, Faculty of Medical & Health Sciences, Tel-Aviv
15	⁷ Brain Institute, Tel-Aviv Sourasky Medical Center, Faculty of Medical & Health Sciences, Sagol School of Neurosciences, Tel-Aviv University, Tel-Aviv, Israel.

*Corresponding author. Email: aisrael@leumit.co.il.

Abstract:

5

Parkinson's disease (PD) is characterized by progressive neurodegeneration in the autonomic and central nervous systems, manifesting with hallmark symptoms of resting tremor, bradykinesia and rigidity. The etiology of PD remains elusive, and currently available treatments do not halt disease progression.

In this study, conducted within a national health provider, we examined the impact of vaccination and medication purchase on PD occurrence and severity, using an innovative

10 machine learning algorithm to track disease progression.

Our findings reveal a significant reduction in PD occurrence following anti-tetanus vaccination, with a time-dependent association between the elapsed time since vaccination and both the rate and progression of PD. These results are supported by

15 evidence that antimicrobial treatments significantly alter disease severity, suggesting the actual involvement of *Clostridium Tetani* in PD pathology.

Thus, tetanus vaccination and *C. tetani* eradication could be promising strategies for preventing PD and slowing its progression, pending controlled clinical trials.

Parkinson's disease (PD) is a neurodegenerative disorder affecting over six million individuals worldwide.¹ PD is characterized by the progressive loss of neurons throughout the peripheral autonomic and central nervous systems. Current diagnosis of PD is based on the presence of resting tremor, bradykinesia, rigidity and postural response abnormalities which are tightly associated with dopaminergic loss. The etiology of PD remains poorly understood, although aging, genetic and environmental factors have been identified as risk factors. Currently, only exercise has shown an effect on disease progression and no approved treatment targets the basic mechanism of the disease or its direct cause.

5

20

- Clostridia are gram-negative obligate anaerobe bacteria that are prevalent in the environment. Several members of the Clostridium species produce highly potent toxins ². Notably, *C. tetani* can produce Tetanus neurotoxin, the causative agent of tetanus.^{3,4} In tetanus, the neurotoxin is internalized into signaling endosomes and transported retrogradely to the neuronal soma, interfering with the release of neurotransmitters,
- notably glycine and gamma-aminobutyric acid, blocking inhibitory impulses.⁵ Lacking inhibitory neurotransmission, stimulation of motor neurons increases, producing rigidity, unopposed muscle contraction and spasm.⁶

To protect from the severe risks associated with tetanus disease, the combined Tetanus and Diphtheria toxoid (TD) vaccine is routinely administered to infants, children and adolescents as part of the standard immunization programs.^{7,8} Adults usually receive antitetanus booster vaccination upon clinical indication, when they present in clinic with a wound susceptible to be infected with *C. tetani*, and there is no record of vaccination in the last ten years (or five years in a dirty wound at high risk for contamination). In a

serologic survey performed in the United States between 1988 and 1994, fully protective levels of anti-tetanus and anti-diphtheria antibodies were detected in 91% of individuals aged 6 to 11 years but in only 47% of individuals 20 years or older ^{9,10}. Interestingly, PD happens to be diagnosed at the late adult age, typically after 45, when tetanus antibody protection acquired from childhood and adolescence vaccination is no more protective. Recent studies have revealed significant differences in the gut microbiota composition between patients with Parkinson's Disease (PD) and healthy controls ¹¹. Notably, alterations were observed for Clostridia species.¹² *C. tetani* has been isolated in the feces from adult individuals,¹³ suggesting it might be a common host of the human gut. Given its potential to cause synaptic dysfunction, clostridium neurotoxin might play a role in the neurodegenerative process responsible for PD. If a Clostridium neurotoxin contributes to the pathogenesis of PD, an inverse correlation would be expected between the administration of the tetanus toxoid vaccine and the incidence of PD.

The study cohort

5

10

We performed a large-scale observational study in Leumit Health Services (LHS), a nationwide health organization in Israel with over 22 years of centrally maintained electronic health records (EHR). Using rigorous criteria, we selected 1446 patients who received a PD diagnosis between the ages of 45 and 75, avoiding the rare cases occurring in very young age (<45), where disease is heavily influenced by genetic or environmental factors, and older patients (>75), where non-specific motor symptoms make the clinical diagnosis of PD less reliable. We used the earliest diagnosis date or antiparkinsonian medication purchase as the index date, and selected 7230 control individuals matching in a 5:1 ratio the PD patients. Control individuals, were assigned the same index dates as

their respective cases, so that the same depth of recorded EHR history was available in the two groups.

Table 1 presents the demographic and clinical comparison of the two at the index date. The age, gender and socioeconomic distribution of the two groups are very similar. In both groups 55.6% were male, with an average age of 65.7 ± 7.2 . Clinical characteristics also appear to be similar, with the notable exception of smoking status, with PD patients

being less likely to be smokers: Odds Ratio for current smoking (OR=0.59; P<0.001),

consistent with literature.^{14,15}

5

20

Vaccination is associated with decreased risk of PD occurrence.

The last line of Table 1 displays a striking difference between the groups for TD vaccination status: only 1.64% of patients with PD had a record of vaccination before the index date, compared to 3.19% in the control group (OR=0.50, P=0.001). A characteristic feature of a disease occurring because of waning antibody protection is a progressive increase in the disease rate with time elapsing since vaccination, as our group has shown for SARS-CoV-2.^{16,17} We therefore assessed the relationship between time elapsed since last vaccination and PD risk.

PD risk in vaccinated patients is associated with time elapsed since vaccination

Figure 1 displays comparisons between the case and control groups according to the timing of vaccination. As the forest plot shows clearly, there is a time-dependent protection effect of the TD vaccination. The odds-ratio for PD occurring within two years after last TD vaccination is 0.00 (P=0.006), within 5 years, it is 0.17 (P=0.003). The odds increase to 0.26 between 5 to 10 years (P=0.004). Between 10- and 15-years post-vaccination, there is still a trend for protection, but the odds difference loses

statistical significance. After 15 years post-vaccination the trend is for increased risk, not statistically significant.

If not getting a TD vaccination was to be a confounder of some early symptoms of PD (e.g. people with early signs of PD, before diagnosis, may decrease their activity and hence reduce opportunities to get wounded, and subsequently get vaccination), then one would expect that this pattern would continue and even amplify after the index date. But this does not occur: instead, we observe an opposite trend, of increased odds for PD among those who got a vaccine after the index date (OR=1.33, P=0.39). This reversing time relationship strongly indicates that lack of TD vaccination is unlikely to be a

10 confounder for early PD.

5

20

Having shown that PD occurrence is highly associated with time elapsed since last TD vaccination, we proceeded to verify, in the few patients who were either diagnosed with PD after vaccination, or were vaccinated once PD was diagnosed, whether disease course was affected by vaccination.

15 Vaccinated PD patients have slower rates of disease progression

In order to follow the disease course, we developed a method to accurately assess disease severity during the follow-up period. Parkinson's disease is characterized by progressive course of motor and functional deterioration as shown by Hoehn & Yahr's clinical staging of the laterality and axial symptoms¹⁸ and by the Unified Parkinson's Disease Rating Scale (UPDRS).¹⁹ Symptoms are typically treated by antiparkinsonian medications that are increased over time as the disease progresses in dosage and potency. Having records of medications purchased by each patient, we could use the annual medication consumption to train a machine learning model, that would assess disease

severity for each patient and year of disease. For this purpose, we trained a gradient boosting model by cross-validation folds, utilizing yearly medication consumption per catalog entry as training variables, together with the gender of the patient (since disease course is affected by gender), to estimate time elapsed since disease onset, set as the target variable. The output of the model is a PD severity score expressed in a scale analogous to years of disease (e.g. a patient with a severity score of 6 in a given year has purchased medications that are typically used by a PD patient in the 6th year of the disease). Using this model, we computed severity scores for 8,793 PD patient years of disease, of which 201 were from patients with prior vaccination record.

5

20

Figure 2A displays a kernel density plot of these scores vs. the actual disease duration, along with a linear regression line. The calculated severity scores correlate nicely with the actual year of the disease. The Pearson correlation coefficient, r is 0.443, with a Pvalue under 10^{-200} . Of note, a quadratic regression model (a polynomial model of degree 2) was slightly more informative, exhibiting a lower Akaike Information Criterion (AIC) and a higher R² compared to the linear model (see Fig. 2B).

Figure 2C displays the severity scores over time, according to vaccination status. Disease severity in vaccinated patients (orange curve) was significantly lower than in non-vaccinated patients. Linear regression models show vaccination status is significantly associated with reduced severity, both in a univariable model (P<0.001, Table S1), and in a multivariable model adjusted for age, gender and smoking status (P<0.001, Table S2). Here again the quadratic regression is more informative (Table S3), so we use a quadratic model to plot disease severity according to time elapsed since PD diagnosis: Figure 2D shows that, even though disease severity was roughly similar at disease onset, disease

progression was significantly slower in the vaccinated group. Their regression curve shows that the average severity in vaccinated patients at the seventh year of the disease (red line) was similar to the average severity score of unvaccinated patients who were at the third year of the disease.

5 We see here, that even among PD patients, disease course was less severe postvaccination than in PD patients with no prior vaccination record.

Disease progression is associated with time elapsed since vaccination

10

15

20

We have shown that PD progresses more slowly in patients who have been vaccinated than in those who did not. A further indication that TD vaccine affects PD progression would be to find, focusing on the vaccinated patients, an effect that is more pronounced when closer to the date of vaccination, even after correcting for time since diagnosis. For this analysis, which requires comparison of patients who have experienced varying durations of the disease, we calculated "relative PD severity scores", which are obtained by subtracting the mean severity score of patients at the same disease year from the severity scores.

Figure 2E displays these relative PD severity scores according to time since diagnosis, divided by vaccination status, showing once again that vaccinated individuals have lower disease severity (P<0.001, Table S4). Figure 2F follows the relative disease severity of vaccinated patients according to time elapsed since last vaccination. The relative disease severity decrease is correlated with time elapsed since vaccination (r=0.19, P=0.03, Tables S5 and S6 for crude and adjusted regression analyses).

Disease severity is affected by antimicrobial treatments

If C. tetani present in the patients' microbiome are actually involved in PD pathology, then PD disease severity should be affected by antimicrobial treatments that kill these bacteria. Having calculated PD disease severity scores, we can use them to assess 5 whether antimicrobial courses are actually associated with changes in disease severity. For this analysis, we identified forty classes of medications for which we observed varying consumption rates between PD patients and control patients in the years prior to the index date, which is a potential signal for an effect on disease occurrence.²⁰ For each of these classes, and for each year of disease in a PD patient, we calculated a variable that 10 reflects whether medications of the class were purchased by the patient over the three preceding years. Starting with a multivariable regression model that includes age, gender, and smoking status, we employed a stepwise approach to incrementally select the ten medication classes most strongly associated with the relative PD severity scores. Table 2 presents the results of this regression. Not surprisingly, the tetanus toxoid vaccine had the 15 strongest effect on disease severity, purchase of the vaccine in the preceding three years significantly reduced the relative disease severity (-1.11, P=0.0003).

Several antibiotic agents displayed a substantial and significant effect in reducing disease severity: benzathine penicillin, which is an intramuscular formulation of penicillin, displayed a significant reducing effect on disease severity (-0.69, P=0.003). Benzathine penicillin is the classical treatment for tetanus,²¹ its sustained release formulation is likely to deplete the clostridial population producing the toxin, explaining its beneficial effect on disease severity. Likewise, Cefadroxil, a first generation cephalosporine structurally close to penicillin displayed a significant decreasing effect on disease severity (-1.02,

20

P=0.003). Two macrolide antibiotics also displayed a beneficial effect on disease severity: erythromycin (-0.74, P=0.0002) and roxithromycin (-0.13, P=0.008), as well as two compounds of the fluoroquinolone family: ofloxacin (-0.45, P<0.001), and levofloxacin (-0.22, P=0.026).

- 5 On the other hand, Clindamycin was associated with a significant and substantial increase in disease severity (+0.40, P=0.002). Clindamycin is primarily used to treat infections caused by susceptible anaerobic bacteria, but it is notorious for causing *Clostridium difficile* colitis, a dangerous condition in which Clostridium bacteria, inherently resistant to clindamycin, colonize the human colon.²² The observed effect of increasing severity of
- PD is consistent with a competitive advantage granted by this antibiotic to clostridia, over susceptible anaerobes present in the microbiome, enabling proliferation of *C. tetani* in a manner similar to *C. difficile* proliferation following Clindamycin treatment.
 Interestingly, we also observed strong apposing effects for two laxative agents commonly used to treat constipation in PD patients. Macrogol (polyethylene glycol), was
- significantly associated with increased disease severity (+0.34, P<0.001), while ispaghula (Psyllium), a dietary fiber, was associated with decreased disease severity (-0.42, P=0.0004). Since these two compounds primary affect the gut environment, these opposite effects are strongly suggestive of a pivotal role played by an organism which resides in the intestinal microbiome.
- 20 We tested whether a dose-response relationship could be detected, by looking for a correlation between the number of purchases by each patient during the three preceding years and the relative PD severity score. **Figure 3** display scatter plots and correlation analyses for the identified compounds. For most compounds we observe a significant

dose-response relationship strengthening the direction of the association detected by the regression. Since the regression was based only on a binary variable for each compound, and did not account for the number of purchases, a significant dose-response relationship provides an independent corroboration of the effect of these medications.

5 Discussion

10

This large-scale, real-life population study provides compelling evidence for the protective effect of diphtheria-tetanus (DT) toxoid vaccination on PD risk and its rate of progression. The significant associations observed between time elapsed since vaccination and both PD occurrence and rate of disease progression are all supporting the interpretation that the anti-tetanus vaccine provides protection against PD, and that this protection wanes over time. In addition, disease progression is significantly associated, in a dose-dependent manner, to antibiotics that affect clostridium growth negatively (betalactam, macrolides, fluroquinolones), or positively (clindamycin), strongly supporting the involvement of actual clostridium bacteria as the causative agent of disease.

Clostridia are known to reside in the microbiome, and several studies have reported differential prevalence of clostridial species in the gut microbiome of PD patients.^{11,12} There is strong indication that *C. tetani* is a common host of the human microbiome: a study performed in Vietnam incidentally reported *C. tetani* presence in the feces of 42% of control subjects.¹³ *C. tetani* are also known to be present in the saliva, and
contamination of a wound with saliva in an unvaccinated individual is indication for treatment with tetanus immunoglobulin.²³ *C. Tetani* spores are prevalent in the human environment, extremely resistant: properties that should allow this organism to progressively colonize the human flora. *C. tetani* toxin secreted in the gut would

primarily affect the mesenteric neurons that regulate peristalsis. Remarkably, impairment of the autonomic nervous system of the gut, manifested as constipation and bloating, happens to be a very common prodromal non-motor symptom of PD.²⁴

- The subsequent spreading of *C. tetani* to other microbiome niches would explain how the pathology of PD is further observed in multiple foci. *C. tetani*, present in the nasopharynx would damage by its neurotoxin the olfactory neurons, affecting smell.²⁵ Once significant damage is incurred to dopamine secreting neurons within the basal ganglia via retrograde axonal transport,²⁶ the disease manifests by the characteristic movement disturbances of PD,²⁷ while nonspecific neurodegeneration presents as depression and dementia.²⁸ In the
- 10 mouth, C. tetani would affect the innervation of the salivary glands (causing drooling); invasion of the sebaceous glands would dysregulate sweating, a common PD symptom. The oropharynx and the paranasal sinuses, which lie in close proximity to the CNS, are the primary sites we suspect to harbor C. tetani in PD. In these sensitive sites, even trace amounts of the extremely potent tetanus neurotoxin would be expected to cause over time 15 synaptic dysfunction and subsequent damage to neighboring neuronal tissue. The α synuclein aggregates that are the hallmark of PD could hence be a marker of synaptic dysfunction triggered by the toxin rather than the original cause of neurodegeneration. Given its prevalence in the environment, presence of *C. tetani* in the microbiome, even after loss of the humoral response that could neutralize its toxin, is certainly not the sole 20 factor of PD occurrence. Genetic and environmental factors, as well as lifestyle and comorbidities, likely influence the ability of C. tetani to colonize susceptible niches and harm adjacent neuronal tissue. Interestingly, cigarette smoking has been shown to inhibit

Clostridium growth,²⁹ hence, mere inhibition of C. *tetani* proliferation can explain the decreased risk observed for PD in smokers.^{14,15}

In conclusion, the findings presented in this study strongly support a causative role played by *C. tetani* in the pathological process leading to diffuse and progressive neurodegeneration observed in PD. Based on solid epidemiological data, we propose here a new paradigm that offers a mechanistic explanation of PD pathogenesis and its wide diversity of symptoms progression, in the form of neurotoxin mediated damage caused by *C. tetani* colonizing the microbiome at sensitive sites.

5

This study suggests that vaccination against tetanus neurotoxin (active or passive),

- 10 possibly combined with treatments aimed at eradicating *C. tetani* from body reservoirs, could offer promising avenues to prevent Parkinson's Disease (PD) occurrence and slow disease progression. Our findings make a strong case for a large-scale prospective randomized controlled trial to test and validate the protective effect of anti-tetanus vaccination on PD progression in early and advanced stages of PD, as well as in subjects
- at risk. Additionally, further research should explore factors affecting *C. tetani* proliferation and colonization potential, particularly in relation to medication, diet, food
 supplements, hygiene products, and lifestyle habits.

Methods

Ethics Statement

The Institutional Review Board (IRB) of Leumit Health Services gave ethical approval for this work (LEU-0001-24), with a waiver of informed consent, since data were analyzed retrospectively and anonymously.

Study design

5

10

15

20

This research was conducted as an observational cohort study in Leumit Health Services (LHS), one of the four national health providers in Israel, providing comprehensive healthcare services to approximately 720,000 members. All Israeli citizens are entitled for comprehensive health insurance and receive a standardized package of health services and medications, as defined by the national "Health Basket" committee. LHS operates a centralized electronic health records (EHR) system, with over two decades of meticulously maintained information on patient demographics, medical diagnoses, healthcare encounters, laboratory test results, and records of prescribed and purchased medications. Diagnoses are documented during medical encounters by the treating physicians using the International Classification of Diseases, Ninth Revision (ICD-9). Diagnoses can be marked as chronic when they pertain to a chronic condition, and these can be updated or closed by the treating physicians during subsequent patient encounters. The reliability of these chronic diagnosis records in our registry has been previously validated, demonstrating high accuracy.¹⁶

Eligibility for inclusion in the study was defined as any past or current LHS member with at least five years of LHS membership between years 2003 and 2023. Data extraction was

carried out from the LHS central data warehouse in February 2024, encompassing diagnoses, results of laboratory tests, and medication purchases recorded up to December 31, 2023.

5 Cohort definition

10

20

The study cohort comprises eligible PD patients diagnosed for the first time between ages 45 and 75, alongside a control group matched at a 5:1 ratio, of individuals with no documented PD. PD patients were identified by the presence of ICD-9 coded 332 diagnosis "Parkinson's Disease" in the EHR, if recorded by a neurologist or a movement disorder specialist, or recorded by any physician, if accompanied with purchase of Antiparkinsonian medications (listed in Table S7) over a period of more than six months. In order to avoid confusion with overlapping conditions, individuals with a diagnosis

indicative of secondary parkinsonism, schizophrenia, pituitary adenoma, restless legs syndrome, or cerebrovascular accident were excluded from the study (diagnoses listed in

Table S8), as well of individuals with prior purchase of an antipsychotic medication susceptible to induce parkinsonism (medications listed in Table S9).

Controls were precisely matched to the PD patients based on gender, socio-economic status category, and the year of initial enrollment in Leumit Health Services (LHS). For each PD patient, five control individuals were chosen who met these matching criteria and whose birth dates were closest to that of the PD patient, ensuring no individual was duplicated within the cohort.

Diphtheria-Tetanus toxoid vaccinations

We looked for anti-tetanus vaccinations in the medical history of patients from the study cohort, using both vaccine purchase recorded by LHS pharmacies, and vaccine administration records documented in the EHRs. Most anti-tetanus vaccinations

5 performed in LHS for patients in the cohort were with DT IMOVAX from Sanofi-Pasteur, adult dose, 0.5ml.

Data preparation

Data were extracted from electronic health records and prepared for analyses using

10 scripts developed by Leumit Research Institute in Python 3.11 with Pandas library and T-SQL queries. Prior to analysis, patients' data were deidentified and the patient ID number replaced with an identifier internal to the study.

Statistical analyses

15 Statistical analyses were performed in R version 4.3. Unless specified otherwise, Fisher's exact test was used to compare categorical variables and the two-tailed two samples t-test to compare continuous variables across groups. Conditional logistic models were fit to assess the association of vaccination timing and covariates across matched groups. Linear regression models were fit to assess the relationship between linear variables such as disease severity and relative disease severity and explanatory variables. Pearson correlation analyses were performed to assess dose-response relationship. Graphs of this study were produced in Python with *seaborn* and *matplotlib* libraries.

A.I. and E.Me had access to all the original data and serve as guarantor of the study integrity.

Machine Learning Model

5 Gradient boosting model for predicting disease activity score was built with *lightgbm* python library following exploration of model parameters space performed with *FLAML/AutoML* with 5-fold cross-validation. The predicting features of the model are provided as Supplementary Table S10.

10 References and Notes

15

20

- Ray Dorsey, E. *et al.* Global, regional, and national burden of Parkinson's disease, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 17, 939–953 (2018).
- Popoff, M. R. & Bouvet, P. Clostridial toxins. *Future Microbiol* 4, 1021–64 (2009).
- 3. Eisel, U. *et al.* Tetanus toxin: primary structure, expression in E. coli, and homology with botulinum toxins. *EMBO J* **5**, 2495–502 (1986).
- Brüggemann, H. *et al.* The genome sequence of *Clostridium tetani*, the causative agent of tetanus disease. *Proceedings of the National Academy of Sciences* 100, 1316–1321 (2003).
- Pirazzini, M., Montecucco, C. & Rossetto, O. Toxicology and pharmacology of botulinum and tetanus neurotoxins: an update. *Arch Toxicol* 96, 1521–1539 (2022).

- 6. Hassel, B. Tetanus: Pathophysiology, Treatment, and the Possibility of Using Botulinum Toxin against Tetanus-Induced Rigidity and Spasms. Toxins (Basel) 5, 73-83 (2013).
- 7. Centers for Disease Control and Prevention. Diphtheria, Tetanus, and Pertussis Vaccine Recommendations. https://www.cdc.gov/vaccines/vpd/dtap-tdaptd/hcp/recommendations.html https://www.cdc.gov/vaccines/vpd/dtap-tdaptd/hcp/recommendations.html.
 - 8. Israeli Ministry of Health. Tetanus Vaccine. https://www.gov.il/en/pages/tetanusvaccine.
- 10 9. Gergen, P. J. et al. A Population-Based Serologic Survey of Immunity to Tetanus in the United States. New England Journal of Medicine 332, 761–767 (1995).
 - 10. McQuillan, G. M., Kruszon-Moran, D., Deforest, A., Chu, S. Y. & Wharton, M. Serologic Immunity to Diphtheria and Tetanus in the United States. Ann Intern *Med* **136**, 660 (2002).
- 15 11. Li, Z. et al. Gut bacterial profiles in Parkinson's disease: A systematic review. CNS Neurosci Ther 29, 140–157 (2023).
 - Keshavarzian, A. et al. Colonic bacterial composition in Parkinson's disease. 12. *Movement Disorders* **30**, 1351–1360 (2015).
 - 13. Hao, N. Van et al. The Role of the Gastrointestinal Tract in Toxigenic Clostridium tetani Infection: A Case-Control Study. Am J Trop Med Hyg 105, 494–497 (2021).
 - Mappin-Kasirer, B. et al. Tobacco smoking and the risk of Parkinson disease. 14. *Neurology* **94**, (2020).

5

	15.	Kandinov, B., Giladi, N. & Korczyn, A. D. Smoking and tea consumption delay
		onset of Parkinson's disease. Parkinsonism Relat Disord 15, 41-46 (2009).
	16.	Israel, A. et al. Elapsed time since BNT162b2 vaccine and risk of SARS-CoV-2
		infection: test negative design study. The BMJ 375, e067873 (2021).
5	17.	Israel, A. et al. Large-Scale Study of Antibody Titer Decay following BNT162b2
		mRNA Vaccine or SARS-CoV-2 Infection. Vaccines (Basel) 10, 64 (2021).
	18.	Hoehn, M. M. & Yahr, M. D. Parkinsonism. Neurology 17, 427–427 (1967).
	19.	Ramaker, C., Marinus, J., Stiggelbout, A. M. & van Hilten, B. J. Systematic
		evaluation of rating scales for impairment and disability in Parkinson's disease.
10		Movement Disorders 17, 867–876 (2002).
	20.	Israel, A. et al. Identification of drugs associated with reduced severity of covid-19
		– a case-control study in a large population. <i>Elife</i> 10 , (2021).
	21.	Smith, J. W. G. Penicillin in Prevention of Tetanus. Br Med J 2, 1293 (1964).
	22.	Duffy, C. R. et al. Clindamycin, Gentamicin, and Risk of Clostridium difficile
15		Infection and Acute Kidney Injury During Delivery Hospitalizations. Obstetrics &
		<i>Gynecology</i> 135 , 59–67 (2020).
	23.	Centers for Disease Control and Prevention. Epidemiology and Prevention of
		Vaccine-Preventable Diseases.
	24.	Xu, J., Wang, L., Chen, X. & Le, W. New Understanding on the Pathophysiology

20 and Treatment of Constipation in Parkinson's Disease. *Front Aging Neurosci* **14**, (2022).

- 25. Keller, A., Sweeney, S. T., Zars, T., O'Kane, C. J. & Heisenberg, M. Targeted expression of tetanus neurotoxin interferes with behavioral responses to sensory input in *Drosophila*. *J Neurobiol* **50**, 221–233 (2002).
- 26. Ovespian, S. V., Bodeker, M., O'Leary, V. B., Lawrence, G. W. & Oliver Dolly, J. Internalization and retrograde axonal trafficking of tetanus toxin in motor neurons and trans-synaptic propagation at central synapses exceed those of its C-terminalbinding fragments. *Brain Struct Funct* **220**, 1825–1838 (2015).
 - 27. Blandini, F., Nappi, G., Tassorelli, C. & Martignoni, E. Functional changes of the basal ganglia circuitry in Parkinson's disease. *Prog Neurobiol* **62**, 63–88 (2000).
- Bagetta, G. & Nisticò, G. Tetanus toxin as a neurobiological tool to study mechanisms of neuronal cell death in the mammalian brain. *Pharmacol Ther* 62, 29–39 (1994).
 - Wang, X. *et al.* Active Smoking Induces Aberrations in Digestive Tract Microbiota of Rats. *Front Cell Infect Microbiol* 11, (2021).

15

5

End Notes: Acknowledgements:

We thank Alejandro A. Schäffer for his assistance in preparation of this manuscript.

The content of this publication does not necessarily reflect the views or policies of the
 Department of Health and Human Services, nor does mention of trade names,
 commercial products, or organizations imply endorsement by the US Government.
 Funding: This study was funded internally by Leumit Health Services. NG reports grants
 from the Aufzien Family Charity Fund and the Michael J Fox Foundation, stock holding

and consulting Vibrant, Longevity AI, GaitBetter, Consulting fee from Biogen, Bial and Pharma2B.

Author contributions:

Conceptualization: AI, EMa

5 Methodology: AI, EMe

Investigation: AI, NG

Writing – original draft: AI, NG

Writing - review & editing: AI, EMa, EMe, ER, SV, NG

Competing interests: Authors declare that they have no competing interests.

10 **Supplementary Materials:** Supplementary Information is available for this paper.

Materials & Correspondence: Correspondence and requests for materials should be addressed to AI.

		PD patients	Control	D	Odds
		N=1446	N=7230	P	Ratio
Gender	Female	655 (45.3 %)	3,275 (45.3 %)	1.000	1
	Male	791 (54.7 %)	3,955 (54.7 %)	1.000	1
Age (years)		65.7 ± 7.2	65.6 ± 7.3	0.537	
	45-49	47 (3.25 %)	243 (3.36 %)	0.873	0.97
Age category	50-59	247 (17.08 %)	1,249 (17.28 %)	0.879	0.99
1.20 0000201	60-69	591 (40.87 %)	3,081 (42.61 %)	0.232	0.93
	70-	561 (38.80 %)	2,657 (36.74 %)	0.136	1.09
Weight (kg)		77.1 ± 15.8	78.1 ± 15.7	0.026	
() orgine (ing)	missing	29 (2.01 %)	109 (1.51 %)		
Height (cm)		165 ± 12	165 ± 11	0.264	
	missing	53 (3.67 %)	182 (2.52 %)		
BMI (kg/m2)		28.3 ± 5.1	28.6 ± 5.1	0.035	
	missing	55 (3.80 %)	188 (2.60 %)		
	<18.5				
	Underweight	17 (1.22 %)	55 (0.78 %)	0.110	1.57
BMI category	18.5-24.9 Normal	330 (23.72 %)	1,618 (22.98 %)	0.554	1.04
	25-29.9				
	Overweight	581 (41.77 %)	2,951 (41.91 %)	0.929	0.99
	>=30 Obese	463 (33.29 %)	2,418 (34.34 %)	0.458	0.95

Table 1. Demographic and clinical characteristics of the study cohort at index date

	missing	55 (3.80 %)	188 (2.60 %)	0.014	1.48
BP systolic		134 ± 25	133 ± 21	0.246	
(mmHg)	missing	4 (0.28 %)	41 (0.57 %)		
BP diastolic		78.2 ± 9.6	78.0 ± 9.5	0.491	
(mmHg)	missing	5 (0.35 %)	45 (0.62 %)		
	Non-smoker	1,155 (79.88 %)	5,478 (75.77 %)	0.001	1.27
Smoking status	Past smoker	43 (2.97 %)	275 (3.80 %)	0.145	0.78
6	Smoker	132 (9.13 %)	1,051 (14.54 %)	< 0.001	0.59
	missing	116 (8.02 %)	426 (5.89 %)	0.003	1.39
Socio-economic		9.72 ± 3.40	9.73 ± 3.40	0.948	
status (1-20)	missing	74 (5.12 %)	370 (5.12 %)		
eGFR MDRD		79.4 ± 20.2	80.6 ± 24.5	0.089	
(mL/min/1.73m2)	missing	132 (9.13 %)	844 (11.67 %)		
Glucose (mg/dL)		111 ± 32	109 ± 33	0.107	
	missing	127 (8.78 %)	807 (11.16 %)		
Hemoglobin A1c		6.22 ± 1.10	6.22 ± 1.16	0.945	
(%)	missing	418 (28.9 %)	2,490 (34.4 %)		
Record of prior Tetanus-Diphtheria toxoid vaccination		22 (1.52 %)	216 (2.99 %)	0.001	0.50

Table 2. Multivariable regression model examining the impact of medication purchased in the three preceding years on relative PD disease severity

ATC code		Estimate	95% Confidence Interval	P-value
	Age	-0.00	[-0.01 to 0.00]	0.157
	Smoker	0.18	[0.05 to 0.32]	0.006
J01MA01	Ofloxacin	-0.45	[-0.55 to -0.34]	7.08.10-17
A06AD15	Macrogol	0.34	[0.23 to 0.44]	3.09.10-10
J01FA01	Erythromycin	-0.74	[-1.12 to -0.35]	0.0002
A06AC01	Ispaghula (Psyllium)	-0.42	[-0.65 to -0.19]	0.0004
J07AM51	Tetanus-Toxoid vaccine	-1.11	[-1.71 to -0.51]	0.0003
J01CE08	Benzathine	-0.69	[-1.15 to -0.24]	0.0028
	Benzylpenicillin			
J01DB05	Cefadroxil	-1.02	[-1.70 to -0.35]	0.0029
J01FF01	Clindamycin	0.40	[0.15 to 0.64]	0.0017
J01FA06	Roxithromycin	-0.13	[-0.23 to -0.04]	0.0076
J01MA12	Levofloxacin	-0.22	[-0.41 to -0.03]	0.0263





Ā

5

	PD	Control	-	Odds ratio
Vaccination timing	N=1446	N=7230	Р	[95% CI]
Vaccination 15-20 years before	8 (0.55 %)	29 (0.40 %)	0.381	1.38 [0.54 to 3.11]
Vaccination 10-15 years before	8 (0.55 %)	49 (0.68 %)	0.722	0.82 [0.33 to 1.74]
Vaccination 5-10 years before	4 (0.28 %)	75 (1.04 %)	0.003	0.26 [0.07 to 0.71]
Vaccination 0-5 years before	2 (0.14 %)	61 (0.84 %)	0.002	0.16 [0.02 to 0.62]

В

Vaccination 0-2 years before		32 (0.43 %)	0.007	0.00 [0.00 to 0.61]
Vaccination after	13 (0.90 %)	47 (0.65 %)	0.297	1.39 [0.69 to 2.61]
No recorded vaccination	1,411 (97.58 %)	6,965 (96.33 %)	0.018	1.53 [1.07 to 2.26]

	Univariable model		Multivariable model		
Vaccination	Odds ratio [95% CI]	Р	Adj. OR [95% CI]	Р	
Vaccination 15-20 years before	1.45 [0.66 to 3.18]	0.359	1.37 [0.62 to 3.02]	0.439	
Vaccination 10-15 years before	0.82 [0.39 to 1.74]	0.602	0.79 [0.37 to 1.67]	0.537	
Vaccination 5-10 years before	0.26 [0.09 to 0.71]	0.009	0.26 [0.09 to 0.71]	0.008	
Vaccination 0-5 years before	0.16 [0.04 to 0.68]	0.012	0.17 [0.04 to 0.70]	0.014	
Vaccination after	1.31 [0.70 to 2.43]	0.396	1.33 [0.71 to 2.43]	0.368	
No recorded vaccination	Ref.	Ref.	Ref.	Ref.	

Figure 2. Severity kernel density plots and regression analyses





 ${f C}$ PD Severity score over time in vaccinated vs. others







B PD Severity score over time, quadratic regression (k=2)



D PD Severity score over time in vaccinated vs. others (k=2)



 ${\bf F}_{\rm R}$ Relative PD Severity vs. time since last vaccination



(A) PD Severity score over time, kernel density plot and linear regression

- (B) PD Severity score over time, kernel density plot and quadratic regression (k=2)
- (C) PD Severity score over time, kernel density plot and linear regression in vaccinated
- 5 patients vs. others

(D) PD Severity score over time, kernel density plot and quadradic regression (k=2) in

vaccinated vs. others

(E) Relative PD Severity over time since PD diagnosis, kernel density plot and linear regression in vaccinated vs. others

10 (F) Relative PD Severity vs. time since last vaccination in vaccinated



Fig. 3. Dose-response effect associated with purchase of selected antimicrobial agents and food supplements

Dose-response effect to Roxithromycin

В

- r = -0.127 P = 0.002 Relative PD severity Relative PD severity -1 -2 -2 -3 -3 3 4 5 6 Number of Roxithromycin purchases over 3 years F G Dose-response effect to Clindamycin r = 0.152 P = 0.094 Relative PD severity Relative PD severity -1 -2 -2 -3 10 12 14 Number of Clindamycin purchases over 3 years (B) Dose-response effect to Roxithromycin
- (E) Dose-response effect to Cefadroxil
- (F) Dose-response effect to Clindamycin



- (C) Dose-response effect to Levofloxacin
- (G) Dose-response effect to Psyllium



- (D) Dose-response effect to Ofloxacin
- (H) Dose-response effect to Macrogol



B PD Severity score over time, quadratic regression (k=2)









В

С



D



H Dose-response effect to Macrogol

