1 Title: Serum metabolome indicators of early childhood development in the Brazilian National Survey

2 on Child Nutrition (ENANI-2019)

- 3 Marina Padilha, PhD¹; Victor N. Keller, PhD¹; Paula Normando, PhD¹; Raquel M. Schincaglia, PhD¹;
- 4 Nathalia C. Freitas-Costa, PhD¹; Samary S. R. Freire, MSc¹; Felipe M. Delpino, PhD¹; Inês R. R. de Castro²,
- 5 PhD; Elisa M. A. Lacerda, PhD³; Dayana R. Farias, PhD¹; Zachary Kroezen, BSc⁴; Meera
- 6 Shanmuganathan, PhD⁴; Philip Britz-Mckibbin, PhD⁴; Gilberto Kac, PhD¹
- 7
- 8 ¹ Department of Social and Applied Nutrition, Federal University of Rio de Janeiro, Josué de Castro
- 9 Nutrition Institute, Rio de Janeiro, RJ, Brazil.
- ² Department of Social Nutrition, Institute of Nutrition, State University of Rio de Janeiro, Rio de Janeiro,
- 11 Brazil
- 12 ³ Department of Nutrition and Dietetics, Federal University of Rio de Janeiro, Josué de Castro Nutrition
- 13 Institute, Rio de Janeiro, RJ, Brazil.
- ⁴ Department of Chemistry and Chemical Biology, McMaster University, Hamilton, Canada.
- 15
- 16 Corresponding Author: Gilberto Kac, Avenida Carlos Chagas Filho, 373, J2, room 29, Cidade Universitária
- 17 Rio de Janeiro RJ 21941902, phone: 55 21 39386595, e-mail: gilberto.kac@gmail.com

18 Abstract

19	Background: The role of circulating metabolites on child development is understudied. We investigated
20	associations between children's serum metabolome and early childhood development (ECD).
21	Methods: Untargeted metabolomics was performed on serum samples of 5,004 children aged 6-59 months,
22	a subset of participants from the Brazilian National Survey on Child Nutrition (ENANI-2019). ECD was
23	assessed using the Survey of Well-being of Young Children's milestones questionnaire. The graded
24	response model was used to estimate developmental age. Developmental quotient (DQ) was calculated as
25	the developmental age divided by chronological age. Partial least square regression was used to select
26	metabolites with a variable importance projection ≥ 1 .
27	Results: Twenty-eight top-ranked metabolites were included in linear regression models adjusted for
28	child's nutritional status, diet quality and age. Interaction between these metabolites and child age was
29	tested. Cresol sulfate (β = -0.07; adjusted-p < 0.001), hippuric acid (β = -0.06; adjusted-p < 0.001),
30	phenylacetylglutamine (β = -0.06; adjusted-p < 0.001), and trimethylamine- <i>N</i> -oxide (β = -0.05; adjusted-p
31	= 0.002) showed inverse associations with DQ. We observed opposite directions in the association of DQ
32	for creatinine (for children aged -1 SD: β = -0.05; p =0.01; +1 SD: β = 0.05; p =0.02) and methylhistidine
33	(-1 SD: β = - 0.04; p =0.04; +1 SD: β = 0.04; p =0.03).
34	Conclusion: Serum biomarkers, including dietary and microbial derived metabolites involved in the gut-

brain axis, may potentially be used to track children at risk for developmental delays.

36 Funding: Supported by the Brazilian Ministry of Health and Brazilian National Research Council.

37 Keywords: early childhood development, infancy, metabolites, metabolomic, child development, SWYC

38 Introduction

39 The early years of life are characterized by remarkable growth and neurodevelopment (United Nations 40 Children's Fund (UNICEF), 2017). Child development encompasses many dimensions of a child's well-41 being. It is generally described into specific streams or domains of development, including motor 42 development, speech and language progression, cognitive abilities, and socio-emotional skills (Brown et 43 al., 2020).

44 Neurogenesis starts in the intrauterine environment, continuing to shape brain morphology and plasticity 45 after birth (Doi et al., 2022). The interval from birth to eight years represents a unique and critical period 46 in which the development of a child's brain can be significantly shaped. This phenomenon encompasses 47 special sensitivity to experiences that promote cognitive, social, emotional, and physical development 48 (United Nations Children's Fund (UNICEF), 2017). The acquisition of developmental skills results from 49 an interplay between the development of the nervous system and other organ systems (Brown et al., 2020). 50 Optimal brain development requires a stimulating environment, adequate nutrients, and social interaction 51 with attentive caregivers (Britto et al., 2017).

The early childhood development (ECD) impacts long-term individual and population health outcomes, including the ability to learn, achievements in school and later life, citizenship, involvement in community activities, and overall quality of life (van den Heuvel, 2019). An estimated 250 million children under five years old in low-income and middle-income countries are at risk of not attaining their developmental potential, leading to an average deficit of 19.8% in adult annual income (Black et al., 2017). In 2015, the importance of ECD was recognized and incorporated into the "United Nations Sustainable Development Goals".

59 Studies have demonstrated that early child metabolome disturbances may be implicated in the pathogenesis
60 of non-typical neurodevelopment, including autism spectrum disorder (ASD) (Sotelo-Orozco et al., 2023;
61 Zhu et al., 2022), communication skills development (Kelly et al., 2019), and risk of impaired
62 neurocognitive development (Moreau et al., 2019).

63 Children diagnosed with neurodevelopmental delays tend to experience more favorable treatment outcomes 64 when these conditions are identified and addressed earlier (Clark et al., 2018; Li et al., 2023). Therefore, 65 biomarkers are urgently needed to predict an infant's potential risk for developmental issues while gaining 66 new insights into underlying disease mechanisms. Although child development has been a focal point of 67 research for decades, studies in low- to middle-income countries on the potential role of circulating

68 metabolites in early childhood development (ECD) remain limited. The present study aims to identify 69 associations between children's serum metabolome and ECD. Identifying the relationships between 70 metabolic phenotypes and ECD outcomes can elucidate pathways and targets for potential interventions, 71 such as serum metabolites associated with food consumption in infancy (Bruce et al., 2023).

72

73 Results

74 In total, 14,558 children under five years were evaluated and 12,598 children aged 6-59 months were 75 eligible for blood collection, of whom 8,829 (70%) had the biological material collected. Infants < 6 months 76 of age were not included due to the greater difficulty with venipuncture, increased risk of post-blood draw 77 complications (e.g., bruises), and the lack of reference values for diagnosing micronutrient deficiencies. 78 Due to the costs involved in the metabolome analysis, it was necessary to further reduce the sample size. 79 Then, samples were stratified by age groups (6 to 11, 12 to 23, and 24 to 59 months) and health conditions 80 related to iron metabolism, such as anemia and nutrient deficiencies. The selection process aimed to 81 represent diverse health statuses, including those with no conditions, with specific deficiencies, or with 82 combinations of conditions. Ultimately, through a randomized process that ensured a balanced 83 representation across these groups, a total of 5,004 children were selected for the final sample (Figure 1). 84 The mean age was 34 months, and 48.9% of the participants were between 36 and 59 months. Most children 85 were normal weight, and 25% were at risk or with excessive weight. The prevalence of MDD was 40%. 86 Most children lived in households with a monthly income \geq the Brazilian minimum wage of USD 248.7 87 (72.4%) and had mother/caregiver with at least ≥ 11 years of education (51%) (Table 1). The DQ mean 88 (95% CI) was 0.98 (0.97; 0.99). Overall, children had lower DQs if they were male [t (4970) = 7.61, p < 89 0.001], older (r = -0.31; p < 0.001), from the northern region [F (4, 4999) = 17.77, p < 0.001], had a lower 90 monthly family income (r = 0.05, p < 0.001), and a mother/caregiver with fewer years of education [F (3, 91 5000 = 30.95, p < 0.001] (Supplementary Table 1).



92

- 93 Figure 1. Flow chart of children included in the metabolome analysis of ENANI-2019 (the Brazilian
- 94 National Survey on Child Nutrition).
- 95 Note: COVID-19, coronavirus disease 2019.

96

97 Table 1. Characteristics of children 6-59 months evaluated in a subset sample of ENANI-2019 (the

98 Brazilian National Survey on Child Nutrition) (n = 5004).

Variables	Mean; Frequency (%)	95% CI	
Age (months)	34.0	33.6; 34.5	
Age groups (months)			
6 - 23	29.6	28.3; 30.8	
24 - 35	21.5	20.4; 22.7	
36 - 59	48.9	47.5; 50.3	
Sex			
Male	51.1	49.7; 52.5	
Female	48.9	47.5; 50.3	
Weight for length/height (z-score)	0.3	0.29; 0.36	
Weight for length/height categories ^{1,2}			

Underweight (z-score < -2)	1.6	1.3; 2.0
Normal $(-2 \le z \text{-score} \le 1)$	73.3	72.1; 74.5
Overweight risk $(1 < z$ -score $\leq 2)$	17.3	16.3; 18.4
Excessive weight (z-score > 2)	7.3	6.6; 8.0
Mother/caregiver education (years)		
0 to 7	25.5	24.4; 26.8
8 to 10	23.5	22.3; 24.7
≥11	51.0	49.6; 52.4
Mode of delivery		
Vaginal	53.7	52.3; 55.1
Elective c-section	19.0	18.0; 20.2
Non-elective c-section	27.3	26.1; 28.5
Monthly family income (USD) ³		
< 62.20	3.8	3.3; 4.4
62.20 to 124.40	4.5	4.0; 5.1
124.50 to 248.70	19.2	18.2; 20.3
> 248.70	72.4	71.2; 73.7
Minimum dietary diversity (MDD) ⁴		
\geq 5 food groups	59.3	39.4; 42.1
< 5 food groups	40.7	57.9; 60.6

99 Note: CI: Confidence interval; USD: United States dollar.

¹ 24 missing values; ² Reference WHO child growth standards, 2006 (World Health Organization (WHO), 2006); ³ Estimated from the Brazilian minimum wage (R\$ 998.00) and converted to the USD exchange rate (R\$ 4.013 = USD 1) in 2019 (Brazilian minimum wage = USD 248.70). ⁴ MDD: frequency of children who received ≥ 5 or < 5 out of eight food groups on the day before the interview, [food groups: (1) breast milk; (2) grains, roots and tubers; (3) beans, nuts and seeds; (4) dairy products; (5) flesh foods; (6) eggs; (7) vitamin A-rich fruits and vegetables; and (8) fruits and vegetables] (World and Health Organization, 2021) (World health Organization (WHO) & United Nations Children's Fund (UNICEF), 2021).

Supplementary Table 1. Means and confidence intervals (CI) for developmental quotient (DQ) by sociodemographic variables.

1	1	0	
•		-	

Variables	Mean	95% CI
Region		
North	0.91	0.89; 0.93
Midwest	1.00	0.98; 1.02
Southeast	1.00	0.98; 1.02
Northeast	1.01	0.99; 1.04
South	1.02	0.99; 1.04
Sex		
Male	0.95	0.93; 0.96
Female	1.02	1.01; 1.04

¹⁰⁷

6 - 21	1.14	1.12; 1.16
22 - 35	0.99	0.97; 1.01
36 - 47	0.91	0.90; 0.93
47 - 59	0.88	0.86; 0.89
Maternal/caregiver education (years)		
0 to 7	0.92	0.90; 0.94
8 to 10	0.98	0.96; 1.00
≥11	1.07	1.00; 1.03
Monthly family income (USD) ¹		
< 62.2	0.97	0.92; 1.02
62.2 - 124.4	0.92	0.88; 0.97
124.4 - 248.7	0.94	0.91; 0.96
> 248.7	1.00	0.99; 1.01

111 Note: CI: Confidence interval; USD: United States dollar.

112 ¹ Estimated from the Brazilian minimum wage (R\$ 998.00) and converted to the USD exchange rate (R\$

113 4.013 = USD 1) in 2019. 114

Age quartiles (months)

115

116 Pearson's correlation between serum metabolites and DQ indicated 26 negative and 2 positive statistically

117 significant correlations (Supplementary Figure 1). Correlations were found for phenylacetylglutamine

118 (PAG, r = -0.16; p < 0.001), cresol sulfate (CS, r = -0.15; p < 0.001), hippuric acid (HA, r = -0.14; p < -0.14; p <

119 0.001), creatinine (Crtn, r = -0.13; p < 0.001), trimethylamine-*N*-oxide (TMAO, r = -0.10; p < 0.001),

120 citrulline (Cit, r = -0.09; p < 0.001), deoxycarnitine or g-butyrobetaine (dC0, r = -0.09; p < 0.001), and

121 methylhistidine (MeHis, r = -0.07; p < 0.001).

122



123 124

125

Supplementary Figure 1. Volcano plot for Pearson correlations between serum metabolome and
 developmental quotients of children 6-59 months evaluated in ENANI-2019 (the Brazilian National
 Survey on Child Nutrition) (n = 5004).

129

134 In the PLSR, the training data suggested that 3 components best predicted the data, while the test data

showed a slightly more predictive model with 4 components (**Supplementary Figure 2**). The model with

- 136 3 components was used for parsimony and to avoid overfitting. The serum metabolites that had the highest
- 137 loads on the components were the branched-chain amino acids, including leucine (Leu), isoleucine (Ile),
- 138 and valine (Val) on component 1, the uremic toxins, CS and PAG on component 2 and betaine and amino
- acids, mainly glutamine (Gln) and asparagine (Asn) on component 3 (Supplementary Figure 3). The 3
- 140 components accounted for 39.8% of total metabolite variance (Supplementary Figure 4) and 4.3% of the

Note: The Y-axis and X-axis represent the negative logarithm (base-10) of adjusted-p-values of each
 correlation coefficient and the correlation coefficient, respectively. Dashed line is the Benjamini-Hochberg
 adjusted-p-value for 72 metabolites comparisons. Statistically significant metabolites are labeled on plots.

- 141 DQ variance (Supplementary Figure 2). Twenty-eight metabolites showed a VIP ≥ 1 (Figure 2). These
- 142 metabolites were entered into the multiple linear regression adjusted for the child's diet quality (MDD),
- 143 nutritional status (w/h z-score), and age (months).
- 144



- 145 146
- 147

Supplementary Figure 2. Residual means squared error (RMSEP) and R squared across Partial Least
 Square Regression (PLSR) models with different numbers of components.

150 151 Note: Training and test datasets were done with 80% (n = 4003) and 20% (n = 1001) of observations, 152 respectively.

153



154 155

156

157 Supplementary Figure 3. Partial Least Square Regression (PLSR) loading plots for components 1, 2, and
 158 3.

159

160 Note: Labeled points are metabolites with loading greater than 0.2 or lower than -0.2 on any of the components in the plot.

161 component 162



163

164

Supplementary Figure 4. Partial Least Square Regression (PLSR) score scatterplots for components 1,
 2, and 3.

167

Note: Percentages in parentheses indicate the total metabolite variance explained by each component. Dots
were colored according to the developmental quotient. The child development was based on the Survey of
Well-being of Young Children – the Brazilian version of the milestones questionnaire which estimates the
child's developmental age considering developmental milestones achieved. The developmental quotient
(DQ) was calculated as the child's developmental age divided by the chronological age (Sheldrick & Perrin,
2013). DQ = 1 indicates that expected age milestones have been attained; DQ <1 and > 1 suggest that the
attainment of specific age milestones occurred below or above expectations, respectively.



Figure 2. Variable importance projection (VIP) ranking scores from Partial Least Squares Regression (PLSR) analysis were performed to select the most relevant metabolites that explain the developmental quotient variability.

Note: Metabolites with higher VIP scores are the most contributory variables, whereas smaller VIP scores
provide less contribution to explain the developmental quotient variability. Only metabolites with VIP
scores higher than one are displayed.

180

185 We found inverse associations for serum concentrations of CS ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA ($\beta = -0.07$; adjusted-p < 0.001), HA (β

186 0.06; adjusted-p < 0.001), PAG (β = -0.06; adjusted-p < 0.001), and TMAO (β = -0.05; adjusted-p = 0.002)

187 with the DQ of children, even after including the interaction between child age and the metabolite in the

188 regression analysis (Table 2). Considering the interactions between serum metabolites and child age, we

189 observed associations for Crtn (β -interaction = 0.05; adjusted-p = 0.003), HA (β -interaction = 0.04;

adjusted-p = 0.041), MeHis (β -interaction = 0.04; adjusted-p = 0.018), PAG (β -interaction = 0.04; adjusted-

- 191 p = 0.018), TMAO (β -interaction = 0.05; adjusted-p = 0.003), and Val (β -interaction = 0.04; adjusted-p = 0.018)
- 192 0.039) (**Table 2**).
- 193

	Mai	in effect without interaction ter	child age m ¹	Main effect with child age interaction term ^{1, 2} Interaction ter(Metabolites: Chil		term ild age) ^{1, 2}			
Serum Metabolites	β	95% CI	adjusted-p- value	β	95% CI	adjusted- p-value	β	95% CI	adjusted- p-value
Asymmetric dimethylarginine	< 0.01	-0.03; 0.02	0.942	< 0.01	-0.03; 0.02	0.984	< 0.01	-0.02; 0.03	0.965
Alanine	-0.02	-0.04; 0.01	0.431	-0.02	-0.04; 0.01	0.414	< 0.01	-0.03; 0.02	0.928
Carnitine	< 0.01	-0.03; 0.03	0.954	< 0.01	-0.03; 0.03	0.985	< 0.01	-0.02; 0.03	0.965
Citrulline	-0.03	-0.05; 0.00	0.164	-0.03	-0.05; 0.00	0.176	0.02	0.00; 0.05	0.186
Creatinine	-0.01	-0.03; 0.02	0.935	< 0.01	-0.03; 0.03	0.985	0.05	0.02; 0.08	0.003
Cresol sulfate	-0.07	-0.10; -0.04	< 0.001	-0.06	-0.09; -0.03	< 0.001	0.03	0.01; 0.06	0.059
Deoxycarnitine	-0.03	-0.06; -0.01	0.065	-0.03	-0.06; -0.01	0.073	0.02	-0.01; 0.05	0.227
Hippuric acid	-0.06	-0.09; -0.04	< 0.001	-0.06	-0.09; -0.03	< 0.001	0.04	0.01; 0.06	0.041
Histidine	-0.02	-0.05; 0.00	0.183	-0.02	-0.05; 0.00	0.183	0.02	0.00; 0.05	0.156
Hydroxyproline	-0.01	-0.03; 0.02	0.912	-0.01	-0.03; 0.02	0.976	-0.02	-0.05; 0.01	0.227
Isoleucine	-0.03	-0.05; 0.00	0.164	-0.03	-0.05; 0.00	0.176	0.02	-0.01; 0.04	0.312
Leucine	-0.02	-0.05; 0.00	0.183	-0.02	-0.05; 0.00	0.203	0.03	0.00; 0.05	0.144
Lysine	< 0.01	-0.03; 0.02	0.935	< 0.01	-0.03; 0.02	0.984	0.02	0.00; 0.05	0.194
Methionine	-0.03	-0.05; 0.00	0.164	-0.03	-0.05; 0.00	0.176	0.02	-0.01; 0.05	0.227
Methylhistidine	< 0.01	-0.02; 0.03	0.952	< 0.01	-0.03; 0.03	0.985	0.04	0.02; 0.07	0.018
Phenylacetylglutamine	-0.06	-0.09; -0.04	< 0.001	-0.06	-0.09; -0.03	0.001	0.04	0.01; 0.07	0.018
Phenylalanine	-0.03	-0.06; 0.00	0.164	-0.03	-0.05; 0.00	0.176	0.03	0.00; 0.05	0.129
Threonine	< 0.01	-0.03; 0.02	0.953	< 0.01	-0.03; 0.02	0.985	0.01	-0.02; 0.03	0.836
TMAO	-0.05	-0.08; -0.02	0.002	-0.04	-0.07; -0.02	0.010	0.05	0.02; 0.07	0.003
Tyrosine	-0.01	-0.04; 0.02	0.816	-0.01	-0.04; 0.02	0.810	0.02	0.00; 0.05	0.194

Table 2. Association between serum metabolites and developmental quotient (DQ) of children with and without child age interaction term for children 6-59

months evaluated in a subset sample of ENANI-2019 (the Brazilian National Survey on Child Nutrition) (n = 5004).

Unknown 117.0552:1.67:N; C5H10O3	-0.02	-0.05; 0.01	0.317	-0.02	-0.05; 0.01	0.322	< 0.01	-0.03; 0.03	0.94 95
Unknown 135.0293:1.71:N; C4H8O5	0.01	-0.02; 0.04	0.721	0.01	-0.02; 0.04	0.653	-0.01	-0.04; 0.01	0.52497 197
Uric acid	-0.01	-0.03; 0.02	0.924	< 0.01	-0.03; 0.02	0.984	0.02	0.00; 0.05	0.1 598
Valine	-0.03	-0.05; 0.00	0.164	-0.02	-0.05; 0.00	0.183	0.04	0.01; 0.06	0.0399
2-Hydroxybutyric acid	-0.01	-0.04; 0.02	0.800	-0.01	-0.04; 0.02	0.653	0.03	0.00; 0.06	0.1201
3-Hydroxybutyric acid	-0.03	-0.05; 0.00	0.164	-0.03	-0.05; 0.00	0.176	< 0.01	-0.03; 0.03	0.9002
3-Methyl-2-oxovaleric acid	< 0.01	-0.03; 0.03	0.957	< 0.01	-0.03; 0.03	0.985	< 0.01	-0.02; 0.03	0.9 &9 3
Xanthine	< 0.01	-0.02; 0.03	0.954	< 0.01	-0.03; 0.03	0.985	< 0.01	-0.03; 0.03	0.965
									205

206 Note: CI: Confidence interval; DQ: Developmental quotient; TMAO: Trimethylamine N-oxide.

¹ Multiple linear regression adjusted for child's diet quality (minimum dietary diversity), child nutritional status (w/h score), and child age (in months).

208 ²Child age (in months) was used as interaction in the regression models.

209 Only metabolites with variable importance projection (VIP) scores > 1 were entered in the regressions (29 metabolites). Adjusted-p-values refer to p-values adjusted for multiple

210 comparisons by the Benjamini-Hochberg method. The child development was based on the Survey of Well-being of Young Children - the Brazilian version of the milestones

211 questionnaire, which estimates the child's developmental age considering developmental milestones achieved. The developmental quotient was calculated as the child's

developmental age divided by the chronological age (Sheldrick & Perrin, 2013). All variables were scaled before running the regression models. Bold values indicate statistically

213 significant adjusted-p-values.

214

215

- 216 Comparing children one standard deviation (SD) above the mean child age with those one standard
- 217 deviation below (49 months vs. 19 months), we observed opposite directions for the association with DQ
- 218 for serum Crtn (for children aged 1 SD: β = 0.05; p = 0.01; + 1 SD: β = 0.05; p = 0.02) and for MeHis (-
- 219 1 SD: $\beta = -0.04$; p = 0.04; + 1 SD: $\beta = 0.04$; p = 0.03) (Figure 3). For TMAO, PAG, Val, and HA, the
- 220 effect size went from a negative value for younger children to a non-significant value for older children
- 221 (Figure 3). No association were found for interactions between child sex and each metabolite on DQ (data
- not shown).
- 223



Child age (months) — + 1 SD — Mean - - 1 SD

224 225 Figure 3. Association between serum metabolites with child age interaction and developmental quotient of 226 children from 6 to 59 months from ENANI-2019 (the Brazilian National Survey on Child Nutrition, 2019) 227 (n = 5004).

228

229 Note: SD: standard deviation; TMAO: Trimethylamine N-oxide.

230 Only statistically significant interactions at an adjusted p-value for multiple comparisons < 0.05 were 231 decomposed into their simple slopes. ** refers to simple slopes with p-value < 0.001; * refers to simple 232 slopes with p-value < 0.05; "NS" refers to "not statistically significant". Regression adjustments used in 233 the models: child diet quality, child age (months), and child nutritional status (w/h score). Mean of children 234 age = 34 months and standard deviation = 15 months. The child development was based on the Survey of 235 Well-being of Young Children - the Brazilian version of milestones questionnaire, which estimates the 236 child's developmental age considering developmental milestones achieved. The developmental quotient

(DQ) was calculated as the child's developmental age divided by the chronological age (Sheldrick and
Perrin, 2013). All variables' data were auto-scaled, before running the regression models.

- 240 Mediation analysis identified that PAG was a mediator for the relationship between mode of delivery 241 (ACME = 0.003, p < 0.001), child's diet quality (ACME = 0.002, p = 0.019), and child fiber intake (ACME 242 = -0.002; p = 0.034) and DQ (Supplementary Figure 5). Serum HA (ACME = -0.004, p < 0.001) and 243 TMAO (ACME = -0.002, p = 0.022) were also mediators for the relationship between child fiber intake 244 and DQ (Supplementary Figure 5). According to the mediation analysis, having a vaginal delivery ($\beta = -$ 245 0.05; p < 0.001), not achieving MDD (β = -0.03; p = 0.019), and greater total fiber intake (β = 0.03; p = 246 0.031) increased the serum PAG concentration, that in turn was inversely associated with DQ. Moreover, a higher fiber intake was directly associated with HA ($\beta = 0.06$; p < 0.001) and TMAO ($\beta = 0.03$; p = 0.018), 247
- which also was inversely associated with DQ.
- 249



Supplementary Figure 5. Effect decomposition plots for mediation analysis of mode of delivery (A),
 child's diet quality (B), and child fiber intake (C) on child's developmental quotient considering metabolites
 as mediators.

254

255 Notes: Dots and lines represent the point estimated and 95% confidence interval based upon the quantiles 256 of bootstrapped distribution. Lines that are across the vertical dashed indicate the absence of statistical 257 significance. ACME = Average causal mediation effect; ADE = Average direct effect; Total effect = the 258 total effect. Only metabolites that showed statistically significant association with a child's developmental 259 quotient were tested in the mediation models. The linear regression models were adjusted for child 260 nutritional status (w/h z-score) and child age (in months). Data were auto-scaled before running the models. 261 The mode of delivery was evaluated as vaginal or c-section; child's diet quality as ≥ 5 or < 5 food groups 262 (minimum dietary diversity), and child fiber intake was given in grams. 263

264 Discussion

265 A limited number of investigations have examined the link between blood, urine or stool metabolites and 266 child development, with most studies focusing on comparing the metabolic profile of patients with 267 developmental disorders against healthy controls (Brydges et al., 2021; Moreau et al., 2019; Needham et 268 al., 2021; Ruggeri et al., 2014; Sotelo-Orozco et al., 2019). This is the first study to explore the association 269 between child serum metabolome and ECD on a population-based level. According to our results, serum 270 concentrations of PAG, CS, HA and TMAO were inversely associated with child DQ. The associations of 271 PAG, HA, TMAO, and Val on DQ were also age-dependent and showed stronger associations for children 272 < 34 months. In addition, inverse associations were found for serum levels of MeHis and Crtn with DQ for 273 children < 34 months, whereas direct associations were found for children > 34 months.

274 PAG is the glutamine conjugate of phenylacetic acid generated from the gut microbial dependent 275 metabolism of phenylalanine (Krishnamoorthy et al., 2023; Poesen et al., 2016). As circulating 276 concentrations of the essential amino acid, phenylalanine, were not directly associated with DQ in our 277 study, this suggests that differences in gut microbiome composition impacting PAG formation among 278 children are likely a major determinant of DQ rather than dietary protein intake. Similarly to our findings, 279 a previous study involving 76 patients with Attention-Deficit/Hyperactivity Disorder (ADHD) and 363 280 healthy children aged 1-18 years identified an inverse relationship between urinary PAG and ADHD (Tian 281 et al., 2022). While the specific pathways contributing to such disorders remain to be fully elucidated, it is 282 known that PAG is structurally similar to catecholamines and can activate adrenergic receptors (Huvnh, 283 2020). The stimuli of adrenergic receptors may have broader implications on behavioral responses, 284 potentially influencing neurological activities (Connors et al., 2005; Pliszka et al., 1996). 285 Likewise, in our study, circulating TMAO levels were inversely associated with child DQ. Elevated

286 concentrations of TMAO in plasma and cerebrospinal fluid are also implicated in age-related cognitive

287 dysfunction, neuronal senescence, and synaptic damage in the brain (Praveenraj et al., 2022). In addition, 288 its increased levels have been associated with activation of inflammatory pathways (Seldin et al., 2016) and 289 neurodegenerative diseases (Mudimela et al., 2022). Previous studies reported that TMAO can activate 290 astrocytes and microglia and trigger a cascade of inflammatory responses in the brain, induce oxidative 291 stress, superoxide production, and mitochondrial impairment, and cause inhibition of mTOR signaling in 292 the brain (Mudimela et al., 2022; Praveenraj et al., 2022). In this context, dysregulation of the mTOR 293 signaling pathway may lead to substantial abnormalities in brain development, contributing to a wide array 294 of neurological disorders, including ASD, seizure, learning impairments, and intellectual disabilities 295 (Altomare & Gitto, 2015; Lee, 2015).

296 HA is a glycine conjugate derived from exposure to benzoic acid (i.e., preservative in processed foods), or 297 generated via intestinal microbial fermentation of dietary polyphenols and phenylalanine (Assem et al., 298 2018). Similar to other co-metabolized species, circulating HA concentrations depend on dietary exposures, 299 host metabolism, and early life gut microbiota colonization and maturation in young children from in utero 300 to after birth (Jian et al., 2021). Khan et al. (Khan et al., 2022) in a case-control study involving 65 children 301 with ASD and 20 children with typical development, reported that urinary HA was significantly higher in 302 the ASD group, which corroborates with the inverse association found with DQ in our study. However, the 303 effect of HA on metabolic health is still controversial as it has been proposed as a potential dietary 304 biomarker for fruit and vegetable consumption in healthy children and adolescents (Krupp et al., 2012; 305 Pallister et al., 2017). HA also inhibits the Organic Anion Transporter (OAT) 3 function and contribute to 306 the toxic action of other compounds, including indoxyl sulfate (Ticinesi et al., 2023), which may affect 307 cognitive function by disrupting the brain barrier (Lin et al., 2019).

308 CS is a product of tyrosine fermentation in the gut involving more than 55 p-cresol producing bacteria prior 309 to hepatic sulfate conjugation (Saito et al., 2018). CS was inversely associated with DQ in our study, and it 310 has been studied in the early stages of life, particularly concerning conditions such as ASD (Guzmán-Salas 311 et al., 2022; Persico & Napolioni, 2012). Urinary p-cresol and CS have been found elevated in ASD-312 diagnosed children < 8 years (Altieri et al., 2011; Persico & Napolioni, 2012). Animal models have shown 313 that CS is a gut derived neurotoxin that can impact neuronal cell structural remodeling even at low doses 314 via oxidative stress and secretion of brain-derived neurotrophic factor (Tevzadze et al., 2022). Indeed, p-315 cresol might impact developmental processes since it is related to impaired dendritic development,

316 synaptogenesis, and synapse function in hippocampal neurons, which are crucial for cognitive and neural

development in children (Guzmán-Salas et al., 2022).

318 Prior investigations have identified PAG, CS, HA, and TMAO as products of gut microbiota metabolism 319 (Pallister et al., 2017; Reichard et al., 2022). Specifically, dietary aromatic amino acids are metabolized by 320 gut microbiota in the large intestine, converting phenylalanine into PAG and HA and tyrosine into CS 321 (Reichard et al., 2022; Ticinesi et al., 2023). In contrast, TMAO originates from trimethylamine (TMA), 322 which is produced from betaine compounds, including g-butyrobetaine (dC0), choline, and carnitine via 323 gut microbiota co-metabolism that is subsequently oxidized to TMAO in the liver (Reichard et al., 2022). 324 Once these microbiota-derived compounds enter the bloodstream, they may elicit physiological responses 325 influencing the central nervous system through direct passage across the blood-brain barrier or indirectly 326 through vagus nerve stimulation (Morais et al., 2021). Such dynamics underscore the complex interactions 327 within the microbiota-gut-brain axis (Carabotti et al., 2015; Connell et al., 2022) that can be mediated by 328 environmental exposures early in life, such as mode of birth and child's diet (Azad et al., 2018).

Overall, PAG, HA, and TMAO showed a significant average causal mediation effect with dietary fiber intake that was inversely associated with DQ. However, the interpretation of mediation effects is limited by the observational nature of the data and third variables may explain unexpected relationships between the variables in the analysis. Mediation analysis cannot determine causality or rule out spurious effects and measurement procedures, such as a single measure of dietary intake. These analyses only provide context for the main findings and point to potential future directions.

335 The age-dependent associations observed in our study are consistent with the age-related changes in 336 metabolic profile reported by previous studies (Chiu et al., 2016; Gu et al., 2009; Psihogios et al., 2008; 337 Tian et al., 2022). Increased urinary TMAO and betaine levels were found in children aged six months, 338 whereas creatine and Crtn levels increased significantly after six months (Chiu et al., 2016). Similar 339 findings were reported by Gu et al. in a study including children from newborn to 12 years of age. The 340 urinary Crtn increased with age, whereas glycine, betaine/TMAO, citrate, succinate, and acetone decreased 341 (Gu et al., 2009). These changes may reflect a physiological age-dependent process related to the rapid 342 growth occurring in early life (Chiu et al., 2016), besides the dynamic process of gut microbiota maturation 343 during the first years of life (Derrien et al., 2019).

Interestingly, we observed that the child's age changed the direction of the association between Crtn andMeHis with DQ. Crtn is generated non-enzymatically from creatine, and is related to energy production

within skeletal muscle tissue, whereas MeHis is related to protein turn-over and has been evaluated as a biomarker for the rate of skeletal muscle breakdown (Kreider & Stout, 2021; Wang et al., 2012). For example, plasma and urinary MeHis are temporally associated with changes to a health-promoting Prudent diet in contrast to a Western diet, whose concentrations are positively correlated with greater self-reported daily intake of protein (Wellington et al., 2019). We hypothesize that higher serum concentrations of Crtn and MeHis in older children (> 49 months) may be due to the greater physical activity/mobility needed through the first years (Chiu et al., 2016).

353 Our study provided valuable insights into the potential role of serum metabolome on ECD for children aged 354 6-59 months. One of the strengths of this study is the large sample size, which allows for a more 355 comprehensive representation of the population on a national level. Furthermore, our study employed a 356 quantitative targeted and exploratory untargeted metabolomics method. This high-throughput 357 metabolomics platform is strengthened by implementing rigorous quality control measures and batch-358 correction algorithms, ensuring the high accuracy and reproducibility needed for large-scale 359 epidemiological studies. We used the DQ as a variable for evaluating ECD, which consists of a continuous 360 parameter that integrates developmental milestones attained with the child's chronological age at its 361 achievement. Some limitations are worth mentioning. First, the analysis of hydrophobic/water-insoluble 362 lipids was not included in this study limiting overall metabolome coverage. Also, the inherent limitations 363 of a cross-sectional study prevent us from making causal inferences concerning the temporal relationship 364 between serum metabolic phenotypes and ECD trajectories. Moreover, birth weight and breastfeeding 365 practices were available only for a limited number of participants and were not included in the regression 366 adjustments. Concerning the child's diet assessment, we estimated dietary diversity and fiber intake based 367 on one-day food intake reports, with the MDD specifically measuring dietary diversity within diet quality. 368 In conclusion, this manuscript represents a pioneering effort in Brazil, a population-based survey targeting 369 children from 6-59 months of age that incorporated metabolome and ECD analysis. We found that serum 370 PAG, HA, CS, and TMAO were inversely associated with ECD and that age can modify the effect of PAG, 371 HA, TMAO, Crtn, and MeHis on development. These results suggest that a panel of circulating metabolites 372 might offer a preliminary warning of developmental risk and potentially be used as a screening tool to help 373 identify children at risk for developmental delays at early stages of life.

Further prospective longitudinal studies, including microbiome analysis, are warranted to validate our
findings and establish targeted intervention biomarkers besides providing further insights into possible
mechanistic pathways.

377

378 Materials and Methods

379 Study design and participants

This cross-sectional study uses data from the Brazilian National Survey on Child Nutrition (ENANI-2019). ENANI-2019 is a population-based household survey with national coverage and representativeness of children aged < 5 years that has investigated dietary intake, anthropometric status, and micronutrient deficiency. Details of the ENANI-2019 sample design, study completion, and methodology have been published previously (Alves-Santos et al., 2021; Castro et al., 2021; Vasconcellos et al., 2021). ENANI-2019 data collection took place from February 2019 and ended in March 2020 due to the COVID-19 pandemic.

387

388 Covariates

389 Trained interviewers administered a structured questionnaire to collect socio-demographic, health and 390 anthropometric data (Alves-Santos et al., 2021). The variables included in this study were: the child's age 391 (in months), sex (male or female), educational level of the mother/caregiver of the child (0–7, 8–10, and \geq 392 11 completed years of education), mode of delivery (vaginal or c-section), monthly familiar income (< 62.2, 393 62.2-24.4, 124.5-248.7, > 248.7 USD). Body weight (kg) and length or height (m) were used to calculate 394 the weight for length/height z-scores (w/h z-scores). Also, the w/h z-scores were classified considering the 395 age and sex of the child, according to World Health Organization (WHO) standards (World Health 396 Organization (WHO), 2006).

The child's diet quality was assessed using the minimum dietary diversity (MDD) indicator proposed by the WHO (World health Organization (WHO) & United Nations Children's Fund (UNICEF), 2021). MDD requires consumption of foods from at least five of eight food groups during the previous day. The eight food groups are 1) breast milk; 2) grains, roots, tubers, and plantains; 3) pulses (beans, peas, lentils), nuts, and seeds; 4) dairy products (milk, infant formula, yogurt, cheese); 5) flesh foods (meat, fish, poultry, organ meats); 6) eggs; 7) vitamin-A rich fruits and vegetables; and 8) other fruits and vegetables. The variable was dichotomized as children who had consumed ≥ 5 or < 5 food groups. Data to produce this indicator

101	was derived from the ENANU's structured questionnaire related to foods consumed the day before the first
404	was derived from the Environ's structured questionnane related to roods consumed the day before the first

405 interview (Lacerda et al., 2021) Furthermore, in ENANI-2019 caregivers fulfilled one 24-hour food recall

- 406 (R24h) reporting all children's food and beverage intake in the day before the interview. Child fiber intake
- 407 (grams) was obtained from the R24h.
- 408

409 Assessment of ECD

410 The Survey of Well-being of Young Children (SWYC) milestones questionnaire was used to assess ECD.

411 This questionnaire inquiries about motor, language, and cognitive milestones appropriate for the age range

412 of the form (Whitesell et al., 2015). It is recognized by the American Academy of Pediatrics and is a widely

- disseminated screening tool for identifying developmental delays in children aged 1 to 65 months (Lipkinet al., 2020).
- 415 The SWYC milestones questionnaire was developed and validated by Sheldrick and Perrin (Sheldrick & 416 Perrin, 2013), and a version of the SWYC (SWYC-BR) has been translated, cross-culturally adapted, and 417 validated for use in Brazilian children (Moreira et al., 2019). A recently published study evaluated the 418 internal consistency of the SWYC-BR milestones questionnaire using the ENANI-2019 data and 419 Cronbach's alpha, which showed adequate performance (0.965; 95% CI: 0.963-0.968) (Freitas-Costa et al., 420 2023). SWYC-BR comprises 12 distinct forms, each aligned with the recommended age for routine 421 pediatric wellness visits from 1 to 65 months (specifically at 1-3, 4-5, 6-8, 9-11, 12-14, 15-17, 18-22, 422 23-28, 29-34, 35-46, 47-58, and 59-65 months). Each form is a 10-item questionnaire. For each item, a 423 parent/caregiver can choose one of three answers that best describe their child ("not yet", "somewhat", or 424 "very much").
- 425 The ENANI-2019 data collection system automatically selected the appropriate set of developmental 426 milestones according to the child's age. The corrected age was used to select the proper set of developmental 427 milestones for children under two years who were born preterm (< 37 gestational weeks) (Gould et al., 428 2021).

429

430 Developmental quotient

431 The item response theory and graded response models were used to estimate development age (Samejima,
432 1997). The analysis used the full information method and incorporated the complex sample design in the
433 Mplus software version 7 (Los Angeles, EUA) (Muthén & Muthén, n.d.). The estimated model allowed the

434 construction of an item characteristic curve (ICC) for each milestone, representing the change in the 435 probability of a given response (sometimes or always) and the discrimination of each milestone 436 development by age, estimating the development age (Freitas-Costa et al., 2023). The ICC and its 437 coefficients were used to estimate developmental age according to the developmental milestones reached 438 by each child. This methodology has been previously used to assess ECD with the SWYC (Freitas-Costa 439 et al., 2023; Sheldrick et al., 2019; Sheldrick & Perrin, 2013) and the Denver Test (Drachler et al., 2007). 440 The developmental quotient (DQ) was calculated by dividing the developmental age by the chronological 441 age (Freitas-Costa et al., 2023; Sheldrick & Perrin, 2013). DQ equals to 1 indicates that the expected age 442 milestones are attained. DQ values < 1 or > 1 suggest attaining age milestones below or above expectations, 443 respectively. This method allows analyzing the outcome as a continuous variable.

444

445 Blood collection

446 Details of the procedures adopted for blood collection and laboratory analyses have been previously 447 described (Castro et al., 2021). Fasting was not required, and changes in medication were not necessary to 448 draw the blood sample. Briefly, 8 mL of blood sample were drawn and distributed in a trace tube (6 mL) 449 and EDTA tube (2 mL) and transported in a cooler with a controlled temperature (from 2 °C to 8 °C) to a 450 partner laboratory. Aliquots from the trace tube were centrifugated and the serum was transferred to a 451 second trace tube and stored at freezing temperature (-20 °C) until laboratory analyses were performed. 452 Serum samples with sufficient volume were stored in a biorepository (-80 °C) prior to untargeted 453 metabolome analysis.

454

455 Serum processing and metabolome analysis

456 Untargeted metabolomic analysis was performed on serum samples using a high-throughput platform based 457 on multisegment injection-capillary electrophoresis-mass spectrometry (MSI-CE-MS). Samples were first 458 thawed slowly on ice, where 50 μ L were aliquoted and then diluted four-fold to a final volume of 200 μ L 459 in deionized water with an internal standard mix containing 40 µM 3-chlorotyrosine, 3-fluorophenylalanine, 460 2-fluorotyrosine, trimethylamine-N-oxide[D9], g-amino butyrate[D6], choline[D9], creatinine[D3], 461 ornithine [15N2], histidine [15Na], carnitine [D3], 3-methylhistidine [D3] and 2 mM glucose [13C6]. Diluted 462 serum samples were then transferred to pre-rinsed Nanosep ultracentrifuge devices with a molecular weight 463 cutoff of 3 kDa (Cytiva Life Sciences, Malborough, USA), and centrifuged at 10,000 g for 15 min to remove

464 proteins. Following ultrafiltration, 20 μ L of diluted serum filtrate samples were transferred to CE-465 compatible polypropylene vials and analyzed using MSI-CE-MS. A pooled QC was also prepared to 466 evaluate technical precision throughput the study using 50 μ L aliquots collected from the first batch of 979 467 serum samples processed. Overall, serum specimens were prepared and run as three separate batches of 468 979, 1990, and 2035 samples over an eighteen-month period. A QC-based batch correction algorithm was 469 applied to reduce long-term system drift and improve reproducibility with QC samples analyzed in a 470 randomized position within each analytical run (Wehrens et al., 2016).

471 MSI-CE-MS was performed using an Agilent 6230B time-of-flight mass spectrometer (Agilent, Santa 472 Clara, USA) with an electrospray ion source coupled to an Agilent G7100A capillary electrophoresis (CE) 473 instrument (Agilent, Santa Clara, USA). The metabolome coverage comprises primarily 474 cationic/zwitterionic and anionic polar metabolites when using full-scan data acquisition under positive and 475 negative ionization modes. Given the isocratic separation conditions with steady-state ionization via a 476 sheath liquid interface, MSI-CE-MS increases sample throughput using a serial injection format where 12 477 samples and a pooled QC are analyzed within a single analytical run. Instrumental and data preprocessing 478 parameters have been previously described (Saoi et al., 2019; Shanmuganathan et al., 2021).

479 The technical precision for serum metabolites measured in pooled QC samples had a median CV of 10.5% 480 with a range from 2.7 to 31% (n=422), which were analyzed in every run throughout data acquisition by 481 MSI-CE-MS following batch correction. Overall, seventy-five circulating polar metabolites were measured 482 in most samples (frequency > 50%) with adequate technical precision (CV < 30%) with the exception of 483 symmetric dimethylarginine that was removed. Most metabolites were identified by spiking (i.e., co-484 migration with low mass error < 5 ppm) and quantified with authentic standards, except for 13 unknown 485 metabolites that were annotated based on their accurate mass (m/z), relative migration time (RMT), 486 ionization mode (N or P), and most likely molecular formula. The metabolite distributions were severely 487 asymmetric (average skewness = 40) and leptokurtic (average kurtosis = 1810). Therefore, a log 488 transformation was performed on each metabolite, which reduced average skewness to 2.4 and kurtosis to 489 20.8. Metabolite z-scores > 5 or < -5 were considered outliers and were removed (0.12% of the data).

490 Missing data were treated following the procedures recommended by Wei et al. (Wei et al., 2018) with one 491 modification. Instead of using the "80% rule" of excluding metabolites with < 80% non-missing cases (> 492 20% missing cases) in all dependent variable categories, a less stringent 50% rule was applied to reduce the 493 risk of excluding relevant serum metabolites. For the sole purpose of performing the exclusions, the DQ

494 was recoded as a categorical variable ($DQ \ge 1$ as "within or above expectations", and DQ < 1 as "below 495 expectations") to avoid removing metabolites that had a missingness pattern associated with DQ. Cysteine-496 S-sulfate and an unknown anionic metabolite (209.030:3.04:N; C₆H₁₀O₈) had > 50% missing cases in both 497 DQ categories and were thus excluded.

498 Of the remaining seventy-two serum metabolites that satisfied the above selection criteria, 12.5% of the 499 data were missing due to matrix interferences, and 1.5% were missing due to non-detection (i.e., below 500 method detection limit). Missing data due to matrix interference were imputed using the random forest (RF) 501 method, and non-detection missing data were imputed using quantile regression imputation of left-censored 502 data (QRILC) (Wei et al., 2018). The RF method used all serum metabolome data to predict what value 503 the missing cases would likely have taken.

504

505 Statistical analysis

We carried out descriptive and inferential analyses. The descriptive analyses were based on frequency with
a 95% confidence interval (95% CI) and *Student* t-test or ANOVA were used to compare DQ in groups.

508 The Pearson correlation was first used to explore the correlations between circulating metabolites 509 (exposure) and DQ (outcome). We employed partial least squares regression (PLSR) analysis to compare 510 the metabolites within a single model (Worley & Powers, 2013). The PLSR reduces the metabolites to 511 orthogonal components, which are maximally predictive of the outcome and generate an indicator of how 512 much each metabolite contributes to predicting the outcome, called the variable importance projection 513 (VIP). Because our goal was not to determine the components that are maximally predictive of DQ but to 514 rank the metabolites on their contribution to predicting the outcome, we only used the VIPs from this 515 analysis.

The PLSR was trained on 80% of the data, and the remaining 20% were used as test data. Training and test data were randomly allocated. The model with the optimal number of components considering predictive value and parsimony was used to generate VIP values. Metabolites were selected if they had a VIP \geq 1.

519 The subsequent step was to disentangle the selected metabolites from confounding variables. A Directed

520 Acyclic Graph (DAG) was used to identify potential confounding variables of the association between

521 metabolites and DQ (Breitling et al., 2021). The DAG was produced considering variables related to the

522 exposure (metabolome) and outcome (DQ) based on evidence reported by systematic reviews or meta-

523 analysis (Figure 4). Birth weight, breastfeeding, child's diet quality, the child's nutritional status, and the

- 524 child's age were the minimal adjustments suggested by the DAG. Birth weight was a variable with high
- 525 missing data, and indicators of breastfeeding practice data (referring to exclusive breastfeeding until 6
- 526 months and/or complemented until 2 years) were collected only for children aged 0–23 months. Therefore,
- 527 those confounders were not included as adjustments. Child's diet quality was evaluated as MDD, the child's
- 528 nutritional status as w/h z-score, and the child's age in months.
- 529



530

Figure 4. Directed acyclic graph (DAG) of the association between child metabolome and early childhood
 development.

533 Note: IUGR: intrauterine growth restriction.

Breastfeeding refers to breastfeeding practice exclusively until 6 months and/or complemented until 2
years. Minimum adjustments suggested by the DAG: Birth weight, breastfeeding, child age, child diet
quality, child nutritional status (w/h z-score). Blue rectangles: ancestor of the outcome; pink rectangles:
ancestor of exposure and outcome; white circles: non observed variables in the ENANI-2019.

- 538
- 539 Multiple linear regression between each metabolite and DQ were performed and adjusted for the

540 confounders. Since the child diet and metabolism may change by child's age and that neurodevelopmental

- 541 disorders occur more frequently in boys than in girls, interactions between serum metabolites and child age
- (in months) and between metabolites and child sex were also tested to evaluate a possible modification
- 543 effect of these variables in the models.
- 544 We employed mediation analyses to explore the potential role of serum metabolites as mediators in the
- relationship between certain exposure variables related to the microbiome establishment in early life, such

546 as mode of delivery (Reyman et al., 2019), child's diet quality (Baldeon et al., 2023), as well as child fiber 547 intake (Cronin et al., 2021) and DQ. We adopted the approach proposed by Tingley et al. (Tingley et al., 548 2014) which provides independent estimates for the average causal mediation effect (ACME - the effect of 549 the exposure variable on DQ that is mediated by the metabolite), the average direct effect (ADE - controlling 550 for metabolite concentrations) of the exposure variable on DQ, and the total effect of the exposure variable 551 on DQ (mediation plus direct effect). Bootstrap tests using 5,000 iterations evaluated whether the effects 552 were statistically significant. Due to the exploratory nature of the mediation analysis, significance was not 553 corrected for multiple testing. The child's age (in months) and w/h z-score were entered as covariates. 554 Results were considered statistically significant at an adjusted- $p \le 0.05$ after the Benjamini-Hochberg 555 correction for multiple comparisons. Statistical analyses were carried out using the R programming 556 language (R Core Team; http://www.R-project.org), through JupyterLab, using the following packages: 557 ggplot2 (http://ggplot2.org), interactions (https://cran.r-project.org), dplyr (https://cran.r-project.org), 558 tidyverse 2007), (www.tidyverse.org), pls (Mevik & Wehrens, plsVarSel 559 (https://github.com/khliland/plsVarSel), mediation (Tingley et al., 2014). 560

561 Ethical aspects

The ENANI-2019 was approved by the Research Ethics Committee of the Clementino Fraga Filho University Hospital of the Federal University of Rio de Janeiro (UFRJ) under the number CAAE 89798718.7.0000.5257. Data were collected after a parent/caregiver of the child authorized participation in the study through an informed consent form and following the principles of the Declaration of Helsinki.

566

567 Contributors

568	MP contributed to conceptualization, data curation, formal analysis, investigation, methodology, and
569	writing - original draft. VNK contributed to data curation, formal analysis, investigation, methodology,
570	software, and writing - original draft. PN contributed to conceptualization, investigation, methodology,
571	project administration and writing - original draft. RMS contributed to data curation, formal analysis,
572	investigation, methodology, validation, and writing - original draft NCFC contributed to data curation,
573	methodology, and writing - original draft. SSRF contributed to conceptualization, data curation, and
574	writing- review & editing. FMD contributed to formal analysis, validation, and writing - original draft.
575	IRRC, EMAL, DRF contributed to conceptualization, investigation, methodology, validation, visualization,
576	and writing- review & editing. ZK, MS, and PBM contributed to data curation, formal analysis,
577	investigation, methodology, visualization, and writing- review & editing. GK contributed to
578	conceptualization, funding acquisition, investigation, methodology, project administration, resources,
579	supervision, validation, visualization, writing - original draft, writing- review & editing. All authors read
580	and approved the final manuscript.

581

582 Competing interests

583 All authors declare that they have no competing interests.

584

585 Acknowledgements

- 586 We thank the participating families who made this study possible and the Brazilian National Survey on
- 587 Child Nutrition (ENANI-2019) team collaborators for their support in the fieldwork and organization of the
- 588 database. We also thank the Brazilian Ministry of Health and Brazilian National Research Council (CNPq;

589 process n. 440890/2017-9).

590

591 Data Sharing Statement

- 592 Data described in the manuscript, code book, and analytic code will be available upon publication at
- 593 https://dataverse.nutricao.ufrj.br/privateurl.xhtml?token=e44dcdad-6ae3-4d9d-848e-898cbd6eb411.

15

594 References

- 595 Altieri, L., Neri, C., Sacco, R., Curatolo, P., Benvenuto, A., Muratori, F., Santocchi, E., Bravaccio, C., 596 Lenti, C., Saccani, M., Rigardetto, R., Gandione, M., Urbani, A., & Persico, A. M. (2011). Urinary 597 p-cresol is elevated in small children with severe autism spectrum disorder. *Biomarkers*, 16(3), 598 252-260. https://doi.org/10.3109/1354750X.2010.548010 599 Altomare, D., & Gitto, S. (2015). Recent insights into the pathophysiology of mTOR pathway 600 dysregulation. Research and Reports in Biology, 6, 1-16. https://doi.org/10.2147/rrb.s57088 601 Alves-Santos, N. H., de Castro, I. R. R., dos Anjos, L. A., de Aquino Lacerda, E. M., Normando, P., de 602 Freitas, M. B., Farias, D. R., Boccolini, C. S., De Vasconcellos, M. T. L., Do Nascimento Silva, P. 603 L., & Kac, G. (2021). General methodological aspects in the brazilian national survey on child 604 nutrition (enani-2019): A population-based household survey. Cadernos de Saude Publica, 37(8). 605 https://doi.org/10.1590/0102-311X00300020 606 Assem, M., Lando, M., Grissi, M., Kamel, S., Massy, Z. A., Chillon, J. M., & Hénaut, L. (2018). The impact of uremic toxins on cerebrovascular and cognitive disorders. Toxins, 10, 1-25. 607 608 https://doi.org/10.3390/toxins10070303 609 Azad, M. B., Robertson, B., Atakora, F., Becker, A. B., Subbarao, P., Moraes, T. J., Mandhane, P. J., 610 Turvey, S. E., Lefebvre, D. L., Sears, M. R., & Bode, L. (2018). Human milk oligosaccharide 611 concentrations are associated with multiple fixed and modifiable maternal characteristics, 612 environmental factors, and feeding practices. Journal of Nutrition, 148, 1733-1742. 613 https://doi.org/10.1093/jn/nxy175 614 Baldeon, A. D., McDonald, D., Gonzalez, A., Knight, R., & Holscher, H. D. (2023). Diet quality and the 615 fecal microbiota in adults in the American Gut Project. Journal of Nutrition, 153, 2004–2015. 616 https://doi.org/10.1016/j.tjnut.2023.02.018 617 Black, M. M., Walker, S. P., Fernald, L. C. H., Andersen, C. T., DiGirolamo, A. M., Lu, C., McCoy, D. 618 C., Fink, G., Shawar, Y. R., Shiffman, J., Devercelli, A. E., Wodon, Q. T., Vargas-Barón, E., & 619 Grantham-McGregor, S. (2017). Advancing early childhood development: from science through the 620 life course. The Lancet, 389(10064), 77-90. https://doi.org/10.1016/S0140-6736(16)31389-621 7.Advancing 622 Breitling, L. P., Duan, C., Dragomir, A. D., & Luta, G. (2021). Using dagR to identify minimal sufficient 623 adjustment sets and to simulate data based on directed acyclic graphs. International Journal of 624 Epidemiology, 50(6), 1772-1777. https://doi.org/10.1093/ije/dyab167 625 Britto, P. R., Lye, S. J., Proulx, K., Yousafzai, A. K., Matthews, S. G., Vaivada, T., Perez-Escamilla, R., 626 Rao, N., Ip, P., Fernald, L. C. H., MacMillan, H., Hanson, M., Wachs, T. D., Yao, H., Yoshikawa, 627 H., Cerezo, A., Leckman, J. F., & Bhutta, Z. A. (2017). Nurturing care: promoting early childhood 628 development. The Lancet, 389, 91-102. https://doi.org/10.1016/S0140-6736(16)31390-3 629 Brown, K. A., Parikh, S., & Patel, D. R. (2020). Understanding basic concepts of developmental 630 diagnosis in children. Translational Pediatrics, 9(Suppl 1), S9–S22. 631 https://doi.org/10.21037/TP.2019.11.04 632 Bruce, C. Y., Shanmuganathan, M., Azab, S. M., Simons, E., Mandhane, P., Turvey, S. E., Subbarao, P., 633 Azad, M. B., Britz-McKibbin, P., Anand, S. S., de Souza, R. J., & Stearns, J. C. (2023). The 634 relationship between diet, gut microbiota, and serum metabolome of South Asian infants at 1 year. 635 The Journal of Nutrition, 153, 470-482. https://doi.org/10.1016/j.tjnut.2022.12.016 636 Brydges, C. R., Fiehn, O., Mayberg, H. S., Schreiber, H., Dehkordi, S. M., Bhattacharyya, S., Cha, J., 637 Choi, K. S., Craighead, W. E., Krishnan, R. R., Rush, A. J., Dunlop, B. W., Kaddurah-Daouk, R., 638 Penninx, B., Binder, E., Kastenmüller, G., Arnold, M., Nevado-Helgado, A., Blach, C., ... Rinaldo, 639 P. (2021). Indoxyl sulfate, a gut microbiome-derived uremic toxin, is associated with psychic 640 anxiety and its functional magnetic resonance imaging-based neurologic signature. Scientific
- 641 *Reports*, *11*, 21011. https://doi.org/10.1038/s41598-021-99845-1

642 643	Carabotti, M., Scirocco, A., Maselli, M. A., & Severi, C. (2015). The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems. <i>Annals of Gastroenterology</i> , 28, 203–209.
644	Castro, I. R. R., Normando, P., Alves-Santos, N. H., Bezerra, F. F., Citelli, M., Pedrosa, L. de F. C.,
645	Jordão, A. A., Lira, P. I. C., Kurscheidt, F. A., Silva, P. R. P., Salvatte, K., Lacerda, E. M. A.,
646	Anjos, L. A., Boccolini, C. S., & Kac, G. (2021). Methodological aspects of the micronutrient
647	assessment in the Brazilian National Survey on Child Nutrition (ENANI-2019): A populationbased
648	household survey. <i>Cadernos de Saude Publica</i> , <i>37</i> (8), e00301120. https://doi.org/10.1590/0102-
649	311X00301120
650	Chiu, C. Y., Yeh, K. W., Lin, G., Chiang, M. H., Yang, S. C., Chao, W. J., Yao, T. C., Tsai, M. H., Hua,
651	M. C., Liao, S. L., Lai, S. H., Cheng, M. L., & Huang, J. L. (2016). Metabolomics reveals dynamic
652	metabolic changes associated with age in early childhood. <i>PLoS ONE</i> , 11(2), e0149823.
653	https://doi.org/10.1371/journal.pone.0149823
654	Clark, M. L. E., Vinen, Z., Barbaro, J., & Dissanayake, C. (2018). School Age Outcomes of Children
655	Diagnosed Early and Later with Autism Spectrum Disorder. <i>Journal of Autism and Developmental</i>
656	<i>Disorders</i> , 48(1), 92–102. https://doi.org/10.1007/s10803-017-3279-x
657	Connell, E., Le Gall, G., Pontifex, M. G., Sami, S., Cryan, J. F., Clarke, G., Müller, M., & Vauzour, D.
658	(2022). Microbial-derived metabolites as a risk factor of age-related cognitive decline and
659	dementia. <i>Molecular Neurodegeneration</i> , 17, 43. https://doi.org/10.1186/s13024-022-00548-6
660	Connors, S. L., Crowell, D. E., Eberhart, C. G., Copeland, J., Newschaffer, C. J., Spence, S. J., &
661	Zimmerman, A. W. (2005). β2-Adrenergic receptor activation and genetic polymorphisms in
662	autism: data from dizygotic twins. <i>Journal of Child Neurology</i> , 20, 876–884.
663	https://doi.org/10.1177/08830738050200110401
664	Cronin, P., Joyce, S. A., O'toole, P. W., & O'connor, E. M. (2021). Dietary fibre modulates the gut
665	microbiota. <i>Nutrients</i> , 13, 1655. https://doi.org/10.3390/nu13051655
666	Derrien, M., Alvarez, A. S., & de Vos, W. M. (2019). The gut microbiota in the first decade of life.
667	<i>Trends in Microbiology</i> , 27(12), 997–1010. https://doi.org/10.1016/j.tim.2019.08.001
668	Doi, M., Usui, N., & Shimada, S. (2022). Prenatal Environment and Neurodevelopmental Disorders.
669	Frontiers in Endocrinology, 13, 1–6. https://doi.org/10.3389/fendo.2022.860110
670	Drachler, M. D. L., Marshall, T., & Leite, J. C. C. (2007). A continuous-scale measure of child
671	development for population-based epidemiological surveys: A preliminary study using Item
672	Response Theory for the Denver Test. <i>Paediatric and Perinatal Epidemiology</i> , 21, 138–153.
673	https://doi.org/10.1111/j.1365-3016.2007.00787.x
674 675 676 677 678	Freitas-Costa, N. C., Andrade, P. G., Normando, P., Nunes, K. S. S., Raymundo, C. E., Castro, I. R. R. de, Lacerda, E. M. de A., Farias, D. R., & Kac, G. (2023). Association of development quotient with nutritional status of vitamins B6, B12, and folate in 6–59-month-old children: Results from the Brazilian National Survey on Child Nutrition (ENANI-2019). <i>American Journal of Clinical Nutrition</i> , 118(1), 162–173. https://doi.org/10.1016/j.ajcnut.2023.04.026
679	Gould, J. F., Fuss, B. G., Roberts, R. M., Collins, C. T., & Makrides, M. (2021). Consequences of using
680	chronological age versus corrected age when testing cognitive and motor development in infancy
681	and intelligence quotient at school age for children born preterm. <i>PLoS ONE</i> , 16(9), e0256824.
682	https://doi.org/10.1371/journal.pone.0256824
683 684 685	 Gu, H., Pan, Z., Xi, B., Hainline, B. E., Shanaiah, N., Asiago, V., Gowda, G. A. N., & Raftery, D. (2009). 1H NMR metabolomics study of age profiling in children. <i>NMR in Biomedicine</i>, 22, 826–833. https://doi.org/10.1002/nbm.1395
686	Guzmán-Salas, S., Weber, A., Malci, A., Lin, X., Herrera-Molina, R., Cerpa, W., Dorador, C., Signorelli,
687	J., & Zamorano, P. (2022). The metabolite p-cresol impairs dendritic development, synaptogenesis,
688	and synapse function in hippocampal neurons: Implications for autism spectrum disorder. <i>Journal</i>
689	of Neurochemistry, 161, 335–349. https://doi.org/10.1111/jnc.15604

690	Huynh, K. (2020). Novel gut microbiota-derived metabolite promotes platelet thrombosis via adrenergic
691	receptor signalling. <i>Nature Reviews Cardiology</i> , 17, 265. https://doi.org/10.1038/s41569-020-0367-
692	y
693	Jian, C., Carpén, N., Helve, O., de Vos, W. M., Korpela, K., & Salonen, A. (2021). Early-life gut
694	microbiota and its connection to metabolic health in children: Perspective on ecological drivers and
695	need for quantitative approach. <i>EBioMedicine</i> , 69, 103475.
696	https://doi.org/10.1016/j.ebiom.2021.103475
697 698 699 700	 Kelly, R. S., Boulin, A., Laranjo, N., Lee-Sarwar, K., Chu, S. H., Yadama, A. P., Carey, V., Litonjua, A. A., Lasky-Su, J., & Weiss, S. T. (2019). Metabolomics and communication skills development in children: evidence from the Ages and Stages Questionnaire. <i>Metabolites</i>, 9(1–17). https://doi.org/10.3390/metabo9030042
701 702 703 704	Khan, Z. U. N., Chand, P., Majid, H., Ahmed, S., Khan, A. H., Jamil, A., Ejaz, S., Wasim, A., Khan, K. A., & Jafri, L. (2022). Urinary metabolomics using gas chromatography-mass spectrometry: potential biomarkers for autism spectrum disorder. <i>BMC Neurology</i> , 22(101), 1–8. https://doi.org/10.1186/s12883-022-02630-4
705	Kreider, R. B., & Stout, J. R. (2021). Creatine in health and disease. <i>Nutrients</i> , <i>13</i> , 447.
706	https://doi.org/10.3390/nu13020447
707 708 709 710	 Krishnamoorthy, N. K., Kalyan, M., Hediyal, T. A., Anand, N., Kendaganna, P. H., Pendyala, G., Yelamanchili, S. V., Yang, J., Chidambaram, S. B., Sakharkar, M. K., & Mahalakshmi, A. M. (2023). Role of the gut bacteria-derived metabolite phenylacetylglutamine in health and diseases. <i>ACS Omega</i>, <i>9</i>, 3164–3172. https://doi.org/10.1021/acsomega.3c08184
711	Krupp, D., Doberstein, N., Shi, L., & Remer, T. (2012). Hippuric acid in 24-hour urine collections is a
712	potential biomarker for fruit and vegetable consumption in healthy children and adolescents. <i>The</i>
713	<i>Journal of Nutrition</i> , 142(7), 1314–1320. https://doi.org/10.3945/jn.112.159319
714	Lacerda, E. M. A., Boccolini, C. S., Alves-Santos, N. H., Castro, I. R. R., Anjos, L. A., Crispim, S. P.,
715	Costa, T. H. M., Ferreira, M. M. M., Farias, D. R., Carneiro, L. B. V., Berti, T. L., & Kac, G.
716	(2021). Methodological aspects of the assessment of dietary intake in the Brazilian National Survey
717	on Child Nutrition (ENANI-2019): a population-based household survey. <i>Cadernos de Saude</i>
718	<i>Publica</i> , 37(8), e00301420. https://doi.org/10.1590/0102-311X00301420
719 720	Lee, D. Y. (2015). Roles of mTOR signaling in brain development. <i>Experimental Neurobiology</i> , 24(3), 177–185. https://doi.org/10.5607/en.2015.24.3.177
721 722 723 724	Li, K., Bertrand, K., Naviaux, J. C., Monk, J. M., Wells, A., Wang, L., Lingampelly, S. S., Naviaux, R. K., & Chambers, C. (2023). Metabolomic and exposomic biomarkers of risk of future neurodevelopmental delay in human milk. <i>Pediatric Research</i> , 93(6), 1710–1720. https://doi.org/10.1038/s41390-022-02283-6
725	Lin, Y. T., Wu, P. H., Liang, S. S., Mubanga, M., Yang, Y. H., Hsu, Y. L., Kuo, M. C., Hwang, S. J., &
726	Kuo, P. L. (2019). Protein-bound uremic toxins are associated with cognitive function among
727	patients undergoing maintenance hemodialysis. <i>Scientific Reports</i> , <i>9</i> , 20388.
728	https://doi.org/10.1038/s41598-019-57004-7
729 730 731 732 733 734	 Lipkin, P. H., Macias, M. M., Hyman, S. L., Levy, S. E., Spooner, S. A., Rodgers, B., Kuo, D. Z., Apkon, S., Davidson, L. F., Ellerbeck, K. A., Foster, J. E. A., Hyman, S. L., Noritz, G. H., O'Connor Leppert, M., Saunders, B. S., Stille, C., Yin, L., Brei, T., Davis, B. E., High, P. C. (2020). Promoting optimal development: Identifying infants and young children with developmental disorders through developmental surveillance and screening. <i>Pediatrics</i>, <i>145</i>(1), e20193449. https://doi.org/10.1542/PEDS.2019-3449
735	Mevik, B. H., & Wehrens, R. (2007). The pls package: principal component and partial least squares
736	regression in R. <i>Journal of Statistical Software</i> , 18(2), 1–24.
737	https://doi.org/https://doi.org/10.18637/jss.v018.i02.

18

738	Morais, L. H., Schreiber, H. L., & Mazmanian, S. K. (2021). The gut microbiota–brain axis in behaviour
739	and brain disorders. <i>Nature Reviews Microbiology</i> , 19, 241–255. https://doi.org/10.1038/s41579-
740	020-00460-0
741	Moreau, G. B., Ramakrishnan, G., Cook, H. L., Fox, T. E., Nayak, U., Ma, J. Z., Colgate, E. R.,
742	Kirkpatrick, B. D., Haque, R., & Petri, W. A. (2019). Childhood growth and neurocognition are
743	associated with distinct sets of metabolites. <i>EBioMedicine</i> , 44, 597–606.
744	https://doi.org/10.1016/j.ebiom.2019.05.043
745	Moreira, R. S., Magalhães, L. D. C., Siqueira, C. M., & Alves, C. R. L. (2019). Adaptação transcultural
746	do instrumento de vigilância do desenvolvimento infantil "Survey of Wellbeing of Young Children
747	(SWYC)" no contexto brasileiro. <i>Journal of Human Growth and Development</i> , 29(1), 28–38.
748	https://doi.org/10.7322/jhgd.145001
749 750 751 752	Mudimela, S., Vishwanath, N. K., Pillai, A., Morales, R., Marrelli, S. P., Barichello, T., & Giridharan, V. V. (2022). Clinical significance and potential role of Trimethylamine N-oxide in neurological and neuropsychiatric disorders. <i>Drug Discovery Today</i> , 27(11), 1–17. https://doi.org/10.1016/j.drudis.2022.08.002
753	Muthén, L. K., & Muthén, B. O. (n.d.). Mplus User's Guide (8th ed., p. 944). https://doi.org/DOI
754 755 756 757	Needham, B. D., Adame, M. D., Serena, G., Destanie, R. R., Preston, G. M., Conrad, M. C., Campbell, A. S., Donabedian, D. H., Fasano, A., Ashwood, P., & Mazmanian, S. (2021). Plasma and fecal metabolite profiles in Autism Spectrum Disorder. <i>Biological Psychiatry</i> , 89(5), 451–462. https://doi.org/10.1016/j.biopsych.2020.09.025
758	Pallister, T., Jackson, M. A., Martin, T. C., Zierer, J., Jennings, A., Mohney, R. P., MacGregor, A.,
759	Steves, C. J., Cassidy, A., Spector, T. D., & Menni, C. (2017). Hippurate as a metabolomic marker
760	of gut microbiome diversity: modulation by diet and relationship to metabolic syndrome. <i>Scientific</i>
761	<i>Reports</i> , 7, 13670. https://doi.org/10.1038/s41598-017-13722-4
762 763	Persico, A. M., & Napolioni, V. (2012). Urinary p-cresol in autism spectrum disorder. <i>Neurotoxicology and Teratology</i> , 36, 82–90. https://doi.org/10.1016/j.ntt.2012.09.002
764	Pliszka, S. R., McCracken, J. T., & Maas, J. W. (1996). Catecholamines in attention-deficit hyperactivity
765	disorder: current perspectives. <i>Journal of the American Academy of Child and Adolescent</i>
766	<i>Psychiatry</i> , 35(3), 264–272. https://doi.org/10.1097/00004583-199603000-00006
767	Poesen, R., Claes, K., Evenepoel, P., De Loor, H., Augustijns, P., Kuypers, D., & Meijers, B. (2016).
768	Microbiota-derived phenylacetylglutamine associates with overall mortality and cardiovascular
769	disease in patients with CKD. <i>Journal of the American Society of Nephrology</i> , 27(11), 3479–3487.
770	https://doi.org/10.1681/ASN.2015121302
771 772 773 774 775	Praveenraj, S. S., Sonali, S., Anand, N., Tousif, H. A., Vichitra, C., Kalyan, M., Kanna, P. V., Chandana, K. A., Shasthara, P., Mahalakshmi, A. M., Yang, J., Pandi-Perumal, S. R., Sakharkar, M. K., & Chidambaram, S. B. (2022). The role of a gut microbial-derived metabolite, Trimethylamine N-Oxide (TMAO), in neurological disorders. <i>Molecular Neurobiology</i> , <i>59</i> , 6684–6700. https://doi.org/10.1007/s12035-022-02990-5
776	Psihogios, N. G., Gazi, I. F., Elisaf, M. S., Konstantin, I. S., & Bairaktari, E. T. (2008). Gender-related
777	and age-related urinalysis of healthy subjects by NMR-based metabonomics. <i>NMR in Biomedicine</i> ,
778	21, 195–207. https://doi.org/10.1002/nbm.1176
779	Reichard, C. A., Naelitz, B. D., Wang, Z., Jia, X., Li, J., Stampfer, M. J., Klein, E. A., Hazen, S. L., &
780	Sharifi, N. (2022). Gut microbiome-dependent metabolic pathways and risk of lethal prostate
781	cancer: prospective analysis of a PLCO Cancer Screening Trial Cohort. <i>Cancer Epidemiology</i>
782	<i>Biomarkers & Prevention</i> , 31, 192–199. https://doi.org/10.1158/1055-9965.EPI-21-0766
783 784	Reyman, M., van Houten, M. A., van Baarle, D., Bosch, A. A. T. M., Man, W. H., Chu, M. L. J. N., Arp, K., Watson, R. L., Sanders, E. A. M., Fuentes, S., & Bogaert, D. (2019). Impact of delivery mode-

785 786	associated gut microbiota dynamics on health in the first year of life. <i>Nature Communications</i> , 10(4997), 1–12. https://doi.org/10.1038/s41467-019-13014-7
787 788 789	Ruggeri, B., Sarkans, U., Schumann, G., & Persico, A. M. (2014). Biomarkers in autism spectrum disorder: the old and the new. <i>Psychopharmacology</i> , 231, 1201–1216. https://doi.org/10.1007/s00213-013-3290-7
790 791 792	Saito, Y., Sato, T., Nomoto, K., & Tsuji, H. (2018). Identification of phenol- and p-cresol-producing intestinal bacteria by using media supplemented with tyrosine and its metabolites. <i>FEMS</i> <i>Microbiology Ecology</i> , 94(9), 1–11. https://doi.org/10.1093/femsec/fiy125
793 794 795	Samejima, F. (1997). Graded Response Models. In W. J. van der Linden & R. K. Hambleton (Eds.), Handbook of Item Response Theory (pp. 85–100). Springer. https://doi.org/10.1201/9781315119144-6
796 797 798 799	Saoi, M., Li, A., McGlory, C., Stokes, T., von Allmen, M. T., Phillips, S. M., & Britz-Mckibbin, P. (2019). Metabolic perturbations from step reduction in older persons at risk for sarcopenia: plasma biomarkers of abrupt changes in physical activity. <i>Metabolites</i> , 9, 1–19. https://doi.org/10.3390/metabo9070134
800 801 802 803	Seldin, M. M., Meng, Y., Qi, H., Zhu, W. F., Wang, Z., Hazen, S. L., Lusis, A. J., & Shih, D. M. (2016). Trimethylamine N-oxide promotes vascular inflammation through signaling of mitogen-activated protein kinase and nuclear factor-ĸb. <i>Journal of the American Heart Association</i> , 5, e002767. https://doi.org/10.1161/JAHA.115.002767
804 805 806 807 808	Shanmuganathan, M., Kroezen, Z., Gill, B., Azab, S., de Souza, R. J., Teo, K. K., Atkinson, S., Subbarao, P., Desai, D., Anand, S. S., & Britz-McKibbin, P. (2021). The maternal serum metabolome by multisegment injection-capillary electrophoresis-mass spectrometry: a high-throughput platform and standardized data workflow for large-scale epidemiological studies. <i>Nature Protocols</i> , 16, 1966–1994. https://doi.org/10.1038/s41596-020-00475-0
809 810 811	Sheldrick, R. C., & Perrin, E. C. (2013). Evidence-based milestones for surveillance of cognitive, language, and motor development. <i>Acta Paediatrica</i> , 13(6), 577–586. https://doi.org/10.1016/j.acap.2013.07.001
812 813	Sheldrick, R. C., Schlichting, L. E., Berger, B., Clyne, A., Ni, P., Perrin, E. C., & Vivier, P. M. (2019).
814	Establishing new norms for developmental milestones. <i>Pediatrics</i> , <i>144</i> (6), e20190374. https://doi.org/10.1542/peds.2019-0374
814 815 816 817	 Establishing new norms for developmental milestones. <i>Pediatrics</i>, <i>144</i>(6), e20190374. https://doi.org/10.1542/peds.2019-0374 Sotelo-Orozco, J., Hertz-Picciotto, I., Abbeduto, L., & Slupsky, C. M. (2019). Metabolomics analysis of children with autism, idiopathic-developmental delays, and Down syndrome. <i>Translational Psychiatry</i>, <i>9</i>, 1–15. https://doi.org/10.1038/s41398-019-0578-3
814 815 816 817 818 819 820 821	 Establishing new norms for developmental milestones. <i>Pediatrics</i>, <i>144</i>(6), e20190374. https://doi.org/10.1542/peds.2019-0374 Sotelo-Orozco, J., Hertz-Picciotto, I., Abbeduto, L., & Slupsky, C. M. (2019). Metabolomics analysis of children with autism, idiopathic-developmental delays, and Down syndrome. <i>Translational Psychiatry</i>, <i>9</i>, 1–15. https://doi.org/10.1038/s41398-019-0578-3 Sotelo-Orozco, J., Schmidt, R. J., Slupsky, C. M., & Hertz-Picciotto, I. (2023). Investigating the urinary metabolome in the first year of life and its association with later diagnosis of autism spectrum disorder or non-typical neurodevelopment in the MARBLES study. <i>International Journal of Molecular Sciences</i>, <i>24</i>, 1:20. https://doi.org/10.3390/ijms24119454
 814 815 816 817 818 819 820 821 822 823 824 825 	 Establishing new norms for developmental milestones. <i>Pediatrics</i>, <i>144</i>(6), e20190374. https://doi.org/10.1542/peds.2019-0374 Sotelo-Orozco, J., Hertz-Picciotto, I., Abbeduto, L., & Slupsky, C. M. (2019). Metabolomics analysis of children with autism, idiopathic-developmental delays, and Down syndrome. <i>Translational Psychiatry</i>, <i>9</i>, 1–15. https://doi.org/10.1038/s41398-019-0578-3 Sotelo-Orozco, J., Schmidt, R. J., Slupsky, C. M., & Hertz-Picciotto, I. (2023). Investigating the urinary metabolome in the first year of life and its association with later diagnosis of autism spectrum disorder or non-typical neurodevelopment in the MARBLES study. <i>International Journal of Molecular Sciences</i>, <i>24</i>, 1:20. https://doi.org/10.3390/ijms24119454 Tevzadze, G., Barbakadze, T., Kvergelidze, E., Zhuravliova, E., Shanshiashvili, L., & Mikeladze, D. (2022). Gut neurotoxin p-cresol induces brain-derived neurotrophic factor secretion and increases the expression of neurofilament subunits in PC-12 cells. <i>AIMS Neuroscience</i>, <i>9</i>(1), 12–23. https://doi.org/10.3934/NEUROSCIENCE.2022002
 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 	 Establishing new norms for developmental milestones. <i>Pediatrics</i>, <i>144</i>(6), e20190374. https://doi.org/10.1542/peds.2019-0374 Sotelo-Orozco, J., Hertz-Picciotto, I., Abbeduto, L., & Slupsky, C. M. (2019). Metabolomics analysis of children with autism, idiopathic-developmental delays, and Down syndrome. <i>Translational Psychiatry</i>, <i>9</i>, 1–15. https://doi.org/10.1038/s41398-019-0578-3 Sotelo-Orozco, J., Schmidt, R. J., Slupsky, C. M., & Hertz-Picciotto, I. (2023). Investigating the urinary metabolome in the first year of life and its association with later diagnosis of autism spectrum disorder or non-typical neurodevelopment in the MARBLES study. <i>International Journal of Molecular Sciences</i>, <i>24</i>, 1:20. https://doi.org/10.3390/ijms24119454 Tevzadze, G., Barbakadze, T., Kvergelidze, E., Zhuravliova, E., Shanshiashvili, L., & Mikeladze, D. (2022). Gut neurotoxin p-cresol induces brain-derived neurotrophic factor secretion and increases the expression of neurofilament subunits in PC-12 cells. <i>AIMS Neuroscience</i>, <i>9</i>(1), 12–23. https://doi.org/10.3934/NEUROSCIENCE.2022002 Tian, X., Liu, X., Wang, Y., Liu, Y., Ma, J., Sun, H., Li, J., Tang, X., Guo, Z., Sun, W., Zhang, J., & Song, W. (2022). Urinary metabolomic study in a healthy children population and metabolic biomarker discovery of Attention-Deficit/Hyperactivity Disorder (ADHD). <i>Frontiers in Psychiatry</i>, <i>13</i>, 1–12. https://doi.org/10.3389/fpsyt.2022.819498

833	Tingley, D., Yamamoto, T., Hirose, K., Keele, L., & Imai, K. (2014). Mediation: R Package for Causal
834	Mediation Analysis. <i>Journal of Statistical Software</i> , 59(5), 1–38.
835	https://doi.org/https://doi.org/10.18637/jss.v059.i05
836 837	United Nations Children's Fund (UNICEF). (2017). Early Moments Matter for every child. United Nations Children's Fund (UNICEF).
838	van den Heuvel, M. (2019). Metabolomics, stunting and neurodevelopment. <i>EBioMedicine</i> , 44, 10–11.
839	https://doi.org/10.1016/j.ebiom.2019.05.067
840	Vasconcellos, M. T. L., Silva, P. L. N., Castro, I. R. R., Boccolini, C. S., Alves-Santos, N. H., & Kac, G.
841	(2021). Sampling plan of the Brazilian National Survey on Child Nutrition (ENANI-2019): A
842	populationbased household survey. <i>Cadernos de Saude Publica</i> , 37(8), e00037221.
843	https://doi.org/10.1590/0102-311X00037221
844	Wang, H., Hu, P., & Jiang, J. (2012). Measurement of 1- and 3-methylhistidine in human urine by ultra
845	performance liquid chromatography-tandem mass spectrometry. <i>Clinica Chimica Acta</i> , 413, 131–
846	138. https://doi.org/10.1016/j.cca.2011.09.007
847 848 849 850	 Wehrens, R., Hageman, J. A., van Eeuwijk, F., Kooke, R., Flood, P. J., Wijnker, E., Keurentjes, J. J. B., Lommen, A., van Eekelen, H. D. L. M., Hall, R. D., Mumm, R., & de Vos, R. C. H. (2016). Improved batch correction in untargeted MS-based metabolomics. <i>Metabolomics</i>, <i>12</i>, 1–12. https://doi.org/10.1007/s11306-016-1015-8
851	Wei, R., Wang, J., Su, M., Jia, E., Chen, S., Chen, T., & Ni, Y. (2018). Missing value imputation
852	approach for mass spectrometry-based metabolomics data. <i>Scientific Reports</i> , 8, 1–10.
853	https://doi.org/10.1038/s41598-017-19120-0
854	Wellington, N., Shanmuganathan, M., De Souza, R. J., Zulyniak, M. A., Azab, S., Bloomfield, J., Mell,
855	A., Ly, R., Desai, D., Anand, S. S., & Britz-McKibbin, P. (2019). Metabolic trajectories following
856	contrasting prudent and western diets from food provisions: Identifying robust biomarkers of short-
857	term changes in habitual diet. <i>Nutrients</i> , 11, 2407. https://doi.org/10.3390/nu11102407
858 859 860 861 862	 Whitesell, N. R., Sarche, M., Trucksess, C., Ayoub, C. C., Barnes-Najor, J., Ferron, C., Fitzgerald, H. E., Lyon, K., Lundy, C., Meyer, A. L., Myra Parker, J. D., Salvador, M., Walls, M., & Williams, S. (2015). The survey of well-being of young children: Results of a feasibility study with American Indian and Alaska Native communities. <i>Infant Mental Health Journal</i>, <i>36</i>(5), 483–505. https://doi.org/10.1002/imhj.21526
863	World Health Organization (WHO). (2006). WHO Child Growth Standards: Length/Height-forAge,
864	Weight-for-Age, Weight-for-Length, Weight -for-Height and Body Mass Index-for-Age: Methods
865	and Development. In Developmental Medicine & Child Neurology.
866 867 868	World health Organization (WHO), & United Nations Children's Fund (UNICEF). (2021). Indicators for assessing infant and young child feeding practices: definitions and measurement methods. In <i>World Health Organization</i> .
869	Worley, B., & Powers, R. (2013). Multivariate Analysis in Metabolomics. Current Metabolomics, 1(1),
870	92–107. https://doi.org/10.2174/2213235x11301010092
871	Zhu, J., Hua, X., Yang, T., Guo, M., Li, Q., Xiao, L., Li, L., Chen, J., & Li, T. (2022). Alterations in gut
872	vitamin and amino acid metabolism are associated with symptoms and neurodevelopment in
873	children with autism spectrum disorder. <i>Journal of Autism and Developmental Disorders</i> , 52, 3116–
874	3128. https://doi.org/10.1007/s10803-021-05066-w
875	

(Serum metabolome and childhood development)