Endovascular thrombectomy and perioperative stroke

1	
2	Endovascular thrombectomy:
3	an effective and safe therapy for perioperative ischemic stroke
4	
5	Feng Wang ^{1,*} , Xiaoping Xu ^{1,*} , Ling Zheng ¹ , Jiawei Zhong ¹ , En Wang ¹ , Yang Liu ^{1,2,#} , and
6	Shaofa Ke ^{1,#}
7	
8	¹ Department of Neurology, Taizhou Hospital of Zhejiang Province, Affiliated to Wenzhou
9	Medical University, Linhai, Zhejiang, China
10	⁵ Department of Neurology, Saarland University, Homburg, Germany
11	*: These two authors equally contribute to this study
12	#: These two authors share the senior authorship.
13	
14	Running title: Endovascular thrombectomy and perioperative stroke
15	
16	Correspondence should be addressed to:
17	Dr. Yang Liu, Department of Neurology, Saarland University, 66421 Homburg, Germany. Tel:
18	+49-6841-1624260; Fax: +49-6841-1624175; E-mail: <u>a.liu@mx.uni-saarland.de</u> ; ORCID:
19	0000-0002-7614-4233
20	or
21	Dr. Shaofa Ke, Department of Neurosurgery, Taizhou Hospital of Zhejiang Province affiliated
22	to Wenzhou Medical University, 317000 Linhai, Zhejiang province, China. Tel: +86-576-
23	85133031; E-mail: kesf@enzemed.com
24	
25	Total word count of the manuscript: 6134 NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Endovascular thrombectomy and perioperative stroke

1 Abstract

Background: Perioperative ischemic stroke is a rare but devastating complication.
Mechanical thrombectomy is a promising therapeutic method, but very little data is available
on its efficacy and safety. This study aims to answer this question by comparing the clinical
outcomes of perioperative and community-onset stroke patients after endovascular therapy.

Methods: A retrospective cohort study was conducted on a total of 35 perioperative and 584
community-onset acute ischemic stroke patients who underwent endovascular thrombectomy
at our hospital over the past 3.5 years. The recanalization rate, clinical recovery and cerebral
hemorrhage within 90 days after therapy were compared between these two patient groups.

Results: Endovascular thrombectomy provided perioperative and community-onset ischemic 10 stroke patients with comparable rates of successful reperfusion (mTICI \geq 2b grade) (97.1% 11 vs. 97.3%; P = 0.967) and favorable functional recovery (mRS ≤ 2) (51.4% vs. 43.3%, P =12 0.348), with no increase in severe intracranial hemorrhage (0% vs. 2.6% and 1.0%, for 13 hematoma \geq 30% of infarcted tissue and intraventricular hemorrhage, respectively) within 90 14 15 days. In addition, perioperative stroke patients had higher prevalence of atrial fibrillation (42.9 % vs. 26.7 %; P = 0.038) and intracranial cerebral artery stenosis without clear 16 embolism (17.1 % vs. 3.8 %; P < 0.001) than community-onset stroke patients. 17

18 Conclusions: Endovascular thrombectomy is an effective and safe therapeutic approach for 19 patients with perioperative ischemic stroke, although the results need to be validated by 20 further studies with larger populations. Atrial fibrillation and large artery stenosis may 21 contribute to the pathogenesis of perioperative ischemic stroke.

22

Keywords: Perioperative stroke, community-onset stroke, endovascular thrombectomy,
 intracranial hemorrhage, prognosis

Endovascular thrombectomy and perioperative stroke

1 Introduction

2 Perioperative stroke is defined as any embolic, thrombotic or hemorrhagic cerebrovascular incident within 30 days of an operation¹. Similar to community-onset stroke, 3 the majority of perioperative strokes are ischemic rather than hemorrhagic 2 . The incidence of 4 perioperative stroke ranges from approximately 0.075% to 1.9% in non-cardiac/major 5 vascular/neurological operations²⁻⁵ to almost 10% in individuals undergoing high-risk 6 cardiovascular surgery ⁶. In patients undergoing cardiac surgery, emboli are frequently 7 identified as a primary cause of perioperative stroke ^{1 3}. In patients undergoing non-8 cardiac/vascular surgery, the cause of stroke is less clear. Suspected mechanisms of 9 perioperative stroke include cerebral hypotension/low-flow states and previously undetected 10 large-artery stenosis ¹. It is known that intracranial atherosclerotic stenosis is more common 11 in the Asian people than in the Western population ⁷. The mechanism of large artery stenosis 12 in perioperative stroke should be given special attention in Chinese hospitals. 13

Perioperative stroke is a devastating complication for both patients and the treating surgeons/anesthesiologists ³ ⁶. The 30-day mortality rate for patients who suffer a perioperative stroke after non-cardiac/major-vascular/neurologic surgery is up to 8 times higher than for stroke patients unrelated to surgery ³. The length of hospitalization and the likelihood of discharge to a long-term care facility are also increased in patients with a perioperative stroke. An efficient therapy is urgently needed.

Intravenous thrombolysis is usually avoided in cases of perioperative stroke due to surgery itself being a contraindication for thrombolysis, leading to an elevated risk of bleeding ². Mechanical endovascular thrombectomy (EVT) has become a standard treatment for stroke caused by large vessel occlusion ⁸, and also emerged as a novel therapeutic approach for perioperative stroke. Its use has been increasing over the past 15 years ⁹. A retrospective study showed that EVT can achieve comparable success in cerebral reperfusion

Endovascular thrombectomy and perioperative stroke

1 in patients with perioperative stroke and stroke unrelated to surgery, although mortality within 3 months after EVT is higher in the former than in the latter $(33.3\% \text{ vs. } 4.2\%)^{10}$. 2 However, in a retrospective study comparing patients with in-hospital and community-onset 3 4 stroke, EVT resulted in a lower recanalization rate and poorer recovery in in-hospital stroke patients ¹¹. We remain optimistic about EVT therapy in perioperative stroke as surgical 5 6 procedures continue to evolve and minimally invasive procedures in particular is becoming more common, which means that the pathophysiology of current perioperative stroke patients 7 may be different from that in the past. Further research is needed to evaluate the benefits and 8 9 safety of EVT in perioperative stroke.

In our research, we conducted a retrospective analysis of all 697 patients with acute ischemic stroke (AIS) who underwent EVT at our hospital over the past 3.5 years, and compared cerebral reperfusion post therapy, clinical recovery within 3 months, and incidence of cerebral bleeding between patients with perioperative and community-onset stroke. We found that EVT was an effective and safe therapeutic approach for perioperative stroke.

15

16 Methods

17 Study Design

Our project was a retrospective cohort study. The study protocol was approved by the 18 Ethics Committee of Taizhou Hospital, Zhejiang Province, China (Registration number: 19 K20181204). Written consent was waived due to the retrospective design. Between January 20 21 2020 and June 2023, total 697 patients aged ≥ 18 years suffering from AIS (including 35) patients with perioperative stroke) at our hospital received EVT according to the international 22 and Chinese guidelines ^{12 13}. Perioperative stroke was defined as AIS occurring during or 23 within 30 days following the operative procedure ⁶. Inclusion criteria were as follows: Age 24 >18 years; diagnosis of acute ischemic stroke with proven arterial occlusion confirmed by 25

Endovascular thrombectomy and perioperative stroke

computed tomography (CT) or magnetic resonance imaging (MRI); at least one attempt of
 EVT; EVT performed within 24 hours of symptom onset. Exclusion criteria was the presence
 of severe medical conditions, such as advanced heart failure, end-stage kidney disease, or
 terminal illness, which needed more urgent management.

5

EVT and post-operative management

6 Endovascular therapy included stent retriever, aspiration catheter, balloon angioplasty and combination of different techniques. Balloon angioplasty was performed as the 7 therapeutic method of first choice for recanalization in AIS patients who exhibited the "first-8 9 pass effect" of the microcatheter, in which blood was already flowing through the occluded vessel when the microcatheter was first passed and then withdrawn to the proximal side of 10 the occluded vessel during EVT ¹⁴. The number of passes of catheter needed to achieve 11 recanalization or until the end of procedure and the location of endovascular therapy were 12 documented. At the end of EVT, the interventional neurologist determined a modified 13 Thrombolysis in Cerebral Infarction (mTICI) score¹⁵. Twenty-four hours after EVT or at any 14 time within 24 hours when neurologic deficits progressed, patients were reexamined by head 15 CT and scored using the National Institute of Health Stroke Scale (NIHSS). In the absence of 16 hemorrhagic side effects, regular treatment consisted of antiplatelet agents and anticoagulants 17 started 24 hours after EVT. Other treatment strategies included the therapy for causative 18 surgeries in perioperative AIS patients, and statins, blood glucose and blood pressure control 19 or combinations of these treatments according to the Chinese guidelines for the early 20 treatment of AIS patients¹³. 21

22 Data Collection

The aim of this study was to compare the efficacy and safety of EVT in patients with perioperative and community-onset ischemic stroke. The primary efficacy outcome was the recovery of functional outcome as shown in the modified Rankin Scale (mRS) at 90 days

Endovascular thrombectomy and perioperative stroke

after EVT. Good functional outcome was defined as mRS score ≤2. The secondary efficacy
outcome was the attenuation of neurological deficits as assessed by NIHSS scores at 24
hours, and 7 days after EVT (or at the hospital discharge). The third efficacy outcome was
recanalization of the occluded cerebral artery. Successful reperfusion was defined as mTICI
score of 2b, 2c or 3 and unsuccessful revascularization as mTICI score of 0, 1 or 2a after
EVT ¹⁵.

7 The primary safety outcome was intracranial hemorrhage as reflected by CT scanning 8 or MRI, which were typically conducted within 7 days after EVT, with the timing determined 9 by the individual patient's situation and the examination capacity of our neuroradiology 10 department. Upon neurological deterioration, a head CT was immediately performed. The 11 Heidelberg Bleeding Classification was used to categorize the hemorrhage ¹⁶. The second 12 safety outcome was all-cause mortality within 90 days after EVT.

Baseline demographic, clinical information, and laboratory findings within 24 hours after EVT were collected for all enrolled patients. Stroke subtypes were classified according to the Trial of ORG 10172 in the Acute Stroke Treatment classification (TOAST) ¹⁷. The Alberta stroke program early CT score (ASPECTS) was performed for both anterior and posterior circulations on the CT scan at admission ^{18 19}.

18 Statistical analysis

19 The statistical analysis was conducted using SPSS software for Windows (Version 20 26.0, IBM, Armonk, USA). The data for continuous variables were described as mean \pm SD, 21 and categorical variables were presented as frequencies. Continuous variables were compared 22 between independent groups by *T*-test or Mann-Whitney *U* test depending on whether the 23 continuous variables were normally distributed. Categorical variables were compared by 24 Pearson χ^2 test. The association between the reduction of NIHSS score 24 hours after EVT 25 compared with the score before EVT, and various clinical and laboratory parameters was first

Endovascular thrombectomy and perioperative stroke

1 investigated by Pearson correlation test and then linear regression analysis using 2 bootstrapping. P < 0.05 was considered statistically significant.

- 3
- 4 **Results**

5 Patient data

6 From January 2020 to June 2023, 35 perioperative AIS patients and 662 communityonset AIS patients underwent EVT in our hospital according to the international and Chinese 7 guidelines ¹² ¹³. The causative surgeries of 35 perioperative stroke patients are listed in 8 9 Supplementary Table 1, of whom 28 (80.0%) developed ischemic stroke minimally 1 day (1 -28 days with a median duration of 2 days) after surgery and 10 (28.6%) had undergone 10 11 cardiovascular surgery. After discharge from hospital, all AIS patients were followed up for 12 more than 3 months by face-to-face examination or by telephone. Of community-onset AIS 13 patients, 47 patients underwent EVT 24 hours after symptom onset due to the presence of a 14 penumbra on CT or MRI imaging; and 31 AIS patients dropped out during the 90-day follow-15 up period. These 78 patients were not included in this study. Finally, 35 perioperative and 584 community-onset AIS patients were analyzed in this study (see Supplemental Figure S1, 16 17 flowchart).

The demographic and baseline characteristics of perioperative and community-onset 18 AIS patients were shown in Table 1. The atrial fibrillation was more frequent in perioperative 19 group than in community-onset group (42.9% vs. 26.7%; χ^2 (1) = 4.305, P = 0.038). 20 21 However, both the systolic and diastolic blood pressures were lower in perioperative than in community-onset stroke patients (P < 0.001). The number of monocytes, concentrations of C-22 reactive protein and fibrinogen in the blood were higher in perioperative than in community-23 onset AIS patients (P < 0.05). Due to the causative surgeries, significantly more perioperative 24 AIS patients than community-onset patients received anticoagulant or together with 25

Endovascular thrombectomy and perioperative stroke

1 antiplatelet therapy after EVT ($\chi 2$ (3) = 54.721, P < 0.001). All other analyzed parameters, 2 including age, sex, pre-stroke risk factors (e.g., hypertension, diabetes mellitus, coronary heart disease, previous stroke and current smoking), laboratory findings 24 hours after EVT 3 4 (e.g., platelets, D-dimer, glucose, creatinine, cholesterol, et al.), and TOAST classification were not different between these two groups. Of note, there were no significant differences 5 6 between the perioperative and community-onset groups in the neurological severity of stroke patients as shown by NIHSS scores before EVT (median value 17.4 ± 10.1 vs. 15.9 ± 7.6 ; 7 Mann-Whitney-U-Test, Z = -0.553, P = 0.580). Thus, these 2 comparable groups of AIS 8 9 patients were suitable for analysis of the efficacy and safety of EVT treatments in perioperative stroke. 10

11

EVT provided perioperative AIS patients with a comparable reperfusion rate and 3month rehabilitation, and even a faster recovery within 24 hours compared to community-onset AIS patients

15 There was no significant difference in the rate of successful reperfusion $[mTICI \ge 2b]$ grade]) between perioperative and community-onset AIS patients (Table 2; 97.1% vs. 97.3%; 16 χ^2 (1) = 0.002, P = 0.967). Similarly, the distribution of mTICI scores (1 = mTICI 0, 1 or 2a, 17 2 = mTICI 2b, and 3 = mTICI 3), which categorize the level of successful reperfusion, did 18 not show any significant differences between these two AIS patient groups (Supplementary 19 Fig. S2; $\chi 2$ (2) = 0.035, P = 0.983). It was also found that the causative surgeries did not 20 complicate the EVT procedure, as the number of catheter passes required for recanalization 21 did not vary significantly between these two groups of patients (Table 2; 1.6 ± 0.9 vs. $1.4 \pm$ 22 0.8; Mann-Whitney-U-Test, Z = -1.436, P = 0.151). In addition, perioperative and 23 community-onset AIS patients did not differ in operative time of EVT and the site of arterial 24 occlusion (Table 2). 25

Endovascular thrombectomy and perioperative stroke

It should be noted that there were more patients in the perioperative stroke group than in the community-onset stroke group (Table 2; 17.1% vs. 3.8%; χ^2 (1) = 13.680, *P* < 0.001), who had the "first-pass effect" of the microcatheter, which suggests severe stenosis in the intracranial cerebral arteries and not severe embolism-related vascular occlusion ²⁰, and whose cerebral blood flow could be sufficiently improved by balloon angioplasty alone.

6 Possibly due to the professional medical care in the hospital, patients with perioperative stroke were much easier to recognize than patients with community-onset 7 stroke. As shown in Table 2, the time from onset of symptoms to puncture was significantly 8 shorter (202.5 \pm 217.3 vs. 448.6 \pm 279.7 minutes; Mann-Whitney-U-Test, Z = -6.553, P < 9 0.001), and the ASPECT score was significantly higher (9.9 \pm 0.7 vs. 8.5 \pm 1.8; Mann-10 11 Whitney-U-Test, Z = -5.454, P < 0.001) in perioperative AIS patients than in the community-12 onset AIS controls. The improvement in neurological deficits shown by the reduction in NIHSS score 24 hours after EVT compared with NIHSS score before EVT was significantly 13 more pronounced in perioperative stroke patients than in community-onset stroke patients 14 (Table 2; Δ NIHSS score: 5.5 ± 8.9 vs. 1.56 ± 6.0; Mann-Whitney-U-Test, Z = -3.052, P = 15 0.002). Unfortunately, this favorable recovery after EVT was lost in perioperative AIS 16 patients compared to community-onset AIS patients in the following 7 days, as the NIHSS 17 scores did not differ between perioperative and community-onset AIS patients 7 days after 18 EVT (Table 2; 9.4 ± 9.4 vs. 11.0 ± 12.0 ; Mann-Whitney-U-Test, Z = -1.271, P = 0.204). 19

To investigate the underlying mechanisms hindering further recovery of perioperative AIS patients, we combined perioperative and community-onset AIS patients and performed a correlation analysis to identify variables related to NIHSS score reduction within 24 hours after EVT (see Supplementary Table 2). We then performed a regression analysis with NIHSS reduction as the dependent variable and the correlated parameters as independent variables. We found that perioperative stroke classification, smoking and NIHSS score at admission

Endovascular thrombectomy and perioperative stroke

positively predicted, while diabetes, white blood cell count, D-dimer, creatine, systolic blood pressure, puncture-to-recanalization time and location of occluded artery with MCA as the reference location negatively predicted NIHSS score reduction (Supplementary Table 3). In combination with the high levels of monocytes and C-reactive protein in perioperative AIS patients compared to community-onset stroke patients (Table 1), inflammatory activation could be an influential factor affecting clinical recovery.

7 To assess the functional recovery of all AIS patients under study, mRS scores were evaluated 90 days after EVT. The analysis revealed no significant differences in the 8 9 distribution of mRS scores between patients with perioperative and community-onset AIS (Figure 1; χ^2 (6) = 5.923, P = 0.432). The percentage of patients achieving a favorable 10 11 outcome (mRS \leq 2) in perioperative group was comparable with that in community-onset 12 AIS group (Table 2; 51.4% vs. 43.3%, χ^2 (1) = 0.882, P = 0.348). The mortalities during the entire course of the study were also not significantly different between perioperative and 13 community-onset AIS patients (Table 2; 11.4% vs. 17.8%, $\gamma 2$ (1) = 0.933, P = 0.334). 14

15

16 EVT did not increase a hemorrhagic risk for perioperative AIS patients compared with 17 community-onset AIS patients

community-onset AIS patients

Hemorrhagic transformation is the most important adverse event after EVT therapy in 18 AIS patients. The percentage of patients diagnosed with overall intracranial hemorrhage 19 20 within 7 days after EVT was comparable in the perioperative and community-onset AIS 21 groups (Table 3; 25.7% vs. 19.5%; χ^2 (1) = 0.796, P = 0.372). After dividing the AIS patients into subgroups according to the Heidelberg Bleeding Classification ¹⁶, we found that 22 perioperative and community-onset AIS patients differed significantly in terms of intracranial 23 hemorrhage (Table 3; $\chi 2$ (5) = 16.299, P = 0.004). However, the difference was limited in the 24 subgroups of HI1 (11.4% vs. 3.8; χ^2 (1) = 4.817, P = 0.028) and PH1-type (8.6% vs. 2.4%; 25

Endovascular thrombectomy and perioperative stroke

χ2 (1) = 4.713, P = 0.030) hemorrhages, and subarachnoid hemorrhage (5.7% vs. 0.9%; χ2
 (1) = 6.970, P = 0.008) (Table 3). There were no perioperative AIS patients in the subgroups
 of PH2 and intraventricular hemorrhage, perhaps due to the limited sample size (Table 3).

4

5 Discussion

6 Intravenous thrombolysis is often not feasible in perioperative AIS patients as it 7 increases the risk of bleeding ²⁶. EVT is the most useful method for recanalization of cerebral 8 arteries in perioperative AIS patients and is increasingly practiced in many hospitals ⁹. 9 However, the efficacy of EVT in improving patient outcomes remains uncertain. Our study 10 showed that EVT in perioperative AIS patients resulted in a similar reperfusion rate and 3-11 month recovery without an increased risk of bleeding and death compared to patients with 12 community-onset stroke.

In our study, we used the entire cohort of community-onset stroke patients instead of 13 selecting certain matched AIS patients as a control group for perioperative AIS patients, 14 15 which we believe can limit the sampling bias and better represent the real-world information. The NIHSS score is higher in perioperative AIS patients than in community-onset AIS 16 patients, even though there is no statistical difference, which demonstrates the efficacy of 17 EVT in perioperative stroke. We found that atrial fibrillation was significantly more common 18 in perioperative AIS patients than in community-onset AIS patients, which is consistent with 19 20 previous studies that perioperative or postoperative atrial fibrillation was associated with an 21 increased risk of both early and long-term ischemic stroke, especially in patients undergoing non-cardiac surgery ^{21 22}. Our study seems to show that undetected stenosis of the intracranial 22 arteries is a pathogenic mechanism of perioperative stroke. We observed more AIS patients in 23 the perioperative stroke group than in the community-onset stroke group (17.1 % vs. 3.8 %) 24 who had atherosclerotic stenosis in the cerebral arteries without clear thrombi or emboli, in 25

Endovascular thrombectomy and perioperative stroke

whom balloon angioplasty alone was able to restore blood flow to the brain tissue.
Intracranial atherosclerotic stenoses pose a challenge for EVT in AIS patients as they increase
intraprocedural re-occlusion ²³. A recent study in AIS patients with intracranial
atherosclerosis-related large vessel occlusion showed that balloon angioplasty as a firstchoice recanalization strategy has a higher efficiency in recanalization and better functional
outcomes at 90 days compared to thrombectomy ¹⁴.

7 Our study showed that the recanalization rate after EVT did not differ between perioperative and community-onset AIS stroke patients, which corroborates previous studies 8 on stroke patients after both cardiovascular and non-cardiovascular surgeries ^{10 24}. 9 Importantly, the time from symptom onset to groin puncture in our study is shorter in 10 11 perioperative stroke patients than in community-onset AIS stroke patients (median time, 239 12 vs. 433 minutes). The symptom onset-to-groin puncture time is strongly associated with better clinical outcome, e.g., functional independence at discharge ²⁵. We found that the 13 14 reduction in NIHSS scores within 24 hours of EVT was significantly greater in perioperative AIS patients than in community-onset AIS patients, although this difference disappeared 15 when patients were discharged from our hospital. We observed that: 1) the increase in white 16 blood cell count was negatively associated with NIHSS score reduction after EVT; and 2) 17 monocyte counts and C-reactive protein amount were higher in perioperative patients than in 18 19 community-onset AIS patients. It therefore stands to reason that inflammatory activation impedes functional recovery in perioperative AIS patients. It has been reported that the 20 increase in C-reactive protein 26 , or the inflammatory index C-reaction protein \times 21 neutrophil/lymphocyte count ²⁷ predicts an unfavorable functional outcome in AIS patients 90 22 day after EVT. 23

Our study also showed that EVT provided perioperative and community-onset AIS patients with comparable functional recovery and mortality 3 months after EVT, which is

Endovascular thrombectomy and perioperative stroke

1 different from a previous observation that perioperative AIS patients had a higher rate of death within 3 months of EVT than community-onset AIS patients ¹⁰. Not surprisingly, the 2 underlying diseases requiring surgeries and comorbidities influence the outcome of 3 perioperative AIS patients after EVT. In the previous study ¹⁰, there were 68% perioperative 4 AIS patients receiving cardiovascular surgery and 12% receiving neurosurgery, while we had 5 only 29% patients in the perioperative AIS group, who had undergone cardiovascular 6 procedure. In studies of the therapeutic efficacy of EVT in in-hospital and community-onset 7 AIS patients, the former generally had poorer recovery and higher mortality, which was 8 9 correlated with the modified Charlson Comorbidity Index (mCCI), incorporating 7 comorbidities, age, diabetes, anemia, active cancer, myocardial infarction, congestive heart 10 disease, and ulcer disease into the model ^{11 28}. We further suppose that the widely used mini-11 invasive surgery in our studying cohort may favor the functional recovery of perioperative 12 AIS patients. Compared to open surgical procedures such as sternotomy or open thoracotomy, 13 minimally invasive techniques offer advantages to patients, such as a lower risk of surgical 14 and postoperative complications, shorter recovery times and a reduction in postoperative 15 pain, which can reduce systemic inflammation and hemodynamic changes ²⁹. 16

Consistent with previous studies ^{10 11 24 28}, EVT did not lead to an increase in total 17 intracranial hemorrhage in perioperative AIS patients compared to community-onset AIS 18 patients. However, when hemorrhages with different subtypes according to the Heidelberg 19 hemorrhage classification were considered, there were significantly more perioperative AIS 20 21 patients with HI1 and PH1 type hemorrhages and subarachnoid hemorrhage than communityonset AIS patients. Nevertheless, these types of intracranial hemorrhage were generally 22 thought to have little impact on patient outcome. There were no perioperative AIS patients 23 with PH2-type hemorrhage and intravascular hemorrhage, which often lead to symptomatic 24 intracranial hemorrhage and poorer prognosis ^{16 30}. 25

Endovascular thrombectomy and perioperative stroke

1 Obviously, our study has a limitation. The study population was recruited from a 2 single institution with a limited number of patients. The patients cannot be further subdivided 3 to evaluate the therapeutic efficacy and safety of EVT in perioperative AIS patients with 4 cardiovascular or non-cardiovascular procedures. The incidence of PH2-type intracranial 5 hemorrhage and intravascular hemorrhage in perioperative AIS patients after EVT also 6 remains unclear.

7

8 Conclusion

9 Our study shows that endovascular thrombectomy or balloon angioplasty is an 10 effective and safe therapeutic method for the treatment of perioperative strokes with large 11 vessel occlusions. It is helpful for clinicians to make treatment decisions for perioperative 12 stroke patients. Our study also supports the hypothesis that atrial fibrillation and intracranial 13 cerebral artery stenosis contribute to the occurrence of perioperative stroke. However, our 14 results need to be validated by further studies with larger populations.

15

Endovascular thrombectomy and perioperative stroke

1 Acknowledgments

- 2 The authors thank all patients for contributing to this study.
- 3

4 Funding

- 5 This work was supported by grants from Zhejiang Provincial Basic and Public Welfare
- 6 Research Program (No. GF21H020023); Zhejiang Provincial Medicine and Health Research
- 7 Foundation (grant number: 2021RC141).
- 8

9 Data availability statement

10 The data that support the findings of this study are available on request from the 11 corresponding author, Y.L. and S.K. The data are not publicly available because it contains 12 information that could compromise the privacy of the research participants.

13

14 Contributions

15 Study idea and design: Feng Wang, Yang Liu and Shaofa Ke; Coordination and organization:

- 16 Feng Wang and Shaofa Ke; Participants enrollment: En Wang; Data collection and statistical
- 17 analysis: Xiaoping Xu, Ling Zheng, and Yang Liu; Writing, editing and revision: Feng Wang
- 18 and Yang Liu.
- 19

20 Ethics declarations

21 Conflict of interest

- The author(s) declared no potential conflicts of interest with respect to the research,authorship, and/or publication of this article.
- 24

25 Ethical standards

- 26 The studies involving human participants were reviewed and approved by the Ethics
- 27 Committee of Taizhou Hospital, Zhejiang Province, China (K20181204).
- 28

Endovascular thrombectomy and perioperative stroke

1 References

- Benesch C, Glance LG, Derdeyn CP, et al. Perioperative Neurological Evaluation and Management to Lower the Risk of Acute Stroke in Patients Undergoing Noncardiac, Nonneurological Surgery: A Scientific Statement From the American Heart Association/American Stroke Association. *Circulation* 2021;143(19):e923-e46. doi: 10.1161/CIR.00000000000968
- 7 2. Mashour GA, Moore LE, Lele AV, et al. Perioperative care of patients at high risk for stroke during or after non-cardiac, non-neurologic surgery: consensus statement from the Society for Neuroscience in Anesthesiology and Critical Care*. J Neurosurg Anesthesiol 2014;26(4):273-85. doi: 10.1097/ANA.0000000000000087
- Mashour GA, Shanks AM, Kheterpal S. Perioperative stroke and associated mortality after noncardiac, nonneurologic surgery. *Anesthesiology* 2011;114(6):1289-96. doi: 10.1097/ALN.0b013e318216e7f4
- 4. Vasivej T, Sathirapanya P, Kongkamol C. Incidence and Risk Factors of Perioperative
 Stroke in Noncardiac, and Nonaortic and Its Major Branches Surgery. J Stroke *Cerebrovasc* Dis 2016;25(5):1172-76. doi:
 10.1016/j.jstrokecerebrovasdis.2016.01.051
- 18 5. Bateman BT, Schumacher HC, Wang S, et al. Perioperative acute ischemic stroke in noncardiac and nonvascular surgery: incidence, risk factors, and outcomes.
 20 Anesthesiology 2009;110(2):231-8. doi: 10.1097/ALN.0b013e318194b5ff
- 21 6. Selim M. Perioperative stroke. N Engl J Med 2007;356(7):706-13. doi:
 22 10.1056/NEJMra062668
- 7. Kim JS, Kim YJ, Ahn SH, et al. Location of cerebral atherosclerosis: Why is there a
 difference between East and West? *Int J Stroke* 2018;13(1):35-46. doi:
 10.1177/1747493016647736
- 8. Maingard J, Foo M, Chandra RV, et al. Endovascular Treatment of Acute Ischemic Stroke.
 Curr Treat Options Cardiovasc Med 2019;21(12):89. doi: 10.1007/s11936-019-0781-9 [published Online First: 20191211]
- 9. Shah S, Hatfield J, Fuller M, et al. Mechanical thrombectomy for perioperative ischemic
 stroke following elective inpatient surgery in the United States. *J Clin Neurosci*2022;101:100-05. doi: 10.1016/j.jocn.2022.05.009
- 32 10. Premat K, Clovet O, Frasca Polara G, et al. Mechanical Thrombectomy in Perioperative
 33 Strokes: A Case-Control Study. Stroke 2017;48(11):3149-51. doi:
 34 10.1161/STROKEAHA.117.018033
- Monch S, Lehm M, Maegerlein C, et al. Worse endovascular mechanical recanalization
 results for patients with in-hospital onset acute ischemic stroke. *J Neurol* 2018;265(11):2525-30. doi: 10.1007/s00415-018-9035-0
- 12. Powers WJ, Rabinstein AA, Ackerson T, et al. Guidelines for the Early Management of
 Patients With Acute Ischemic Stroke: 2019 Update to the 2018 Guidelines for the
 Early Management of Acute Ischemic Stroke: A Guideline for Healthcare
 Professionals From the American Heart Association/American Stroke Association. *Stroke* 2019;50(12):e344-e418. doi: 10.1161/STR.000000000000211 [published
 Online First: 20191030]
- Liu L, Chen W, Zhou H, et al. Chinese Stroke Association guidelines for clinical management of cerebrovascular disorders: executive summary and 2019 update of clinical management of ischaemic cerebrovascular diseases. *Stroke Vasc Neurol* 2020;5(2):159-76. doi: 10.1136/svn-2020-000378 [published Online First: 20200618]

Endovascular thrombectomy and perioperative stroke

- 14. Zhang L, He X, Li K, et al. Balloon angioplasty as first-choice recanalization strategy for intracranial atherosclerosis-related emergent large vessel occlusion with small clot burden. *Neuroradiology* 2024;66(3):399-407. doi: 10.1007/s00234-023-03278-8
- 4 15. Higashida RT, Furlan AJ, Roberts H, et al. Trial design and reporting standards for intraarterial cerebral thrombolysis for acute ischemic stroke. *Stroke* 2003;34(8):e109-37.
 6 doi: 10.1161/01.STR.0000082721.62796.09
- 7 16. von Kummer R, Broderick JP, Campbell BC, et al. The Heidelberg Bleeding
 8 Classification: Classification of Bleeding Events After Ischemic Stroke and
 9 Reperfusion Therapy. *Stroke* 2015;46(10):2981-6. doi:
 10 10.1161/STROKEAHA.115.010049
- 17. Adams HP, Jr., Bendixen BH, Kappelle LJ, et al. Classification of subtype of acute 11 12 ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of 13 Org 10172 in Acute Stroke Treatment. Stroke 1993;24(1):35-41. doi: 10.1161/01.str.24.1.35 14
- 18. Barber PA, Demchuk AM, Zhang J, et al. Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. ASPECTS Study Group. Alberta Stroke Programme Early CT Score. *Lancet* 2000;355(9216):1670-4. doi: 10.1016/s0140-6736(00)02237-6
- Puetz V, Sylaja PN, Coutts SB, et al. Extent of hypoattenuation on CT angiography source
 images predicts functional outcome in patients with basilar artery occlusion. *Stroke* 2008;39(9):2485-90. doi: 10.1161/STROKEAHA.107.511162
- 20. Yi TY, Chen WH, Wu YM, et al. Microcatheter "First-Pass Effect" Predicts Acute
 Intracranial Artery Atherosclerotic Disease-Related Occlusion. *Neurosurgery* 24 2019;84(6):1296-305. doi: 10.1093/neuros/nyy183
- 25 21. Siontis KC, Gersh BJ, Weston SA, et al. Association of New-Onset Atrial Fibrillation
 26 After Noncardiac Surgery With Subsequent Stroke and Transient Ischemic Attack.
 27 JAMA 2020;324(9):871-78. doi: 10.1001/jama.2020.12518
- 28 22. Lin MH, Kamel H, Singer DE, et al. Perioperative/Postoperative Atrial Fibrillation and
 29 Risk of Subsequent Stroke and/or Mortality. *Stroke* 2019;50(6):1364-71. doi:
 30 10.1161/STROKEAHA.118.023921
- 31 23. Tsang ACO, Orru E, Klostranec JM, et al. Thrombectomy Outcomes of Intracranial
 32 Atherosclerosis-Related Occlusions. *Stroke* 2019;50(6):1460-66. doi:
 33 10.1161/STROKEAHA.119.024889
- 24. Bay B, Gloyer NO, Remmel M, et al. Mechanical thrombectomy in ischemic stroke after
 cardiovascular procedures: a propensity-matched cohort analysis. *J Neurointerv Surg* 2023;15(e1):e129-e35. doi: 10.1136/jnis-2022-019152
- 25. Jahan R, Saver JL, Schwamm LH, et al. Association Between Time to Treatment With
 Endovascular Reperfusion Therapy and Outcomes in Patients With Acute Ischemic
 Stroke Treated in Clinical Practice. *JAMA* 2019;322(3):252-63. doi:
 10.1001/jama.2019.8286
- 26. Finck T, Sperl P, Hernandez-Petzsche M, et al. Inflammation in stroke: initial CRP levels
 can predict poor outcomes in endovascularly treated stroke patients. *Front Neurol*2023;14:1167549. doi: 10.3389/fneur.2023.1167549
- 27. Du M, Xu L, Zhang X, et al. Association Between Inflammatory Burden Index and Unfavorable Prognosis After Endovascular Thrombectomy in Acute Ischemic Stroke.
 J Inflamm Res 2023;16:3009-17. doi: 10.2147/JIR.S419087
- 47 28. Jung JW, Kim KH, Yun J, et al. Effectiveness of endovascular treatment for in-hospital
 48 stroke vs. community-onset stroke: a propensity score-matched analysis. *Journal of* 49 *Neurology* 2024 doi: 10.1007/s00415-024-12232-4

Endovascular thrombectomy and perioperative stroke

- 29. Mohiuddin K, Swanson SJ. Maximizing the benefit of minimally invasive surgery. J Surg
 Oncol 2013;108(5):315-9. doi: 10.1002/jso.23398
- 30. Hall E, Ullberg T, Andsberg G, et al. Incidence of intracranial hemorrhagic complications
 after anterior circulation endovascular thrombectomy in relation to occlusion site: a
 nationwide observational register study. *J Neurointerv Surg* 2023 doi: 10.1136/jnis 2023-020768
- 7
- 8 Table 1, Characteristics of the patients

	Total	Perioperative	Community-		
Variables	(n = 619)	stroke	onset stroke	<i>P</i> -value	
Demographic characteristics		(n = 35)	(n = 584)		
Female n (%)	237 (38 3)	14 (40 0)	223 (38.2)	0.830	
Age mean \pm SD vears	237(30.3)	14(40.0) 65.83 + 12.21	69.05 ± 12.05	0.050	
Rick factors for stroke	00.07 ± 12.07	05.05 ± 12.21	07.05 ± 12.05	0.125	
Atrial fibrillation $n(0/2)$	171 (27.6)	15 (42.0)	156 (267)	0.028	
Attraction $n (\%)$	1/1(27.0)	13(42.9)	130(20.7)	0.030	
Diskatas a (9()	507 (59.5) 112 (19.2)	21(00.0)	540 (59.2) 107 (19.2)	0.950	
Diabetes, n (%)	113 (18.3)	6 (17.1) 2 (5.7)	107 (18.3)	0.861	
Coronary heart disease, n (%)	35 (5.7)	2 (5.7)	33 (5.7)	0.987	
Previous stroke, n (%)	124 (20.0)	5 (14.3)	119 (20.4)	0.382	
Smoking, n (%)	186 (30.0)	7 (20.0)	179 (30.7)	0.182	
Laboratory findings					
White blood cells, median (P25, P75), 10 ⁹ /L	9.3 (7.5, 11.5)	10 (8.4, 12.9)	9.3 (7.4, 11.3)	0.119	
Monocytes, median (P25, P75), 10 ² /L	0.4 (0.3, 0.6)	0.6 (0.4, 0.7)	0.4 (0.3, 0.6)	0.002	
Neutrophils, median (P25, P75), 10 ⁹ /L	7.7 (5.8, 9.8)	8.3 (6, 11.5)	7.6 (5.8, 9.7)	0.110	
Lymphocytes, median (P25, P75), 10 ⁹ /L	1 (0.7, 1.4)	1.1 (0.8, 1.4)	1 (0.7, 1.4)	0.404	
Platelets, median (P25, P75), 10 ⁹ /L	191 (152, 227)	188 (147, 241)	191 (152, 226)	0.927	
C-reactive protein, median (P25, P75), mg/L	6.8 (2.9, 18.7)	15.4 (4.9, 54.4)	6.4 (2.7, 17.4)	< 0.001	
Fibrinogen, median (P25, P75), g/L	3.1 (2.6, 3.7)	3.7 (3.1, 4.8)	3.1 (2.6, 3.7)	< 0.001	
D-dimer, median (P25, P75), mg/L	1.3 (0.7, 2.5)	1.5 (0.6, 2.3)	1.3 (0.7, 2.5)	0.594	
Cr, median (P25, P75), µmoI/L	66 (54, 81)	63 (50, 86)	66 (55, 80)	0.506	
BUN, median (P25, P75), mmoI/L	5.0 (3.9, 6.4)	5.5 (4.8, 6.7)	5.0 (3.9, 6.3)	0.069	
Glucose, median (P25, P75), mmoI/L	6.7 (5.6, 8.4)	6.6 (5.6, 8.5)	6.7 (5.5, 8.4)	0.938	
TG, median (P25, P75), mmoI/L	1.0 (0.7, 1.5)	1.1 (0.7, 1.8)	1.0 (0.7, 1.5)	0.911	
TC, median (P25, P75), mmoI/L	4.0 (3.4, 4.7)	4.1 (3.1, 4.7)	4.0 (3.4, 4.7)	0.806	
HDL, median (P25, P75), mmoI/L	1.2 (1.0, 1.4)	1.1 (1.0, 1.4)	1.2 (1.0, 1.4)	0.913	
LDL, median (P25, P75), mmoI/L	2.2 (1.7, 2.7)	2.2 (1.5, 2.6)	2.2 (1.7, 2.7)	0.642	
Hemoglobulin, median (P25, P75), g/L	126 (114, 136)	126 (114, 133)	126 (114, 136)	0.637	
Clinical findings					
NIHSS at admission, median (P25, P75)	15 (11, 19)	16 (11, 20)	15 (11, 19)	0.581	
Systolic blood pressure, mean \pm SD, mmHg	151 ± 25	132 ± 21	152 ± 24	< 0.001	
Diastolic blood pressure, median (P25, P75),	87 (76, 98)	79 (70, 84)	87 (77, 99)	< 0.001	
TOAST classification				0.056	
Cardioembolism	254 (41.0)	14 (40 0)	240 (41 1)	0.000	
Large-artery atherosclerosis	341 (55 1)	17 (48 6)	324 (55 5)		
Stroke of undetermined or other determined	511 (55.1)	1, (10.0)	52 (55.5)		
etiology	24 (3.9)	4 (11.4)	20 (3.4)		
Postoperative medication				< 0.001	

Endovascular thrombectomy and perioperative stroke

No antiplatelet and anticoagulant drugs, n (%)	72 (11.6)	2 (5.7)	70 (12.0)
Antiplatelet drugs, n (%)	512 (82.7)	24 (68.6)	488 (83.6)
Anticoagulant drugs, n (%)	8 (1.3)	5 (14.3)	3 (0.5)
Antiplatelet and anticoagulant drugs, n (%)	27 (4.4)	4 (11.4)	23 (3.9)

Note: Continuous variables with normal distribution were expressed as mean \pm SD and non-normally distributed data were expressed as median (IQR). Categorical variables were presented as frequencies. Cr, serum creatinine; BUN, blood urea nitrogen; TG, triglyceride; TC, total cholesterol; HDL, high density lipoprotein; LDL, low density lipoprotein; NIHSS, National Institute of Health stroke scale. Continuous variables were compared between independent groups by *T*-test or Mann-Whitney *U* test depending on whether the continuous variables were normally distributed. Categorical variables were compared by Pearson χ^2 test.

Endovascular thrombectomy and perioperative stroke

1 Table 2, Characteristics of endovascular therapy procedure

Variables	Total (n = 619)	Perioperative stroke	Community- onset stroke	<i>P</i> -value
		(n = 35)	(n = 584)	
ASPECT score, median (P25, P75)	9 (8, 10)	10 (10, 10)	9 (8, 10)	< 0.001
Fibrinolysis	189 (30.5)	2 (5.7)	187 (32.0)	0.001
Endovascular therapy				
Onset-to-puncture time, median (P25, P75), minutes	420 (300, 655)	239 (178, 299)	433 (315, 666)	< 0.001
Puncture-to-recanalization time, median (P25, P75), minutes	66 (45, 100)	60 (38, 90)	66 (45, 102)	0.129
Successful reperfusion (mTICI $\ge 2b$ grade)	602 (97.3)	34 (97.1)	568 (97.3)	0.967
Numbers of passes of catheter, median (P25, P75)	1 (1, 2)	1 (1, 2)	1 (1, 2)	0.151
Location of occluded artery				0.222
Terminus internal carotid artery, n (%)	114 (18.4)	10 (28.6)	104 (17.8)	
Middle cerebral artery, n (%)	316 (51.1)	20 (57.1)	296 (50.7)	
Anterior cerebral artery, n (%)	16 (2.6)	1 (2.9)	15 (2.6)	
\geq 2 arteries in anterior circulation, n (%)	58 (9.4)	0 (0)	58 (9.9)	
Basilar artery, n (%)	80 (12.9)	3 (8.6)	77 (13.2)	
Vertebral artery, n (%)	25 (4.0)	0 (0)	25 (4.3)	
Posterior cerebral artery, n (%)	10 (1.6)	1 (2.9)	9 (1.5)	
Endovascular therapy procedure				< 0.001
Stent retriever, n (%)	423 (68.3)	24 (68.6)	399 (68.3)	
Balloon angioplasty, n (%)	28 (4.5)	6 (17.1)	22 (3.8)	
Stent retriever and balloon angioplasty, n (%)	127 (20.5)	1 (2.9)	126 (21.6)	
Aspiration catheter, n (%)	17 (2.7)	3 (8.6)	14 (2.4)	
Aspiration catheter or stent retriever, n (%)	24 (3.9)	1 (2.9)	23 (3.9)	
Clinical findings				
NIHSS at 24 hours after EVT, median (P25, P75)	13 (7, 19)	8 (5, 18)	13 (7, 19)	0.094
Reduction of NIHSS at 24 hours compared to NIHSS before EVT, median (P25, P75)	0 (0, 5)	5 (0, 10)	0 (0, 4)	< 0.001
NIHSS at 7 days after EVT, median (P25, P75)	7 (2, 15)	3 (1, 14)	7 (2, 15)	0.204
mRS \leq 2 at 90 days after EVT, n (%)	271 (43.8)	18 (51.4)	253 (43.3)	0.348
Mortality at the end of study, n (%)	108 (17.4)	4 (11.4)	104 (17.8)	0.933

Note: Non-normally distributed data were expressed as median (IQR). Categorical variables were presented as frequencies. Continuous variables were compared between independent groups by Mann-Whitney U test. Categorical variables were

6

compared by Pearson $\chi 2$ test.

Endovascular thrombectomy and perioperative stroke

1 Table 3, Characteristics of intracranial hemorrhage

Variables	Total (n = 619)	Peri-operative stroke (n = 35)	Community- onset stroke (n = 584)	χ2 test P-value
Intracranial hemorrhages	123 (19.9)	9 (25.7)	114 (19.5)	0.372
The Heidelberg Bleeding Classification				0.004
0. No hemorrhages	496 (80.1)	26 (74.3)	470 (80.5)	
1. Hemorrhagic transformation of infarcted brain tissue				
1a (HI1), Scattered small petechiae, no mass effect	26 (4.2)	4 (11.4)	22 (3.8)	
1b (HI2), Confluent petechiae, no mass effect	54 (8.7)	0 (0)	52 (8.9)	
1c (PH1), Hematoma within infarcted tissue, occupying <30%, no substantive mass effect	16 (2.6)	3 (8.6)	14 (2.4)	
2. Intracerebral hemorrhage within and beyond infarcted brain tissue				
PH2, Hematoma occupying 30% or more of the infarcted tissue, with obvious mass effect	14 (2.3)	0 (0)	15 (2.6)	
3. Intracerebral hemorrhage outside the infarcted b	3. Intracerebral hemorrhage outside the infarcted brain tissue			
3a Parenchymal hematoma remote from infarcted brain tissue	13 (2.1)	0 (0)	1 (0.2)	
3b Intraventricular hemorrhage		0 (0)	6 (1.0)	
3c Subarachnoid hemorrhage		2 (5.7)	5 (0.9)	

Endovascular thrombectomy and perioperative stroke

1 Figure legends

2 Figure 1. Endovascular thrombectomy achieves comparable functional recovery in perioperative and community-onset stroke patients. Functional recovery of patients with 3 4 acute ischemic stroke was assessed using the modified Rankin Scale (mRS) at 90 days after endovascular thrombectomy. The mRS scores range from 0 to 6, with score 0 indicating no 5 symptoms, to score 5 indicating severe disability and score 6 indicating death. The 6 7 distribution of patients with different mRS scores did not differ between the perioperative and community-onset stroke patient groups. χ^2 test, n = 35 and 584 for perioperative and 8 9 community-onset stroke groups, respectively.

10



(Pearson χ^2 test, *P*=0.432)