

Neuroblastoma in Adults: A Scoping Review of Presentations, Genetics and Therapies

Bader H. Alsaikhan, MD, MSc, FRCSC^{1,2,3}; Basmah Alwahhabi, MBBS, FRCP(UK)⁴; Abdullah Alshalan, MD⁵; Alex Koziarz, MD, MSc⁶; Abdullah M. Alkhayal, MD, FRCSC^{1,2,3}; Khalid Alrabeeah, MD, FRCSC^{1,2,3}

1. College of Medicine, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia.
2. King Abdullah International Medical Research Centre (KAIMARC), Riyadh, Saudi Arabia.
3. Division of Urology, Department of Surgery, King Abdulaziz Medical City, National Guard Health Affairs (MNGHA), Riyadh, Saudi Arabia.
4. Department of Medicine, King Abdullah bin Abdulaziz University Hospital, Riyadh, Saudi Arabia.
5. Department of Surgery, Division of Urology, King Abdulla bin Abdulaziz University Hospital, Riyadh Saudi Arabia.
6. Department of Radiology, McMaster University, Hamilton, Ontario, Canada.

Take home messages

- 679 cases of adult neuroblastoma have been published in the literature.
- Compared to pediatrics, adult cases have more somatic mutations (eg: *ALK*, *ATRX*).
- Registry data showed that adult neuroblastoma has a 5-year survival of 36%.

Co-Investigators:

Bader H. Alsaikhan, MD, MSc, FRCSC
Alex Koziarz, MD, MSc

Correspondence:

Bader H. Alsaikhan, MD, MSc, FRCSC
Assistant Professor
Division of Urology
King Saud bin Abdulaziz University for Health Sciences
King Abdullah International Medical Research Centre
National Guard Health Affairs (MNGHA)
Riyadh, Saudi Arabia
Tel: +966 56 309 2222
Email: alsaikhanba@mngha.med.sa

Acknowledgements: none.

Conflicts of interest: none.

Financial disclosures: none.

Funding/support: none.

ABSTRACT

Purpose: As a scoping review, evaluate the literature on the presentations, genetics, and therapies for neuroblastoma in adult patients.

Methods: We searched four databases for studies reporting adults with neuroblastoma. Cohort studies, case series, and case reports were synthesized qualitatively. Progression-free and overall survival were compared amongst cohort studies.

Results: Of 2287 unique records, 136 studies published in 141 articles were included. A total of 679 patients were included. On review of individual patient-level data, the adrenal gland and retroperitoneum were the most common primary site (47.3%). *MYCN* was rarely amplified: seven studies reported zero patients with *MYCN* amplified, two studies with a single patient, and one study with 3/7 patients. Adult patients appear to show a high frequency of somatic mutations, specifically *ALK* (42%) and *ATRX* (58%). Registry data of included studies showed 5-year overall survival to be 36.3% in adults aged ≥ 20 years.

Conclusion: Of nearly 700 cases of adult neuroblastoma published in the literature, the most common primary site is the adrenals or retroperitoneum. Relative to pediatric cases, adult cases demonstrate a considerable rate of somatic mutations such as *ALK* and *ATRX*. Registry data showed 5-year survival of 36%. Future studies evaluating targeted therapies in larger samples are needed.

Keywords: adult; cancer; genetics; kidney; neuroblastoma; survival

Introduction

Neuroblastoma is the most common extracranial solid tumor of childhood [1]. Unfortunately, over half of the children have metastatic disease on presentation. Neuroblastoma arises from cells of the neural crest that form the sympathetic ganglia and adrenal medulla. Tumors may occur anywhere along the sympathetic chain within the neck, thorax, retroperitoneum, or pelvis, or in the adrenal gland. The most common primary site includes the adrenal gland [1].

The occurrence of neuroblastoma is common in children however very rare in adults. Neuroblastoma is highly malignant, usually characterized by invasive growth and propensity for organ metastasis. The prognosis of adult patients with neuroblastoma is generally poor, and management is similar compared to the pediatric population.

The literature describing neuroblastoma is scattered with few cohort studies and countless case reports. Previously literature reviews include only a small fraction of the published cases, which may produce inaccurate summaries of the presentation, management, and prognosis. Therefore, we performed a scoping review with the aim to summarize all published reports of neuroblastoma in adults to inform clinicians of the available evidence of this heterogeneous disease.

Methods

We performed this scoping review in an attempt to summarize the available literature on the presentations, genetics, and therapies for neuroblastoma in adult patients. All patients were required to be ≥ 18 years old and have a pathologic diagnosis of neuroblastoma. Studies were excluded if the primary site was intracranial, cutaneous, or in the extremities. We also excluded cases of olfactory neuroblastoma (esthesioneuroblastoma) and if only ganglioneuroblastoma were reported. If studies included a combination of neuroblastoma and ganglioneuroblastoma,

we attempted to describe them separately. The following databases were searched for randomized controlled trials, observational studies, case series, and case reports: Medline, Embase, CINAHL, Web of Science. We used a combination of key word and medical subject headings such as ‘neuroblastoma’, ‘adult’, ‘geriatric’ from inception to 4 October 2022 (Medline strategy found in **Table A1**). There was no limitation on language or publication type (full text article or published conference abstract). The search was current until 7 October 2022. References of all included studies were reviewed to identify studies not captured in the search of databases. Research ethics board approval was not required as all data was published and publicly available.

Studies were screened by a single investigator and reviewed by a second investigator for inclusion. All studies were reported qualitatively and categorized according to study design (cohort study, case series, case report). We aimed to identify the total number of published unique cases of adult neuroblastoma. We also aimed to report patient ages, genders, and primary tumor sites. For cohort studies with sufficient data, we reported relevant genetic abnormalities (eg, *MYCN* amplification, *ALK* or *ATRX* mutations), management strategies, and progression-free and overall survival.

In addition, we described a case of neuroblastoma in a 24-year-old male patient managed at King Abdullah University Hospital (Riyadh, Saudi Arabia).

Results

Scoping Review

Characteristics of included studies (number, study design, age, sex)

Our initial literature search identified 2287 unique articles, of which 101 were included following full text review. Forty articles were identified through manual searching of references

of included articles. Accordingly, a total of 136 studies published in 141 articles were included in the scoping review (**Figure 1; Table 1**). **Table 2** details the genetic characteristics of neuroblastoma tumors of included studies. There were 6 cohort studies [2-7], 28 case series (**Table 3**) [11-38], and 102 case reports (**appendix Table A2**) [appendix references A1-A102]. The total number of adult neuroblastoma cases was 679. Five articles included overlapping patients [8-10,39,40], but were still included as they provided different information. For cohort studies, male proportion ranged between 44% [6] to 59% [2]. Rogowitz and colleagues [5] reported the largest study, documenting 216 adult neuroblastoma cases sourced from the Surveillance, Epidemiology, and End-Results (SEER) registry (1973-2010). Of these 216 patients, 35 (16%) were over 60 years old.

Primary site of tumor

In the largest series [5], the most common primary tumor site in patients aged 18-60 years was the central nervous system (39%, not reported if this included sympathetic ganglia), followed by the retroperitoneum (17%), and endocrine tissues including thymus (14%). Suzuki and colleagues reported a series of 44 patients from Italy, whereby the most common primary site was adrenal gland (26/44, 59%), followed by the abdominal paraspinal region (11/44, 25%), and mediastinum (5/44, 11%). Tang and colleagues [2] reported a series of 27 patients published in 1975, of which 15 patients had a primary site identified. The most common primary site was the mediastinum (7/15, 47%), followed by the retroperitoneum (5/15, 33%). Notably, 12 patients were unable to have a primary site identified given multiple sites involved. Sorrentino and colleagues [6] report a series of 16 patients from Italy, of which the adrenal gland (5/16, 31%), and abdomen (5/16, 31%) were the commonest primary site, followed by the pelvis (4/16, 27%), and thorax (2/16, 13%). When totaling patients included in included case series (n=131), the

most commonly reported site included the retroperitoneum not otherwise specified (n=38, 29%), followed by adrenal gland (n=24, 18%), pelvis (n=16, 12%), abdomen not otherwise specified (n=12, 9%), and mediastinum (n=11, 8%). If the primary sites of adrenal gland and retroperitoneum were combined, this site would total 47.3% (62/131) of patients.

Genetics

MYCN amplification occurred rarely in included studies of tested adult neuroblastoma patients: 0/22 patients in Kushner and colleagues [29], 0/5 in Podda and colleagues [31], 1/7 in Berbegall and colleagues [33], 0/40 in Suzuki and colleagues [7], 0/9 in Polishchuk and colleagues [3], 0/3 in Mazzocco and colleagues [36], and 1/10 in Sorrentino and colleagues [6]. The patient with *MYCN* amplification in Sorrentino and colleagues [6] was a 30-year-old female who presented with adrenal neuroblastoma with metastasis to the liver, and underwent radical tumor resection followed by cisplatin and three courses of peptichemio; she unfortunately died 10 months after initial presentation. Also in this study, one of five patients presented with 1p chromosome deletion. This was a 27-year-old male who presented with a neuroblastoma primary in the thorax with metastasis to bone. He underwent four courses of carboplatin and etoposide chemotherapy, and four courses of etoposide alone. He had a minor response but showed local and bone progression at 13 months and died 38 months following initial presentation.

Suzuki and colleagues reported a series of 44 patients [7], whereby somatic *ATRX* mutations or deletions were identified in 58% (15/26) of tested tumors and somatic *ALK* mutations were identified in 42% (10/16) of tested tumors. A proportion of this cohort underwent targeted exome sequencing at initial diagnosis (3/11) or relapse (8/11), which found a greater variety of mutations and total number of mutations at relapse. One patient had targeted exome sequencing performed both at initial diagnosis and at relapse: initial sequencing showed *CSF1R*

V32G mutation, relapse sequencing showed *CRLF2* deletion (Xp22.33) in addition to two other mutations (*FAT1* deletion (4q35.2), *NUP93* S654G).

Management and outcomes

The management of neuroblastoma varied according to extent of disease. Sorrentino and colleagues [6] reported a cohort of 16 patients: 4 patients were stage 1 or 2 and all underwent surgery except for 1 patient who also had chemotherapy; 6 patients were stage 3, and underwent various combinations of surgery, chemotherapy, and radiation, one of whom underwent surgery only but had bone relapse at 5 months; 6 patients were stage 4, all of whom received chemotherapy, 3 of whom also received surgery and 2 also received radiation [6]. Suzuki and colleagues [7] reported a series of 44 patients. Eight patients had locoregional neuroblastoma (five stage 1, one stage 2, five stage 3). According to International Neuroblastoma Response Criteria [A41]: all (5/5) patients with stage 1 disease had complete response to induction therapy; the only patient (1/1) with stage 2 disease had partial response; 80% (4/5) of stage 3 patients had complete response and 20% (1/5) had progressive disease; 21% (7/33) of stage 4 patients had complete response, 21% (7/33) had partial response, 36% (12/33) had no response, and 21% (7/33) had progressive disease [7].

Polishchuk and colleagues [3] reported a cohort of 16 adult patients with recurrent or refractory neuroblastoma managed with ¹³¹I-MIBG monotherapy. Five of these patients had ganglioneuroblastoma according to Shimada histologic classification [A42]. Treatment toxicities included thrombocytopenia in 69% (20/29; 4 patients grade 3, 16 patients grade 4) [A43]; neutropenia in 69% (20/29; 7 patients grade 3; 13 patients grade 4); need for granulocyte colony-stimulating factor support in 45% (13/29); need for autologous hematopoietic stem cell support in 25% (7/28); 13% (2/15) late-onset grade 2 thyroid abnormalities; 6% (1/16) had second

hematologic malignancies. Stiefel and colleagues [10] reported 14 adult neuroblastoma patients with somatic *ALK* mutations, 7 of whom were treated with FDA-approved *ALK* inhibitors. Adverse events included nausea/vomiting (86%, 6/7) and neurologic side effects (43%, 3/7; hallucinations [1/3], drowsiness [1/3], dizziness [1/3]). Median overall survival since *ALK* inhibitor treatment was 46.5 (range 17-74) months.

Discussion

This scoping review provides a detailed overview of adult neuroblastoma cases published in the literature from 1927 to present. We provide several examples of the heterogeneous nature of neuroblastoma, such as an 86-year-old presenting with a mediastinal mass. Our total count of 679 cases provides the literature with an update, as previous literature reviews quoted numbers ranging up to 200 cases. In addition, we detail patient survival according to modern risk stratification guidelines, and introduce recently published outcomes for targeted therapies.

In pediatrics, a considerable proportion of neuroblastoma cases have been associated with mutations of the *ALK* or *PHOX2B* gene [44,45], with somatic acquisitions *ALK* in up to 15% of cases [46]. *MYCN* amplification is reported in approximately 20% of pediatric neuroblastoma patients [44], and has been used as a biomarker to risk stratify patients for poorer prognosis. In our review, we identified seven studies detailing *MYCN* amplification testing, which reported 0 to 14% [7,33] of patients in respective series to test positive for *MYCN* amplification. In fact, the majority of studies reported 0% of patients harboring such a mutation [3,7,29,31,36]. However, Suzuki and colleagues [7] identified that somatic mutations in *ALK* occurred in 42% of patients, and *ATRX* mutations or deletions occurred in 58%. Accordingly, adult neuroblastoma cases may be more associated with progressive accumulation of somatic mutations [7,47], whereas childhood neuroblastoma cases may be more related to genetic susceptibility [44].

Studies of neuroblastoma in pediatrics versus adults show conflicting conclusions [4,7]. Conter and colleagues report in a conference abstract from University of Texas MD Anderson Cancer Center (1994-2012) of 118 adult and 112 pediatric patients no significant difference in survival for L1- ($p=0.4$), L2- ($p=0.54$), and M-stage ($p=0.73$) disease [4]. Esiashvilli and colleagues using SEER data reported a 5-year overall survival to be 84.6% in infants, 47.8% in children 1-9 years old, 46.2% in children 10-19 years old, and 36.3% in adults aged ≥ 20 years [8]. Navalke and colleagues also used SEER data (1975-2006) and showed a decreasing survival rate from patients aged 0-4 years to patients aged 15-19 years; there were too few patients in age groups 20-24 and 25-29 years to produce stable survival rates [9]. Accordingly, whether the true survival of adult neuroblastoma patients is inferior to pediatric patients or whether the overall differences are secondary to different disease stage distributions of each population is unknown; however, the consistent reduction in overall survival in the aforementioned studies suggests that adults generally fare worse.

The history of adult neuroblastoma has improved due to advancements of diagnostic and therapeutic techniques. Our literature search spanning nearly a century demonstrates the progression of initially relying solely on basic imaging techniques such as radiography [A4,A6] to routinely using magnetic resonance imaging and MIBG (iodine-123 labeled with metaiodobenzylguanidine) [34,A74]. The use of targeted therapies has also expanded, with studies reporting immunotherapy such as anti-GD2 [7], which was previously used exclusively in high risk children [48]. Future studies would contribute to the literature by providing outcomes of additional adult patients undergoing targeted therapies.

This scoping review has several strengths. First, our literature search was thorough given several methods of identifying articles: searching four databases, reviewing literature searches of

previous reviews, and reviewing the references list of all 142 included studies. This increases the confidence of readers that the included number of cases approaches the true number of published cases of adult neuroblastoma. This new count of 679 cases can be quoted in the future to provide clinicians a more accurate impression of the probabilities of encountering this rare entity.

Second, this review outlined the presentations, management strategies, and outcomes using cohort studies. Cohort studies generally provide a more balanced representation of the average adult neuroblastoma case. This contrasts with case reports, which have the aim of showcasing unusual presentations and outcomes. Using our scoping review, readers can appreciate both the general prognosis of adult neuroblastoma and how it varies according to risk stratification, as well as unusual cases we described in our table of case reports.

Nevertheless, this review is not without limitations. As thorough as literature searches may be, they are inherently imperfect, so some cases may have been missed. Second, we were unable to meta-analyze survival data to provide updated estimates. This was due to a lack of individual patient data, as well as suspected difficulty in ascertaining whether patient stages were similarly categorized. Third, there is a potential for overlapping cases if case reports were published both by themselves as well as included in neuroblastoma registries. We aimed to overcome this issue with case series and cohort studies by reviewing the data sources such as the SEER registry. Our total count included only the studies with the highest number of patients of registries to avoid patient duplication. Whereas we agree that some case reports may not be representative of the typical adult neuroblastoma patient, neuroblastoma is a disease regarded for its clinical heterogeneity, so awareness of such obscure cases broadens the differential of the clinician met with this enigmatic disease.

Conclusion

This scoping review documents the variable presentations, management strategies, and outcomes of 679 adult patients with neuroblastoma. Adult neuroblastoma is a rare disease entity that may present at any age, and appears to show considerable rates of somatic mutations. Future studies evaluating targeted therapies in larger samples are needed.

Acknowledgements: none.

Author contributions: BHA, BA, AA, AK conceived study idea. AK designed the study protocol. BHA and AK were involved in data extraction and synthesis of results. All authors were involved in manuscript writeup and critical review of the manuscript.

References

1. Pizzo PA, Poplack DG. Principles and practice of pediatric oncology. Lippincott Williams & Wilkins; 2015.
2. Tang CK, Hajdu SI. Neuroblastoma in adolescence and adulthood. *New York State Journal of Medicine*. 1975;75:1434-8.
3. Polishchuk AL, DuBois SG, Haas-Kogan D. Response, survival, and toxicity after iodine-131-metaiodobenzylguanidine therapy for neuroblastoma in preadolescents, adolescents, and adults. *Cancer*. 2011;117:4286-93.
4. Conter HJ, Gopalakrishnan V, Ravi V, Ater J, Patel S, Araujo DM. Are adult and pediatric neuroblastoma clinically different entities? *Journal of Clinical Oncology*;31(suppl):10049.
5. Rogowitz E, Babiker HM, Kanaan M, Millius RA, Ringenberg QS, Bishop M. Neuroblastoma of the elderly, an oncologist's nightmare: case presentation, literature review and SEER database analysis. *Experimental Hematology & Oncology*. 2014;3:1-7.
6. Sorrentino S, Gigliotti AR, Sementa AR, et al. Neuroblastoma in the adult: the Italian experience with 21 patients. *Journal of Pediatric Hematology/Oncology*. 2014;36:e499-505.
7. Suzuki M, Kushner BH, Kramer K, et al. Treatment and outcome of adult-onset neuroblastoma. *International journal of cancer*. 2018;143:1249-58.
8. Esiashvili N, Goodman M, Ward K, Marcus Jr RB, Johnstone PA. Neuroblastoma in adults: incidence and survival analysis based on SEER data. *Pediatric blood & cancer*. 2007;49:41-6.
9. Navalkele P, O'Dorisio MS, O'Dorisio TM, Zamba GK, Lynch CF. Incidence, survival, and prevalence of neuroendocrine tumors versus neuroblastoma in children and young adults: nine standard SEER registries, 1975–2006. *Pediatric blood & cancer*. 2011;56:50-7.
10. Stiefel J, Kushner BH, Basu EM, Roberts SS, Modak S. ALK inhibitors for treatment of adult-onset neuroblastoma. *Pediatric Blood and Cancer*. 2021;68(suppl5):S226
11. Fortner J, Nicastrì A, Murphy ML. Neuroblastoma: natural history and results of treating 133 cases. *Annals of Surgery*. 1968;167:132.
12. Mackay B, Luna MA, Butler JJ. Adult neuroblastoma. Electron microscopic observations in nine cases. *Cancer*. 1976;37:1334-51.
13. Dosik GM, Rodriguez V, Benjamin RS, Bodey GP. Neuroblastoma in the adult. Effective combination chemotherapy. *Cancer*. 1978;41:56-63.
14. Lopez R, Karakousis C, Rao U. Treatment of adult neuroblastoma. *Cancer*. 1980;45:840-4.
15. Feinstein RS, Gatewood OM, Fishman EK, Goldman SM, Siegelman SS. Computed tomography of adult neuroblastoma. *Journal of computer assisted tomography*. 1984;8:720-6.

16. Aleshire SL, Glick AD, Cruz VE, Bradley CA, Parl FF. Neuroblastoma in adults. Pathologic findings and clinical outcome. *Archives of pathology & laboratory medicine*. 1985;109:352-6.
17. Allan SG, Cornbleet MA, Carmichael J, Arnott SJ, Smyth JF. Adult neuroblastoma report of three cases and review of the literature. *Cancer*. 1986;57:2419-21.
18. Kaye JA, Warhol MJ, Kretschmar C, Landsberg L, Iii EF. Neuroblastoma in adults. Three case reports and a review of the literature. *Cancer*. 1986;58:1149-57.
19. Gohji K, Nakanishi T, Hara I, Hamami G, Kamidono S. Two cases of primary neuroblastoma of the kidney in adults. *The Journal of urology*. 1987;137:966-8.
20. Hoefnagel CA, Klumper A, Voute PA. Single photon emission tomography using meta-[131 I] iodobenzylguanidine in malignant pheochromocytoma and neuroblastoma. *Journal of Medical Imaging*. 1987;1:57-60.
21. Heyman S, Evans AE, D'Angio GJ. I-131 metaiodobenzylguanidine: diagnostic use in neuroblastoma patients in relapse. *Medical and pediatric oncology*. 1988;16:337-40.
22. Behr, A., Uiks, RRJ & Phillips JI. Neuroblastoma in adults-a report of 3 cases. *South African Medical Journal*. 1989;75:333-5.
23. Krikke AP, van der Jagt EJ. Adult neuroblastoma: a report of two cases. *Rofo: Fortschritte auf dem Gebiete der Rontgenstrahlen und der Nuklearmedizin*. 1989;150:138-41.
24. Prestidge BR, Donaldson SS. Treatment results among adults with childhood tumors: a 20-year experience. *International Journal of Radiation Oncology Biology Physics*. 1989;17:507-14.
25. Raina V, Kumar R, Sharma SK, Sarkar C, Bal CS. Neuroblastoma in thoracic region in adults: a report of five cases. *Annals of Saudi Medicine*. 1994;14:338-41.
26. Moody AM, Norman AR, Tait D. Paediatric tumours in the adult population: the experience of the Royal Marsden Hospital 1974–1990. *Medical and Pediatric Oncology*. 1996;26:153-9.
27. Argani P, Erlandson RA, Rosai J. Thymic neuroblastoma in adults: report of three cases with special emphasis on its association with the syndrome of inappropriate secretion of antidiuretic hormone. *American journal of clinical pathology*. 1997;108:537-43.
28. Cotterill SJ, Parker L, More L, Craft AW. Neuroblastoma: Changing incidence and survival in young people aged 0–24 years. A report from the North of England Young Persons' Malignant Disease Registry. *Medical and Pediatric Oncology*. 2001;36:231-4.
29. Kushner BH, Kramer K, LaQuaglia MP, Modak S, Cheung NK. Neuroblastoma in adolescents and adults: The Memorial Sloan-Kettering experience. *Medical and pediatric oncology*. 2003;41:508-15.
30. Tateishi U, Hasegawa T, Makimoto A, Moriyama N. Adult neuroblastoma: radiologic and clinicopathologic features. *Journal of computer assisted tomography*. 2003;27:321-6.
31. Podda MG, Luksch R, Polastri D, Gandola L, Piva L, Collini P, Cefalo G, Terenziani M, Ferrari A, Casanova M, Spreafico F. Neuroblastoma in patients over 12 years old: a 20-year experience at the Istituto Nazionale Tumori of Milan. *Tumori Journal*. 2010;96:684-9.
32. Sartelet H, Decaussin M, Latour M, et al. Neuroblastoma in adults: A case series. *Virchows Arch*. 2013;463:101-352
33. Berbegall AP, Villamón E, Tadeo I, et al. Neuroblastoma after childhood: prognostic relevance of segmental chromosome aberrations, ATRX protein status, and immune cell infiltration. *Neoplasia*. 2014;16:471-80.
34. Jrebi NY, Iqbal CW, Joliat GR, Sebo TJ, Farley DR. Review of our experience with neuroblastoma and ganglioneuroblastoma in adults. *World journal of surgery*. 2014;38:2871-4.
35. Mahkamova K, Harrison BJ. Functional Retroperitoneal Neuroblastoma and Ganglioneuroblastoma in Adults. *World Journal of Endocrine Surgery*. 2014;6:89-95.
36. Mazzocco K, Defferrari R, Sementa AR, et al. Genetic abnormalities in adolescents and young adults with neuroblastoma: a report from the Italian Neuroblastoma group. *Pediatric blood and cancer*. 2015;62:1725-32.
37. Duan K, Dickson BC, Marrano P, Thorner PS, Chung CT. Adult-onset neuroblastoma: Report of seven cases with molecular genetic characterization. *Genes, Chromosomes and Cancer*. 2020;59:240-8.

38. Jin QY, Du SB, Yuan XJ. Exploring the prognosis of neuroblastoma in adolescents and adults: a case series and literature review. *Neoplasma*. 2022.
39. Franks LM, Bollen A, Seeger RC, Stram DO, Matthay KK. Neuroblastoma in adults and adolescents: an indolent course with poor survival. *Cancer: Interdisciplinary International Journal of the American Cancer Society*. 1997;79:2028-35.
40. Hasegawa T, Hirose T, Ayala AG, et al. Adult neuroblastoma of the retroperitoneum and abdomen: clinicopathologic distinction from primitive neuroectodermal tumor. *The American journal of surgical pathology*. 2001;25:918-24.
41. Brodeur GM, Pritchard J, Berthold F, et al. Revisions of the international criteria for neuroblastoma diagnosis, staging, and response to treatment. *J Clin Oncol* 1993;11:1466–77.
42. Shimada H, Chatten J, Newton WA, et al. Histopathologic prognostic factors in neuroblastic tumors. *J Natl Cancer Inst*. 1984;73:405-416.
43. US Department of Health and Human Services. Common Terminology Criteria for Adverse Events (CTCAE). Version 3. Washington, DC: National Institute of Health, National Cancer Institute; 2006.
44. Maris JM. Recent advances in neuroblastoma. *New England Journal of Medicine*. 2010 Jun 10;362:2202-11.
45. Mossé YP, Laudenslager M, Longo L, Cole KA, Wood A, Attiyeh EF, Laquaglia MJ, Sennett R, Lynch JE, Perri P, Laureys G. Identification of ALK as a major familial neuroblastoma predisposition gene. *Nature*. 2008;455:930-5.
46. George RE, Sanda T, Hanna M, Fröhling S, Ii WL, Zhang J, Ahn Y, Zhou W, London WB, McGrady P, Xue L. Activating mutations in ALK provide a therapeutic target in neuroblastoma. *Nature*. 2008;455:975-8.
47. Kennedy SR, Loeb LA, Herr AJ. Somatic mutations in aging, cancer and neurodegeneration. *Mech Ageing Dev* 2012;133:118–26.
48. Ploessl C, Pan A, Maples KT, Lowe DK. Dinutuximab: an anti-GD2 monoclonal antibody for high-risk neuroblastoma. *Annals of Pharmacotherapy*. 2016;50:416-22.

Figure legends

Figure 1: PRISMA flow diagram

Table legends

Table 1: Characteristics of 6 included cohort studies

Table 2: Genetic characteristics of neuroblastoma tumors in included studies

Table 3: Characteristics of 28 included case series

Table A1: Sample search strategy using medline database

Table A2: Characteristics of 102 included case reports

Identification of studies via databases

Identification

Records identified from:
Embase + Medline (n=1768)
CINAHL (n=184)
Web of Science (n=1216)

Records removed *before screening*:
Duplicate records removed (n=881)

Screening

Records screened (n=2287)

Records excluded (n=2156)

Reports sought for retrieval
(n=131)

Reports not retrieved
(n=0)

Reports assessed for eligibility
(n = 131)

Reports obtained from
reference review of
included studies (n=40)

Reports excluded: (n=30)
Wrong patient population (n=23)
Duplicate study (n=7)

Included

Studies included in review
(n=136)
Reports of included studies
(n=141)

Table 1: Characteristics of 6 included cohort studies

Study	Country and Data Source	Patient cohort	Primary Tumor Site	Management	Outcome
Tang 1975 [2]	United States: Memorial Hospital 1948-1972	N=27. 16 males (59%).	Mediastinum (7), retroperitoneum (5), pelvis (1), extremity (1), neck (1), multiple sites (12).	Surgical resection in 26 patients. Combined chemotherapy and radiation in 18. Radiation only in 7.	67% 1-year survival. 28% 2-year survival. 12% lived over 5 years.
Polishchuk 2011 [3]	United States: University California San Francisco (recurrent or refractory cases)	N=16. 8 males (50%)	Abdomen (10), thorax (3), thoracoabdominal (1), pelvic (1), unknown (1).	¹³¹ I-MIBG monotherapy	Median survival 36 months, estimated 3-year survival 43.2%.
Conter 2013 [4]	United States: MD Anderson Cancer Center 1994-2012	N=118. Mean age 47 (range 18-82) years.	NA	Specific management not available.	Adults similar survival to pediatrics (L1 p=0.40, L2 p=0.54, M p=0.73). L1 disease actuarial survival 94% at 3 years, 90% at 5 years, 69% at 10 years. L2 disease estimated survival 83% at 3 years, 73% at 5 years, 41% at 10 years. M actuarial survival 68% at 1 year, 33% at 2 years, 13% at 5 years. INRG stage was prognostic (p<0.001).
Rogowitz 2014 [5]	United States National Cancer Surveillance, Epidemiology, and End-Results registry (SEER)	N=216. 35 patients >60 years (19%). 107 male (50%)	Most common site CNS (71) in patients 18-60 years. Most common site soft tissue including heart (21) in patients >60 years.	Specific management not available.	In patients >60 years, 5 year disease-specific survival was 40%. In patients with mediastinal, tracheal, and other resp neuroblastomas, 5 year disease-specific survival was 44% for 9 adult patients.
Sorrentino 2014 [6]	Italy: Istituto Giannina Gaslini	N=16. Mean age 26 (range 18-37). 7 male (44%).	Abdomen (5), adrenal (5), thorax (2), pelvis (4). 6 patients metastatic disease on diagnosis.	Stage 1/2 (4): All surgery, of which 1 also chemotherapy. Stage 3 (6): Various combinations of surgery (3), chemotherapy (4), radiation (2). Stage 4 (6): All chemotherapy, 3 also surgery and 2 radiation.	Stage 1/2: 2 died at 25 and 48 months. 1 alive with disease at 107 months. 1 remission at 130 months. Stage 3: 5 patients died of disease at mean 70 months; 1 patient remission at 142 months. Stage 4: 6 patients died of disease at mean 46 months.

Suzuki 2018 [7]	United States: Memorial Sloan Kettering Cancer Center 1979-2015	N=44. Median age 25 (range 18-71). 25 males (57%).	Adrenal gland (26), abdominal paraspinal region (11), mediastinum (5); bone or bone marrow (2).	<p>Locoregional neuroblastoma (11): All surgery, and 8 additionally received chemotherapy.</p> <p>Stage 4 (33): 27 high-dose chemotherapy. 1 patient crizotinib for ALK-mutation. Surgery was attempted in 11 patients.</p> <p>Of 7 stage 4 patients with complete response, 5 received autologous stem cell transplant, 1 received autologous stem cell transplant and anti-GD2 immunotherapy with 3F8. 7 with incomplete response received anti-GD2 immunotherapy: 4 with m3F8 + GM-CSF, 2 with 3F8 alone, 1 with 3F8.</p>	<p>Locoregional neuroblastoma: 10-year progression-free 35.4% and overall survival was 61.4%</p> <p>Stage 4: 7 (21%) achieved complete response (CR) after induction chemotherapy and/or surgery. 5-year progression-free survival 9.7%. 10- year overall survival 19%. Patients who achieved complete response after induction had superior survival.</p>
-----------------	---	--	---	--	--

*Three studies not detailed given overlapping patients [8-10]

Table 2: Genetic abnormalities of neuroblastoma tumors in cohort studies and case series

Study	Genetic testing	Genetic outcomes
Polishchuk 2011 [3]	NA.	0/9 MYCN amplifications.
Sorrentino 2014 [6]	NA.	1/10 tested patients MYCN amplification. 1/5 tested patients 1p chromosome deletion.
Suzuki 2018 [7]	FISH; whole exome sequencing; DNA sequencing for <i>ALK</i> ; targeted exome sequencing using Foundation One or MSK-IMPACT	0/40 tumor <i>MYCN</i> amplification. 15/26 (58%) <i>ATRX</i> mutations or deletions. 10/16 (42%) <i>ALK</i> mutations. Various somatic mutations in 11 patients with targeted exome sequencing including <i>BRCA1</i> , <i>IKZF2</i> , <i>NF2</i> , <i>MUTYH</i> , <i>TP53</i> , <i>CSF1R</i> , <i>CRLF2</i> , <i>FAT1</i> , <i>NUP93</i> , <i>BLM</i> , <i>PTPRD</i> , <i>RAC1</i> , <i>EZH2</i> , <i>SMARCA4</i> , <i>ERBB3</i> , <i>NOTCH3</i> , <i>PAK1</i> , <i>RBM10</i> , <i>BCL6</i> , <i>KIT</i> , <i>RASA1</i> .
Kushner 2003 [29]	NA.	0/22 tumor <i>MYCN</i> amplification.
Podda 2010 [31]	NA.	0/5 tumor <i>MYCN</i> amplification.
Berbegall 2014 [33]	aSNP, MLPA, and FISH in tumors with adequate DNA quality; FISH used for MYCN	1/7 tumor heterogenous <i>MYCN</i> amplification. 1/6 17q gain. 1/6 11q deletion. 2/4 focal segmental chromosome aberration (+5p; +20p). 2/4 copy-neutral loss of heterozygosity (4q, 11p, 11p; 12q, 19p; 6p).
Mazzocco 2015 [36]	Double color FISH on interphase nuclei using <i>MYCN/LAF</i> probe for chromosomes 2p24 and 2q11; <i>ALK</i> assessed in exons 20 to 28 of the <i>ALK</i> gene; <i>ATRX</i> assessed in five exons (10, 16, 19, 30, 32) and five portions of exon 9 and flanking splice junctions of <i>ATRX</i>	0/3 tumor <i>MYCN</i> amplification. 0/3 <i>ALK</i> mutations. 1/3 <i>ATRX</i> mutations (c.6572A>C p.D2191A). 1 patient 1p deletion, 9p deletion. 1 patient 2p gain. 1 patient 1p deletion, 9p deletion, 11q deletion, 17q gain.
Sartelet 2013 [32]	FISH for <i>MYCN</i> amplification.	0/4 <i>MYCN</i> amplification.
Mahkamova 2014 [35]	NA.	0/1 <i>MYCN</i> amplification.
Duan 2019 [37]	FISH used for MYCN copy number, chromosome 1p loss and 17q gain;	MYCN: 4/7 no amplification, 1/7 focal amplification, 1/7 diffuse amplification, 1/7 gain.

	detecting alternate lengthening of telomeres (<i>ALT</i>); and for detecting telomerase gene <i>TERT</i> rearrangement.	2/6 <i>ALK</i> mutation using IHC, of which no break detected using FISH. 4/5 <i>ATRX</i> mutation using IHC, of which 1 positive in 3.8% of cells using FISH. 1/5 1p deletion. 4/4 17q gain. 1/5 <i>TERT</i> rearrangement (diffuse amplification of 200x) detected using FISH.
--	---	--

NA: not available; M: male; F: female; mo: month. MSK-IMPACT: Memorial Sloan Kettering Integrated Mutation Profiling of Actionable Cancer Targets; FISH: fluorescence in situ hybridization; IHC: immunohistochemistry;

Table 3: Characteristics of 28 included case series

Study	Age	Sex	Location of Primary
Fortner 1968 [11]	7 patients (2 male), locations not reported		
Mackay 1976 [12]	54	Female	Right parotid gland
	72	Male	Left paravertebral area, buttocks, right inguinal area
	60	Male	Left subdigastric region, posterior cervical area
	35	Male	Extending throughout retroperitoneum, left ankle
	25	Male	Right liver lobe
	37	Female	Retroperitoneum involving pancreas
Dosik 1978 [13]	37	Female	Liver, retroperitoneum involving pancreas
	20	Male	Posterior abdominal mass extending into pelvis
	23	Male	Posterior abdominal mass extending into pelvis
Lopez 1979 [14]	33	Female	Left adrenal extending to left upper quadrant of abdomen
	52	Female	Mediastinum, metastatic to abdomen, retroperitoneum, brain
Feinstein 1984 [15]	40	Female	Posterior mediastinum
	20	Male	Pelvis with another mass right chest above diaphragm
	19	Male	Right adrenal involving retrocaval lymph nodes and encasing IVC
	20	Male	Left hemithorax extending laterally to soft tissues, T3-T5 vertebral bodies, acetabulum, femur
	28	Male	Left para-aortic region and left renal vein
Aleshire 1985 [16]	75	Male	Soft tissue of neck
	57	Female	Parotid
	32	Male	Iliac fossa
	26	Male	Soft tissue of neck
	65	Female	Soft tissue of neck
	29	Female	Adrenal gland
	53	Male	Retroperitoneum
	68	Male	Soft tissue of neck
Allan 1986 [17]	26	Female	Right pelvis, previous ovarian cystectomy
	34	Female	Right thorax infiltrating intervertebral foramina
	25	Male	Pelvis encasing IVC
Kaye 1986 [18]	26	Female	Right lower neck, anterior mediastinum, bones
	38	Female	Right adrenal mass metastatic to bone
	20	Female	Pelvis involving uterus, and omentum, mesentery, liver
Gohji 1987 [19]	35	Male	Lower pole of right kidney
	29	Male	Right kidney infiltrating IVC wall and psoas muscle
Hoefnagel 1987 [20]	18	NA	NA
	22	Male	Abdomen, mediastinum
	32	NA	NA
	52	NA	NA
Heyman 1988 [21]	22	Male	NA
	30	Male	NA
Behr 1989 [22]	47	Female	Below right scapula
	26	Female	Retroperitoneum infiltrating bladder and right superior mediastinum
Krikke 1989 [23]	25	Female	Left adrenal gland metastatic to abdominal viscera and bones
	23	Female	Right paravertebral area T12-L2
Prestridge 1989 [24]	18	Male	Hemi-abdomen, pelvis
	19	Female	Hemi-abdomen
	19	Male	Hemi-abdomen
	25	Female	Hemi-abdomen
	34	Female	Hemi-abdomen, pelvis
Raina 1994 [25]	18	Female	Left infra-axillary region involving 4 th rib
	21	Male	Anterosuperior mediastinum
	18	Male	Right upper lung lobe, mediastinum
	35	Female	Left paravertebral region extending into right paravertebral region, left hemithorax

Moody 1996 [26]	33	Male	Retroperitoneum with multiple liver metastases
	24	Male	Metastasis to bone, primary location unknown
	24	Female	Adrenal gland metastasis to bone
Argani 1997 [27]	80	Male	Thymus
	71	Male	Anterior mediastinum
Cotterill 2001 [28]	3 patients, unknown genders and neuroblastoma locations.		
Kushner 2003 [29]	18	Male	Right adrenal, cortical bone, liver, thoracic
	23	Female	Left adrenal, bone marrow, cortical bone
	23	Male	Bone marrow, cortical bone
	29	Male	Left adrenal, bone marrow, cortical bone, liver
	31	Female	Left adrenal, thoracic, left supraclavicular
	20	Male	Retroperitoneum
	24	Female	Retroperitoneum, bone marrow, cortical bone, thoracic
	19	Female	Retroperitoneum, bone marrow, cortical bone, thoracic
	24	Female	Pelvis, bone marrow, cortical bone
	31	Female	Retroperitoneum, cortical bone, thoracic
	32	Male	Retroperitoneum, bone marrow, cortical bone
	33	Male	Left adrenal, bone marrow, cortical bone
	41	Male	Retroperitoneum, pelvis, sphenoid
	18	Female	Thoracoabdominal
	18	Male	Thoracoabdominal, left supraclavicular
	21	Female	Retroperitoneum, bone marrow, thoracic, supraclavicular
	24	Female	Bone marrow, cortical bone
	27	Female	Thoracic, left supraclavicular
Tateishi 2003 [30]	6 patients, mean age 49, tumor sites include retroperitoneum (n=2), pelvis (n=2), anterior mediastinum (n=1)		
Podda 2010 [31]	69	Male	Adrenal gland, no metastasis
	18	Female	Thorax, no metastasis
	30	Male	Thorax, no metastasis
	36	Male	Thorax, no metastasis
	32	Male	Adrenal gland, no metastasis
	19	Male	Thorax, metastasis to lymph node
	18	Male	Abdomen (not adrenal gland), metastasis to lymph node and bone marrow
	20	Female	Abdomen (not adrenal gland), metastasis to lymph node
	21	Female	Abdomen (adrenal gland), metastasis to skeleton
	25	Female	Abdomen (not adrenal gland), metastasis to bone marrow, skeleton, liver
	19	Male	Abdomen (not adrenal gland), metastasis to lymph node
	23	Female	Adrenal gland, metastasis to lymph node
	Sartelet 2013 [32]	4 patients (2 male); mean age 34 (range 22-53) years; 3 tumors in retroperitoneum, 1 tumor in adrenal gland.	
Berbegall 2014 [33]	18	Male	Pelvis
	19	Female	Adrenal gland
	21	Male	Pelvis
	24	Male	Adrenal gland
	36	Female	Adrenal gland
	39	Female	Adrenal gland
	60	Female	NA
Jrebi 2014 [34]	25	Female	Pelvis
	19	Female	Mediastinum
	22	Male	Mediastinum
	31	Male	Retroperitoneum
	50	Female	Groin, pelvis
	28	Female	Pelvis
	20	Male	Abdomen
66	Female	Abdomen, chest	

	51	Female	Adrenal gland
	21	Female	Pelvis
Mahkamova 2014 [35]	36	Male	Retroperitoneum: encasing aorta and IVC, infiltrating posterior abdominal wall
	21	Male	Retroperitoneum: extending from upper pole of right kidney to aortic bifurcation
Mazzocco 2015 [36]	22	Male	NA
	18	Male	NA
	19	Male	NA
Duan 2019 [37]	9 patients (3 males); median age 24 (range 18-30) years; tumor locations include retroperitoneum (n=8), neck (n=1)		
Jin 2022 [38]	20	Female	Retroperitoneum, metastasis to bone and bone marrow
	25	Female	Adrenal gland

*Two studies not detailed given overlapping patients [39, 40]

Appendix: Neuroblastoma in Adults: A Systematic Review of Presentations, Genetics and Therapies

Table A1: Sample search strategy using medline database

Database: Embase <1974 to 2022 October 04>, OVID Medline Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE(R) Daily and Ovid MEDLINE(R) 1946 to Present	
Search Strategy:	

1	neuroblastoma.ti,ab. (86935)
2	adult*.ti,ab. (3411525)
3	geriatrics/ (63893)
4	geriatric*.ti,ab. (142867)
5	octogenarian.ti,ab. (3901)
6	nonagenarian.ti,ab. (1547)
7	2 or 3 or 4 or 5 or 6 (3565808)
8	1 and 7 (4349)
9	limit 8 to english language (4111)
10	limit 9 to humans (3221)
11	limit 10 to "all adult (19 plus years)" [Limit not valid in Embase; records were retained] (2531)
12	limit 11 to animal studies [Limit not valid in Ovid MEDLINE(R),Ovid MEDLINE(R) Daily Update,Ovid MEDLINE(R) PubMed not MEDLINE,Ovid MEDLINE(R) In-Process,Ovid MEDLINE(R) Publisher; records were retained] (763)
13	11 not 12 (1768)

Table A2: Characteristics of 102 included case reports

Study	Age	Sex	Location of Neuroblastoma
Capaldi 1927 [A1]	44	Female	Sympathetic chain left C7-T1
Frost 1936 [A2]	38	Male	Right lung apex
Balas 1959 [A3]	31	Female	Lung parenchyma
Buthker 1964 [A4]	67	Female	Anterior mediastinum
Eklof 1967 [A5]	27	Male	Chest
Griff 1968 [A6]	21	Male	Posterior mediastinum
Hutchinson 1968 [A7]	51	Male	Anterior mediastinum
Hale 1970 [A8]	26	Female	Right adrenal gland
Baumgartner 1975 [A9]	56	Female	Left kidney
Chadhury 1975 [A10]	29	Female	Retroperitoneum
Attal 1975 [A11]	25	Male	Left suprarenal region
Rowe 1979 [A12]	29	Female	Right adrenal gland
Reynolds 1981 [A13]	20	Female	Retrouterine pouch
Berkel 1982 [A14]	36	Female	Left adrenal gland metastatic to spine
Blijham 1982 [A15]	19	Male	Right sub-hepatic space extending to iliac crests
Raje 1982 [A16]	20	Male	Pelvis with extension to abdominal viscera, parietal peritoneum, and diaphragm
Holgerson 1983 [A17]	29	NA	Thoracic region metastatic to lumbosacral spine
Block 1984 [A18]	22	Female	Right ovary metastatic to retroperitoneum, mediastinum, and supraclavicular nodes
Grubb 1984 [A19]	58	Female	Right submandibular salivary gland
Yashiro 1984 [A20]	26	Female	Left adrenal gland
Mackay 1987 [A21]	23	Male	Right scapula, possibly femur
Mir 1987 [A22]	29	Male	Left paraspinal region
Hoover 1988 [A23]	57	Male	Posterior mediastinum
Onishi 1988 [A24]	34	Female	Right adrenal gland with tumor thrombus extending into right renal vein, IVC, RA
Roberts 1988 [A25]	22	Male	Left ilium, suprasellar region, right anterior clinoid process
Bullock 1989 [A26]	35	Male	Left orbit
Cheah 1989 [A27]	24	Male	Left retroperitoneum involving suprarenal region and left liver lobe
Urios 1989 [A28]	30	Male	Right paraspinal region at T6-7 level
McDonnell 1990 [A29]	31	Male	Entire peritoneal cavity
Rossitti 1991 [A30]	39	Male	Left side of lumbosacral plexus
Chaloupka 1992 [A31]	31	Male	Anterior and posterior vertebral elements of T10 with encasement of spinal cord
Salter 1995 [A32]	80	Female	Anterior mediastinum
Varona 1996 [A33]	18	Female	Right paravertebral region
Cowan 1997 [A34]	28	Male	Left kidney, left pelvis, left upper quadrant of abdomen
Wester 1997 [A35]	35	Female	Thoracolumbar region, metastasis to suprasellar and pontine cisterns
Hirokawa 1998 [A36]	56	Female	Retroperitoneum extending into pelvic cavity
Custodio 1999 [A37]	32	Female	Right adrenal gland region
Yamamoto 2000 [A38]	21	Male	Right thoracic paravertebral region
Kawakami 2001 [A39]	43	Female	Retroperitoneum
Genc 2003 [A40]	30	Female	Left adrenal gland
Shimizu 2003 [A41]	33	Male	Posterior mediastinum, 8-9 ribs, sympathetic chain
McLean 2004 [A42]	39	Male	Right adrenal gland
Sucker 2004 [A43]	21	NA	Left upper lung, left neck
Genc 2005 [A44]	52	Male	Left adrenal gland
Schalk 2005 [A45]	51	Male	Right adrenal gland
Ayadi-Kaddour 2007 [A46]	61	Female	Left posterior mediastinum
Loeser 2007 [A47]	39	Female	Cervix, with several recurrences afterwards
Hadi 2008 [A48]	25	Female	Left adrenal gland, infiltrating kidney and pancreas
Refaat 2008 [A49]	25	Female	Right adrenal gland
Jarius 2009 [A50]	23	Female	Para-aortic region adjacent to left adrenal gland
Ogawa 2009 [A51]	60	Male	Anterior mediastinum
Zerrweck-Lopez 2009 [A52]	31	Male	Bilateral retroperitoneal mass displacing aorta and kindeys

Jost 2010 [A53]	67	Female	Left adrenal gland, metastasis to paravertebral soft tissues, humerus, femur, pelvis
Panovska-Stavridis 2010 [A54]	38	Male	Thoracic para-spinal region
Singh 2010 [A55]	38	Female	Bilateral adnexa, lymph nodes
Soares 2010 [A56]	23	Female	T3-T5 paravertebral region extending into spinal canal, right parietal lobe
Then 2010 [A57]	55	Male	Possibly mediastinum; vertebral bodies T6-T10
Venat-Bouvet 2010 [A58]	25	Male	Retroperitoneum
Kajbafzadeh 2011 [A59]	52	Male	Rectovesical pouch
Ohtaki 2011 [A60]	64	Male	Superior aspect of mediastinum
Marin Oyaga 2011 [A61]	29	Male	Adrenal gland
Selcukbiricik 2011 [A62]	53	Male	Right neural foramina extending to spinal duct at T4-6 levels
Shi 2011 [A63]	42	Female	Anterior mediastinum
Abdou 2012 [A64]	22	Female	Pancreatic head
Bayrak 2012 [A65]	37	Female	Lower pole of right kidney
Pellegrino 2012 [A66]	79	Female	Anterior mediastinum
Shobha 2012 [A67]	26	Female	Right ovary
Ueda 2012 [A68]	65	Female	Anterior mediastinum
Gupta 2013 [A69]	47	Male	Right adrenal region with infiltration into upper pole of right kidney
Koumarianou 2013 [A70]	30	Male	Right adrenal gland
Smith 2013 [A71]	39	Female	Paraspinal, with metastasis to axial and appendicular skeleton, liver, lymph nodes.
Tiu 2013 [A72]	79	Female	Lower pole of left kidney.
Imvrios 2014 [A73]	26	Female	Right lumbar region
Skoura 2014 [A74]	33	Male	Right adrenal gland
Stevens 2014 [A75]	29	Female	Right pelvis with lytic lesion of left iliac crest
Thomas 2014 [A76]	79	Female	Adrenal glands bilaterally
Rogowitz 2014 [A77]	86	Male	Mediastinum
Febrero 2015 [A78]	39	Male	Left infrarenal space
Liu 2015 [A79]	22	Female	Upper pole of left kidney
Wiesel 2015 [A80]	62	Male	Anterior mediastinum
Daboussi 2016 [A81]	21	Male	Posterior mediastinum
Kaluzna-Oleksy 2016 [A82]	21	Female	Lumbar spine
Kurokawa 2016 [A83]	62	Male	Left adrenal gland region
Rais 2016 [A84]	40	Male	Right adrenal gland
He 2016 [A85]	28	Female	Cervix
Jain 2017 [A86]	53	Female	Right renal fossa
Malik 2017 [A87]	23	Male	Left lumbar region
Huang 2018 [A88]	41	Female	Right kidney with retrocaval lymphadenopathy and right renal vein thrombus
Ma 2018 [A89]	24	Male	Posterior mediastinum
Manoharan 2018 [A90]	32	Female	Left adrenal gland
Wu 2018 [A91]	20	Female	Left retroperitoneal space
Martinez-Ciarpaglini 2019 [A92]	40	Male	Left inguinal region
Naeem 2019 [A93]	29	Male	Sacrum S1-S4, with mass surrounding sacral vertebrae extending into central canal
Ramsignh 2019 [A94]	22	Female	Left adrenal gland
Tan 2019 [A95]	46	Female	Left dorsal pleura
Yanik 2019 [A96]	40	Male	Anterior mediastinum
Zhang 2019 [A97]	75	Female	Left adrenal gland
Bukhari 2020 [A98]	31	Female	T12 vertebra
Findakly 2020 [A99]	63	Male	NA
Nawata 2021 [A100]	63	Male	Left kidney with enlargement of intra-abdominal lymph nodes
Xu 2021 [A101]	40	Female	Left adrenal gland
Gomez 2022 [A102]	24	Female	Adrenal gland, retroperitoneum

References of included case reports:

- A1. Capaldi, B. Zwei Falle von Sympathicoblastom. Frankfurt Z. Path 1927;35:83-100.
A2. Frost TF, Wolpaw SE. Intra-thoracic Sympathoblastoma. Am. J. Cancer 1936;26:483.

- A3. Balas A. Neurogerier Lungentumor (Sympathoblasom). *Der Chirurg* 1959;30:325.
- A4. Büthker W, Feltkamp-Vroom T, Groen AS, Wieberdink J. Sympathicoblastoma in the anterior mediastinum. *Diseases of the Chest*. 1964;46:531-6.
- A5. Eklof OL, Gooding CA. Intrathoracic neuroblastoma. *American Journal of Roentgenology*. 1967;100:202-7.
- A6. Griff LC, Griff RE. Neuroblastoma emphasis on the mediastinal neuroblastoma. *American Journal of Roentgenology*. 1968;103:19-24.
- A7. Hutchinson III JE, Nash AD, McCord CW. Neuroblastoma of the anterior mediastinum in an adult. *The Journal of Thoracic and Cardiovascular Surgery*. 1968;56:147-52.
- A8. Hale JE. A case of neuroblastoma in an adult with raised catecholamines. *Journal of British Surgery*. 1970;57:551-3.
- A9. Baumgartner GC, Gaeta J, Wajzman Z, Merrin C. Neuroblastoma presenting as renal cell carcinoma in an adult. *Urology*. 1975;6:376-8.
- A10. Chadhury A. Abdominal sympathoblastoma in an adult with lung abscess. *J Indian Med Assoc* 1975;64:240-242.
- A11. Attal HC, Grover S, De'Souza E. Neuroblastoma presenting as disseminated osseous metastasis (a case report). *The Journal of the Association of Physicians of India*. 1975;23:531-4.
- A12. Rowe PH, Oram JJ, Scott GW. Neuroblastoma in adults. *Postgraduate Medical Journal*. 1979;55:579-80.
- A13. Reynolds CP, Smith RG, Frenkel EP. The
Cancer. 1981;48:2088-94.
- A14. Berkel MV, Rijn HJ, Lammers HA. Adrenal neuroblastoma sympathicum in a 36-year-old woman. *European Journal of Nuclear Medicine*. 1982;7:278-9.
- A15. Blijham GH, Barlogie B, Richman S, Latreille J, Vellekoop L. Medulloblastoma and neuroblastoma in adults. *Netherlands Journal of Medicine*. 1982;25(4):94-9.
- A16. Raje D, Shah DJ, Lofters WS. Neuroblastoma in an adult. *West Indian medical journal*. 1982;149-52.
- A17. Holgersen LO, Santulli TV, Schullinger JN, Berdon WE. Neuroblastoma with intraspinal (dumbbell) extension. *Journal of Pediatric Surgery*. 1983;18:406-11.
- A18.
term survival. Case report and review of the literature. *Cancer*. 1984;54:590-5.
- A19. Grubb BP, Thant M. Neuroblastoma in an adult. *Southern Medical Journal*. 1984;77:1180-2.
- A20. Yashiro N, Yoshida H, Kuwajima S, Nomura T, Akima M. Adult adrenal neuroblastoma with extension into inferior vena cava. *Radiation Medicine*. 1984;2:234-6.
- A21. Mackay B, Ordóñez NG. Adult neuroblastoma of bone: A case report. *Ultrastructural Pathology*. 1987;11:455-64.
- A22. Mir R, Phillips SL, Schwartz G, Mathur R, Khan A, Kahn LB. Metastatic neuroblastoma after 52 years of dormancy. *Cancer*. 1987;60:2510-4.
- A23. Hoover EL, Hsu HK, Dressler C, Fani K, Webb H, Ketosugbo A, Kharma B. Neuroblastoma: a rare primary intrathoracic neurogenic tumor in adults. *Texas Heart Institute Journal*. 1988;15:107.
- A24. Onishi S, Shibahara N, Yasuta H, et al. Adult neuroblastoma of the adrenal with intraatrial tumor thrombus: report of a case. *Hinyokika kyo. Acta Urologica Japonica*. 1988;34:1195-9.
- A25. Roberts JL, Magid D, Miller NR, Kuhajda FP, Fishman EK. Case report 493. *Skeletal Radiol*. 1988;17:375-377.
- A26. Bullock JD, Goldberg SH, Rakes SM, Felder DS, Connelly PJ. Primary orbital neuroblastoma. *Archives of Ophthalmology*. 1989;107:1031-3.
- A27. Cheah PL, Jayalakshmi P, Jeyamalar R, Kuperan P. Adult neuroblastoma: a case report. *The Malaysian Journal of Pathology*. 1989;11:69-71.
- A28. Urios JI, Garceran LR, Rosell TV. Neuroblastoma in an adult causing spinal cord compression: report of a case and review of the literature. *Spinal Cord*. 1989;27:394-401.

- A29. McDonnell III CH, McLeod M, Baker ME. Primary peritoneal neuroblastoma: computed tomography findings. *Clinical imaging*. 1990;14(1):41-3.
- A30. Rossitti SL, Matos PS, Balbo RJ, Zuiani AR, Thomaz IR. Adult peripheral neuroblastoma. *Arquivos de Neuro-Psiquiatria*. 1991;49:357-61.
- A31. Chaloupka JC, Castillo M. Primary neuroblastoma of a vertebra: an unusual location for a variant of primitive neuroectodermal tumor of bone. *AJR. American journal of roentgenology*. 1992;159:1130.
- A32. Salter JE, Gibson D, Ordóñez NG, Mackay B. Neuroblastoma of the anterior mediastinum in an 80-year-old woman. *Ultrastructural Pathology*. 1995;19:305-10.
- A33. Varona L, Ruiz J. Adult-onset neuroblastoma with intraspinal (dumbbell) extension. *Neurologia (Barcelona, Spain)*. 1996;11:88-9.
- A34. Cowan JM, Dayal Y, Schwaitzberg S, Tischler AS. Cytogenetic and immunohistochemical analysis of an adult anaplastic neuroblastoma. *The American journal of surgical pathology*. 1997;21:957-63.
- A35. Wester S, Mork S, Smievoll AI. Neuroblastoma and venous sinus thrombosis in an adult patient. *British journal of neurosurgery*. 1997;11:241-4.
- A36. Hirokawa S, Yamashita I, Kuroki Y, et al. Neuroblastoma in an adult with a high serum level of carbohydrate antigen, CA125: report of a case. *Surgery today*. 1998;28:349-54.
- A37. Custodio CM, Semelka RC, Balci NC, Mitchell KM, Freeman JA. Adrenal neuroblastoma in an adult with tumor thrombus in the inferior vena cava. *Journal of Magnetic Resonance Imaging*. 1999;9:621-3.
- A38. Yamamoto R, Tada H, Kishi A, Tojo T, Asada H, Inoue T. Neuroblastoma resection in an adult with a 10-year history of chest-mass shadow. *The Japanese Journal of Thoracic and Cardiovascular Surgery*. 2000;48:809-11.
- A39. Kawakami M, Koda M, Matsunaga N, et al. Adult-type neuroblastoma originated in retroperitoneum beginning with obstructive jaundice. *Clinical imaging*. 2001;25:284-7.
- A40. Genc FA, Aksoy M, Kapran Y, et al. Adrenal neuroblastoma in an adult: Report of a case. *Surgery today*. 2003;33:879-81.
- A41. Shimizu M, Shimizu T, Adachi T, Nishiuchi M, Kanzaki M, Onuki T. Long-term survival in adult mediastinal neuroblastoma. *The Japanese Journal of Thoracic and Cardiovascular Surgery*. 2003;51:326-9.
- A42. McLean TW, Iskandar SS, Shimada H, Hall MC. Neuroblastoma in an adult. *Urology*. 2004;64:1232.
- A43. Sucker C, Dölken G, Lorenz G, Fiedler W, Stockschräder M. Thoracic neuroblastoma in a young adult. *Deutsche Medizinische Wochenschrift (1946)*. 2004;129:249-52.
- A44. Genc H, Hacıyanlı M, Hacıyanlı SG, Gelal F, Uçarsoy AA, Dag F. An adult adrenal neuroblastoma: a case report. *Acta Chirurgica Belgica*. 2005;105:673-6.
- A45. Schalk E, Mohren M, Koenigsmann M, Buhtz P, Franke A, Jentsch-Ullrich K. Metastatic adrenal neuroblastoma in an adult. *Oncology Research and Treatment*. 2005;28:353-5.
- A46. Ayadi-Kaddour A, Chaabouni S, Mlika M, Kilani T, Hassine RB, El Mezni F. Neuroblastoma of the posterior mediastinum in a 61-year-old woman. *Respiratory Medicine Extra*. 2007;3(3):114-6.
- A47. Loeser A, Gerharz EW, Riedmiller H. Recurrent pelvic neuroblastoma in an adult patient. *Gynecologic oncology*. 2007;106:257-8.
- A48. Hadi AM, Khalaf IJ, Abbass YI. An adult adrenal neuroblastoma presented as a right shoulder pain: A case report. *The Medical Journal of Tikrit University*. 2008;2(142).
- A49. Refaat MM, Idriss SZ, Blaszkowsky LS. Case report: an unusual case of adrenal neuroblastoma in pregnancy. *The Oncologist*. 2008;13:152-6.
- A50. Jarius S, Arnold S, Linke R, et al. Long term survival in anti-Hu associated adult neuroblastoma. *Journal of the neurological sciences*. 2009;284:205-8.

- A51. Ogawa F, Amano H, Iyoda A, Satoh Y. Thymic neuroblastoma with the syndrome of inappropriate secretion of antidiuretic hormone. *Interactive Cardiovascular and Thoracic Surgery*. 2009;9:903-5.
- A52. Zerrweck-López C, Quijano-Orvañanos F, Montañez-Ramírez H, et al. Neuroblastoma in the adult. Case report. *Cirugia y Cirujanos*. 2009;77:397-401.
- A53. Jost G, Frank S, Fischer N, Taub E, Mariani L. An epidural neuroblastoma causing spinal cord compression in a 67-year-old woman. *Rare tumors*. 2010;2(2):74-7.
- A54. Panovska-Stavridis I, Ivanovski M, Hadzi-Pecova L, et al. A case report of aggressive adult neuroblastoma mimicking acute leukemia with fulminant course and fatal outcome. *Prilozi*. 2010;31:349-59.
- A55. Singh M, Arora P, Singh S, et al. Adult neuroblastoma of the ovary: a rare tumor with review of literature. *Journal of Cancer Research and Therapeutics*. 2010;6:367.
- A56. Soares PM, Quirino Filho S, Pereira DE, Bonan PF, Martelli-Júnior H. Neuroblastoma in an adult: Case report. *Rev Med Chile*. 2010;138:1131-4.
- A57. Then C, Ebelt K, Langer A, Mayr D, Schmidmaier R, Oduncu F. Neuroblastoma in a 55-year-old patient: a case report. *Case Reports in Oncology*. 2010;3:458-62.
- A58. Venat-Bouvet L, Le Brun-Ly V, Martin J, Gasnier O, Falkowsky S, Tubiana-Mathieu N. Long-term survival in adult neuroblastoma with multiple recurrences. *Case Reports in Oncology*. 2010;3:45-8.
- A59. Kajbafzadeh AM, Sadeghi Z, Esfahani SA, Fisher C, Judson I. Adult pelvic neuroblastoma: long-term survival and review of the literature/Eriskinlerde pelvis nöroblastom: Uzun dönem sagkalim ve literatürün gözden geçirilmesi. *Turkish Journal of Urology*. 2011;37:167.
- A60. Ohtaki Y, Ishii G, Hasegawa T, Nagai K. Adult neuroblastoma arising in the superior mediastinum. *Interactive Cardiovascular and Thoracic Surgery*. 2011;13:220-2.
- A61. Marín Oyaga VA, Cuenca Cuenca JI, Fernández López R, Borrego Dorado I, Vázquez Albertino R. 123I-
journal of nuclear medicine and molecular imaging. 2011;38:1586.
- A62. Selcukbiricik F, Tural D, Esatoglu N, Kocak S, Mandel NM. A very rare adult case with neuroblastoma. *Case Reports in Oncology*. 2011;4:481-6.
- A63. Shi Jr X, Wang X, Yi C, Zhang X, Chen Z. Imaging features of adult neuroblastoma in F-18 FDG PET/CT. *Clinical Nuclear Medicine*. 2011;36:1058-60.
- A64. Abdou AG, Asaad NY, Elkased A, Said H, Dawoud M. Adult pancreatic neuroblastoma, an unusual site and fatal outcome. *Pathology & Oncology Research*. 2012;18:239-43.
- A65. Bayrak O, Seckiner I, Erturhan S, Aydin A, Yagci F. Adult intrarenal neuroblastoma presenting as renal cell carcinoma. *Canadian Urological Association Journal*. 2012;6:E144.
- A66. Pellegrino M, Gianotti L, Cassibba S, Brizio R, Terzi A, Borretta G. Neuroblastoma in the elderly and SIADH: case report and review of the literature. *Case reports in medicine*. 2012;2012.
- A67. Shobha K, Bafna UD, Devi UK, Amirtham U. Primary neuroblastoma of ovary in an adult: A case report and review of literature. *Clinical Cancer Investigation Journal*. 2012 Apr 1;1(2):92.
- A68. Ueda Y, Omasa M, Taki T, et al. Thymic neuroblastoma within a thymic cyst in an adult. *Case reports in oncology*. 2012;5:459-63.
- A69. Gupta P, Maiti A, Aich RK, Deb AR. Adrenal neuroblastoma in an adult. *Journal of Cancer Research and Therapeutics*. 2013;9:96.
- A70. Koumariou A, Oikonomopoulou P, Baka M, et al. Implications of the incidental finding of a MYCN amplified adrenal tumor: a case report and update of a pediatric disease diagnosed in adults. *Case Reports in Oncological Medicine*. 2013;2013.
- A71. Smith L, Minter S, O'Brien P, Kravetka JM, Medina AM, Lazarchick J. Neuroblastoma in an adult: case presentation and literature review. *Annals of Clinical & Laboratory Science*. 2013;43:81-4.
- A72. Tiu A, Latif E, Yau L, Reynolds R, Millar EK, Aslan P. Primary renal neuroblastoma in adults. *Urology*. 2013;82:11-3.

- A73. Imvrios G, Filis D, Fouzas I, et al. A Rare Adult Case with Neuroblastoma. *Journal of Tumor*. 2015;2:278-81.
- A74. Skoura E, Oikonomopoulos G, Vasileiou S, Kyprianou D, Koumakis G, Datseris IE. (18) F-FDG-PET/CT,(123) I-MIBG and (99m) Tc-MDP whole-body scans, in detecting recurrence of an adult adrenal neuroblastoma. *Hellenic Journal of Nuclear Medicine*. 2014;17:58-61.
- A75. Stevens PL, Johnson DB, Thompson MA, Keedy VL, Frangoul HA, Snyder KM. Adult neuroblastoma complicated by increased intracranial pressure: a case report and review of the literature. *Case Reports in Oncological Medicine*. 2014;2014.
- A76. Thomas J, Freedman L, Kumar SC, Hupart KH. Bilateral Adrenal Neuroblastoma in a Geriatric Patient with Elevated Catecholamine Metabolites. *Endocrine Reviews* 2014;35:0770.
- A77. Rogowitz E, Babiker HM, Kanaan M, Millius RA, Ringenberg QS, Bishop M. Neuroblastoma of analysis. *Experimental Hematology & Oncology*. 2014;3:1-7.
- A78. Febrero B, Ríos A, Rodríguez JM, Flores B, Parrilla P. Retroperitoneal neuroblastoma in the adult. An uncommon entity. *Cirugia espanola*. 2015;93:e147-8.
- A79. Liu QL, Liang QL, Ou WT, Li ZY, Zhang XN. Surgical resection and post-operative radiotherapy in an adult renal neuroblastoma patient with multiple bone and joint metastases: A case report. *Oncology Letters*. 2015;9:2591-4.
- A80. Wiesel O, Bhattacharyya S, Vaitkevicius H, Prasad S, McNamee C. Ataxia induced by a thymic neuroblastoma in the elderly patient. *World Journal of Surgical Oncology*. 2015;13:1-6.
- A81. Daboussi S, Moetemri Z, M'hamdi S, Aichaouia C, Khadraoui M, Cheikh R. Mediastinal neuroblastoma: Unusual localization with late revelation. *Revue de Pneumologie Clinique*. 2016;72:353-8.
- A82. -Oleksy M, Wachowiak- - Migaj E. Unresectable heart neuroblastoma in an adult: a natural follow-up. *Pol Arch Med Wewn*. 2016;126:365-6.
- A83. Kurokawa S, Mizuno K, Nakane A, et al. Adrenal neuroblastoma in an adult: effect of radiotherapy on local progression after surgical removal. *Case Reports in Urology*. 2016;2657632.
- A84. Rais F, Benhmidou N, Rais G, et al. Neuroblastoma in an adult: a case report of a rare entity with a summary review. *International Journal of Surgery and Medicine*. 2018;4:53.
- A85. He Y, Yao M, Zhang X, Sun P, Gao H. A very rare adult case of cervical neuroblastoma. *International Journal of Clinical and Experimental Pathology*. 2017;10:8746.
- A86. Jain TK, Ashwathanarayana AG, Basher RK. A rare adult renal neuroblastoma better imaged by 18 68 F-FDG than by Ga-dotanoc in the PET/CT scan. *Hellenic Journal of Nuclear Medicine*. 2017;20:100-1.
- A87. Malik D, Jois A, Singh H, Bora GS, Basher RK, Mittal BR. Metastatic neuroblastoma in adult patient, presenting as a super scan on 68Ga-DOTANOC PET/CT imaging. *Clin Nucl Med* 2017;42:697-9.
- A88. Huang MD, Hsu LS, Chuang HC, et al. Adult renal neuroblastoma: a case report and literature review. *Medicine*. 2018;97;1-5.
- A89. Ma X, Yang Y, Wang Z, et al. Surgical removal of adult recurrent neuroblastoma located in the posterior mediastinum and retroperitoneum: a case report. *Medicine*. 2018;97;50.
- A90. Manoharan J, Wiese D, Maurer E, et al. Left side adrenal mass mimicking a pheochromocytoma. *Langenbeck's Archives of Surgery*. 2018;403:903-916.
- A91. Wu XL, Dai YJ, Sun GY, et al. Adult neuroblastoma in the retroperitoneum: a case report. *Medicine*. 2018;97;51;1-4.
- A92. Martinez-Ciarpaglini C, Machado I, Yoshida A, et al. Extra-adrenal adult neuroblastoma with aberrant germ cell marker expression: maturation after chemotherapy as an important clue to a challenging diagnosis. *International Journal of Surgical Pathology*. 2019;27:568-73.
- A93. Naeem M, Maluf H, Baker JC, Jennings JW. Primary osseous sacral neuroblastoma in an adult. *Skeletal Radiology*. 2019;48:985-8.

- A94. Ramsingh J, Casey H, Watson C. Adult neuroblastoma: a rare diagnosis of an adrenal mass. *BMJ Case Reports CP*. 2019;12:e228730.
- A95. Tan YB, Li JF, Li WS, Yang RL. Primary thoracic neuroblastoma in an adult: a rare case report. *Medicine*. 2019;98;1-4.
- A96. Yanik F, Karamustafaoglu YA, Yoruk Y. A rare mediastinal occurrence of neuroblastoma in an adult: case report. *Sao Paulo Medical Journal*. 2018;137:104-6.
- A97. Zhang H, Feng Z. Adrenal neuroblastoma in an elderly adult: a case report and review of the literature. *Journal of Medical Case Reports*. 2019;13:1-4.
- A98. Bukhari N, Harfouch B, Alotaibi MS, Al-Harbi H, Chamdine O. Immediate response to chemotherapy in an adult neuroblastoma patient presenting with cord compression. *Case Reports in Neurological Medicine*. 2020;6401497.
- A99. Findakly D, Nguyen E, Wang J. Clinical features and outcomes of neuroblastomas in adults. *Journal of Clinical Oncology*. 2020;38;15(suppl):e14546.
- A100. Nawata A, Izumi R, Harada K, et al. An elderly-onset neuroblastoma concomitant with minimal change nephrotic syndrome: the first autopsy case report. *CEN Case Reports*. 2021;10:414-21.
- A101. Xu S, Zhang W, Che B, et al. Adult adrenal neuroblastoma: A case report. *Molecular and Clinical Oncology*. 2021;15:1-4.
- A102. Gómez GE, Urbano MA, Martínez V. Adrenal Neuroblastoma Producing Catecholamines Diagnosed in Adults: Case Report. *Case Reports in Oncology*. 2022;15:682-6.