

High acceptability of newborn screening for sickle cell disease among post-natal mothers in western Kenya

John Orimbo¹ * Shehu Shagari Awandu¹, Faith Muhonja², Patrick Owili³, Dickens Omondi¹

¹Department of Biomedical and Nursing Sciences, School of Health Sciences, Jaramogi Oginga Odinga University of Science and Technology, Bondo, Kenya

²Department of Public Health, Amref International University

³African Population and Health Research Centre

*Corresponding Author: John Orimbo. Email: jguda006@gmail.com

Abstract

Background: Sickle cell disease is a genetically inherited blood disorder that manifests early in life with resultant significant health complications. Globally, nearly three quarters of all affected babies are in sub-Saharan Africa. Early identification of babies with sickle cell disease through newborn screening followed by early linkage to care is recommended. However, the program has not been widely adopted in the sub-Saharan Africa. Evidence on acceptability of newborn screening to scale up newborn screening program is scarce. This study assessed factors associated with acceptability of newborn screening among mothers of newborns delivered at Homa Bay county teaching and referral hospital, western Kenya.

Methods: This study employed a cross-sectional design among postnatal mothers at Homa Bay county teaching and referral hospital with 399 postnatal mothers enrolled into the study. A semi-structured questionnaire was used for data collection. Maternal sociodemographic characteristics, knowledge, and perception were assessed. Babies were also screened for sickle cell disease using Sickle SCAN point-of-care test. The acceptability was calculated as percentage of mothers accepting to have their babies screened. Data were analyzed using logistic regression to explore factors associated with acceptability of newborn screening for sickle cell disease.

Results: Ninety-four percent of mothers accepted newborn screening for sickle cell disease. Mother's age and occupation were significantly associated with acceptability of newborn screening for sickle cell disease. Younger mothers (OR=3.01;95%CI=1.16-7.83; $p=0.024$) and being a student (OR= 6.18; 95%CI= 1.18-32.22; $p=0.031$) were significant at bivariate regression analysis. Only being a student (aOR= 25.02; 95% CI=1.29-484.51; $p= 0.033$) was significant at multivariate logistic regression analysis.

Conclusion: The acceptability of newborn screening for sickle cell disease is high in the county. The Homabay county and the Kenyan ministry of health should implement routine newborn screening for sickle cell disease in all level 2-6 hospitals.

Key words: Sickle Cell Disease Acceptability Newborn Screening Postnatal mothers

45 **Introduction**

46
47 Sickle cell disease (SCD) is a major genetically inherited blood disorder, caused by autosomal
48 recessive inheritance, that manifests early in life with resultant significant health complications.
49 Globally, nearly three quarters of all sickle cell disease-affected babies are in sub-Saharan Africa
50 (Uyoga et al., 2019). In 2011, the World Health Organization (WHO) recommended an increased
51 awareness, improved access to health services, and provision of technical support for prevention
52 and management of sickle cell disease (Makani et al., 2013). Programs such as newborn screening
53 for sickle cell disease have shown great success towards preventing and managing the disease
54 (Nnodu et al., 2018). However, only a few countries in Africa have piloted the screening program
55 and none has scaled up the implementation (Inusa et al., 2015; Ohene-Frempong et al., 2008;
56 Tshilolo et al., 2009). Successful implementation of newborn screening for sickle cell disease
57 followed by early initiation of penicillin prophylaxis, pneumococcal vaccines, and hydroxyurea
58 treatment are likely to reduce the disease burden in low- and middle-income countries (LMICs).

59
60 Nonetheless, limited data on the prevalence of sickle cell disease at birth in Africa makes it difficult
61 to quantify the magnitude of the disease burden. Additionally, approximately 90% of the children
62 with sickle cell disease die before their fifth birthday, partly because of late diagnosis (Katamea et
63 al., 2022). This can be solved through adequate implementation of newborn screening for sickle
64 cell disease. This implementation however, heavily depends on its acceptability (Sekhon et al.,
65 2017) and socio-cultural settings (Bediako & Neblett, 2011), which vary from one community or
66 region to another. Unacceptability of the screening program for sickle cell disease by the parents
67 or caregivers or authorities in Africa can greatly hinder effective implementation of the program
68 (Inusa et al., 2015). There is need to assess the acceptability of the program before its
69 implementation which may not only promote public participation but also community ownership
70 of the intervention.

71
72 The prevalence of sickle cell disease varies on the malaria endemicity patterns. The disease burden
73 in Kenya is quite high especially around the lake regions in western as well as the Coastal regions
74 (Ministry of Health, 2020). The prevalence of sickle cell disease around western Kenya is about
75 4.5% for sickle cell disease, and 18% for sickle cell trait (Wanjiku et al., 2019).

76

77 Homa Bay county is yet to implement newborn screening for sickle cell disease. This is despite
78 the high prevalence of sickle cell disease among infants who were accessing maternal and child
79 health services in the county, which was found to be at 9.6% (Wanjiku et al., 2018). Routine
80 newborn screening for sickle cell disease is recommended in regions where sickle cell disease
81 prevalence at birth is 0.05% or more (DeBaun & Galadanci, 2022). There is however, paucity of
82 data on the acceptability of newborn screening in the county. Acceptability is important for
83 successful implementation of the intervention program (Sekhon et al., 2017). Additionally,
84 extrapolating data from another county may not be feasible owing to the variable nature of
85 acceptability (Marsh et al., 2011).

86
87 Here, we assessed the acceptability of newborn screening for sickle cell disease and its associated
88 factors in Homa Bay County, Western Kenya. This is important in promoting the establishment of
89 a systematic newborn screening program for sickle cell disease in Homa Bay County, and
90 subsequently a successful implementation of the intervention (Sekhon et al., 2017). The results of
91 this study are critical to the ministry of health at the county-level for planning of the sickle cell
92 disease screening program in the county.

93 94 **Methods**

95 96 **Study design, study population and study setting**

97 This study employed a cross-sectional design at the postnatal wards in the Homa Bay county
98 teaching and referral hospital, Homa Bay County, western Kenya. Homa Bay county teaching and
99 referral hospital is the county referral hospital with a wider catchment across the eight sub counties
100 in Homa Bay county. The study involved postnatal mothers in the postnatal wards between 14th
101 April 2023 and 15th June 2023. The site has a sickle cell disease clinic however, it lacks newborn
102 screening for sickle cell disease implementation.

103 104 **Sample size and sampling technique**

105 Cochran's formula for calculating the sample when the population is infinite was used. A
106 maximum variability of 50% was used to estimate the maximum sample size required since the
107 acceptability of newborn screening for sickle cell disease was unknown in Homa Bay County. The
108 final sample size was 403 mothers, after adjusting for 5% non-response. Consecutive sampling

109 technique, where all participants meeting eligibility criteria were enrolled until required sample is
110 achieved, was used to select the participants. All postnatal mothers available at the postnatal ward
111 during the study period, and willing to take part in our study and met the inclusion criteria were
112 selected and enrolled. The inclusion criteria were: The postnatal mother giving an informed
113 consent or assent to participate in the study and whose outcome of delivery was an alive infant.
114 Postnatal mothers who were medically unstable to respond to the questionnaire or needed urgent
115 critical care were excluded from participating in the study.

116

117 **Data collection tools**

118

119 Kobo collect tool was used to collect the data using a semi-structured questionnaire with five sub-
120 sections. The first section collected demographic information: Maternal age, marital status,
121 occupation, religion, education level and parity. The second section contained questions on the
122 knowledge (awareness) of the mother about sickle cell disease including ever having heard of or
123 being aware of sickle cell disease, being aware of their haemoglobin status, knowledge about sickle
124 cell disease being a blood disorder, knowledge on transmission of sickle cell disease, when it can
125 be diagnosed and knowledge on the existence of newborn screening for sickle cell disease
126 programs. The third section contained the questions on the perceptions of the newly delivered
127 mothers. The perception was measured by asking question on their opinion on to when screening
128 for sickle cell disease should be done, ethical issues and the perceived effectiveness of the newborn
129 screening for sickle cell disease. The perception was considered good when in line with the
130 national government newborn screening policy. The fourth section of the questionnaire explored
131 the possible intervening variables mainly, the cultural issues around newborn screening and the
132 role of the male partner in making the decision as to whether to accept newborn screening for
133 sickle cell disease. The fifth section collected data on their intent/willingness to have their
134 newborns screened and those who were willing had their newborns screened using sickle SCAN
135 kit (BioMedomics Inc, NC, USA). The sickle SCAN test kit is capable of identifying Hemoglobin
136 A, S and C variants in blood samples. The sex of the newborns was also captured here.

137

138 **Enrollment and Data collection**

139 A total of 399 participants were enrolled into the study, representing 99% response rate. Eligible
140 participants provided written informed consent for mothers who were 18 years and above. Minor

141 mothers also were provided with a signed assent. The participants were provided with a copy of
142 the signed consent form and another copy retained by the investigator. After consenting, a semi-
143 structured questionnaire was administered to each participant by the research assistants. The
144 research assistants comprised of 1 clinical officer/physician assistant, 2 nurses, 1 laboratory
145 technician and 2 counsellors. The principal investigator trained the research assistants prior to the
146 start of the study and supervised data collection. The mothers were finally questioned on their
147 intent/willingness to have their newborns screened. Willing participants had their babies screened
148 for sickle cell disease using the Sickle SCAN® point-of-care test following manufacturer's
149 instructions (BioMedomics Inc.,NC, United States). The results were interpreted as sickle cell trait
150 with 3 lines visible at the control, HbA and HbS; sickle cell disease when two lines appeared at
151 the control and HbS or no haemoglobinopathy when two lines at the control and HbA. The
152 laboratory technician conducted the screening test by the bedside following the manufacturer's
153 instructions. The newborns' heels were pricked using 1.2mm lancet pricker, and 2 drops of blood
154 collected directly into the device. The results were ready in 3-5 minutes. The counsellors offered
155 psychological counselling to all the mothers pretest and to mothers whose babies had sickle cell
156 trait or sickle cell disease post-test. The mothers whose newborns screened positive for sickle cell
157 disease were referred to the sickle cell disease comprehensive clinic and also received
158 psychological counselling from the study team.

159

160 **Statistical analyses**

161 The data were entered into excel for cleaning before analysis using STATA version 16.0
162 (StataCorp, 2019). The level of acceptability of newborn screening for sickle cell disease was
163 analyzed using descriptive statistics only. Even though acceptability is a multidimensional
164 construct, in this study it was defined as the willingness of the mother to accept newborn screening
165 for sickle cell disease. The percentage acceptability, calculated as the percentage of mothers who
166 agreed to have their newborns screened for sickle cell disease, was calculated.

167 The factors associated with acceptability of newborn screening for sickle cell disease were
168 analyzed using both descriptive statistics and inferential statistics. Descriptively, proportions of
169 acceptability among various factors were calculated and represented in a table. Inferentially, Chi-
170 square and Fisher's exact test (where at least a cell has a count less than 5) were used to assess the
171 association between the sociodemographic, maternal knowledge and maternal perception

172 characteristics and the acceptability of newborn screening for sickle cell disease. The factors that
173 had statistically significant association with acceptability were further analyzed using logistic
174 regression analysis to establish the point estimate and the magnitude of association. After the
175 unadjusted analysis, statistically significant variables, were fit into a multivariate logistics
176 regressions model using forward stepwise regression approach. Both unadjusted and adjusted odds
177 ratio (OR) and 95% confidence intervals (95% CI) were reported. The results of the newborn
178 screening were used to calculate the prevalence of sickle cell disease and sickle cell trait among
179 the newborns.

180 **Results**

181 **Summary of participant characteristics**

182 The mean age of the mothers was 26.2 years (SD 6.07) with 25-34 years being modal age-group,
183 comprising 44.1% of the respondents. Majority (300/399; 75%) of respondents were married;
184 59.7% (238/399) had one or 2 children; 41.4% (165/399) had attained secondary level education
185 while, 70% were engaged in either informal or no employment. Only 2 mothers (0.5%) reported
186 to have had a previous child with Sickle cell disease. Majority (371/399; 93%) of the respondents
187 had ever heard about sickle cell disease; only 4% knew their sickle cell disease status with only
188 2% having ever been screened for sickle cell disease before; over 90%(362/399) knew the cause
189 of sickle cell disease; only 7%(28/399) of respondents believed that sickle cell disease was as a
190 result of God's will with less than 2%(5/399) believing sickle cell disease was due to God's
191 punishment or evil spirits and majority (386/399;96.7%) belived the best timing for sickle cell
192 disease testing was during the postnatal period (Table 1).

Table 1: Characteristics of Postnatal Mothers (N=399) at Homay Bay County Teaching and Referral Hospital, western Kenya

Variable Characteristic	Frequency	Percentage
Age group		
11-24	175	43.9
25-34	176	44.1
≥35 years	48	12.0
Education		
Tertiary	120	30.1
Secondary	165	41.4
Primary	112	28.1
No Education	2	0.5


Marital Status

Married	300	75.2
Widowed	5	1.3
Separated	21	5.3
Never married	73	18.3
Occupation		
Student	72	18.1
Formal employment	40	10.0
Informal employment	226	56.6
No employment	61	15.3
Religion		
Roman Catholic	73	18.3
Protestant	206	51.6
Pentecostal	55	13.8
Muslim	6	1.5
Others	59	14.8
Parity		
One child	141	35.3
Two children	97	24.3
Three children	80	20.1
Four children and above	81	20.3
Child with sickle cell disease		
Yes	2	0.5
No	397	99.5
Death of child under 2 years		
Yes	29	7.3
No	370	92.7

193
194
195
196
197
198
199
200

Level of acceptability of newborn screening for sickle cell disease among postnatal mothers at Homabay County Teaching and Referral Hospital, western Kenya

Out of the total 399 mothers interviewed, 375/399 (94.0%) agreed to have their newborns screened for sickle cell disease using the sickle scan point of care kit (Fig. 1). The level of acceptability was therefore 94.0%.



201
202 *Figure 1: Pie chart showing acceptability of New Born Screening for Sickle Cell Disease among*
203 *postnatal mothers in Homa Bay County Teaching and Referral Hospital, western Kenya*
204

205 **Association between sociodemographic characteristics and acceptability of newborn**
206 **screening for Sickle cell disease among postnatal mothers at Homabay County Teaching and**
207 **Referral Hospital, western Kenya**

208 Chi-square method was used to assess association between the sociodemographic characteristics
209 of the respondents and their acceptability of newborn screening for sickle cell disease. Maternal
210 occupation ($p= 0.026$) and maternal age ($p= 0.023$) were the only factors that had statistically
211 significant association with the acceptability of newborn screening for sickle cell disease.
212 Respondents who were students and those younger than 25 years were more likely to accept
213 newborn screening for sickle cell disease. The student as an occupation category comprised
214 mothers who were school going and therefore not in any other occupation. Characteristically in
215 this study, about 90% (65/72) were younger than 25 years. Majority of the students 91%(66/72)
216 had post primary education, with the modal group in secondary schools. The odds of accepting
217 newborn screening for sickle cell disease among student postnatal mothers was higher than those
218 in formal employment (OR=6.18; 95%CI=1.18-32.22; $p= 0.031$). Mothers younger than 25 years
219 of age had higher odds of accepting newborn screening for sickle cell disease than older mothers

220 (OR=3.01;95%CI=1.16-7.83; $p= 0.024$). No significant association was found between
 221 acceptability of newborn screening for sickle cell disease and other sociodemographic
 222 characteristics. (Table 2).

223

224

Table 2: Association between acceptability of newborn screening for sickle cell disease and Sociodemographic characteristics of postnatal mothers at Homabay County Teaching and Referral Hospital, western Kenya

Characteristic	NBS YES	NBS NO	X ² value	df	p-value
Marital Status*					
Married	279(93.00)	21(7.00)			
Widowed	5(100.00)	0(0.00)		3	0.181
Separated	19(90.48)	2(9.52)			
Never married	72(98.63)	1(1.37)			
Education (highest completed)					
Tertiary	110(91.67)	10(8.33)			
Secondary	158(95.76)	7(4.24)	2.06	1	0.357
Primary	107(93.86)	7(6.14)			
Religion*					
Roman Catholic	69(94.52)	4(5.48)			
Protestant	192(93.20)	14(6.80)		4	0.341
Pentecostal	51(92.73)	4(7.23)			
Muslim	5(83.33)	1(16.67)			
Others	58(98.31)	1(1.69)			
Parity					
One child	135(95.74)	6(4.26)			
Two children	92(94.85)	5(5.15)		3	0.369
Three children	72(90.00)	8(10.00)			
Four children and above	76(93.83)	5(6.17)			
Age group*					
11-24	169(96.57)	7(3.43)			
25-34	159(90.34)	17(9.66)		2	0.023
≥35	47(97.52)	1(2.48)			
Previous Child with sickle cell disease*					
Yes	2(100.00)	0(0.00)		1	1.000
No	373(93.95)	24(6.05)			
Occupation*					
Student	70(97.22)	2(2.78)			

Formal employment	34(85.00)	6(15.00)	3	0.026
Informal employment	211(93.36)	15(6.64)		
Unemployed	60(98.36)	1(1.64)		
Previous Death of child under 2 years*				
Yes	28(96.55)	1(3.45)	1	0.546
No	347(93.78)	23(6.22)		

225 *Fisher's exact method used where cell contain value <5, p-value<0.05

226

227 **Association between maternal knowledge and acceptability of newborn screening for sickle**
 228 **cell disease among postnatal mothers at Homabay County Teaching and Referral Hospital,**
 229 **western Kenya**

230 Fisher's exact method was used to assess the association between maternal knowledge, measured
 231 by finding what they know about sickle cell disease as shown in table 3 below, and the acceptability
 232 of newborn screening for sickle cell disease since the criteria for chi-square was not met. Most of
 233 the cells had values less than 5. No statistically significant association was found between maternal
 234 knowledge dimensions and acceptability of newborn screening for sickle cell disease. Most of the
 235 mothers had ever heard about sickle cell disease and knew it is a non-contagious blood disorder.
 236 However, only few mothers had ever been screened for sickle cell disease nor knew their sickle
 237 cell disease status. Most had good knowledge of the inheritance patterns of sickle cell disease and
 238 nearly two thirds had never heard about newborn screening for sickle cell disease before (Table
 239 3).

240

Table 3: Association between acceptability of newborn screening for sickle cell disease and knowledge of postnatal mothers at Homabay County Teaching and Referral Hospital, western Kenya

Characteristics	NBS YES	NBS NO	P-value
Ever heard about SCD			1.000
Yes	348(93.80)	23 (6.20)	
No	27(96.40)	1(3.60)	
Knowledge of SCD status			0.612
Yes	16(100.00)	0(0.00)	
No	359(93.70)	24(6.30)	
Ever screened for SCD			1.000
Yes	8(100.00)	0(0.00)	
No	367(93.90)	24(6.10)	
SCD is blood disorder			1.000
Yes	340(93.10)	22(6.10)	
No	35(94.60)	2(5.40)	
SCD is contagious			1.000
Yes	6(100.00)	0(0.00)	

No	369(93.90)	24(6.10)	
Maternal inheritance only			1.000
Yes	2(100.00)	0(0.00)	
No	373(94.00)	24(6.00)	
Paternal inheritance only			1.000
Yes	1(100.00)	0(0.00)	
No	374(94.00)	24(6.00)	
Heard about SCD screening			0.079
Yes	129(97.00)	4(3.00)	
No	246(92.30)	20(7.70)	

241 *Fisher's exact method used, p-value <0.05*

242

243 **Association between acceptability of newborn screening for sickle cell disease and perception**
 244 **of postnatal mothers at Homabay County Teaching and Referral Hospital, western Kenya**

245

246 Fisher's exact method was used to assess the association between maternal perception and the
 247 acceptability of newborn screening for sickle cell disease since the criteria for chi-square test was
 248 not met. Most of the cells had values less than 5. No statistically significant association was found
 249 between maternal perceptions and acceptability of newborn screening for sickle cell disease.
 250 Postnatal mothers however had good perception about sickle cell disease. Most of the postnatal
 251 mothers had good perception of what causes sickle cell disease, with most believing that the best
 252 time to screen for sickle cell disease is during the newborn period (0-6 weeks). Majority of the
 253 postnatal mothers believed newborn screening for sickle cell disease was morally right and would
 254 be useful in controlling the disease (Table 4).

255

Table 4: Association between acceptability of newborn screening for sickle cell disease and perception of postnatal mothers at Homabay County Teaching and Referral Hospital, western Kenya

Characteristics	NBS YES	NBS NO	p-value
SCD occur due to God's will			1.000
True	27(96.43)	1(3.57)	
False	348(93.80)	23(6.20)	
SCD occur due to God's punishment			0.268
True	4(80.00)	1(20.00)	
False	371(94.16)	23(5.84)	
SCD occur due to evil spirit			0.268
True	4(80.00)	1(20.00)	

False	371(94.16)	23(5.84)	
Best timing for SCD Screening			0.180
Prenatally	4(100.00)	0(0.00)	
0-6 weeks	364(94.30)	22(5.70)	
> 6 weeks	7(77.78)	2(22.22)	
NBS for SCD is morally right			0.312
True	370(94.15)	23(5.85)	
False	5(83.33)	1(16.67)	
NBS is useful for controlling SCD			1.000
True	360(93.75)	24(6.25)	
False	15(100.00)	0(0.00)	

256 *Fisher's exact method used, p-value < 0.05*

257

258

259 **Multivariate logistic regression analysis of factors with significant association with**
 260 **acceptability of newborn screening for sickle cell disease at bivariate analysis**

261 Only being a student (aOR= 25.02; 95% CI=1.29-484.51; $p= 0.033$) was significant at multivariate
 262 logistic regression analysis, though with a wide confidence interval. Male partner support was
 263 found to be a strong modifying factor with positive effect (coef=7.50; std error=1.31;95%CI=4.94-
 264 10.07; $p<0.001$). In the multivariate model, the coefficient of determination(r^2) was found to be
 265 63.9% (Table 5).

266

Table 5: Bivariate and Multivariate logistic regression analysis model of acceptability of newborn screening for sickle cell disease

Characteristic	Unadjusted model			Adjusted model		
	OR	95%CI	P-value	aOR	95%CI	P-value
Occupation						
Formal employment	REF			REF		
student	6.18	1.18-32.22	0.031	25.02	1.29-484.51	0.033
Informal employment	2.48	0.90-6.84	0.079	10.14	1.67-61.36	0.012
No employment	10.59	1.22-91.67	0.032	19.24	0.97-383.14	0.053
Age category						
24-35	REF			REF		
≤24 years	3.01	1.16-7.83	0.024	0.44	0.07-2.64	0.365
≥35 years	5.03	0.65-38.76	0.121	0.65	0.06-6.89	0.720
Cultural support						
No	REF			REF		
yes	4.32	1.13-16.49	0.032	0.24	0.01-5.89	0.380

Male support	partner			partner		
No	REF			REF		
Yes	559.5	105.46-	<0.00	1816.9	139.34-	<0.00
	0	2968.41	1	1	23691.24	1

267 *Table legend: model variable significant at 95% CI; p= 0.05; R²= 63.9%*

268
269
270

271 Discussion

272 The acceptability of newborn screening for sickle cell disease observed in the current study sample
273 was relatively high (94%). Comparatively, a similar study conducted at the postnatal wards at
274 Kisumu county referral hospital, Kenya, previously observed newborn screening for sickle cell
275 disease acceptability of 99.4% (Kuta et al., 2019). This informed successful implementation of
276 routine NBS for SCD in Kisumu County. The two studies are both hospital-based with similar
277 design and thus the comparable acceptability. Studies from Nigeria (Oluwole *et al.* 2020;Nnodu
278 *et al.* 2018) observed 86% acceptability of NBS which is below the acceptability in this current
279 study but is still comparable. The difference between the current study and that of Nnodu and team
280 in Nigeria was target population of postnatal mothers while that in Nigeria was conducted in health
281 institutions and University with diverse target groups.

282

283 Similar studies conducted in Gabon (Mombo et al., 2021) showed low acceptability rates of about
284 30% despite satisfactory knowledge among the respondents, demonstrating wide variabilities
285 (Marsh et al., 2011). The reason for the huge difference in the level of acceptability was that most
286 mothers in Gabon feared blood collection from their newborns (Mombo et al., 2021). This was not
287 the case in Homabay. There were obvious variations across socio-economic and demographic
288 stages, with younger single mothers being more receptive. Awareness and knowledge levels as
289 well as availability and access to services in high burden areas have been observed to influence
290 acceptability (Nnodu et al., 2018).

291

292 Despite the high willingness of the service recipient, other factors not explored in this study such
293 as cost barriers and service availability remain import factors in the overall evaluation of the
294 intervention acceptability. It is important for policy development to make the intervention

295 universally available and accessible in the county (Gates et al., 2021). Such a policy is currently
296 not in place in the county even though health is a devolved function.

297

298 Awareness about the intervention or the existence of the intervention helps improve acceptability
299 since those with low knowledge or awareness are less likely to accept the interventions. It will be
300 important that prior to implementation of the newborn screening programme in the county,
301 adequate awareness creation is created, and cost barriers eliminated. This has improved
302 acceptability in the HIV care set up (Izizag *et al.*, 2018) and can be extrapolated to sickle cell
303 disease care.

304

305 Primary interventions such as newborn screening programme are highly recommended to avert
306 disease complications and early deaths. Acceptance of health technologies is necessary for
307 progressive adoption and continued use by consumers as well as for its implementation at scale
308 (Kuta et al., 2019). Despite the relatively high acceptability rate, which might imply higher
309 willingness for newborn screening by the recipients, recent studies in Africa show that acceptance
310 for newborn screening is diverse and variable over time across different population groups as well
311 as the implementation contexts (Sekhon *et al.*, 2017; Katamea *et al.*, 2022). This might be
312 accounted for by the fact that health technology acceptance is a dynamic concept, and evolves
313 sequentially through a continuum of stages including, pre-use acceptability; initial use
314 acceptability and sustained use acceptability. However, few studies exist on the pre-use health
315 technology acceptance in Africa (Sekhon et al., 2017). The current study examined pre-use
316 acceptability. Whether high acceptability across these stages necessarily translates to scalability
317 remains a multidimensional challenge.

318

319 This study found out that occupation was strongly associated with acceptability of newborn
320 screening for sickle cell disease. Those in formal employment were less likely to accept newborn
321 screening for sickle cell disease when compared to the students and those in other occupations.
322 This is comparable to the findings of Nnodu et al. in a multicenter survey in Nigeria (Nnodu et al.,
323 2018). The similarities in both studies was the cross sectional design. The study in Nigeria was
324 however a multicenter study both in health institutions and universities. This provides a diverse
325 context to confirm the finding. However, contrasting findings were found by Katamea et al. in DRC

326 (Katamea et al., 2022) and Ahmed et al. in Nigeria (Ahmed et al., 2021) where occupation was not
327 associated with acceptability. This current study differed with the DRC study in that this was
328 hospital-based study with only postnatal mothers responding to the questionnaire while that of
329 DRC had all adults participate and it was a community survey. This could have contributed to the
330 differences in findings since not all those adults interviewed were having children. Either, most
331 students are usually younger and thus they may have been left out of the survey disproportionately.
332 The students in this study were predominantly younger and younger age has been associated with
333 increased acceptability of newborn screening for sickle cell disease.

334
335 Maternal age was found to be a strong predictor of acceptability of newborn screening for sickle
336 cell disease. Mothers who were younger than 25 years were found to be 3 times more likely to
337 accept newborn screening than those older. This finding is comparable to the findings of Mombo
338 et al. in a study done in Gabon where mothers younger than 29 years were more likely to accept
339 newborn screening than those older than 29 years (Mombo et al., 2021). Similarly, a study by
340 Nnodu et al. in Nigeria found that mothers younger than 21 years were more likely to accept
341 newborn screening for SCD than older ones (Nnodu et al., 2018). Younger women were thought
342 to be less culturally bound and therefore more likely to accept newborn screening for SCD
343 (Mombo et al., 2021). All the 3 studies above had an aspect of hospital-based interviews thus likely
344 the similarity in finding about the age.

345
346 Male partner support was found in this study to improve the odds of acceptability of newborn
347 screening for SCD across the various occupational categories. Male partner support may reflect
348 not only mutual consensus on decisions regarding the health interventions but also, potential
349 influence (Dsouza et al., 2022). It is therefore important to involve males in planning of
350 implementation of health interventions such as this (Mboane & Bhatta, 2015).

351
352 The multivariate model in this study had the significant factors explaining up to 63% of the variable
353 relationships. The unexplained 37% is likely due to unexplored factors. Review of previous similar
354 studies conducted (Katamea et al., 2022; Mombo et al., 2021) shows diverse factors may be
355 involved but there is no consensus on how different factors interact to influence or potentially
356 affect maternal acceptability of the newborn screening for SCD.

357
358 The study also found that the confidence intervals for most of the odds ratios of the associated
359 factors were very wide. This typically occur due to reasonably high variability in the factor
360 characteristics within the community of study (Marsh et al., 2011) and likely lower sample size,
361 even though this was above the minimum calculated sample size. However, this may also, result
362 from study limitations in that, in as much as acceptability is a multidimensional aspect, only
363 willingness was assessed.

364
365 The study limitations included sampling from one health facility within the county thus the mothers
366 coming from near the health facility could have contributed disproportionately in numbers.
367 However, conducting the study at the county referral hospital was to allow for a representation
368 from all other 8 sub counties within the county with each subcounty contributing at least 5% of the
369 total sample. The assessment of acceptability in a unidimensional approach as opposed to the
370 multidimensional approach was another limitation. However, this was done to allow the
371 comparison with other similar studies.

372
373

374 **Conclusion**

375 The acceptability of Newborn screening for Sickle cell disease among the postnatal mothers in
376 Homa Bay County was high, mainly among younger mothers. Male partner support was a strong
377 motivating factor for newborn screening acceptability with positive effect. The high acceptability
378 is indicative of a willingness for screening for sickle cell and is necessary for scaled up
379 programming. Implementation of routine newborn screening program for sickle cell disease is
380 highly recommended in view of the good user acceptability and high prevalence of sickle cell
381 disease at birth in the county. Additionally, there is need to explore other determinants of
382 acceptability and how they impact on acceptability and actual uptake.

383

384 **List of abbreviations**

385 aOR: Adjusted Odds Ratio
386 CI: Confidence Interval
387 EC: Ethics Committee
388 JOOUST: Jaramogi Oginga Odinga Teaching University of Science and Technology

389 NACOSTI: National Commission Of Science, Technology and Innovation

390 NBS: Newborn Screening

391 OR: Odds Ratio

392 SCD: Sickle Cell Disease

393 SCT: Sickle Cell trait

394 SSA: Sub-Saharan Africa

395 **Declarations**

396 **Ethical approval and consent to participate**

397 The proposal was approved by the board of postgraduate studies of Jaramogi Oginga Odinga
398 University of Science and Technology. Ethical approval for the study was obtained by JOOUST
399 Ethics Committee (ERC 36/02/23-33) and the NACOSTI permit obtained the National
400 Commission of Science, Technology and Innovation (Research License 101836). Using the
401 approvals from the Ethic Committee (EC) and NACOSTI, the County authorization was obtained
402 from the County Health Management Team and the County Director of Medical Services and
403 notice issued to the Hospital management and the relevant department. The research assistants
404 signed a confidentiality agreement form before starting the data collection and were adequately
405 trained to minimize any potential harm to the participants. Informed consent was obtained from
406 the study participants prior to any study procedure. The data collected was kept confidential and,
407 in a password-protected computer. Data was entered directly into the password-protected computer
408 with limited access. The access of the research assistants was terminated immediately data
409 cleaning was complete. No paper data was intended for the study. Key personal identifying
410 information such as name of the participant were not captured in order to deidentify the participant
411 and minimize chances of harm from breach of confidentiality. Mothers of newborns who were
412 willing to have their newborns screened were offered psychological counselling pre and post the
413 test and those whose newborns screened positive and need advanced psychological counseling
414 were referred to the counselling department of the Homa Bay County Teaching and Referral
415 hospital. The newborns who screened positive for SCD were linked with the SCD clinic of the
416 Homa Bay County Teaching and Referral Hospital for early initiation of comprehensive care,
417 usually initiated at the age of 2 months. The mothers of the newborns who were found to have
418 SCT were offered genetic counselling. The risk to the newborns screened for SCD included
419 minimal pain, swelling and bruise at the sample collection site due to the prick but this was

420 minimized by having the blood sample collected by adequately trained and professional research
421 assistants with experience in neonatal blood sampling from the heels. The heel prick was done
422 while maintaining the infection prevention control measures like adequate sterilization of the prick
423 site to minimize chances of prick site infection.

424

425

426 **Consent for publication**

427 Not applicable

428

429 **Competing interests**

430 The authors declare they have no competing interests

431

432 **Acknowledgment**

433 We acknowledge the postnatal mothers who consented to take part in the study which may improve
434 the policy on newborn screening and the care of patients with sickle cell disease and the research
435 assistants who helped with data collection.

436

437 **Funding**

438 This data collection for Msc program was funded by The European and Developing Countries
439 Clinical Trials Partnership (EDCTP) through the African Population and Health Research Centre
440 (APHRC), Lund University, Jaramogi Oginga Odinga University of Science and Technology and
441 Amref International University.

442

443 **Availability of data and materials**

444 The datasets used and/or analysed during the current study are available from the corresponding
445 author on reasonable request.

446

447 **Authors Contributions**

448 JO conceptualized the idea, drafted the protocol, collected data, did data analysis and wrote the
449 manuscript.

450 SA did critical review of the protocol and guided its writing, reviewed data collection tools, did
451 data analysis and reviewed the manuscript.

452 FM did critical review of the protocol and guided its writing, reviewed data collection tools, did
453 data analysis and reviewed the manuscript.

454 PO did critical review of the protocol and guided its writing, reviewed data collection tools, did
455 data analysis and reviewed the manuscript.

456 DO did critical review of the protocol and guided its writing, reviewed data collection tools, did
457 data analysis and reviewed the manuscript.

458

459

460 **References**

461

462 Bediako, S. M., & Neblett, E. W. (2011). Optimism and Perceived Stress in Sickle-Cell Disease:
463 The Role of an Afrocultural Social Ethos. *Journal of Black Psychology*, 37(2), 234–253.
464 <https://doi.org/10.1177/0095798410385681>

465 DeBaun, M. R., & Galadanci, N. A. (2022). *Sickle cell disease in sub-Saharan Africa*. 39.

466 Dsouza, J. P., Van Den Broucke, S., Pattanshetty, S., & Dhoore, W. (2022). Factors explaining
467 men's intentions to support their partner's participation in cervical cancer screening. *BMC*
468 *Women's Health*, 22(1), 443. <https://doi.org/10.1186/s12905-022-02019-y>

469 Gates, A., Gates, M., Rahman, S., Guitard, S., MacGregor, T., Pillay, J., Ismail, S. J., Tunis, M.
470 C., Young, K., Hardy, K., Featherstone, R., & Hartling, L. (2021). A systematic review of
471 factors that influence the acceptability of vaccines among Canadians. *Vaccine*, 39(2), 222–
472 236. <https://doi.org/10.1016/j.vaccine.2020.10.038>

473 Inusa, B. P., Juliana Olufunke, Y. D., & John Dada, L. (2015). Sickle Cell Disease Screening in
474 Northern Nigeria: The Co-Existence of \hat{I} '-Thalassemia Inheritance. *Pediatrics &*
475 *Therapeutics*, 05(03). <https://doi.org/10.4172/2161-0665.1000262>

476 Izizag, B. B., Situakibanza, H., Mbutiwi, T., Ingwe, R., Kiazayawoko, F., Nkodila, A., Mandina,
477 M., Longokolo, M., Amaela, E., & Mbula, M. (2018). Factors associated with acceptability

478 of HIV self-testing (HIVST) among university students in a Peri-Urban area of the
479 Democratic Republic of Congo (DRC). *Pan African Medical Journal*, 31.
480 <https://doi.org/10.11604/pamj.2018.31.248.13855>

481 Katamea, T., Mukuku, O., Mpoy, C. W., Mutombo, A. K., Luboya, O. N., & Wembonyama, S. O.
482 (2022). Factors Associated with Acceptability of Newborn Screening for Sickle Cell
483 Disease in Lubumbashi City, Democratic Republic of the Congo. *Global Journal of*
484 *Medical, Pharmaceutical, and Biomedical Update*, 17, 5.
485 https://doi.org/10.25259/GJMPBU_7_2022

486 Kuta, E. S. M., Tenge, C. N., Ganda, B. K. O., & Njuguna, F. M. (2019). Newborn screening for
487 sickle cell disease at Kisumu county hospital, Kisumu -Kenya. *East African Medical*
488 *Journal*, 96(2), 2419–2429. CABDirect.
489 <https://www.ajol.info/index.php/eamj/article/view/198133>

490 Makani, J., Ofori-Acquah, S. F., Nnodu, O., Wonkam, A., & Ohene-Frempong, K. (2013). Sickle
491 Cell Disease: New Opportunities and Challenges in Africa. *The Scientific World Journal*,
492 2013, 1–16. <https://doi.org/10.1155/2013/193252>

493 Marsh, V. M., Kamuya, D. M., & Molyneux, S. S. (2011). ‘All her children are born that way’:
494 Gendered experiences of stigma in families affected by sickle cell disorder in rural Kenya.
495 *Ethnicity & Health*, 16(4–5), 343–359. <https://doi.org/10.1080/13557858.2010.541903>

496 Mboane, R., & Bhatta, M. P. (2015). Influence of a husband’s healthcare decision making role on
497 a woman’s intention to use contraceptives among Mozambican women. *Reproductive*
498 *Health*, 12(1), 36. <https://doi.org/10.1186/s12978-015-0010-2>

499 *Ministry of Health 2020 (Sickle Cell Disease Guidelines—Kenya)*.
500

- 501 Misati Akuma, J. (2015). Socio-cultural and family change in Africa: Implications for adolescent
502 socialization in Kisii County, South Western, Kenya. *Les Cahiers d'Afrique de L'Est*, 50,
503 80–98. <https://doi.org/10.4000/estafrica.296>
- 504 Mombo, L. E., Makosso, L. K., Bisseye, C., Mbacky, K., & Edou, A. (2021). *Acceptability of*
505 *neonatal sickle cell disease screening among parturient women at the Paul Moukambi*
506 *Regional Hospital in rural Eastern Gabon, Central Africa*. 6.
- 507 Nnodu, O. E., Adegoke, S. A., Ezenwosu, O. U., Emodi, I. I., Ugwu, N. I., Ohiaeri, C. N., Brown,
508 B. J., Olaniyi, J. A., Isa, H., Okeke, C. C., Bene, B. A., Balongun, M. T., Okocha, E. O.,
509 Aneke, J. C., Lawson, J. O. J., Usman, A. M., Diaku-Akinmumi, I. N., Okolo, A. A., Israel-
510 Aina, Y. T., ... Adekile, A. D. (2018). A Multi-centre Survey of Acceptability of Newborn
511 Screening for Sickle Cell Disease in Nigeria. *Cureus*. <https://doi.org/10.7759/cureus.2354>
- 512 Odunvbun, M. E., Okolo, A. A., & Rahimy, C. M. (2008). Newborn screening for sickle cell
513 disease in a Nigerian hospital. *Public Health*, 122(10), 1111–1116.
514 <https://doi.org/10.1016/j.puhe.2008.01.008>
- 515 Ohene-Frempong, K., Oduro, J., Tetteh, H., & Nkrumah, F. (2008). SCREENING NEWBORNS
516 FOR SICKLE CELL DISEASE IN GHANA. *Pediatrics*, 121(Supplement_2), S120–S121.
517 <https://doi.org/10.1542/peds.2007-2022UUU>
- 518 Oluwole, E. O., Adeyemo, T. A., Osanyin, G. E., Odukoya, O. O., Kanki, P. J., & Afolabi, B. B.
519 (2020). Feasibility and acceptability of early infant screening for sickle cell disease in
520 Lagos, Nigeria—A pilot study. *PLOS ONE*, 15(12), e0242861.
521 <https://doi.org/10.1371/journal.pone.0242861>

- 522 Sekhon, M., Cartwright, M., & Francis, J. J. (2017). Acceptability of healthcare interventions: An
523 overview of reviews and development of a theoretical framework. *BMC Health Services*
524 *Research*, 17(1), 88. <https://doi.org/10.1186/s12913-017-2031-8>
- 525 Tshilolo, L., Aissi, L. M., Lukusa, D., Kinsiama, C., Wembonyama, S., Gulbis, B., & Vertongen,
526 F. (2009). Neonatal screening for sickle cell anaemia in the Democratic Republic of the
527 Congo: Experience from a pioneer project on 31 204 newborns. *Journal of Clinical*
528 *Pathology*, 62(1), 35–38. <https://doi.org/10.1136/jcp.2008.058958>
- 529 Uyoga, S., Macharia, A. W., Mochamah, G., Ndila, C. M., Nyutu, G., Makale, J., Tendwa, M.,
530 Nyatichi, E., Ojal, J., Otiende, M., Shebe, M., Awuondo, K. O., Mturi, N., Peshu, N., Tsofa,
531 B., Maitland, K., Scott, J. A. G., & Williams, T. N. (2019). The epidemiology of sickle cell
532 disease in children recruited in infancy in Kilifi, Kenya: A prospective cohort study. *The*
533 *Lancet Global Health*, 7(10), e1458–e1466. [https://doi.org/10.1016/S2214-](https://doi.org/10.1016/S2214-109X(19)30328-6)
534 [109X\(19\)30328-6](https://doi.org/10.1016/S2214-109X(19)30328-6)
- 535 Wanjiku, C. M., Njuguna, F., Asirwa, F. C., Njuguna, C., Roberson, C., & Greist, A. (2018).
536 Validation and Feasibility of a Point of Care Screening Test for Sickle Cell Disease in a
537 Resource Constrained Setting—A New Frontier. *Blood*, 132(Supplement 1), 2229–2229.
538 <https://doi.org/10.1182/blood-2018-99-118876>
- 539