

A CROSS-SECTIONAL STUDY OF THE PSYCHOLOGICAL IMPACT OF COVID-19 ON MEDICAL STUDENTS DURING & AFTER RECOVERY FROM COVID-19

AUTHOR NAMES AND AFFILIATIONS:

1. Dr. Anushka Misra, Undergraduate student, K.S. Hegde Medical Academy, Nitte (Deemed to be University), Mangalore, India
2. Dr. Damini Rane, Junior Resident, Department of Psychiatry, K.S. Hegde Medical Academy, Nitte (Deemed to be University), Mangalore, India
3. Dr. Arun Bhat P., Assistant Professor, Department of Psychiatry, K.S. Hegde Medical Academy, Nitte (Deemed to be University), Mangalore, India

ABSTRACT

Introduction: Mental health of medical students has been an utmost topic of discussion. It is important to know whether the upcoming doctors are mentally equipped to handle the pressure that comes with the profession. There is substantial literature indicating negative impact on the mental health during COVID-19 pandemic. Studies regarding the same on frontline workers seems scarce. Therefore, it is essential to assess the prevalence of symptoms anxiety and depression among medical students during and after the recovery from COVID-19 infection.

Materials and methods: It is a cross-sectional time bound study on 32 medical undergraduate students with active COVID-19 infection. They were screened using Self-reporting Questionnaire 20 (SRQ). Hamilton Anxiety Rating Scale (HAM-A) and Hamilton Depression Rating Scale (HAM-D) were administered on those who screened positive on SRQ-20 to assess the severity of anxiety and depressive symptoms. These were administered during and after COVID-19 infection on virtual platform.

Results: In our study, 59.37% population was screened positive for mental health disorder during COVID-19 infection. 89.5% of the positively screened respondents had mild anxiety symptoms and 10.5% had moderate anxiety symptoms. 47.4% of

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

screened respondents had mild depressive symptoms, 31.6% reported moderate depressive symptoms and 21.1% had severe depressive symptoms. Following recovery, there was a reduction in severity of anxiety and depressive symptoms which was statistically significant ($p < 0.001$).

Conclusion: Medical students have a propensity to develop anxiety and depressive symptoms while suffering from COVID-19 infection. Some may continue to have residual anxiety and depressive symptoms after recovery but the severity significantly reduces.

Keywords: Anxiety, COVID-19, Depression, Pandemic, Students, SARS-CoV-2.

A CROSS-SECTIONAL STUDY OF THE PSYCHOLOGICAL IMPACT OF COVID-19 ON MEDICAL STUDENTS DURING & AFTER RECOVERY FROM COVID-19

BACKGROUND

Coronavirus disease (COVID-19) was identified as an atypical form of severe acute respiratory syndrome (SARS) in Wuhan, China in December 2019 caused by novel coronavirus-2 (SARS-CoV-2)^{1,2}. The rapid spread of COVID-19 pandemic was a global health concern that can potentially impact the citizens of all nations due to its devastating consequences³.

Although ample clinical attention and research focus has been towards the treatment of the physical effects of the disease, the psychological fallout of the virus has been not been thoroughly studied on the affected individuals during and after recovery from the infection. As observed during previous viral outbreaks like SARS and MERS, viral infection and the isolation and quarantine swiftly culminate into anxiety, depression and sleep disturbances^{4,5}. Various psychiatric symptoms have been reported among SARS survivors, including post-traumatic stress disorder (PTSD), depression, panic disorder, and obsessive-compulsive disorder (OCD) up to 1 to 50 months of follow-up⁶. Previous prospective studies suggested that the psychological distress due to any illness was a predictor of future health and disease outcomes⁷. However, the current belief is that the psychological fallout of the COVID-19 pandemic may surpass the previous outbreaks mainly due to the widespread of misinformation by the social media which further perpetuates the sense of fear caused by the virus^{8,9}.

The SARS-CoV-2 pandemic poses an additional risk for medical students because they serve as frontline workers. They are more likely to have stress and anxiety and hence are more prone to develop depression, anxiety and sleep disorders¹⁰. Studies on junior doctors and medical students revealed anxiety and depression amongst the frontline workers. Some highlighted a higher propensity among females than males whereas few studies did not show a gender variation^{11,12,13}. Suicidal thoughts were prevalent in school going students during active COVID-19 infection and even after the recovery. Many of them reported of marked depression, anxiety, stress,

loneliness, social anxiety, lack of self-control and many more negative perceptions during this time¹⁴. Furthermore, there is increased worry and fear of their health and their loved ones, disturbed sleep patterns, trouble concentrating, dwindling social interaction due to quarantine and social distancing which contributed to their other worries during the COVID-19 pandemic¹⁵.

These medical students are naturally concerned, not only because of their health but also about the impact of infectious disease on their peers and the potential impact on their studies. Targeted interventions provided after diagnosis of anxiety or depression can improve the mental health of these students and can potentially alleviate the psychological impact of the novel coronavirus. To date, there have been limited studies on the psychological impact of the SARS-CoV-2 outbreak on the mental well-being of medical students during and after recovery from the illness. This study aims to measure the levels of anxiety and depressive symptoms among medical students during and after the recovery from COVID-19 infection.

MATERIALS AND METHODOLOGY

Participants

This was a cross-sectional study done on 32 students who were diagnosed with COVID-19 infection. It was a time-bound study for 9 months conducted in a tertiary care teaching institute on all medical students from Phase 1 to 4. A sample size was not pre-determined and was established upon active participation during the course of the study. Clearance from Institutional Ethics Committee (IEC) was obtained. Students who completed the online interview were recruited in the study. Those students who did not respond to the questionnaire or responded incompletely were excluded from the study.

Tools

A semi-structured questionnaire was devised to obtain socio-demographic and COVID-19 related details of the students. Self-reporting Questionnaire 20 (SRQ-20)¹⁶ is a self-administered scale which was provided to all students who had an active COVID-19 infection and also consented for the study. This is a screening

tool for assessing anxiety and depressive symptoms which was circulated on an online platform. Hamilton Anxiety Rating Scale (HAM-A)¹⁷ and Hamilton Depression Rating Scale (HAM-D)¹⁸ were administered on those who screened positive on SRQ-20 to assess the severity of anxiety and depressive symptoms. These were administered during and 6 months after recovery from COVID-19 infection on virtual platform.

Procedure

Medical students from Phase 1 to 4 who were diagnosed with COVID-19 taken up for the study. Verbal and written informed consent were taken before including them in the study. All the data was collected on an online video-calling platform. Socio-demographic and COVID-19 related details were obtained through the proforma. Self-administered SRQ-20 scale was provided to screen for anxiety and depressive symptoms at the time of active infection. Those who scored 8 or more on SRQ-20 were screened positive and were further assessed for severity of anxiety and depressive symptoms using HAM-A and HAM-D during and 6 months after recovery from COVID-19 infection. On the basis of the available information, inference was drawn regarding the difference in levels of anxiety and depressive symptoms during and after COVID-19 infection.

Statistical analysis

Statistical analysis was performed using Statistical Package for the Social Sciences (SPSS) version 23.0. The collected data was entered on a Microsoft Excel Sheet. Data was expressed in descriptive statistics using frequency and percentage. The pre/post comparison of HAM-A and HAM-D was done using Wilcoxon signed-rank test. P-value <0.05 was considered statistically significant.

RESULTS

The questionnaire was distributed via virtual platform among 600 medical students. A total of 32 students responded of whom 53.1% were females. More than half (65.6%) were less than 23 years of age. Out of the respondents, majority were from third and fourth year which was 31.3% in either year. Most of them did not have a

family history of psychiatric illness and had never consulted a psychiatrist. 4 of 32 participants had taken treatment for anxiety or depression in the past. Only 1 participant had a past history of COVID-19 infection. Majority of the respondents did not require in-patient care during current COVID19 infection. The socio-demographic details have been summarized in **Table 1**.

In our study, 59.4% were screened positive for mental health disorder with the mean age of 22 years (standard deviation [SD] =1.0) while 40.6% did not report of any mental health problems, mean age being 22.8 years (SD=1.67). Majority of the respondents were female (53.1%) from which 70.5% were screened positive for mental health disorder using the SRQ scale. Out of the remaining 46.9% male participants, less than half (46.6%) were screened positive for anxiety and depressive symptoms using SRQ scale. 15.8% of the participants with a positive SRQ score reported of a prior mental health disorder resulting in worsening of the symptoms. This data has been depicted in **Table 2**.

The subset of respondents that were screened positive for mental health related symptoms at the time of COVID-19 infection were assessed for severity of anxiety symptoms using HAM-A scale. 89.5% of the positively screened respondents had mild anxiety symptoms and 10.5% had moderate anxiety symptoms. Following recovery these respondents were reassessed to observe the change in severity of anxiety. Post recovery, 50% of these participants were found to have mild anxiety symptoms. This comparative data has been depicted in **Table 3** and **Figure 1**.

Similarly, the respondents that were screened positive for mental health related symptoms were also assessed for severity of depressive symptoms using HAM-D scale. At the time of active infection, it was found that 47.4% of screened respondents had mild depressive symptoms, 31.6% reported moderate depressive symptoms and 21.1% had severe depressive symptoms. Additionally, they were reassessed for change in severity of depression after recovery. 10.5% participants reported to have mild depressive symptoms, 5.3% had moderate depressive symptoms and 5.3% had severe depressive symptoms respectively. This comparative data has been depicted in **Table 4** and **Figure 2**.

The anxiety and depressive symptoms that was precipitated during COVID-19 infection reduced after recovery from the infection and this reduction was found to be

statistically significant ($z=3.66$, $N=32$, $p<0.001$). Anxiety symptoms reduced significantly with median HAM-A score during the infection being 5 [Interquartile range (IQR): 3-8] to post-infection score being 2 (IQR 1-8). Depressive symptoms also reduced significantly ($z=3.83$, $N=32$, $p<0.001$) with HAM-D score being 14 (IQR 11-17) during the infection and reducing to 9 (IQR 4-13) after recovery from infection.

DISCUSSION:

This cross-sectional study investigated mental health status of medical students during COVID-19 pandemic. The primary purpose was to investigate mental health status of medical students during and after recovery from COVID-19 infection and to determine their psychological wellbeing during the second wave of SARS-CoV2 virus.

Our study comprised of both genders but showed a higher number of women as compared to men. An Indian survey done in 2020-21 indicated that a higher percentage of women enrolled in medical science as compared to the men for higher education¹⁹. Furthermore, there has been an increase in female population in higher education from 2014 to 2020^{19,20}. Hence our study population did not show a deviation from the general trend. The students in our study were within the age range of 20-26 years. Other studies showed a higher prevalence of anxiety and depression among younger population²¹. A survey looking into depression and anxiety among medical students indicated that both anxiety and depression were higher among third year medical students which was consistent with our study findings²².

There are studies done on Indian medical students which showed a predominance of anxiety and depression which was also seen in our study highlighting that 12.5% of the medical students had a previous history of psychiatric consultation²³. Some of the participants in our study reported of anxiety and depressive symptoms during COVID-19 infection. Those who had a past history of mental health disorder reported of higher anxiety and depression during COVID-19 as compared to those who did not have a past history of psychiatric disorder. Studies contributing to our findings concluded that individuals with a history of mental health disorder have a higher prevalence of anxiety and depression²⁴.

In our study, more than half the population with COVID-19 infection had anxiety and depressive symptoms which was found to be in line with the available literature suggesting that individuals suffering from COVID-19 infection experienced more adverse mental health outcomes than their non-COVID-19 counterparts^{25,26}. Our study found an improvement in mental health status among undergraduate students after recovery from COVID-19 infection as compared to when they had COVID-19 infection. Our findings concerning the level of anxiety and depression symptoms among medical students were reported to be higher in females which complimented other studies which showed that females had significantly higher median anxiety and depression scores than the male participants²⁷. With our study we noticed statistically significant decrease in anxiety and depression. Residual mental health consequences were noted even after recovery from the infection. This was in line with another study done on COVID-19 recovered patients from general population which highlighted that one-third of the population continued to experience anxiety and depression 6 months after the discharge²⁸. Whereas, a similar study done on undergraduate students suggested no difference in mental health status before and after COVID-19 infection²⁹.

LIMITATIONS

The sample population for this study included undergraduate students pursuing Bachelor of Medicine, Bachelor of Surgery. Therefore, the results obtained cannot be generalized to general population. Due to the nature of the study being a time-bound study, the sample size obtained was small making it further difficult to generalize the findings of our study.

CONCLUSION

This study highlighted that undergraduate medical students have propensity to developing anxiety and depressive symptoms during COVID-19 infection. Certain socio-demographic groups were found of have been more at risk for mental health disorder; like female gender, respondents with past history of mental health disorder, students from third and fourth year of study. Moreover, respondents that had mental health disorder during COVID-19 infection had residual anxiety and depressive

symptoms when reassessed after recovery. Nonetheless, there was a significant reduction in severity of their anxiety and depressive symptoms after recovering from COVID-19 infection.

REFERENCES:

1. Lotfi, M. , Hamblin M.R. & Rezaei N.. 2020. COVID-19: transmission, prevention, and potential therapeutic opportunities. *Clin. Chim. Acta* 508: 254–266.
2. Zhai, P. , Ding Y., Wu X., *et al* 2020. The epidemiology, diagnosis and treatment of COVID-19. *Int. J. Antimicrob. Agents* 55: 105955.
3. Rajkumar RP. Ayurveda and COVID-19: where psychoneuroimmunology and the meaning response meet. *Brain Behav Immun.* 2020.
4. Khan, S. , Siddique R., Li H., *et al* 2020. Impact of coronavirus outbreak on psychological health. *J. Glob. Health* 10: 010331.
5. Deng J, Zhou F, Hou W, Silver Z, Wong CY, Chang O et al., The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: a meta-analysis. *Ann N Y Acad Sci.* 2021 Feb;1486(1):90-111.
6. Wu Y., Xu X., Chen Z. Nervous system involvement after infection with COVID-19 and other coronaviruses. *Brain Behav. Immun.* 2020
7. Turner AI, Smyth N, Hall SJ, Torres SJ, Hussein M, Jayasinghe SU et al. Psychological stress reactivity and future health and disease outcomes: A systematic review of prospective evidence. *Psychoneuroendocrinology.* 2020;114:104599. pmid:32045797
8. Mian, A. & Khan S.. 2020. Coronavirus: the spread of misinformation. *BMC Med.* 18: 89.
9. Zarocostas, J. 2020. How to fight an infodemic. *Lancet* 395: 676.
10. Azad N, Shahid A, Abbas N, Shaheen A, Munir N. Anxiety And Depression In Medical Students Of A Private Medical College. *J Ayub Med Coll Abbottabad.* 2017 Jan-Mar.

11. Pandey U, Corbett G, Mohan S, et al. Anxiety, Depression and Behavioural Changes in Junior Doctors and Medical Students Associated with the Coronavirus Pandemic: A Cross-Sectional Survey [published online ahead of print, 2020 Sep 24]. *J Obstet Gynaecol India*. 2020;71(1):1-5.
doi:10.1007/s13224-020-01366-w
12. Wang X, Hegde S, Son C, Keller B, Smith A, Sasangohar F. Investigating Mental Health of US College Students During the COVID-19 Pandemic: Cross-Sectional Survey Study. *J Med Internet Res*. 2020;22(9):e22817. Published 2020 Sep 17.
doi:10.2196/22817
13. Saraswathi I, Saikarthik J, Senthil Kumar K, Madhan Srinivasan K, Ardhanaari M, Gunapriya R. Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study. *PeerJ*. 2020;8:e10164. Published 2020 Oct 16.
doi:10.7717/peerj.10164
14. Zhu S, Zhuang Y, Lee P, Wong PWC. The changes of suicidal ideation status among young people in Hong Kong during COVID-19: A longitudinal survey. *J Affect Disord*. 2021;294:151-158. doi:10.1016/j.jad.2021.07.042
15. Son C, Hegde S, Smith A, Wang X, Sasangohar F. Effects of COVID-19 on College Students' Mental Health in the United States: Interview Survey Study. *J Med Internet Res*. 2020;22(9):e21279. Published 2020 Sep 3. doi:10.2196/21279
16. Scholte WF, Verduin F, van Lammeren A, Rutayisire T, Kamperman AM. Psychometric properties and longitudinal validation of the self-reporting questionnaire (SRQ-20) in a Rwandan community setting: a validation study. *BMC Med Res Methodol*. 2011;11:116. Published 2011 Aug 16.
doi:10.1186/1471-2288-11-116
17. Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol* 1959; 32:50–55.
18. Hedlund JL, Viewig BW (1979). "The Hamilton rating scale for depression: a comprehensive review". *Journal of Operational Psychiatry*.
19. Ministry of Human Resource Development, (2020-21). All India Survey on Higher Education. New Delhi: Department of Higher Education, MHRD, Government of India.
<https://aishe.gov.in/aishe/viewDocument.action;jsessionid=48A04A2580470998D334B629574037EB.n3?documentId=322>

20. Kraljević G, Čarapina Zovko I, Jelavić B. Sex Ratio of Graduated Medical Students of the School of Medicine in Mostar for the Period 2003-2020. ABCR [Internet]. 2022 Jul. 26 [cited 2023 Jun. 13];1(1):34-8. Available from: <https://abcr-mefmo.org/index.php/abcr/article/view/9>
21. Iqbal S, Gupta S, Venkatarao E. Stress, anxiety and depression among medical undergraduate students and their socio-demographic correlates. *Indian J Med Res.* 2015;141(3):354-357. doi:10.4103/0971-5916.156571
22. Chandavarkar U, Azzam A, Mathews CA. Anxiety symptoms and perceived performance in medical students. *Depress Anxiety.* 2007;24(2):103-111. doi:10.1002/da.20185
23. Raja S, Balasubramanian G, Jamuna Rani R. Prevalence of depression, anxiety and stress among private medical college students in South India: A cross-sectional study. *J Educ Health Promot.* 2022;11:373. Published 2022 Nov 26. doi:10.4103/jehp.jehp_393_22
24. Galindo-Vázquez O, Ramírez-Orozco M, Costas-Muñiz R, Mendoza-Contreras LA, Calderillo-Ruiz G, Meneses-García A. Symptoms of anxiety, depression and self-care behaviors during the COVID-19 pandemic in the general population. Síntomas de ansiedad, depresión y conductas de autocuidado durante la pandemia de COVID-19 en la población general. *Gac Med Mex.* 2020;156(4):298-305. doi:10.24875/GMM.20000266
25. Hossain MM, Tasnim S, Sultana A, et al. Epidemiology of mental health problems in COVID-19: a review. *F1000Res.* 2020;9:636. Published 2020 Jun 23. doi:10.12688/f1000research.24457.1
26. Deng J, Zhou F, Hou W, et al. The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: a meta-analysis. *Ann N Y Acad Sci.* 2021;1486(1):90-111. doi:10.1111/nyas.14506
27. Pandey U, Corbett G, Mohan S, et al. Anxiety, Depression and Behavioural Changes in Junior Doctors and Medical Students Associated with the Coronavirus Pandemic: A Cross-Sectional Survey [published online ahead of print, 2020 Sep 24]. *J Obstet Gynaecol India.* 2020;71(1):1-5. doi:10.1007/s13224-020-01366-w
28. Tizenberg BN, Brenner LA, Lowry CA, et al. Biological and Psychological Factors Determining Neuropsychiatric Outcomes in COVID-19. *Curr Psychiatry Rep.* 2021;23(10):68. Published 2021 Oct 1. doi:10.1007/s11920-021-01275-3

29. Lu P, Yang L, Wang C, et al. Mental health of new undergraduate students before and after COVID-19 in China. *Sci Rep.* 2021;11(1):18783. Published 2021 Sep 22. doi:10.1038/s41598-021-98140-3

Table 1: Sociodemographic data:

SL NO.	SOCIODEMOGRAPHIC DATA	RANGES	FREQUENCY (%)
1.	Age group	≤22 years	21 (65.6)
		>22 years	11 (34.4)
2.	Sex	Male	15 (46.9)
		Female	17 (53.1)
3.	Year of study	First	4 (12.5)
		Second	8 (25.0)
		Third	10 (31.3)
		Fourth	10 (31.3)
4.	Past history of psychiatric illness	Yes	4 (12.5)
		No	28 (87.5)
5.	Family history of psychiatric illness	Yes	1 (3.1)
		No	31 (96.9)
6.	No. of COVID episodes	≤ 1	31 (96.9)
		>1	1 (3.1)
7.	Hospitalization	Yes	4 (12.5)
		No	28 (87.5)

Table 2: Prevalence of mental health problems during COVID-19 infection:

SL NO.	SRQ SCORES	FREQUENCY (%)
1.	<8	13 (40.6)
2.	≥8	19 (59.4)

Table 3: Comparison of severity of anxiety symptoms during and after COVID-19 infection:

SL NO.	HAM A SCORE INTERPRETATION	FREQUENCY (%)	
		DURING COVID INFECTION	POST RECOVERY
1.	Mild	17 (89.5)	19 (100.0)
2.	Moderate	2 (10.5)	None
3.	Severe	None	None

Figure 1: Comparison of severity of anxiety symptoms during and after COVID-19 infection:




Table 4: Comparison of severity of depressive symptoms during and after COVID-19 infection:

SL NO.	HAM D SCORE INTERPRETATION	FREQUENCY (%)	
		DURING COVID INFECTION	POST RECOVERY
1.	Mild	9 (47.4)	2 (40)
2.	Moderate	6 (31.6)	1 (20)
3.	Severe	4 (21.1)	1 (20)

Figure 2: Comparison of severity of depressive symptoms during and after COVID-19 infection:

