Longitudinal Influenza A Virus Screening of Retail Milk from Canadian Provinces (Rolling Updates)

Hannah L. Wallace^{1*}, Jordan Wight^{1*}, Mariana Baz², Barbara Dowding³, Louis Flamand², Tom Hobman⁴, Francois Jean⁵, Jeffrey B Joy^{6,7,8}, Andrew S. Lang⁹, Sonya MacParland¹⁰, Craig McCormick¹¹, Ryan Noyce¹², Rodney S. Russell¹³ Selena M. Sagan⁵, Jumari Snyman⁴, Gabriela J. Rzeszutek¹, Mustafa S. Jafri¹, Isaac Bogoch^{14,15}#, Jason Kindrachuk^{1,16,17}#, Angela L. Rasmussen^{18,19}#, on behalf of the Pan-Canadian Milk Study Network

Corresponding authors: II Bogoch (Isaac.Bogoch@uhn.ca); J Kindrachuk (Jason.Kindrachuk@umanitoba.ca); AL Rasmussen (angela.rasmussen@usask.ca)

¹ Department of Medical Microbiology & Infectious Diseases, Max Rady College of Medicine, University of Manitoba, MB, Canada

² Département de microbiologie-infectiologie et d'immunologie, Faculté de Médecine, Université Laval, QC, Canada

³ Department of Biological Sciences, Faculty of Science, Applied Science, and Engineering, University of New Brunswick – Saint John, NB, Canada

⁴ Department of Cell Biology, University of Alberta, AB, Canada

⁵ Department of Microbiology & Immunology, Faculty of Science, University of British Columbia, BC, Canada

⁶ British Columbia Centre for Excellence in HIV/AIDS, Vancouver, BC, Canada

⁷ Bioinformatics, University of British Columbia, Vancouver, BC, Canada

⁸ Department of Medicine, University of British Columbia, Vancouver, BC, Canada

⁹ Department of Biology, Memorial University, NL, Canada

¹⁰ Schwartz-Reisman Liver Research Centre, Toronto General Research Institute

¹¹ Department of Microbiology & Immunology, Faculty of Medicine, Dalhousie University, NS, Canada

¹² Medical Microbiology and Immunology Department, Faculty of Medicine & Dentistry, University of Alberta, AB, Canada

¹³ Immunology and Infectious Diseases, Division of Biomedical Sciences, Faculty of Medicine, Memorial University, NL, Canada

¹⁴ Division of Infectious Diseases, Toronto General Hospital, Toronto, ON, Canada

¹⁵ Department of Medicine, University of Toronto, Toronto, ON, Canada

¹⁶ Department of Internal Medicine, Max Rady College of Medicine, University of Manitoba, MB, Canada

¹⁷ Manitoba Centre for Proteomics and Systems Biology, Max Rady College of Medicine, University of Manitoba, MB, Canada

¹⁸ Vaccine and Infectious Disease Organization (VIDO), University of Saskatchewan, SK, Canada

¹⁹ Department of Biochemistry, Microbiology, and Immunology, University of Saskatchewan, SK, Canada

^{*}These authors contributed equally

Abstract

Highly pathogenic avian influenza (HPAI) H5N1 has caused the deaths of more than 100 million birds since 2021, and human cases since 1997 have been associated with significant morbidity and mortality. Given the recent detection of HPAI H5N1 in dairy cattle and H5N1 RNA detections in pasteurized retail milk in the United States, we established the Pan-Canadian Milk (PCM) Network. Through our network of collaborators from across Canada, retail milk is being procured longitudinally and sent to a central laboratory for testing for the presence of influenza A virus RNA. To date (20 June 2024), we have tested 84 retail milk samples from all ten Canadian provinces (NL, NS, PEI, NB, QC, ON, MB, SK, AB, and BC) and all have tested negative for influenza A virus RNA. Testing is ongoing and these results will be updated on rolling basis as additional data becomes available. Despite no known HPAI infections of dairy cattle in Canada to date, H5N1 poses a significant threat to the health of both humans and other animals. Routine surveillance of retail milk on a national scale will allow for monitoring of infected dairy cattle on an ongoing basis in a cost-effective, standardized, scalable and easily accessible manner. Our network and testing will act as an early detection system to help inform containment and mitigation activities if positive samples are identified.

Introduction

Influenza A viruses (IAVs) are in the family Orthomyxoviridae which can infect a wide variety of hosts and result in a wide spectrum of illness, from asymptomatic to severe including multiorgan failure resulting in death (1, 2). Highly pathogenic avian influenza (HPAI) is an emerging disease threat with pandemic potential. Most recently arriving in North America in late 2021 (3), HPAI H5N1 clade 2.3.4.4b has infected a broad range of wild and domesticated bird species, and has resulted in the death of more than 100 million birds (4). This virus and derived reassortants have also infected diverse species of marine and terrestrial mammals (5-8), with the most recent infections occurring in dairy cattle in the United States (9-11). Since 1997, spillover of HPAI H5N1 from birds to humans have also been documented. Of the nearly 900 known human infections of H5N1, most have exhibited severe disease symptoms, with a case fatality rate of approximately 50% (12). However, despite the fact that the currently circulating H5N1 clade 2.3.4.4b virus and derived reassortants have caused extensive infections and deaths of birds and mammals, relatively few human cases have been reported, with only 12 cases globally associated with this clade since January 2022 (12, 13). Notably, none of these cases have been fatal, with the majority being reported as exhibiting mild disease symptoms with frequent reports of conjunctivitis. Three of the 12 human cases were reported in the spring of 2024; cases were associated with close interactions with infected dairy cattle. The three cases all had relatively mild symptoms (conjunctivitis), with one also reporting mild respiratory symptoms and all recovered (14–16).

The United States Department of Agriculture (USDA) reported in late April 2024 that dairy cattle on a farm in Texas were exhibiting signs of lethargy, dehydration, mild respiratory symptoms, decreased feed intake, decreased milk production and/or milk with abnormal colour/texture, and further investigation revealed lesions similar to those seen with mastitis (17, 18). There

were also reports of barn cats being found dead on this property (18). Phylogenetic analyses showed that the infection of dairy cattle was the result of a single spillover event of HPAI H5N1 from wild birds in late 2023 (19). This was then followed by subsequent cattle-to-cattle transmission possibly mediated by exposure to contaminated milking equipment, or another currently unknown route (10, 20). Since then, the outbreak has spread to 116 herds in the US as the transport of infected dairy cattle occurred before the extent of the outbreak was known (21). H5N1-infected dairy cattle have now been reported in nine states (21) and recently H5N1 RNA was detected in wastewater from nine cities across Texas (22).

Soon after the initial dairy infections were reported, the FDA began conducting H5N1 surveillance of pasteurized milk and other dairy products and reported that up to 20% of samples tested were positive for H5N1 RNA. Importantly, no viable virus was detected in the retail pasteurized dairy samples (23). Testing of tissue from dairy cattle as well as testing of retail beef is also being performed by the USDA. To date, one muscle sample from a dairy cow known to be infected with H5N1 has shown to be positive for H5N1 RNA and no meat from any dairy cattle has entered the US food supply (24).

To date, no H5N1 infections of dairy cattle have been reported in Canada. Due to current mandatory testing by the USDA before transporting cattle across state boarders to limit interstate spread (25), along with limited importation of dairy cattle from the US into Canada, infected cattle are unlikely to enter Canada and threaten Canadian cattle operations. Additionally, since 29 April 2024, the Canadian Food Inspection Agency (CFIA) has implemented a requirement for proof of negative testing on USDA export certificates for all lactating dairy cattle (26), further protecting Canadian cattle. Furthermore, as this virus still extensively circulates in wild birds, there is the potential for the virus to independently spill over from wild birds into Canadian cattle, as occurred on the index farm in Texas. Therefore, monitoring of cattle in Canada is of vital importance to detect if a potential spillover has occurred as quickly as possible.

Although it is known that cattle can be infected by influenza A viruses (27–29), including H5N1 (30), no viral genomes have been sequenced from cattle, prior to early 2024. Very little is known about the patterns of disease or the tissue types that allow viral replication although decreased milk production had been previously associated with influenza antibodies in dairy cattle (31). A recent pre-print suggested that mammary tissue of dairy cattle express both α -2,3- and α -2,6-sialic acid, associated with avian and human influenza receptors, respectively (32). This suggests the possibility of dairy cattle acting as a "mixing pot" for avian and human influenza virus reassortment, similar to what has been described for swine (10).

In April 2024, we established the Pan-Canadian Milk (PCM) Network, bringing together colleagues from across the country to procure retail milk every two weeks and send samples to a central laboratory for testing. This study will continue longitudinally, with particular interest in testing during the wild bird migration seasons. Recently, the CFIA tested retail milk and found that all samples tested IAV RNA negative; however, these efforts represent a snapshot in time and no detailed plans for longitudinal screening have been announced (33). Ongoing

surveillance in retail milk will not only allow rapid containment measures should any samples test positive but may also further our understanding of dairy cattle susceptibility to various influenza A viruses, not just H5N1.

Methods

Milk Sampling

Pasteurized whole (3.25%) milk was obtained from local stores by collaborators in all Canadian provinces. However, additional detailed data was collected about each retail milk sample though this is not reported here. Samples were collected every two weeks beginning 29 April 2024. Milk cartons were externally disinfected, opened in a biosafety cabinet, and a sample aseptically collected in sterile conical tubes at the laboratories of collaborators. Samples were kept at 4°C and shipped on ice to the Kindrachuk lab at the University of Manitoba for processing. Samples were then stored at -80°C.

RNA Isolation

RNA isolation was performed from 140 μ L of milk using the Qiagen Viral RNA Mini Kit (Qiagen, Product # 52906) as per the manufacturer's instructions. Isolated RNA was stored at -80°C.

Screening for Influenza A Virus

Real-time RT-PCR was used to screen for the presence of IAV matrix gene RNA as per Wight *et al.* (34) using a QuantStudio 6 Flex instrument (Applied Biosystems). Any sample that tests positive for the matrix gene will be subsequently screened for the H5 subtype of the haemagglutinin gene also as per Wight *et al.* (34). In the event of positive sample identification, data will be immediately shared with federal agencies through existing linkages and collaborative connections.

Results (Rolling Updates)

As of 20 June 2024, 84 retail milk samples obtained from each of the Canadian Provinces (NL, NS, PEI, NB, QC, ON, MB, SK, AB, and BC) have been tested. All samples to date have tested negative for IAV matrix RNA (Table 1).

Table 1. Provinces and pan-Canadian totals of retail milk samples tested, the number of IAV RNA positive samples, and the percentage of IAV RNA positive samples. RNA was isolated from milk samples and screened for IAV matrix RNA using real-time RT-PCR.

| Province | Number of Milk Samples Tested | Number of IAV RNA Positive Samples | % of IAV RNA Positive Samples |
|---------------------------|----------------------------------|--|-------------------------------------|
| Newfoundland and Labrador | 8 | 0 | 0% |

| Nova Scotia | 6 | 0 | 0% |
|-------------------------|----|---|----|
| Prince Edward Island | 1 | 0 | 0% |
| New Brunswick | 6 | 0 | 0% |
| Quebec | 15 | 0 | 0% |
| Ontario | 7 | 0 | 0% |
| Manitoba | 8 | 0 | 0% |
| Saskatchewan | 13 | 0 | 0% |
| Alberta | 9 | 0 | 0% |
| British Columbia | 11 | 0 | 0% |
| National Totals | 84 | 0 | 0% |

Discussion

CFIA has recently reported that the 600 milk samples they tested were H5 RNA negative (33) as well as the fact that no outbreaks of disease associated with H5N1 have been reported in dairy cattle in Canada, it was not unexpected that our samples, collected during nearly the same time as CFIA's, also tested negative for influenza.

Given the expansive nature of the H5N1 outbreak in dairy cattle, spillover to other species, and at least three human infections linked to close contact with dairy cattle in the US, continued longitudinal monitoring of milk samples in Canada may allow early detection of HPAI infection of dairy cattle. This will augment and compliment ongoing efforts by government agencies and other academic groups, enabling early containment and preventative measures to inhibit further spread. Sample collection and testing will be ongoing for the foreseeable future. This document will be updated as additional data becomes available. Our results will be provided to federal government agencies in real-time as this live preprint is updated to ensure early, transparent, and collaborative data exchange with key stakeholders and policy makers.

Acknowledgements

We would like to acknowledge all PCM Study collaborators for contributing milk samples from across the country.

Funding

This work was supported by the Canadian Institutes of Health Research (Tier 2 Canada Research Chair, grant number 950-231498 to JK) and by the Natural Sciences and Engineering Research Council Discovery Grant (RGPIN-2018-06036 to JK).

Author Contributions

Conceptualization: HLW, JW, IB, AR, JK

Data Curation: HLW, JW Formal Analysis: HLW, JW Funding Acquisition: JK Investigation: JW, HLW Methodology: JW, HLW

Project Administration: HLW, JW, JK

Resources: JK Supervision: JK

Writing (original draft): HLW, JW

Writing (review and editing): All authors

Competing Interests

The authors declare that they have no competing interests.

References

- Centers for Disease Control and Prevention. 2024. Bird Flu Virus Infections in Humans. Influenza Flu. https://www.cdc.gov/flu/avianflu/avian-in-humans.htm. Retrieved 25 May 2024.
- 2. Canadian Food Inspection Agency. 2024. Animals susceptile to H5N1 highly pathogenic avaian influenza (HPAI). Gov Can. https://inspection.canada.ca/en/animal-health/terrestrial-animals/diseases/reportable/avian-influenza/animals-susceptible-h5n1-hpai. Retrieved 25 May 2024.
- 3. Caliendo V, Lewis NS, Pohlmann A, Baillie SR, Banyard AC, Beer M, Brown IH, Fouchier RAM, Hansen RDE, Lameris TK, Lang AS, Laurendeau S, Lung O, Robertson G, Van Der Jeugd H, Alkie TN, Thorup K, Van Toor ML, Waldenström J, Yason C, Kuiken T, Berhane Y. 2022. Transatlantic spread of highly pathogenic avian influenza H5N1 by wild birds from Europe to North America in 2021. Sci Rep 12:11729.

- 4. Youk S, Torchetti MK, Lantz K, Lenoch JB, Killian ML, Leyson C, Bevins SN, Dilione K, Ip HS, Stallknecht DE, Poulson RL, Suarez DL, Swayne DE, Pantin-Jackwood MJ. 2023. H5N1 highly pathogenic avian influenza clade 2.3.4.4b in wild and domestic birds: Introductions into the United States and reassortments, December 2021–April 2022. Virology 587:109860.
- 5. Lair S, Quesnel L, Signore AV, Delnatte P, Embury-Hyatt C, Nadeau M-S, Lung O, Ferrell ST, Michaud R, Berhane Y. 2024. Outbreak of Highly Pathogenic Avian Influenza A(H5N1) Virus in Seals, St. Lawrence Estuary, Quebec, Canada1. Emerg Infect Dis 30.
- 6. Jakobek BT, Berhane Y, Nadeau M-S, Embury-Hyatt C, Lung O, Xu W, Lair S. 2023. Influenza A(H5N1) Virus Infections in 2 Free-Ranging Black Bears (*Ursus americanus*), Quebec, Canada. Emerg Infect Dis 29.
- 7. Alkie TN, Cox S, Embury-Hyatt C, Stevens B, Pople N, Pybus MJ, Xu W, Hisanaga T, Suderman M, Koziuk J, Kruczkiewicz P, Nguyen HH, Fisher M, Lung O, Erdelyan CNG, Hochman O, Ojkic D, Yason C, Bravo-Araya M, Bourque L, Bollinger TK, Soos C, Giacinti J, Provencher J, Ogilvie S, Clark A, MacPhee R, Parsons GJ, Eaglesome H, Gilbert S, Saboraki K, Davis R, Jerao A, Ginn M, Jones MEB, Berhane Y. 2023. Characterization of neurotropic HPAI H5N1 viruses with novel genome constellations and mammalian adaptive mutations in free-living mesocarnivores in Canada. Emerg Microbes Infect 12:2186608.
- 8. European Food Safety Authority, European Centre for Disease Prevention and Control, European Union Reference Laboratory for Avian Influenza, Fusaro A, Gonzales JL, Kuiken T, Mirinavičiūtė G, Niqueux É, Ståhl K, Staubach C, Svartström O, Terregino C, Willgert K, Baldinelli F, Delacourt R, Georganas A, Kohnle L. 2024. Avian influenza overview December 2023–March 2024. EFSA J 22.
- 9. USDA APHIS. 2024. Federal and State Veterinary Public Health Agencies Share Update on HPAI Detection in Kansas Texas Dairy Herds.
- 10. Nguyen T-Q, Hutter C, Markin A, Thomas M, Lantz K, Killian ML, Janzen GM, Vijendran S, Wagle S, Inderski B, Magstadt DR, Li G, Diel DG, Frye EA, Dimitrov KM, Swinford AK, Thompson AC, Snevik KR, Suarez DL, Spackman E, Lakin SM, Ahola SC, Johnson KR, Baker AL, Robbe-Austerman S, Torchetti MK, Anderson TK. 2024. Emergence and interstate spread of highly pathogenic avian influenza A(H5N1) in dairy cattle https://doi.org/10.1101/2024.05.01.591751.
- 11. Hu X, Saxena A, Magstadt DR, Gauger PC, Burrough E, Zhang J, Siepker C, Mainenti M, Gorden PJ, Plummer P, Li G. 2024. Highly Pathogenic Avian Influenza A (H5N1) clade 2.3.4.4b Virus detected in dairy cattle https://doi.org/10.1101/2024.04.16.588916.
- 12. World Health Organization. 2024. Cumulative number of confirmed human cases for avian influenza A (H5N1) reported to WHO, 2003-2024. https://cdn.who.int/media/docs/default-source/influenza/h5n1-human-case-cumulative-

- table/2024_feb_tableh5n1.pdf?sfvrsn=bccd8c23_1&download=true%22%20%5ct%20%22_blank. Retrieved 9 May 2024.
- 13. Kniss K, Sumner KM, Tastad KJ, Lewis NM, Jansen L, Julian D, Reh M, Carlson E, Williams R, Koirala S, Buss B, Donahue M, Palm J, Kollmann L, Holzbauer S, Levine MZ, Davis T, Barnes JR, Flannery B, Brammer L, Fry A. 2023. Risk for Infection in Humans after Exposure to Birds Infected with Highly Pathogenic Avian Influenza A(H5N1) Virus, United States, 2022. Emerg Infect Dis 29.
- 14. Uyeki TM, Milton S, Abdul Hamid C, Reinoso Webb C, Presley SM, Shetty V, Rollo SN, Martinez DL, Rai S, Gonzales ER, Kniss KL, Jang Y, Frederick JC, De La Cruz JA, Liddell J, Di H, Kirby MK, Barnes JR, Davis CT. 2024. Highly Pathogenic Avian Influenza A(H5N1) Virus Infection in a Dairy Farm Worker. N Engl J Med NEJMc2405371.
- 15. Centers for Disease Control and Prevention. 2024. CDC Reports Second Human Case of H5 Bird Flu Tied to Dairy Cow Outbreak | CDC Online Newsroom | CDC. Press Release.
- 16. Centers for Disease Control and Prevention. 2024. CDC A (H5N1) Bird Flu Response Update May 31, 2024. CDC Influenza Flu. https://www.cdc.gov/flu/avianflu/spotlights/2023-2024/h5n1-response-053124.html#print. Retrieved 3 June 2024.
- 17. Caserta LC, Frye EA, Butt SL, Laverack MA, Nooruzzaman M, Covalenda LM, Thompson A, Prarat Koscielny M, Cronk B, Johnson A, Kleinhenz K, Edwards EE, Gomez G, Hitchener GR, Martins M, Kapczynski DR, Suarez DL, Alexander Morris ER, Hensley T, Beeby JS, Lejeune M, Swinford A, Elvinger F, Dimitrov KM, Diel DG. 2024. From birds to mammals: spillover of highly pathogenic avian influenza H5N1 virus to dairy cattle led to efficient intra- and interspecies transmission https://doi.org/10.1101/2024.05.22.595317.
- 18. Eric R. Burrough, Magstadt DR, Petersen B, Timmermans SJ, Gauger PC, Zhang J, Siepker C, Mainenti M, Li G, Thompson AC, Gorden PJ, Plummer PJ, Main R. 2024. Highly Pathogenic Avian Influenza A(H5N1) Clade 2.3.4.4b Virus Infection in Domestic Dairy Cattle and Cats, United States, 2024. CDC Emerg Infect Dis. Journal, Early Release. https://wwwnc.cdc.gov/eid/article/30/7/24-0508_article. Retrieved 29 April 2024.
- 19. Worobey M, Gangavarapu K, Pekar JE, Joy JB, Moncla L, Kraemer MUG, Dudas G, Goldhill D, Ruis C, Serrano LM, Ji X, Andersen KG, Wertheim JO, Lemey P, Suchard MA, Rasmussen AL, Chand M, Groves N, Pybus OG, Peacock TP, Rambaut A, Nelson MI. 2024. Preliminary report on genomic epidemiology of the 2024 H5N1 influenza A virus outbreak in U.S. cattle (Part 1 of 2) H5N1-global Virological. https://virological.org/t/preliminary-report-ongenomic-epidemiology-of-the-2024-h5n1-influenza-a-virus-outbreak-in-u-s-cattle-part-1-of-2/970. Retrieved 26 May 2024.
- 20. Le Sage V, Campbell AJ, Reed D, Duprex WP, Lakdawala S. 2024. Influenza H5N1 and H1N1 viruses remain infectious in unpasteurized milk on milking machinery surfaces https://doi.org/10.1101/2024.05.22.24307745.

- 21. USDA APHIS. 2024. Highly Pathogenic Avian Influenza (HPAI) Detections in Livestock 2024-06-20. https://www.aphis.usda.gov/livestock-poultry-disease/avian/avian-influenza/hpaidetections/livestock. Retrieved 20 June 2024.
- 22. Tisza MJ, Hanson BM, Clark JR, Wang L, Payne K, Ross MC, Mena KD, Gitter A, Javornik-Cregeen SJ, Cormier J, Avadhanula V, Terwilliger A, Balliew J, Wu F, Rios J, Deegan J, Piedra PA, Petrosino JF, Boerwinkle E, Maresso AW. 2024. Virome Sequencing Identifies H5N1 Avian Influenza in Wastewater from Nine Cities https://doi.org/10.1101/2024.05.10.24307179.
- 23. Spackman E, Jones DR, McCoig AM, Colonius TJ, Goraichuk I, Suarez DL. 2024. Characterization of highly pathogenic avian influenza virus in retail dairy products in the US https://doi.org/10.1101/2024.05.21.24307706.
- 24. USDA APHIS. 2024. Updates on H5N1 Beef Safety Studies. USDA APHIS. https://www.aphis.usda.gov/livestock-poultry-disease/avian/avian-influenza/hpai-detections/livestock/h5n1-beef-safety-studies#:~:text=Meat%20from%20condemned%20cows%20is,cattle%20entered%20the%20food%20supply. Retrieved 25 May 2024.
- 25. USDA APHIS. 2024. Federal Order Requiring Testing for and Reporting of Highly Pathogenic Avian Influenza (HPAI) in Livestock.
- 26. Canadian Food Inspection Agency. 2024. Notice to industry: Highly pathogenic avian influenza (HPAI or H5N1) in dairy cattle in the USA Addendum to export certificate. https://inspection.canada.ca/en/animal-health/terrestrial-animals/diseases/reportable/avian-influenza/latest-bird-flu-situation/notice-industry-2024-04-30. Retrieved 27 May 2024.
- 27. Campbell CH, Easterday BC, Webster RG. 1977. Strains of Hong Kong Influenza Virus in Calves. J Infect Dis 135:678–680.
- 28. Mitchell CA, Walker RVL, Bannister GL. 1953. Preliminary Experiments Relating to the Propagation of Virus in the Bovine Mammary Gland. Comp Med 17:97–104.
- 29. Brown IH, Crawshaw TR, Harris PA, Alexander DJ. 1998. Detection of antibodies to influenza A virus in cattle in association with respiratory disease and reduced milk yield. 23. Vet Rec 143:637–638.
- 30. Kalthoff D, Hoffmann B, Harder T, Durban M, Beer M. 2008. Experimental Infection of Cattle with Highly Pathogenic Avian Influenza Virus (H5N1). Emerg Infect Dis 14:1132–1134.
- 31. Crawshaw TR, Brown IH, Essen SC, Young SCL. 2008. Significant rising antibody titres to influenza A are associated with an acute reduction in milk yield in cattle. Vet J 178:98–102.

- 32. Kristensen C, Jensen HE, Trebbien R, Webby RJ, Larsen LE. 2024. The avian and human influenza A virus receptors sialic acid (SA)- α 2,3 and SA- α 2,6 are widely expressed in the bovine mammary gland https://doi.org/10.1101/2024.05.03.592326.
- 33. Canadian Food Inspection Agency. 2024. Commercial milk sampling and testing for highly pathogenic avian influenza (HPAI) viral fragments in Canada. https://inspection.canada.ca/en/animal-health/terrestrial-animals/diseases/reportable/avian-influenza/latest-bird-flu-situation/commercial-milk-sampling-and-testing-hpai-vira#:~:text=Milk%20sampling%20and%20testing%20results,dairy%20cattle%20detected% 20in%20milk. Retrieved 20 June 2024.
- 34. Wight J, Rahman I, Wallace HL, Cunningham JT, Roul S, Robertson GJ, Russell RS, Xu W, Zhmendak D, Alkie TN, Berhane Y, Hargan KE, Lang AS. 2024. Avian influenza virus circulation and immunity in a wild urban duck population prior to and during a highly pathogenic H5N1 outbreak. preprint. bioRxiv https://doi.org/10.1101/2024.02.22.581693.