

AN EVALUATION OF THE EFFECTIVENESS OF CRITICAL COMPONENTS OF THE CHAIN OF SURVIVAL IN OUT-OF-HOSPITAL CARDIOPULMONARY RESUSCITATION IN NIGERIA

Authors and Affiliations

1. KOLAWOLE, Israel Kayode, (Kolawole I.K), MBBS, DA, PGD. Pall Med. Department of Anaesthesia, University of Ilorin, Ilorin, Nigeria. Email: kolawole.ik@unilorin.edu.ng ORCID ID: 0000-0001-5823-8685
2. ABDUR-RAHMAN, Lukman Olajide, (Abdur-Rahman L.O), MBBS, MPH, MD, Cert. Health Leadership and Management. Department of Surgery, University of Ilorin, Ilorin, Nigeria. Email: bolarjide@yahoo.com ORCID ID:0000-0002-7557-5824
3. KANA, Shehu Abubakar, (Kana S.A), MBBS. Department of Internal Medicine, Aminu Kano Teaching Hospital. Email: shehukana2009@yahoo.com ORCID ID: 0000-0002-9361-2684
4. AJALA, Aisha Oluwabunmi (Ajala A.O), MB.Ch.B. Department of Internal Medicine, University of Port-Harcourt Teaching Hospital, Rivers State, Nigeria. Email: ajalabunmi@gmail.com ORCID ID: 0000-0003-4418-6946
5. AJE. Akinyemi (Aje. A), MBBS. Department of Medicine, University College Hospital, Ibadan, Oyo-State, Nigeria. Email: ajeyemi@yahoo.co.uk ORCID ID: 0000-0002-5943-4220
6. ALAGBE-BRIGGS, Olubusola Temitope, (Alagbe-Briggs O.T), MBBS, MSc Med.Ed., Cert. in LMIH (UW). Department of Anaesthesiology, University of Port-Harcourt. Email: busolabriggs@yahoo.co.uk ORCID NUMBER: 0009-0002-5980-7637
7. AYOOLA, Yekeen Ayodele, (Ayoola Y.A), MBCHB. Department of Internal Medicine, Gombe State University. Email: profyekeen@yahoo.com ORCID ID: 0000000306608458
8. MBADIWE, Nkeiruka Chigekwu, (Mbadiwe N.C), B. Med Sc, MBBS, Cert. Interventional Cardiology (Hyderabad). Department of Internal Medicine, University of Nigeria, Nsuka. Email: nkeiruchimbadiwe@gmail.com ORCID ID: 0000-000-9390-6431
9. GALADANCI, Hadiza Shehu, (Galadanci H.S), MBBS, MSc. Department of Obstetrics and Gynaecology, Aminu Kano Teaching Hospital, Director Africa Centre of Excellence for population Health and Policy, Bayero University, Kano. Email: hgaladanci@yahoo.com ORCID ID: 0000-0001-5810-2884
10. LADIPO, Josephus Kayode, (Ladipo J.K), MBBS. Department of Surgery, University of Ibadan. Email: jkladipo@gmail.com ORCID ID:
11. SALAHU, Dalhat, (Salahu. D), MBBS. Department of Anaesthesia, Aminu Kano Teaching Hospital, Kano. dalhatusalahu@gmail.com ORCID ID: 0000-0003-0215-0873

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

- 1 12. SANUSI, Arinola Adeyoola, (Sanusi A.A), MBBS, DA. Department of Anaesthesia,
2 College of Medicine, University of Ibadan/University College Hospital, Ibadan, Oyo
3 State, Nigeria. Email: arinolasanusi@yahoo.com **ORCID ID: 0000-0003-1725-9097**
4 13. OGUNMODEDE, James Ayodele, (Ogunmodede J.A). MBBS. Department of
5 Medicine, University of Ilorin, Ilorin Nigeria. Email: ogunmodede.ja@unilorin.edu.ng
6 **ORCID ID: 0000000316771232**
7 14. ONYEKA, Tonia Chinyelu, (Onyeka T.C). MBBS, MSc Pain Mgt. Department of
8 Anaesthesia/Pain & Palliative Care Unit, University of Nigeria Teaching Hospital,
9 Ituku-Ozalla, Enugu, IVAN Research Institute, Enugu. Email:
10 tonia.onyeka@unn.edu.ng **ORCID ID:0000-0001-8138-3100**
11 15. YUSUF, Tanimu Sambo, (Yusuf T.S), MBBS. Department of Anaesthesia, Gombe
12 State University. Email: yusufsambo.ys@gmail.com **ORCID ID: 0000-0002-3135-**
13 **0404**
14 16. SOYANNWO, Olaitan A. (Soyannwo O.A), MBBS, DA, M.Med. Centre for
15 Palliative Care, Nigeria. Email: folait2001@yahoo.com **ORCID ID: 0000-0001-**
16 **8388-9105**
17 17. Dr Terfa Kene, MBBS, MSc Epidemiology and Medical Statistics. Ave Health Sense
18 Ltd, Abuja. Email: avehealthsense@gmail.com
19 **ORCID ID: 0000-0002-2110-2407.**

20

21 **Short title: Evaluation of Chain of survival in Nigeria**

22 **Corresponding Author:**

23 **Prof. Israel Kayode KOLAWOLE**

24 Department of Anaesthesia, Faculty of Clinical Sciences, College of Health Sciences,

25 University of Ilorin, Ilorin, Nigeria.

26 Email: kolawole.ik@unilorin.edu.ng

27 **Phone number: +2348033781032.**

28 **WORD COUNTS:**

29 Manuscript length =4,994

30 Abstract =350

31 Number of references =32

32 Number of figures and tables =6

33 **ABSTRACT**

34 **Background:** The “Chain of Survival” (CoS), provides a structure to evaluate an emergency
35 response system for out-of-hospital cardiac arrest (OHCA) in any community. This chain has
36 not been previously evaluated in an African population. The aim of this study was to assess
37 the effectiveness and efficiency of out-of-hospital CPR by evaluating the components of the
38 CoS following sudden cardiac arrest in Nigeria.

39 **Methodology:** This was a prospective multicenter descriptive cross-sectional exploratory
40 survey of epidemiology and outcome of sudden cardiac arrest (SCA) in adult Nigerians. The
41 survey was conducted in six University teaching hospitals in the 6 geopolitical zones.
42 Information concerning the patients were obtained from family member or an accompanying
43 person, and hospital emergency department personnel. Exclusion criteria included age < 18
44 years, incomplete registered data, do not attempt resuscitation (DNAR) decision, and patients
45 in whom resuscitation was not attempted. Primary outcome measures were return of
46 spontaneous circulation (ROSC), survival to hospital admission and discharge.

47 **Results:** A total of 387 OHCA patients aged between 18 and 115 years met the inclusion
48 criteria. They consisted of 62.5% males and 37.5% females, giving a male/female ratio of
49 approximately 1.7:1, and the mean age was 49.9 ± 19.15 years. Most (55.6%) of the arrests
50 occurred at home. Ninety-one-point seven percent (91.7%) of the arrests were witnessed.
51 Calls were made to ambulance services but only in a third (34.6%) did ambulance arrive. The
52 time taken for the ambulance to arrive at the call site ranged between 10 minutes and 60
53 minutes, with a mean time of arrival of 25.6 ± 20.5 minutes. Only 14.3% of ambulances had
54 facilities for both BLS interventions. Pre-hospital resuscitation was attempted in 15.2%. Pre-
55 hospital ROSC was achieved in 3.6% of all the SCA events, but only 1.8% survived to
56 hospital admission and were eventually discharged.

57 **Conclusion:**

58 The emergency response service to OHCA in Nigeria is very poor due to weaknesses in the
59 critical components of the CoS. There is a need for regular evaluation of the effectiveness of
60 the CoS to identify weaknesses, and efforts made to improve performance in every
61 component of the chain, through community wide BLS training.

62 **KEY WORDS:** Out of hospital cardiac arrest, Chain of survival, evaluation, Nigeria

63 **Non-standard Abbreviation**

64 Chain of survival (CoS)

65 **Clinical Perspective:**

66 What is new?

- 67 • This is the first large scale multi-center study in Nigeria, and probably in Africa, to
68 evaluate the effectiveness and efficiency of the components of chain of survival in
69 out-of-hospital cardiopulmonary resuscitation, reflecting the complex interplay of
70 resources, setting and bystander engagement on the clinical outcomes of cardiac
71 arrest.
- 72 • The emergency response service system for OHCA in Nigeria is very poor.
- 73 • This study revealed the shortcomings related to cardiopulmonary resuscitation due to
74 resource-limitation, which bellies the practical difficulties inherent in implementing
75 the scientific recommendations on resuscitation formulated primarily from the
76 perspective of a high-resource environment by the International Liaison Committee
77 on Resuscitation.

78 **Clinical implications**

- 79 • The shortcomings observed in cardiopulmonary resuscitation in this study calls for a
80 well-organized, coordinated deliberate efforts to strengthen all parts of the chain of
81 survival in a manner that emphasizes engagement from the community, and the

82 development of locally implemented, strong pre-hospital chain of care, which is
83 adaptable to the requirements of different cardiac emergencies.

- 84 • There should be a long-term, ongoing evaluation of the CoS to identify weaknesses
85 and efforts made to improve performance in every community through a deliberate
86 effort to ensure training on BLS for everyone (medical and non-medical), and from an
87 early age.

88

89

90

91

92

93

94

95 INTRODUCTION

96 An out-of-hospital sudden cardiac arrest (OHCA) is defined as cessation of cardiac
97 mechanical activity that occurs outside of the hospital setting and is confirmed by the absence
98 of signs of circulation.¹ It remains, a major public health challenge worldwide, with survival
99 ranging between 2 and 11%.^{2,3}

100 The term “Chain of Survival” describes a series of sequentially connected interdependent
101 links of critical interventions designed to reduce the mortality associated with cardiac arrest.
102 The concept was proposed in the 1992 CPR guidelines and provides a structure to evaluate an
103 emergency response system for OHCA patients in any community for optimal outcome.⁴ The
104 six structural components of the Chain include: early recognition and access to emergency
105 medical care, early CPR, early defibrillation, early advanced cardiac life support (ACLS),
106 integrated post-resuscitative care, and long-term rehabilitation. Strict adherence to the Chain
107 is necessary for optimal outcome. However, international resuscitation guidelines are usually
108 based on scientific recommendations, which are formulated typically from the perspective of
109 well-resourced environment, with little consideration of applicability in LMIC.

110 This is the first large scale multi-center study in Nigeria, and probably in Africa, that have
111 assessed the effectiveness and efficiency of the components of CoS in out-of-hospital CPR,
112 reflecting the complex interplay of resources, setting and bystander engagement, on the
113 clinical outcomes of cardiac arrest. The aim was to assess performance and identify gaps in
114 the components of the Chain.

115 METHODS

116 **Study setting:** Nigeria is the most populous country in Africa, and the most populous black
117 nation in the World,⁵ with an estimated population of about 206 million as at 2020.⁶ The
118 country is a multi-ethnic, multi religious and multi-cultural society, officially subdivided into
119 six geopolitical zones (Figure 1). This study was conducted in six purposefully selected

120 University teaching hospitals in the six geopolitical zones. The hospitals are, University of
121 Ilorin Teaching hospital (North Central), Aminu Kano teaching hospital (North-West),
122 University College hospital, Ibadan (South-West), Gombe State University teaching hospital
123 (North-East), University of Nigeria teaching hospital, Ituku-Ozalla, Enugu (South-East) and
124 University of Port-Harcourt teaching hospital (South-South).

125 **Design:** A sub-analysis of the data on OHCA extracted from a prospective multicenter
126 descriptive cross-sectional exploratory survey of epidemiology and outcome of sudden
127 cardiac arrest (SCA) in Nigerian adults was conducted.

128 **Participants:** All adults (≥ 18 years) who suffered OHCA and were transferred to the six
129 selected teaching hospitals from 10th November 2020 – 9th July 2022 were recruited into the
130 study.

131 **Exclusion criteria:** Sudden cardiac arrest in patients that had non-cardiac chronic terminal
132 illness like malignancy that was not in remission, incomplete data, or violation of protocol
133 regarding age limits, and patients with “do not attempt resuscitation” instructions (DNAR),
134 and patients in whom resuscitation was not attempted were excluded.

135 **Main outcome measures**

136 Main outcome measures were ROSC, survival to hospital admission and survival to hospital
137 discharge.

138 **Compliance with Ethical Standards:** The study was approved by the local ethics committee
139 of each participating hospital and the National health research ethics committee
140 (NHREC/01/01/2007-25/08/2020).

141 **STATISTICAL ANALYSIS**

142 Statistical analysis was done using descriptive statistics (frequency, proportions, means and
143 standard deviation) to summarize variables. Analysis of quantitative data was done using
144 IBM-SPSS version 25.0 (SPSS Inc., Chicago, IL, USA. Data was appropriately presented

145 using tables. Continuous variables were summarised using means and standard deviations for
146 normally distributed variables, and any between group comparisons was tested with the
147 student's t-test or ANOVA as indicated. Categorical variables like gender, causes of SCA and
148 treatment types were summarized as number (frequency tables) and percentages, and
149 compared between groups with the Chi-square test and Yate corrected and Fishers' exact
150 tests as indicated. Survival was determined as the percentage of persons who survived to
151 hospital discharge out of the number of persons with SCA. Bi-variate analysis (cross
152 tabulation) was done to have an idea of how some factors relate directly to the outcome and
153 the site of occurrence of the sudden cardiac arrest. All tests were considered significant at
154 $p < 0.05$.

155

156 **RESULTS**

157 A total of 387 OHCA brought to the selected hospitals met the inclusion criteria for analysis.
158 Table I shows the demographic and socio-cultural characteristics of the patients. There were
159 242 (62.5%) males and 145 (37.5) females, giving a male/female ratio of approximately
160 1.7:1. The age ranged between 18 years and 115 years, with a mean age of 49.9 ± 19.15 , and
161 median of 48.0 years. The patients were arbitrarily stratified into 3 age categories. The mean
162 age of males was 49.9 ± 19.35 years and the mean age of females was 49.8 ± 18.86 years.
163 Table II shows the location of arrests, witnesses to the arrest and responses to the patients.
164 Most of the arrests 215 (55.6%) occurred at home, 147 (38.0%) occurred at public places,
165 such as offices, market, roadside, party, place of worship and sports field, while 25 (6.5%)
166 cases occurred in other places, (such as en-route to hospital, peripheral hospital, delivery
167 home and others). Three hundred and fifty-five (91.7%) of the SCA were witnessed. Arrests
168 were not witnessed in 32 (8.3%) of cases. A total of 351 (90.7%) patients possessed mobile
169 phones. Calls were made to ambulance services in 26 (6.7%) of cases, but in only 9 (34.6%)

170 of these cases did ambulance arrive, of which only 3 (42.9%) had facilities for providing life
171 support interventions. The time taken for the ambulance service to arrive to the call site
172 ranged between 10 minutes and 60 minutes with a mean time of arrival of 25.6 ± 20.5 minutes
173 (median =10 minutes). Fifty-five-point six percent of the ambulances arrived between 10 and
174 19 minutes, while the remaining 44.4% arrived after 29 minutes. Ambulance did not arrive at
175 all in 17 (65.4%) of the cases, while calls were not made for ambulance services in 361
176 (93.3%) of the cases. Private or public transportation was the available means of conveying
177 the patients to hospital in 369 (95.3%) of cases, while ambulances were used in 7 (1.8%) of
178 cases, 4 (57.1%) of which had no means of providing life support interventions. Two (28.6%)
179 of the ambulances had facilities for both basic-life support (BLS) interventions and life-
180 saving drugs without a trained health personnel and only one (14.3%) of the ambulances was
181 equipped with facilities for BLS intervention, life-saving drugs, defibrillators, and personnel
182 trained to deal with emergencies. The mode of pre-hospital transportation was not specified
183 in 11 (2.8%) of the cases.

184 Table III shows the pre-hospital medical interventions. Pre-hospital intervention
185 (resuscitation) was attempted, either at the scene or during transportation to health facilities,
186 in 59 (15.2%) cases. The first monitored rhythms in the OHCA were pulseless VT in 2
187 (3.4%) victims, asystole in 1 (1.7%) and bradycardia in another (1.7%). The rhythm status
188 was not monitored for 55 (93.2%) of the cases.

189 Table IV and V show the age and gender distribution of the patients, location of arrest,
190 witnesses to arrest, who instituted resuscitation, place of post cardiac arrest care and
191 outcomes of resuscitation. Pre-hospital ROSC was achieved in 14 (23.7%) of 59 resuscitation
192 attempts, but only 7 (11.9%) patients, accounting for (50.0%) of the pre-hospital ROSC
193 survived to hospital admission and eventually survived to hospital discharge, with an overall
194 hospital discharge survival of 7 out of 387 SCA (1.8%). The remaining 7 (50.0%) pre-

195 hospital ROSC suffered re-arrest en-route hospital. Pre-hospital ROSC and survival to
196 hospital admission and discharge were significantly better in the females than males, ($p <$
197 0.05). Of all the 147 (38.0%) cases of SCA that occurred in the public places, only 23
198 (15.6%) benefited from bystander CPR interventions, which yielded 4 (17.4%) ROSC, none
199 of whom survived to hospital admission. Resuscitation was attempted in 24 (11.2%) of the
200 215 patients who had SCA at home, out of which only 3 (12.5%) achieved ROSC, but only 1
201 survived to hospital admission and was eventually discharged from hospital alive.
202 Resuscitation was attempted in 12 (48.0%) of the 25 patients who had SCA in other places,
203 apart from home and public places, (such as en-route to hospital, peripheral hospital, delivery
204 home and others), out of which 7 (58.3%) achieved ROSC, and 6 (50.0%) survived to
205 hospital admission and were eventually discharged from hospital alive. Proportionately, 12
206 (48.0%) of the arrests that occurred in places such as en-route to hospital, peripheral hospital
207 and delivery homes benefited from pre-hospital intervention compared to those who arrested
208 in other places. Survival was also significantly better in terms of pre-hospital ROSC, survival
209 to hospital admission and survival to eventual discharge from hospital in this category of
210 patients ($p = 0.0001$) compared to the other categories. Majority, 44 of 59 (74.6%) of the pre-
211 hospital CPR intervention were provided by bystanders, while only 15 (25.4) were provided
212 by trained health personnel. However, most (7 or 50.0%) of the pre-hospital ROSC were
213 from patients resuscitated by healthcare personnel and 6 (85.7%) of them survived to hospital
214 admission and were eventually discharged from hospital alive. This is unlike the 7 (50.0%)
215 pre-hospital ROSC resuscitated by bystanders, out of which only 1 (14.3%) survived to
216 hospital admission and was eventually discharged from the hospital alive. These differences
217 were statistically significant for ROSC ($p < 0.05$). The ROSC, survival to hospital admission
218 and discharge were also significantly more in the CPR provided by the trained health
219 personnel ($p < 0.05$).

220 Post cardiac arrest hospital care was done for 4 (57.1%) of the 7 admitted patients with
221 ROSC in the intensive care unit (ICU), 1 (14.3%) in the high dependency unit (HDU) and 2
222 (28.6%) in the open ward. All the patients (100.0%) survived to discharge regardless of the
223 area of post arrest care. The neurological outcome at discharge, assessed using the Cerebral
224 Performance Category (CPC) scale, showed that 4 (57.1%) of the survivors were discharged
225 with a favorable neurological outcome (CPC of 1 or 2), and 3 (42.9%) with unfavorable
226 neurological outcome (CPC of 3).

227

228 **DISCUSSION**

229 In this study, we used three working definitions of survival as described in previous studies.^{7,4}

230 These include (1) return of spontaneous circulation (ROSC), (2) successful hospital
231 admission, defined as sustained ROSC till hospital admission and (3) successful hospital
232 discharge, defined as the patient leaving the hospital alive. However, the survival rate to
233 hospital discharge is regarded as the “gold standard” for determining the effectiveness of
234 treatment of cardiac arrest.⁴

235 The overall outcomes from OHCA in this study are poor both in terms of survival, and
236 neurologic and functional ability. The rate of pre-hospital CPR intervention (15.2%) was low.

237 Despite the 23.7% of ROSC achieved from the pre-hospital CPR intervention, the overall
238 survival to hospital admission and eventual discharge from hospital was only 1.8%. This
239 finding is abysmal compared to the reports from developed countries.^{8,9} This is indicative of
240 the poor quality of pre-hospital CPR intervention and emergency care in our setting. The
241 morbidity and mortality weekly report of the Cardiac Arrest Registry to Enhance Survival
242 (CARES) – an out-of-hospital cardiac arrest surveillance, recorded an overall survival to
243 hospital discharge rate of 9.6%,¹ which according to the authors was higher than in some
244 previous reports.¹⁰ Reports put the median rates of survival to hospital discharge from OHCA

245 at 7.9%.^{11,10} However, in interpreting and comparing the findings of this study with any
246 previous report from another population, we need to consider the context and the standard of
247 care available for out-of-hospital medical emergencies in the different populations. Survival
248 from OHCA varies widely across countries and geographical regions, depending on the
249 characteristics of emergency medical services (EMS) and health care systems in the
250 community, such as treatment availability, training, and care quality.⁸ Reported outcomes of
251 OHCA in low-resource settings are usually lower than those reported in patients from high-
252 resource countries, obviously due to fewer available resources for resuscitation care
253 compared to high-resource regions.³ According to the World Health Organization, “the state
254 of cardiac arrest and trauma resuscitation is a good reflection of the standard of pre-hospital
255 and emergency care in any health system.”¹² Our findings, therefore, reflect the weaknesses
256 in community emergency medical care service system in Nigeria compared to developed
257 world.
258 Nigeria, by the World Bank classification, is a lower middle-income country. The healthcare
259 service provision is cash and carry, based on out-of-pocket payments. Consequently, access
260 to healthcare, including emergencies, remains a very big challenge, severely limited by
261 ability of the citizens to pay for care cost. The country faces major challenges in each of the
262 six areas in the links of the CoS in terms of timing, effectiveness, and efficiency of execution
263 of the components of the chain. There was poor ambulance response time and more
264 importantly, low level of community participation as shown by poor bystander CPR
265 intervention and lack of trained EMS personnel. This is due to lack of organized pre-hospital
266 emergency medical services, and basic infrastructural deficits. In fact, as clearly shown from
267 the findings of this study, it is obvious that the concept of chain of survival is not well rooted
268 in the out-of-hospital emergency medical services in Nigeria.

269 Early access, which is the first link in the Chain of Survival, is regarded as the most essential
270 component of the chain.¹³ However, where the arrest is not witnessed, as happened in 8.3%
271 of cases in this study, the link will obviously be compromised. Previous study reported
272 survival from unwitnessed cardiac arrest to be only 4% compared with 22% for witnessed
273 arrests.¹⁴ There was no record of CPR attempt in cases of unwitnessed arrest in our study.
274 Incidentally, majority (91.7%) of the OHCA events in this study were witnessed. This
275 volume of witnesses is large compared to what was reported in a previous study in South
276 Africa (68%),¹⁵ and in a morbidity and mortality weekly report of CARES in the United
277 States,¹ and in a similar study in Canada.¹⁶

278 The benefit of an arrest being witnessed is the activation of emergency response system. The
279 large volume of witnessed arrest in our study was, therefore, expected to facilitate the
280 activation of the emergency medical system, which is an important part of early access.
281 However, this is only possible where there is a dedicated nationally promoted and well
282 publicized emergency telephone number/code for such communication. Unfortunately,
283 providing toll free emergency response services through a dedicated unique telephone
284 number, like 911 in the US, is a herculean task and almost an impossibility in most
285 communities in Nigeria. This is because, a look at the displays of the emergency telephone
286 number used in many countries throughout the world did not show Nigeria on the list.¹⁷ An
287 area-wide emergency toll-free telephone numbers/code, as obtains in the UK and North
288 America, is either non-existent, or if at all exists in Nigeria, is not well publicized. Enhancing
289 emergency medical response service through a dedicated unique telephone number in all
290 communities should therefore be a top priority for Nigeria. Widespread use of dedicated
291 emergency telephone numbers has simplified and shortened access to emergency assistance
292 in most advanced/developed communities.

293 The home location of the arrests in this study ensured that 68.2% of the witnesses were
294 family members (family bystanders). However, the advantage of this to the survival of the
295 patients, in the form of early access to ensure prompt intervention, was lost in this study in
296 the absence of prompt notification of the cardiac arrest response team, and an inefficient out-
297 of-hospital emergency medical evacuation system (EMS). Nonetheless, the finding reinforces
298 the need to include family members in the education efforts and information dissemination to
299 patients at risk of sudden death in order to ensure timely and effective delivery of
300 interventions by bystanders. The various categories of arrest witnesses in this study may also
301 have contributed to the telephone calls made for ambulance services. Unfortunately, despite
302 the widespread claim of good mobile telephone penetration within the communities in
303 Nigeria, as attested to in this study with 90.7% of the victims possessing mobile phones, calls
304 could only be made for ambulance services in 6.7% of cases.

305 The available emergency medical care system, which can be activated for OHCA in Nigeria
306 are essentially hospital-based ambulance services. These out-of-hospital ambulance services
307 are operated by both public and private hospitals. Furthermore, the ambulance service system
308 in Nigeria is inefficient and bedeviled with numerous structural challenges, such as delayed
309 response time. The fact of this situation was buttressed by the findings of this study in which
310 majority (55.6%) of the ambulances in this study arrived much late; arriving between 10 and
311 19 minutes, while a sizable number (44.4%) arrived after 29 minutes. The median time of
312 ambulance arrival was 10 minutes. This is a period long enough for permanent brain damage
313 to ensue in the absence of any effective CPR intervention by the bystanders. A study in South
314 Africa, which is classified as an upper middle-income country by the World Bank, found that
315 the median response time for OHCA cases was 9 min.¹⁵ Many of the ambulances in Nigeria
316 are mere conveyor or evacuation vehicles and not necessarily designed for life-saving
317 interventions. They are often too poorly equipped and lacked appropriately trained personnel

318 and facilities to provide BLS and AED interventions required for community emergency
319 medical services. A good ambulance services must have adequate facilities for emergency
320 medical intervention. At the least, the service according to the European Resuscitation
321 Council guidelines, must have a driver with a paramedical background and a nurse, both
322 trained and qualified to perform Advanced Cardiac Life Support (ACLS), with expertise to
323 render such services as advanced airway interventions, administration of inotropic drugs and
324 use of a defibrillator.¹⁸

325 The fact that ambulance transportation occurred in only 77.8% of the 9 ambulance responses
326 suggest that, even some of the very few ambulances that responded to calls probably arrived
327 after the patients had been transported to hospital in private or commercial vehicles. In most
328 cases, transportation of patients to health facilities for emergency medical intervention is in
329 Nigeria done through private arrangements using any available vehicle from family, friends
330 or commercial taxis. This is attested to by the findings of this study, in which a high majority
331 (95.3%) of the patients were transported to the hospital by private or public transportation.
332 These vehicles were unlikely to have facilities for medical interventions since they were not
333 designed for such purpose, and the survival of such patients to the hospital cannot be
334 guaranteed. The fact that 50% of patients with pre-hospital ROSC died en-route to hospital
335 before hospital admission demonstrate inadequate facilities / or knowledge and skills of BLS
336 in the ambulance personnel or the accompanying bystanders in the alternative means of
337 transportation used to convey the patients to hospital. In other words, the resources available
338 within the community (ambulance facilities/equipment and personnel) to convey the patients
339 to hospital could not sustain the gain already made at the scene of the event.
340 Bystander CPR is the second link in the CoS and represents the only chance for a victim of
341 sudden cardiac arrest to survive.¹⁹ The time to initiation of bystander CPR, (the time from
342 collapse to initiation of CPR), could not be ascertained in this study because it was not

343 monitored. It has been demonstrated that the probability of survival approximately doubles
344 when bystanders initiate CPR before the arrival of EMS personnel.²⁰ However, the results of
345 the present study show that, the achievement of anything close to this is hampered by a
346 number of challenges in Nigeria due to poorly organized community emergency medical
347 response system, poor bystander intervention and inadequate resources in terms of personnel
348 and facilities to mount a timely and effective intervention.

349 The fact that only 21.8% of public bystanders witnessed arrests benefited from BLS
350 intervention in this study indicates poor knowledge / or willingness to perform this
351 intervention within the community. The rates of bystander CPR remain poor in most low-
352 middle income countries (LMIC).²¹ This reflects the level of community involvement in the
353 treatment of OHCA and forms an important modifiable factor in any efforts to improve the
354 survival of OHCA. It highlights the importance of deliberate, coordinated, and effective
355 community-based CPR training with efforts made to teach community members about the
356 different ways in which out-of-hospital cardiac arrest patients can present, and the need to
357 initiate CPR, and use AEDs in the event of SCA. Countries across the world have tried to
358 address the issue of low bystander intervention within the community by introducing
359 mandatory CPR training at various opportunities, levels and strata of their society.²² The
360 potency of this is verifiable with the experience of Norway in which introduction of CPR
361 training into the national school education curriculum since 1961, has succeeded in raising
362 the average survival from OHCA to about 25%.²³

363 The fear of getting involved because of the distrust of law enforcement, as identified in a
364 previous study in minority low-income neighbourhood community in the United State of
365 America,²⁴ is also relevant and constitute a huge potential barrier to willingness to perform
366 bystander CPR in the LMIC, including Nigeria. To improve bystander action in cases of
367 OHCA in Nigeria therefore, there is need to put in place appropriate laws to absolve the

368 bystander who provides reasonable aid/offer assistance to victims of SCA in the community
369 for the victim's health condition and protect him/her from prosecution if anything goes
370 wrong. Such laws will encourage laypersons to willingly help without any hesitation.
371 Although the CPR knowledge and skill of the family members and bystanders who witnessed
372 the arrest in our study could not be ascertained, the findings of the study demonstrate a poor
373 knowledge and skills of BLS within the community. This is attested to by the fact that, even
374 with the 15.2% pre-hospital CPR interventions, only a small percentage (23.7%) of pre-
375 hospital ROSC was achieved, with an eventual hospital admission and discharge survival of
376 11.9% respectively. The 1.8% overall hospital discharge survival out of the total 387 OHCA
377 events analyzed for the study is abysmally low compared to the overall survival rate of 11.2%
378 reported in the CARES study,¹ and the average survival of 25% in Norway.²³ This shows that
379 even when an arrest is witnessed, the success of CPR, if at all instituted, will depend on the
380 speed of initiation and the knowledge and skills of the operator. Deficient resuscitation skill
381 within the community is a risk factor responsible for poor outcome after out-of-hospital
382 cardiac arrest, even in witnessed arrests. However, it is good enough that some of the
383 bystanders, (family bystanders and public bystanders), initiated the first link in the chain of
384 survival by providing BLS interventions to some of the patients. The efforts of the bystanders
385 (family and public) in this study yielded an overall 15.9% of pre-hospital ROSC in all the
386 cases of their 74.6% CPR interventions. On the other hand, the 25.4% of the trained health
387 personnel-initiated CPR achieved 46.7% pre-hospital ROSC. This clearly shows the benefits
388 of knowledge and skills in achieving good outcomes in CPR. These healthcare workers, apart
389 from participating in the initial resuscitation efforts, may also have assisted in arranging
390 prompt transportation of the patient to hospital.
391 Our study revealed that, the rate of bystander-initiated CPR was substantially lower (10.3%)
392 if the OHCA happened in the patient's home compared with arrest in the public places

393 (21.8%). In fact, the proportion of BLS intervention (21.8%) in public bystander witnessed
394 arrest was twice the proportion of BLS intervention (10.3%) in family bystander witnessed
395 arrest. The low family bystander CPR is similar to the findings of previous studies.^{25,26}
396 However, contrary to their explanation that the lower rate of bystander CPR at home might in
397 some degree be explained by patients living alone, a large proportion of our patients (95.6%)
398 live with others, which is a consequent of communal living arrangement prevalent in our
399 communities. The only plausible reason that could be adduced to lower rate of bystander
400 CPR at home, therefore, would be the fact that the spouses/relatives/living partners were less
401 disposed to initiating CPR, either due to inadequate awareness, fear, or lack of skills. Reason
402 such as this has been documented previously.^{24,26}
403 The third link in the Chain of Survival is early defibrillation. Although, generally, each link
404 in the CoS is important and the relative importance of each of the links has not been clearly
405 demonstrated, rapid defibrillation is often considered to be the single most important factor in
406 determining survival from adult sudden cardiac arrest. Investigators have observed that it is
407 the interaction of early CPR with early defibrillation that is the most powerful intervention
408 for ensuring optimal patient survival.⁴ Vaillancourt et al,¹⁶ identified bystander CPR and early
409 first responder defibrillation as significant contributors to increased survival in OHCA. This
410 is because only defibrillation can reverse a ventricular fibrillation (VF) arrest. Defibrillation
411 is often provided in OHCA by bystanders or trained BLS first responders, if a defibrillator is
412 available, prior to the arrival of EMS providers. However, as revealed in this study,
413 availability of an AED is a rarity for emergency medical response in public places within the
414 community in Nigeria. The ambulances hardly have the necessary facilities and personnel
415 with requisite knowledge to provide emergency medical interventions. Many are hardly
416 equipped with automated defibrillators or never accompanied by paramedics, physicians, or
417 nurses to provide BLS-defibrillation. Of the 9 ambulances that responded for emergency

418 calls, only 3 had life-support interventions. Although one had a defibrillator the ambulance
419 personnel lacked sufficient training to make use of the AED.

420 The initial documented cardiac rhythm at the time of the arrest has been found to be an
421 important determinant of survival, with the VF or VT constituting the bulk of the patients in
422 whom CPR is beneficial following SCA.^{27,28} In a previous study by Tresch et al,²⁸ all the
423 patients who demonstrated VF as the initial out-of-hospital rhythm had the highest immediate
424 successful resuscitation and hospital survival rates with approximately 43% successfully
425 resuscitated and 21% surviving. On the other hand, patients who demonstrated asystole or
426 electromechanical dissociation had the lowest successful resuscitation and survival rates.

427 Unfortunately, rhythm identification was not possible in the large majority of patients in this
428 study, except in (6.8%) of the 59 resuscitation attempts due to lack of resources (personnel
429 and materials) for such within the community. Of these cases in which rhythm identification
430 was possible, only 2 patients (3.4%) had rhythms considered to be shockable (VF), which
431 may explain the low survival rate in our study population. The rhythms in the remaining 2
432 patients, (1.7% each of bradycardia and asystole respectively), were not shockable and have
433 usually been associated with poor prognosis, though some reports have recorded successful
434 treatment with survival to hospital discharge even in asystole cardiac arrests.^{29,30}

435 The fourth critical link in the chain for management of cardiac arrest is early effective ACLS
436 (intubation, intravenous access and intravenous medications), which is commonly provided
437 by paramedics, physicians or nurses at the scene or in-transit. As revealed by the findings of
438 our survey, the present ambulance service system in Nigeria can hardly meet this level of
439 infrastructure, facilities and personnel, which explains why the outcome is so poor. There
440 were no records of the use of definitive airway management in our study, however,
441 intravenous medications were administered in very few cases, suggesting that some patients
442 had intravenous access established for drug administration.

443 The fifth link in the CoS is integrated post-resuscitative care. A person who survives SCA is
444 usually admitted to the hospital for observation and treatment where the heart is monitored
445 closely, medicines are given to reduce the chance of another SCA, and tests are performed to
446 identify the cause of the SCA. The level of care received will depend on the facilities
447 available in the different areas of the hospital. The 2010 ACLS guidelines recommend that all
448 post-cardiac arrest patients should be managed in an ICU.³¹ As revealed in this study, some
449 of the post-cardiac arrest patients were managed in other hospital locations apart from the
450 ICU. This may be as a result of limited ICU bed capacity in some of the study sites.
451 However, it appears that the Nigeria healthcare system is not doing badly in this link of the
452 chain of survival, as all the ROSC delivered to the hospital survived till discharge regardless
453 of the area of the hospital where they received the post cardiac arrest care.
454 Good neurological outcome is generally defined as a Cerebral Performance Category (CPC)
455 ≤ 2 or a modified Ranking scale (mRS) score of ≤ 3 ; both broadly representing the ability to
456 live independently.³² The neurological outcome at discharge in this study, with just 1.0% of
457 all the SCA events achieving a favorable CPC of ≤ 2 and 0.8% CPC of 3, is very poor. This is
458 probably due to the various interplay of a complex of factors, such as delayed initiation of
459 pre-hospital CPR, quality of CPR/defibrillation and unfavorable rhythms of the arrests,
460 resulting in prolonged no flow and low flow phases of the arrests, with significant degrees of
461 brain damage.

462

463 **Limitations:**

464 There were several limitations of this study. First, the study is limited by the fact that it was a
465 descriptive cross-sectional exploratory survey and conducted only in some hospitals that were
466 purposefully selected for representation of the geopolitical zones. This could have introduced
467 some degree of selection bias. Secondly, several time-related events in the chain of survival

468 that may be associated with survival from cardiac arrest were not specifically captured in this
469 study. These time-related events include access time, (the time from collapse to the
470 summoning of emergency aid), time to initiation of cardiopulmonary resuscitation (CPR),
471 (the time from collapse to initiation of CPR) and time to definitive care, (the time from
472 collapse to provision of defibrillation, intubation, or emergency medication). Thus, the
473 prognostic significance of these time-related events could not be tested in this study.

474 **Conclusion:**

475 Despite the various limitations to the study, the strengths of our study lie in the fact that it
476 was a prospective multicenter survey, conducted in tertiary referral hospitals providing
477 inpatient and outpatient services, as well as emergency and critical care services in the six
478 geopolitical zones of the country. The study revealed the shortcomings related to
479 cardiopulmonary resuscitation due to resource-limitation in a low-resource setting like
480 Nigeria. This bellies the practical difficulties inherent in implementing the scientific
481 recommendations on resuscitation formulated primarily from the perspective of a high-
482 resource environment by the International Liaison Committee on Resuscitation (ILCOR).

483

484 **Acknowledgements**

485 We are immensely indebted to late Emeritus Professor Nimi Briggs for his assistance at every
486 stage of the project. We also acknowledge the assistance of the West African College of
487 Surgeons (WACS) and the Endowment Fund Committee (Nigeria) of WACS for the
488 encouragement, moral, technical and logistic support at every stage of the project.

489 **Source of Funding**

490 The project was supported by a grant obtained from the Tertiary Education Trust Fund
491 (TETFund) of the Federal Republic of Nigeria (Grant number TETFUND/DR&D-
492 CE/NRF/2019, BATCH 6).

493 **Disclosures**

494 None

495 **References**

- 496 1. Frieden TR, Harold Jaffe DW, Thacker SB, Moolenaar RL, LaPete MA, Martinroe JC,
497 et al. Morbidity and Mortality Weekly Report Out-of-Hospital Cardiac Arrest
498 Surveillance-Cardiac Arrest Registry to Enhance Survival (CARES), Centers for
499 Disease Control and Prevention MMWR Editorial and Production Staff MMWR
500 Editorial Board. 2011;
- 501 2. Yan S, Gan Y, Jiang N, Wang R, Chen Y, Luo Z, et al. The global survival rate among
502 adult out-of-hospital cardiac arrest patients who received cardiopulmonary
503 resuscitation: A systematic review and meta-analysis. *Crit Care*. BioMed Central Ltd.;
504 2020 Feb 22;24(1):1–13.
- 505 3. Berdowski J, Berg RA, Tijssen JGP, Koster RW. Global incidences of out-of-hospital
506 cardiac arrest and survival rates: Systematic review of 67 prospective studies.
507 *Resuscitation* [Internet]. Elsevier Ireland Ltd; 2010;81(11):1479–87. Available from:
508 <http://dx.doi.org/10.1016/j.resuscitation.2010.08.006>
- 509 4. Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular
510 Care. Part 12: from science to survival: strengthening the chain of survival in every
511 community. The American Heart Association in collaboration with the International
512 Liaison Committee on Resuscitation - PubMed [Internet]. [cited 2023 Jan 26].
513 Available from: <https://pubmed.ncbi.nlm.nih.gov/10966681/>
- 514 5. Population in Africa, by country 2020 | Statista [Internet]. [cited 2023 Jan 26].
515 Available from: [https://www.statista.com/statistics/1121246/population-in-africa-by-](https://www.statista.com/statistics/1121246/population-in-africa-by-country/)
516 [country/](https://www.statista.com/statistics/1121246/population-in-africa-by-country/)
- 517 6. Nigeria Population (2023) - Worldometer [Internet]. [cited 2023 Mar 4]. Available from:
518 <https://www.worldometers.info/world-population/nigeria-population/>
- 519 7. Eisenberg MS, Cummins RO, Damon S, Larsen MP, Hearne TR. Survival rates from
520 out-of-hospital cardiac arrest: recommendations for uniform definitions and data to
521 report. *Ann Emerg Med*. *Ann Emerg Med*; 1990;19(11):1249–59.
- 522 8. Institute of Medicine of the National Academies. Strategies to Improve Cardiac Arrest
523 Survival A Time to Act. Report Brief. 2015.
- 524 9. Vellano K, Crouch A, Rajdev M, Mcnally B. Cardiac Arrest Registry to Enhance
525 Survival (CARES) Report on the Public Health Burden of Out-of-Hospital Cardiac
526 Arrest Institute of Medicine. 2015;
- 527 10. Sasson C, Rogers MAM, Dahl J, Kellermann AL. Predictors of survival from out-of-
528 hospital cardiac arrest: a systematic review and meta-analysis. *Circ Cardiovasc Qual*
529 *Outcomes*. *Circ Cardiovasc Qual Outcomes*; 2010 Jan;3(1):63–81.
- 530 11. Nichol G, Thomas E, Callaway CW, Hedges J, Powell JL, Aufderheide TP, et al.
531 REGIONAL VARIATION IN OUT-OF-HOSPITAL CARDIAC ARREST INCIDENCE
532 AND OUTCOME. *JAMA : the journal of the American Medical Association*. NIH Public
533 Access; 2008 Sep 9;300(12):1423.

- 534 12. Aufderheide TP, Nolan JP, Jacobs IG, Van Belle G, Bobrow BJ, Marshall J, et al.
535 Global health and emergency care: a resuscitation research agenda--part 1. *Acad
536 Emerg Med. Acad Emerg Med*; 2013 Dec;20(12):1289–96.
- 537 13. Cummins R.O, Chamberlain D.A, Abramson N.S et al. Recommended guidelines for
538 uniform reporting of data from out-of-hospital cardiac arrest: the Utstein Style. A
539 statement for health professionals from a task force of the American Heart
540 Association, the European Resuscitation Council, the Heart and Stroke. *Circulation*.
541 1991;84(2):960–75.
- 542 14. Eisenberg MS, Bergner L, Hallstrom A. Cardiac Resuscitation in the Community:
543 Importance of Rapid Provision and Implications for Program Planning. *JAMA*.
544 American Medical Association; 1979 May 4;241(18):1905–7.
- 545 15. Stein C. Out-of-hospital cardiac arrest cases in Johannesburg, South Africa: A first
546 glimpse of short-term outcomes from a paramedic clinical learning database.
547 *Emergency Medicine Journal [Internet]*. 2009 Sep [cited 2023 Jun 27];26(9):670–4.
548 Available from: [https://www.researchgate.net/publication/26760936_Out-of-](https://www.researchgate.net/publication/26760936_Out-of-hospital_cardiac_arrest_cases_in_Johannesburg_South_Africa_A_first_glimpse_of_s hort-term_outcomes_from_a_paramedic_clinical_learning_database)
549 [hospital_cardiac_arrest_cases_in_Johannesburg_South_Africa_A_first_glimpse_of_s](https://www.researchgate.net/publication/26760936_Out-of-hospital_cardiac_arrest_cases_in_Johannesburg_South_Africa_A_first_glimpse_of_s hort-term_outcomes_from_a_paramedic_clinical_learning_database)
550 [hort-term_outcomes_from_a_paramedic_clinical_learning_database](https://www.researchgate.net/publication/26760936_Out-of-hospital_cardiac_arrest_cases_in_Johannesburg_South_Africa_A_first_glimpse_of_s hort-term_outcomes_from_a_paramedic_clinical_learning_database)
- 551 16. Vaillancourt, C., Stiell, IG., et al. Cardiac arrest care and emergency medical services
552 in Canada. *Can J Cardiol*. 2004;20(11):1081–90.
- 553 17. Arntz HR, Bossaert L, Carli P, Chamberlain D, Davies M, Dellborg M, et al. The pre-
554 hospital management of acute heart attacks. Recommendations of a Task Force of
555 the The European Society of Cardiology and The European Resuscitation Council.
556 *Eur Heart J. Eur Heart J*; 1998 Aug;19(8):1140–64.
- 557 18. Soar J, Nolan JP, Böttiger BW, Perkins GD, Lott C, Carli P, et al. European
558 Resuscitation Council Guidelines for Resuscitation 2015: Section 3. Adult advanced
559 life support. *Resuscitation. Elsevier*; 2015 Oct 1;95:100–47.
- 560 19. Digitalcommons@umaine D, Maine M, Jurson C. Incorporation of Civilian Care in
561 Emergency Medicine: Retainment of Training and Familiarization of Resources at the
562 University of Maine.
- 563 20. Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predicting survival from out-
564 of-hospital cardiac arrest: a graphic model. *Ann Emerg Med*. 1993 Nov
565 1;22(11):1652–8.
- 566 21. Tanaka H, Ong MEH, Siddiqui FJ, Ma MHM, Kaneko H, Lee KW, et al. Modifiable
567 Factors Associated With Survival After Out-of-Hospital Cardiac Arrest in the Pan-
568 Asian Resuscitation Outcomes Study. *Ann Emerg Med [Internet]*. *Ann Emerg Med*;
569 2018 May 1 [cited 2022 Dec 22];71(5):608-617.e15. Available from:
570 <https://pubmed.ncbi.nlm.nih.gov/28985969/>
- 571 22. Wissenberg M, Lippert FK, Folke F, Weeke P, Hansen CM, Christensen EF, et al.
572 Association of national initiatives to improve cardiac arrest management with rates of
573 bystander intervention and patient survival after out-of-hospital cardiac arrest. *JAMA*.
574 American Medical Association; 2013 Oct 2;310(13):1377–84.
- 575 23. Lancet T. Out-of-hospital cardiac arrest: a unique medical emergency. *The Lancet*
576 [Internet]. 2018 [cited 2023 Feb 3];391:911. Available from: <http://www.global>

- 577 24. Sasson C, Haukoos JS, Ben-Youssef L, Ramirez L, Bull S, Eigel B, et al. Barriers to
578 Calling 911 and Learning and Performing Cardiopulmonary Resuscitation for
579 Residents of Primarily Latino, High-Risk Neighborhoods in Denver, Colorado. *Ann*
580 *Emerg Med*. Mosby Inc.; 2015 May 1;65(5):545-552.e2.
- 581 25. Fosbøl EL, Dupre ME, Strauss B, Swanson DR, Myers B, McNally BF, et al.
582 Association of neighborhood characteristics with incidence of out-of-hospital cardiac
583 arrest and rates of bystander-initiated CPR: Implications for community-based
584 education intervention. *Resuscitation*. Elsevier Ireland Ltd; 2014 Nov 1;85(11):1512–
585 7.
- 586 26. Swor RA, Jackson RE, Compton S, Domeier R, Zalenski R, Honeycutt L, et al.
587 Cardiac arrest in private locations: Different strategies are needed to improve
588 outcome. *Resuscitation* [Internet]. Elsevier Ireland Ltd; 2003 Aug 1 [cited 2023 Jun
589 19];58(2):171–6. Available from:
590 <http://www.resuscitationjournal.com/article/S0300957203001187/fulltext>
- 591 27. Tresch DD, Thakur RK. Cardiopulmonary resuscitation in the elderly. Beneficial or an
592 exercise in futility? *Emerg Med Clin North Am*. *Emerg Med Clin North Am*;
593 1998;16(3):649–63.
- 594 28. Tresch DD, Thakur R, Hoffmann RG, Brooks HL. Comparison of outcome of
595 resuscitation of out-of-hospital cardiac arrest in persons younger and older than 70
596 years of age. *Am J Cardiol*. 1988 May 1;61(13):1120–2.
- 597 29. Alao DO, Mohammed NA, Hukan YO, Al Neyadi M, Jummani Z, Dababneh EH, et al.
598 The epidemiology and outcomes of adult in-hospital cardiac arrest in a high-income
599 developing country. [cited 2023 Jun 18]; Available from:
600 <https://doi.org/10.1016/j.resplu.2022.100220>
- 601 30. Cope AR, Quinton DN, Dove AF, Sloan JP, Dave SH. Survival from cardiac arrest in
602 the accident and emergency department. *J R Soc Med*. 1987;80(12):746–9.
- 603 31. Peberdy MA, Callaway CW, Neumar RW, Geocadin RG, Zimmerman JL, Donnino M,
604 et al. Part 9: Post-cardiac arrest care: 2010 American Heart Association Guidelines
605 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Vol. 122,
606 *Circulation*. 2010.
- 607 32. Becker LB, Aufderheide TP, Geocadin RG, Callaway CW, Lazar RM, Donnino MW, et
608 al. Primary outcomes for resuscitation science studies: A consensus statement from
609 the american heart association. *Circulation*. 2011 Nov 8;124(19):2158–77.
- 610
- 611
- 612
- 613
- 614
- 615
- 616

617
618

Table I: Demographic and Socio-Cultural Characteristics of the OHCA patients

Parameters	Frequency (N=387)	Percentage
Patient Age		
<40 years	126	32.6
40-64 Years	167	43.1
65+ Years	94	24.3
Patient gender		
Male	242	62.5
Female	145	37.5
Marital status		
Married	271	70.0
Single	76	19.6
Widowed	33	8.5
Divorced/Separated	7	1.8
Type of Residence		
Urban dwellers	319	82.4
Rural dwellers	68	17.6
Living arrangements		
Lives with others	370	95.6
Lives alone	17	4.4
Availability of Family Support		
Good: families around them all the time	225	58.1
Fair: families around them sometimes	62	16.0
Poor: don't have families around them	14	3.6
Unknown	86	22.2
Socio-economic class		
26-29: Upper (I)	37	9.6
16-25: Upper Middle (II)	182	47.0
11-15: Lower Middle (III)	97	25.1
5-10: Upper Lower (IV)	40	10.3
Unknown	31	8.0

619
620
621
622
623
624
625
626
627

628 **Table II: Location of arrest, Witness to the arrest, Possession of mobile phones,**
 629 **activation of Emergency Medical System and means of transportation of patients to**
 630 **health facilities**

Parameters	Frequency (N =387)	Percentage
Location of arrest		
Home	215	55.6
Public places – Office, Market, Roadside, Party, Place of worship, Sports field, Hotel room	147	38.0
En-route to hospital	11	2.8
Peripheral hospital	8	2.1
Delivery home	4	1.0
Unspecified	2	0.5
Arrest witnessed or not witnessed		
Arrest witnessed	355	91.7
Arrest not witnessed	32	8.3
Who witnessed?		
Family bystanders	242	62.5
Public bystanders	87	22.5
Health personnels	26	6.7
Possession of mobile phone (N =387)		
Yes	351	90.7
No	36	9.3
Activation of Emergency Medical System		
Was call made for ambulance service?		
No	361	93.3
Yes	26	6.7
Did any ambulance arrive? (N =26)		
No	17	65.4
Yes	9	34.6
Facilities inside the ambulances (N =7)		
Ambulances without life-support interventions	4	57.1
Ambulances with facilities for both BLS and life-saving drugs, without a trained health personnel	2	28.6
Ambulances with facilities for BLS, life-saving drugs, defibrillators, and trained personnel	1	14.3
Estimated Time of ambulance arrival		
10-19 minutes	5	55.6
20-29	0	-
30-39 minutes	2	22.2
40-49 minutes	0	-
≥50 minutes	2	22.2
Mean time (minutes)	25.6 ±20.5	Median =10
Type of pre-hospital transportation (N =387)		
Private or public transportation	369	95.3
Ambulances	7	1.8
Unspecified	11	2.8

631

632

633

Table III: Pre-hospital medical intervention

Parameters	Frequency	Percentage (%)
Resuscitation (N=387)		
Attempted	59	15.2
Not attempted	328	84.8
Reasons for resuscitation not attempted (n =328)		
Considered futile	134	34.6
Brought in dead	10	2.6
Others	20	5.2
Unknown	164	42.4
Who instituted resuscitation? (N =59)		
Family Bystanders	25	42.4
Public Bystanders	19	32.2
Trained health personnel	15	25.4
Number of people available to institute CPR (N =59)		
Multiple rescuer	30	50.8
Two rescuers	16	27.1
Single rescuer	13	22.0
Equipment used during resuscitation (N=59)		
Bag-mask device (AMBU Bag)	29	49.2
Airway devices:	12	20.3
Oropharyngeal/Nasopharyngeal/Laryngeal mask airway/Tracheal Tube		
ECG monitor/AED/Manual defibrillator	2	3.4
Barrier device – Pocket mask	2	3.4
Unspecified	14	23.7
First monitored rhythm		
Pulse less VT	2	3.4
Asystole	1	1.7
Bradycardia	1	1.7
Unknown	55	93.2

634

635

636

637

638

639

640

641

642

643 **Table IV: Age and gender distribution of OHCA patients, location of arrest, witnesses**
 644 **to arrest and who instituted resuscitation VS outcome**

Parameters	Frequency/Percenta	Pre-hospital	Survival to	Survival to
------------	--------------------	--------------	-------------	-------------

	ge (N =387)	ROSC n =14 (3.6%)	hospital admission n =7 (1.8%)	hospital discharge n =7 (1.8%)
Age in Years				
< 65 years (N =50 or 84.7%)	293(75.7)	11 (22.0)	5 (10.0)	5 (10.0)
≥ 65 Years (N=9 or 15.3%)	94 (24.3)	3 (33.3)	2 (22.2)	2 (22.2)
p Value	Not Significant	0.462	0.296	0.296
Gender				
Male	242 (62.5)	4 (1.7)	1 (0.4)	1 (0.4)
Female	145 (37.5)	10 (6.9)	6 (4.1)	6 (4.1)
p Value	Statistically Significant	0.007	0.023 ^y	0.023 ^y
Location of arrest				
Home N=215 (55.6%)	24 (11.2)	3 (12.5)	1 (4.2)	1 (4.2)
Public places N=147(38.0%)	23 (15.6)	4 (17.4)	0	0
Other places N=25 (6.5%)	12 (48.0)	7 (58.3)	6 (50.0)	6 (50.0)
p Value	Statistically Significant	0.006	0.001 ^y	0.001 ^y
Witnesses to arrest (N =355)				
Family members (N =242)	242 (68.2)	4 (1.7)	1 (0.4)	1 (0.4)
Public bystanders (N =87)	87 (24.5)	3 (3.4)	0 (0.0)	0 (0.0)
Trained health personnel (N =26)	26 (7.3)	7 (26.9)	6 (23.0)	6 (23.0)
p Value	Statistically Significant	0.001	0.001 ^y	0.001 ^y
Who instituted resuscitation (N =59)				
Bystanders (N =25)	44 (74.6)	7(15.9)	1 (2.3)	1 (2.3)
Family members + 19 Public =44)				
Trained health personnel (N =15)	15 (25.4)	7 (46.7)	6 (40.0)	6 (40.0)
p Value	Statistically Significant	0.020	0.001	0.001

645 *^y =Yates correction

646

647

648

649 **Table V: Outcome of resuscitation**

Outcome measures	Frequency	Percentage
Pre-hospital return of spontaneous circulation (ROSC) (N		

=59)	14	23.7
Yes		
No	45	76.3
Survival to hospital admission (N =14)		
Survived	7	50.0
Suffered re-arrest and died during transport to hospital	7	50.0
Place of Post Arrest Care in the hospital (N =7)		
ICU	4	57.1
HDU	1	14.3
Open ward	2	28.6
Overall Hospital discharge survival (N= 387)		
Survived to discharge	7	1.8
Neurological status at discharge (N =7)		
CPC 1 or 2	4	57.1
CPC 3	3	42.9

650 *CPC =Cerebral Performance Category

651



652

Figure 1: Map of Nigeria showing the distribution of participating hospitals in the six geopolitical zones.

● Sites of study.