1	Title	Page

2	Full title: Lifestyles and their relative contribution to biological aging across multiple
3	organ systems: change analysis from the China Multi-Ethnic Cohort Study
4	Yuan Zhang ^{1,#} , Dan Tang ^{2,1,#} , Ning Zhang ¹ , Yi Xiang ¹ , Yifan Hu ¹ , Wen Qian ³ , Yangji
5	Baima ⁴ , Xianbin Ding ⁵ , Ziyun Wang ⁶ , Jianzhong Yin ^{7,*} , Xiong Xiao ^{1,*} , Xing Zhao ¹
6	Affiliations:
7	¹ West China School of Public Health and West China Fourth Hospital, Sichuan
8	University, Chengdu, China;
9	² Xiamen Center for Disease Control and Prevention, Xiamen, China
10	³ Chengdu Center for Disease Control and Prevention, Chengdu, China;
11	⁴ School of Medicine, Tibet University, Lhasa, China;
12	⁵ Chongqing Municipal Centre for Disease Control and Prevention, Chongqing,
13	China;
14	⁶ School of Public Health, the Key Laboratory of Environmental Pollution Monitoring
15	and Disease Control, Ministry of Education, Guizhou Medical University, Guiyang,
16	China;
17	⁷ School of Public Health, Kunming Medical University, Kunming, China;
18	[#] Joint first authors, these authors contributed equally to this work;
19	* Co-corresponding author: Jianzhong Yin, E-mail: yinjianzhong2005@sina.com,
20	Address: No. 1168, Chunrong West Road, Yuhua Street, Chenggong District,
21	Kunming, Yunnan, 650500, China; Xiong Xiao, E-mail: xiaoxiong.scu@scu.edu.cn,
22 NO	Address: No.16, People's South Road, Chengdu, Sichuan, 610041, China. ITE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

23 Funding

This work was primarily supported by the National Natural Science Foundation of China (Grant No. 82273740). The CMEC study was funded by the National Key Research and Development Program of China (Grant No. 2017YFC0907305, 2017YFC0907300). The sponsors had no role in the design, analysis, interpretation, or writing of this article.

29 Ethics approval and consent to participate

This study was approved by the Sichuan University Medical Ethical Review Board [ID: K2016038, K2020022]. All procedures in the study were consistent with the 1964 Declaration of Helsinki and its subsequent amendments. All subjects agreeing to take part in the study signed informed written consent.

34 Abbreviations used

ALB: Albumin; ALP: Alkaline phosphatase; aMED: Alternative Mediterranean diet; 35 AST: Aspartate aminotransferase, BA: Biological age; BMI: Body mass index; CVD: 36 37 Cardiovascular disease; CMEC: China Multi-Ethnic Cohort Study; CA: Chronological age; Cr: Creatinine; FEM: Fixed effects model; GGT: y-Glutamyl 38 transpeptidase; HBA1C: Glycosylated hemoglobin; HLI: Healthy lifestyle index; 39 HDL-CH: High-density lipoprotein cholesterol; IQR: Interquartile range; KDM-BA: 40 Klemera-Doubal method of biological age; LDL-CH: Low-density lipoprotein 41 cholesterol; MCV: Mean corpuscular volume; PEF: Peak expiratory flow; PLT: 42 Platelet count; QGC: Quantile G-computation; SBP: Systolic blood pressure; TG: 43 Triglyceride; WHR: Waist-to-hip ratio. 44

46 Abstract

47 Background

Biological aging exhibits heterogeneity across multi organ systems. However, it
remains unclear how is lifestyle associated with overall and organ-specific aging and
which factors contribute most in Southwest China.

51 **Objective**

52 To examine the associations of healthy lifestyle with comprehensive and organ-

53 specific biological ages and which factors contribute most.

54 Methods

This study involved 8,396 participants who completed two surveys from the China 55 56 Multi-Ethnic Cohort (CMEC) Study. The healthy lifestyle index (HLI) was developed 57 using five lifestyle factors: smoking, alcohol, diet, exercise, and sleep. The comprehensive and organ-specific biological ages (BAs) were calculated using the 58 Klemera-Doubal method based on longitudinal clinical laboratory measurements, and 59 60 validation were conducted to select BA reflecting related diseases. Fixed effects 61 model was used to examine associations between HLI or its components and the 62 acceleration of validated BAs. We further evaluated the relative contribution of lifestyle components to comprehension and organ systems BAs using quantile G-63 64 computation.

65 **Results**

About two-thirds of participants changed HLI scores between surveys. Aftervalidation, three organ-specific BAs (the cardiopulmonary, metabolic, and liver BAs)

68	were identified as reflective of specific diseases and included in further analyses with
69	the comprehensive BA. The health alterations in healthy lifestyle index showed a
70	protective association with the acceleration of all biological ages, with a mean shift of
71	-0.19 (95%CI: -0.34, -0.03) in the comprehensive biological age acceleration. Diet
72	and smoking were the major contributors to overall negative associations of five
73	lifestyle factors with the comprehensive BA and metabolic BA accounting for 24%
74	and 55% respectively.

75 Conclusions

Healthy lifestyle changes were inversely related to comprehensive and organ-specific biological aging in Southwest China, with diet and smoking contributing most to comprehensive and metabolic BA separately. Our findings highlight the potential of lifestyle interventions to decelerate aging and identify intervention targets to limit organ-specific aging in less-developed regions.

Keywords: healthy lifestyle; biological aging; organ-specific aging; change analysis;
Klemera-Doubal method of biological age.

84 Introduction

Aging is a global issue, with about 10 percent of the world population aged 65 years 85 86 and over in 2022, and the proportion expected to be near 16 percent by 2050(1). In China, the proportion has reached 14.2% at the end of 2021(2). Aging leads to 87 88 reduced life expectancy and chronic diseases through dynamic and heterogeneous 89 changes in biological systems(3). While chronological age (CA) is widely used as a marker of aging, biological age (BA) has been constructed as a more accurate 90 indicator for biological aging(4). Aging exhibits variations across and with 91 individuals; people with the same CA can age at different rates, and distinct organ 92 systems within an individual can also age differently(5). Comprehensive aging 93 indicators like epigenetic clocks and PhenoAge, capturing different aging aspects, 94 95 have been developed(6, 7). Recent studies have focused on organ-specific aging markers to understand their aging rate variations(8, 9). The process of biological 96 aging, influenced by genetics, environment, and health behaviors, is modifiable, yet 97 research has predominantly concentrated on comprehensive biological age (10-13). 98 Nevertheless, current research on determinants of multi-organ biological aging 99 remains limited. Further investigation is crucial to deepen understanding of 100 101 comprehensive and organ-specific biological aging and to develop targeted interventions for age-related diseases of each organ system. 102

Lifestyle factors such as smoking, drinking, diet, physical activity, and sleep are recognized as contributors to many age-related diseases(14-16), and researchers have begun to explore how lifestyle factors influence the process of biological aging.

106 Studies on lifestyle and BA can be summarized into two categories, one focusing on the association of a single lifestyle factor with biological aging(17-20) because 107 108 lifestyle factors may be interrelated and have cumulative effects, the other further focusing on the impact of a combined lifestyle on biological aging. Research focusing 109 110 on healthy lifestyles has been conducted in Chinese(21, 22), American(23), and 111 British populations(24) and has found that adherence to a healthy lifestyle is consistently associated with slower biological aging. However, most studies used one-112 time measurements of lifestyle factors as exposure and could not examine the impact 113 114 of lifestyle changes on BA. In addition, research has often focused on epigenetic age and PhenoAge. The Klemera-Doubal method of biological age (KDM-BA) has been 115 shown to perform well in predicting frailty and mortality and is used to identify the 116 117 biological ages of organ systems(8, 25). Therefore, KDM-BA, based on clinical-lab data sets, offers accessible and cost-effective means for aging identification and 118 intervention. Besides, no study has been conducted to analyze further the relative 119 120 contributions of multiple lifestyle changes on various organ systems BA, which could identify important influences on the comprehensive and organ-specific biological 121 aging to prioritize interventions, especially in developing regions with limited 122 resources. Southwest China is characterized by multi-ethnicity, unbalanced internal 123 development, and rapid aging, making identifying priority intervenable aging factors 124 important. 125

126 Therefore, based on the China Multi-Ethnic Cohort Study (CMEC), this study127 aimed to develop potential BAs for organ systems using the KDM algorithm, explore

how lifestyle changes correlate with both comprehensive and organ-specific BAs, and
further investigate the relative contributions of individual lifestyle components in
southwest China.

131 Methods

132 Study Population

133 This study used data from the CMEC baseline and repeated surveys for analysis. The CMEC adopted a multi-stage stratified cluster sampling, with participants covering 134 five provinces and seven ethnic groups (Tibetan, Yi, Miao, Bai, Bouyei, Dong, and 135 136 Han) in Southwest China, representing the local regional and ethnic characteristics. The baseline survey of the CMEC was conducted between May 2018 and September 137 2019, and the repeated survey was conducted between August 2020 and June 2021. 138 139 The CMEC baseline survey included 99,556 individuals, with 10% of the sample population selected for the repeated survey. Details of the CMEC procedures and 140 methodology have been described in our previous study(26). Participants' personal 141 interviews, physical examinations, and clinical laboratory measurements were 142 completed using a uniform standardized operating procedure in both surveys. The 143 study protocol was approved by the Sichuan University Medical Ethical Review 144 Board [ID: K2016038, K2020022]. All participants in the study signed an informed 145 consent form before the investigation. 146

147 In this study, we included participants with available information on lifestyle, 148 measurements for calculating BA required, and covariates. We excluded individuals 149 with missing data to calculate BA and individuals with incomplete data on lifestyle

8

150 and covariates for participants at the baseline and repeated survey. Finally, 8396 individuals aged 30-79 were included as the full set in the longitudinal analysis after 151 152 matching (Supplementary Figure 1).

153

Definition of lifestyle factors

Considering previous studies, the healthy lifestyle index (HLI) was established 154 155 through five lifestyle factors: smoking, alcohol consumption, diet, exercise, and sleep. Data for the lifestyle factors were obtained from the questionnaire, and the simplified 156 FFQ was assessed by trained staff. 157

The definitions of a healthy lifestyle are as follows. Smoking was categorized 158 according to smoking status, and never smoking was defined as healthy. For alcohol 159 consumption, according to the findings of recent research on alcohol consumption in 160 161 Chinese populations, being healthy was defined as being a non-regular drinker (drinking frequency less than once a week)(27). For diet, since the Mediterranean diet 162 (MED) is currently the most internationally recognized healthy dietary pattern and has 163 widely and robustly beneficial effects(28), we used the alternative Mediterranean diet 164 (aMED) to evaluate dietary quality, with details shown in the supplemental material. 165 Since the HLI already contained a drinking component, we removed the drinking item 166 in the aMED, which had a score range of 7-35. We defined individuals with aMED 167 scores \geq population median as healthy diets. More details on the calculation of aMED 168 can be found in previous studies(29). For exercise, based on how often participants 169 participated in physical activity in their leisure time during the past year, regular 170 exercise ("12 times/week", "3-5 times/week," or "daily or almost every day") was 171

172 categorized as healthy(30). For sleep, a sleep duration of 7-8 hours was defined as173 healthy sleep based on previous studies(31).

174 In addition, to comprehensively evaluate the impact of overall healthy lifestyle level on outcomes, we further constructed the HLI in this study. For each lifestyle, we 175 176 assign 1 point to health and 0 points to the opposite. The HLI was calculated by 177 directly adding up the five lifestyle scores, ranging from 0-5, with a higher score representing an overall healthier lifestyle. We then transformed HLI into a 178 dichotomous variable in this study, where a score of 4-5 for HLI was considered a 179 healthy lifestyle, and a score of 0-3 was considered an unfavorable lifestyle that could 180 be improved. 181

182 Covariate Assessment

183 The selection of covariates is based on prior literature review and previously constructed directed acyclic graphs (Supplementary Figure 2). Covariates included 184 age, sex, ethnicity (majority, minority), urbanicity (rural, urban), education (No 185 schooling, primary school, Middle/high school, College/university), occupation 186 (primary industry practitioner, secondary industry practitioner, tertiary industry 187 188 practitioner, unemployed), marital (married/cohabiting, status not married/cohabitating), total energy intake (kcal/day), depression symptoms, anxiety 189 190 symptoms, menopausal status in women (premenopausal, perimenopausal, postmenopausal), beverage intake (never, former consumer, currently consuming), 191 dietary supplement intake, self-reported doctor-diagnosed diseases of diabetes, 192 cardiovascular disease (CVD), and cancer. Among these, sex, ethnicity, urbanicity, 193

and education were considered time-invariant variables, while other variables wereconsidered time-varying.

BA Construction

The comprehensive BA and BAs across multiple organ systems were estimated using 197 198 the Klemera-Doubal method(32), which is well-validated in the Chinese 199 population(33). We selected indicators for constructing BA from clinical-lab data sets measured in the baseline and repeated surveys of the CMEC, filtering based on a 200 missing rate of less than 30%. Based on previous studies, these indicators were 201 202 categorized into five systems based on the organ/system function they represent: cardiopulmonary, metabolic, liver, renal, and immune systems, as detailed in the 203 Supplementary Table 2(8, 9). Thus, we developed a comprehensive BA alongside five 204 205 organ system-specific BAs: cardiopulmonary BA, metabolic BA, liver BA, renal BA, and immune BA. The selection process was then completed following the 206 methodology used in our previous studies, which are detailed in the supplementary 207 208 materials.

After finishing the screening process, 15 measures were used in constructing the
comprehensive BA, which were systolic blood pressure (SBP), waist-to-hip ratio
(WHR), peak expiratory flow (PEF), γ-Glutamyl transpeptidase (GGT), albumin
(ALB), low-density lipoprotein cholesterol (LDL-CH), high-density lipoprotein
cholesterol (HDL-CH), triglyceride (TG), aspartate aminotransferase (AST),
creatinine (Cr), alkaline phosphatase (ALP), urea, mean corpuscular volume (MCV),
glycosylated hemoglobin (HBA1C) and platelet count (PLT). The cardiopulmonary

BA was assessed using SBP and PEF; the metabolic BA was assessed using LDL-CH, HDL-CH, HBA1C, TG and WHR; the liver BA was assessed using AST, GGT, ALP, ALB; the renal BA was assessed using Cr and urea, and the immune BA was assessed using PLT and MCV. Using the above measures, we calculated each BA in the male and female populations separately using the KDM algorithm. To quantify variation in biological aging between participants, we calculated BA acceleration, which is the difference between each BA and CA simultaneously.

BA Validation

To assess the performance of the newly developed BAs of multiple organ systems, we 224 performed a validation analysis to evaluate the relationships between them and organ-225 specific diseases. Given the short follow-up period in the CMEC study, we conducted 226 227 a cross-sectional analysis using baseline data. We separately examined the associations between the comprehension BA and CVD, diabetes, cancer, 228 cardiopulmonary BA and CVD and chronic bronchitis, the metabolic BA and CVD 229 and diabetes, the liver BA and chronic hepatitis or cirrhosis, the immune BA and 230 rheumatoid arthritis. For interpretability, we simultaneously standardized and 231 classified each BA acceleration into categories (BA acceleration < 1, BA acceleration 232 < -1, and BA acceleration > 1). We used logistic regression to assess the associations 233 between each continuous and categorized BA (|BA| acceleration $| \le 1$ serving as the 234 reference group) and the diseases. Since the values of BA acceleration of the 235 cardiopulmonary BA were small, it was not processed as categorical variables. The 236 model was additionally adjusted for five healthy lifestyle factors based on the 237

covariates. The comprehensive BA constructed in the CMEC study has been validated
to better reflect age-related disease and frailty(34). After completing the validation
analysis, we selected BAs reflecting aging and related diseases for further analysis.

241 Statistical analysis

We described changes in lifestyle between the two waves of surveys, where changes 242 243 were categorized as becoming healthier, remaining unchanged, and becoming unhealthier. For single lifestyles, remaining unchanged indicates that the factor has 244 remained healthy or unhealthy at two surveys; becoming healthier suggests that it was 245 246 unhealthy at baseline and became healthy on the repeated survey, and vice versa. For 247 the HLI scores, no change indicates the same score in both surveys, a change to healthier implies a repeated survey score higher than the baseline score, and a shift in 248 249 less healthy indicates a repeated survey score lower than the baseline score. The dichotomous HLI's change categorization was similar to the single lifestyle. The 250 baseline characteristics of participants were described across categories in the 251 252 dichotomized HLI, with continuous variables represented by median (Interquartile range, IQR) and categorical variables by proportions (Table 1). 253

We used the fixed effects model (FEM) to examine associations between each lifestyle factor and the acceleration of validated BAs for people who participated in both surveys. FEM is extensively applied in the analysis of panel data in the fields of sociology, economics, and public health research, where it serves to control for unobserved individual heterogeneity. The detailed methodology of the FEM is outlined in the supplementary materials. Because of the approximately 2-year interval

between the two CMEC waves, we assumed that the short-term effect of lifestyle 260 changes on biological aging was linear during this period. Multivariable-adjusted 261 models were constructed to account for potential confounding, including time-varying 262 and time-invariant variables and CA. We incorporated five lifestyle factors as 263 exposures simultaneously into the model, while the continuous and binary HLI were 264 entered into the model as exposures separately. Since the BAs we analyzed may 265 potentially measure overlapping aspects of human aging, we did not correct for 266 multiple comparisons (Supplementary Figure 3). 267

To further obtain the relative contribution of each lifestyle factor on validated 268 comprehension and organ systems BA, the present study used data from two waves 269 for quantile G-computation (QGC) analysis. QGC is a statistical technique for 270 271 evaluating the effects of mixture exposures, capable of discerning positive and negative influences, and is widely used in environmental epidemiology research(35). 272 Additionally, subgroup analyses for continuous and categorized HLI and each 273 validated BA were conducted across sex (male vs. female), baseline age (<60 vs. 260), 274 ethnicity (majority vs. minority), urbanicity (rural vs. urban), baseline BA acceleration 275 (<0 vs. >0), and baseline disease status (free of diabetes, CVD, and cancer vs. either 276 one). The heterogeneity between strata was assessed using the Q test ($\alpha = 0.1$ was 277 considered significant heterogeneity). 278

We performed sensitivity analyses concerning exposure definitions, confounders, and our analysis method. First, we repeated the analysis of the association between lifestyles and BAs using a standard FEM, with adjustments made only for time-

282 varying variables. Second, we used alternative common health criteria for each lifestyle factor separately, and we repeated the FEM analyses and QGC analyses by 283 284 replacing the health definition of one lifestyle factor at a time. Exposure definitions for the main analyses and sensitivity analyses are shown in the Supplementary Table 285 286 1. Third, we did not adjust for body mass index (BMI) in models because some 287 studies have suggested that BMI may be an intermediate factor between lifestyles and health outcomes. However, in the sensitivity analysis, we additionally adjusted for 288 BMI in the FEM and QGC analyses models. Statistical analyses were performed 289 290 using R Project for Statistical Computing version 4.1.1 (Vienna, Austria).

291 **Results**

292 Lifestyle changes between two waves in CMEC

293 Figure 1 shows the changes in lifestyle factors in the study population from baseline to repeated surveys. For the HLI (range), more than 60 percent of participants 294 experienced a lifestyle change. For the HLI (category), about 30 percent of 295 participants changed lifestyle categories, and the percentages were close for both 296 types. For each lifestyle factor, the proportion of individuals who remained unchanged 297 during the two surveys was higher than that of individuals who changed, with 298 smoking and drinking having the highest proportion of unchanged individuals at more 299 than 90 percent, indicating that these two lifestyle factors are less likely to change. 300 Sleep and diet behaviors had a relatively high percentage of change, accounting for 301 over one-third, and exercise had a slightly lower rate of change, nearly 30 percent. 302

303 Construction and validation of organ systems BAs

304 The results of the cross-sectional analysis between the comprehensive and organ systems BAs and diseases are presented in Supplementary Table 3. Accelerated 305 306 comprehensive BA was positively correlated with an increased risk of CVD and diabetes. The cardiopulmonary BA, the liver BA, and the metabolic BA were each 307 linked to an increased risk of their respective diseases. An increase in 308 309 cardiopulmonary BA acceleration was associated with a higher risk of cardiovascular diseases but not with chronic obstructive pulmonary disease. Per 1-SD increase of the 310 liver BA was associated with a higher risk of chronic hepatitis or cirrhosis (OR 1.28, 311 312 95% CI 1.23-1.33), and per 1-SD increase in metabolic BA corresponded to a higher risk of CVD (OR 1.21, 95% CI 1.19-1.23) and diabetes (OR 3.23, 95% CI 3.12-3.34). 313 However, no expected associations with related diseases were observed for immune 314 315 BA. Finally, we concluded that the cardiopulmonary BA, the metabolic BA, and the liver BA could reflect organ-specific disease and were included in subsequent 316 317 analyses along with comprehensive BA.

318 Characteristics of participants by HLI categories

Table 1 reports the characteristics of populations categorized by dichotomous HLI at baseline and the repeated survey. In this study, 3461 individuals had a healthy lifestyle, and individuals with an unfavorable lifestyle accounted for a higher proportion (n (%) = 4935 (58.8%)) at baseline. There was a slight decrease in the proportion of people with a healthy lifestyle at the time of the repeated survey (n (%) = 3275 (39.0%)), while 5049 (61.0%) had an unfavorable lifestyle.

325 Since the lifestyle categories of participants may change, the distribution of

326 population characteristics between baseline and repeated surveys changes slightly but is generally similar. Participants with healthy lifestyles were more likely to be female, 327 328 in urban areas, Han Chinese, with higher levels of education, less likely to be employed in primary and secondary industries, more likely to be premenopausal and 329 330 with dietary supplement intake, less likely to have anxiety, depression, and younger in 331 biological age, as shown by the lower median of BA and BA acceleration of the comprehensive, cardiopulmonary, metabolic and liver BA. For major diseases, the 332 proportion of people with a healthier lifestyle was closer to that of people with an 333 334 unfavorable lifestyle in both surveys.

Associations of lifestyle factors and HLI changes with BAs acceleration and relative contributions

337 Table 2 shows the associations of individual lifestyle factors and continuous and dichotomous HLI with all validated BAs acceleration. For comprehensive BA, a 338 339 significant negative association was found between HLI and accelerated biological 340 aging, with a mean change of -0.19 (95% CI: -0.34, -0.03) in the BA acceleration for a healthier lifestyle change. Results showed that a change of -0.15 (95% CI: -0.29, -341 342 0.00) in the comprehensive BA acceleration was associated with a shift in diet. For the cardiopulmonary BA, metabolic BA, and liver BA, the HLI was associated with a 343 reduction in each BA acceleration, although some of the results were not statistically 344 significant. Shift in smoking was related to a change in metabolic BA acceleration of -345 0.54 (95%CI: -0.97, -0.11), and the results of individual lifestyle factors and the 346 cardiopulmonary BA as well as the liver BA did not reach statistical significance. 347

348 The relative contribution of each lifestyle factor on the comprehensive and the cardiopulmonary BA, the metabolic BA, and the liver BA acceleration based on the 349 350 QGC method are shown in Figure 2. For comprehensive BA, the relative contribution of diet was 0.24. Although the relative contributions of drinking and exercise were 351 slightly higher than that of diet, they were not statistically significant. For metabolic 352 353 BA, smoking was the major contributor, weighting 0.55. For cardiopulmonary BA and liver BA, exercise and alcohol consumption were the most contributing 354 components in all lifestyle factors separately, although the results were statistically 355 significant. 356

357 Results of subgroup analysis and sensitivity analysis

Figure 3 shows the results of the subgroup analyses of the HLI range and category 358 359 with the comprehensive BA acceleration. For the HLI range, the direction of the association was consistent with the whole population, with statistical significance 360 only in females, the younger, minorities, rural residents, accelerated BA group, and no 361 362 baseline disease populations. The negative association was stronger in minorities (Heterogeneity test P = 0.028). For categorized HLI and AA, the direction of the 363 association was broadly consistent with the whole population. The continued HLI 364 results for cardiopulmonary BA showed a stronger negative association in females 365 (Heterogeneity test P = 0.045, Supplementary Figure 4). For metabolic BA and liver 366 BA, no differences were observed between different subgroups (Supplementary 367 Figure 5 and 6). 368

369 We conducted a series of sensitivity analyses to test the robustness of the results.

370 Firstly, the results of the standard FEM analysis were broadly consistent with the primary analysis (Supplementary Table 4 and Supplementary Figure 7). For the 371 372 comprehensive BA, exercise along with diet was negatively correlated with BA acceleration, and in the main analysis, exercise was borderline statistically significant. 373 While the results of some lifestyles for liver function BA changed, none were 374 375 statistically significant. Secondly, changing the definition of health for any lifestyle factor had negative association estimates with all BAs acceleration for the HLI. 376 Although the effect size of some lifestyle factors changed, most of the results 377 378 remained stable. Full results are reported in Supplementary Table 5 and Supplementary Figure 8-11. Thirdly, with additional adjustment for BMI, the 379 direction and magnitude of the associations were generally consistent with the main 380 381 analysis results (Supplementary Table 6 and Supplementary Figure 12). In summary, the sensitivity analysis results aligned with the main results and were relatively 382 383 robust.

384 Discussion

This study examined the association of lifestyle changes on biological aging in the Southwest Cohort based on two surveys. About two-thirds of the participants were observed to change their HLI scores between the two surveys. The health alterations in HLI showed a protective association between accelerated aging of comprehension and each organ system. The relative contribution of diet was the largest among all lifestyle components for the comprehensive BA, and smoking contributed the most to

the metabolic BA, respectively, suggesting that these factors are particularly crucial inspecifically decelerating the pace of biological aging.

393 In our study, the prevalence of healthy lifestyles was higher than the prevalence reported by research based on the China Kadoorie Biobank study and the China 394 Nutrition and Health Surveillance (36), with those with 4-5 lifestyle factors 395 396 accounting for 41.2% and 39.0% at baseline and repeated surveys, respectively. The relatively high prevalence of healthy lifestyles in this study may be explained by he 397 recent establishment of the CMEC and an overall trend towards healthier lifestyles, 398 399 alongside different lifestyle standards. Furthermore, we noted lifestyle changes in our research and subsequently employed the FEM for analysis based on this observation. 400 Lifestyle may be influenced by environmental factors such as urbanization and 401 402 individual characteristics such as health awareness, and thus, lifestyle may change over time, yet fewer studies have focused on this(37). The high proportion of changes 403 occurring in sleep, exercise, and diet here suggests that they may be more variable and 404 405 relatively easy to intervene. For overall health styles, there was a higher proportion of changes in lifestyle scores but fewer shifts from unhealthy to healthy or healthy to 406 unhealthy. That is, while lifestyle fluctuates over time, it is less likely to change 407 substantially, consistent with the findings of another study(14, 38). 408

409 Our study reveals that healthy lifestyle shifts can slow the pace of comprehensive 410 BA, with diet being a crucial component. Both the HLI range and category showed 411 robust protective associations. Although there are differences in lifestyle definitions 412 and aging indicators, available studies consistently find that healthier lifestyles are

413 associated with lower aging acceleration(21, 23, 39, 40). Despite limited research on how lifestyle components contribute to aging, our study provides additional evidence. 414 415 We used the widely recommended aMED as a healthy dietary criterion, which 416 includes rich plant foods, olive oil, moderate fish, and red wine, some of which may 417 slow aging and promote health(41). In our study, the association between exercise and 418 comprehensive BA was nearly statistically significant. Higher levels of exercise have been shown to be associated with a lower incidence of chronic disease and a longer 419 lifespan(42, 43). Diet and exercise may slow aging through beneficial DNA 420 421 methylation changes(44), modification of gut microbiota(45), and reducing oxidative stress and inflammation(46, 47), collectively enhancing fitness and slowing the aging 422 pace. Combining the previous randomized controlled trial with our findings, the 423 424 evidence suggested that diet and exercise are easily modifiable anti-aging factors in the real world(13). 425

Another contribution of our study is presenting evidence of different lifestyle 426 factors influencing multi-organ BA, which was constructed and validated for 427 Southwest China. The HLI still played a protective role against aging across various 428 429 organ systems, yet the components contributing most varied specifically. Biological aging varies across organ systems, and the associations with lifestyle may differ. 430 Change in smoking played a significant role in metabolic aging. The current study of 431 the British population has also identified smoking as a factor influencing aging across 432 433 multiple organ systems(9). The Coronary Artery Risk Development in Young Adults Study revealed that smoking contributed to epigenetic aging acceleration by 434

435 83.5%(39). The variation in results could be attributed to the fact that current studies offer evidence of cumulative exposure to smoking(17, 21), while our research focuses 436 437 on short-term changes. Short-term smoking behavior changes may be sufficient to impact metabolism aging but do not significantly affect the comprehensive BA. 438 439 Existing research has shown that the metabolic and immune systems can impact 440 various systems through inter-organ aging pathways, making it essential to focus on modifiable factors of metabolic system aging. Hence, we discovered that smoking 441 could serve as an intervention target for organ system aging, holding the potential to 442 443 delay the onset of diseases in specific systems, thereby extending a healthy lifespan.

In subgroup analysis, we discovered that negative associations of HLI and the 444 445 comprehensive BA acceleration varied among ethnic groups. No observed differences 446 in HLI category change between ethnic groups, potentially due to insufficient statistical power. Compared to the majority group, the minority may have lower 447 socioeconomic status and less health awareness, leading to less management of 448 lifestyle factors and, consequently, stronger associations when changes in these 449 factors occur. Given that biological aging is a cause of various long-term outcomes, 450 451 these results highlight the importance of conducting interventions among minority groups in Southwest China. 452

453 Strengths and limitations

Unlike previous studies that used one measurement of lifestyle or biological age, our
study utilized longitudinal data from two waves of the CMEC study, which measured
the longitudinal changes in lifestyle behaviors and BA of the body and multi-organ

457 systems. We also assessed the impact of alterations in healthy lifestyle factors on 458 validated BAs and further characterized the relative contributions. This 459 comprehensive approach offers the latest evidence for the early prevention and 460 targeted behavioral interventions of organ-specific aging and related diseases in 461 southwestern China.

This study has certain limitations. Firstly, our construction of multi-organ BA 462 was limited by the clinical lab data measures, preventing the development of BAs for 463 464 some organ systems like the brain. Additionally, we could not fully capture specific 465 aging processes of the immune system because of few immune-related indicators and the possible omission of critical markers. However, we used routinely detected 466 467 biomarkers to construct the BA, which are cost-effective and easily implemented. 468 Therefore, this multi-organ BA is suitable for broad adoption in primary prevention, especially in developing areas like Southwestern China. Besides, the CMEC had a 469 470 limited range of diseases collected, precluding all BAs validation. Thus, we finally 471 included only those BAs that could be validated for further analysis. Secondly, as an observational study, we established a prospective association between healthy 472 473 lifestyles and accelerated aging, yet it does not confirm a causal relationship. Thirdly, the CMEC was established in 2018 with only data from two survey waves, which 474 475 highlights the drawback of reduced statistical power of FEM. Future research could build on long-term, multi-wave follow-ups, employing FEM analysis based on 476 477 repeatedly collected data to derive more reliable causal effect estimates. Fourthly, there is no gold standard for measuring biological aging. However, the KDM-BA 478

479 method used in this study has the advantage of accurately predicting aging-related 480 outcomes, and it has been validated in our study population(34). Lastly, our study is 481 based on a population from the Southwestern China which could limit the 482 generalizability of our findings. However, the Southwestern China is characterized by 483 its diverse ethnicities and lower developmental level, with few studies addressing 484 biological aging there. This research may provide crucial information for 485 interventions in other less-developed regions.

486 Conclusions

In summary, within the Southwest China population, we observed that healthy lifestyle changes were inversely related to comprehensive and organ-specific accelerated biological aging, with diet and smoking making the most significant contributions to the comprehensive BA and metabolic BA separately. These findings underscore the potential for lifestyle interventions to slow the aging pace and identify priority intervention targets to various organ-specific aging in less-developed regions.

493 Acknowledgements

We sincerely thank all the participants and staff of the CMEC study. We gratefully acknowledge Professor Xiaosong Li, the former principal investigator of CMEC research, for his leadership and tremendous contribution to the establishment of CMEC. Professor Li passed away in 2019.

498 Authors' contributions

DT, JY and XX designed the study. YZ and DT analyzed the data and drafted thepaper. All other authors were involved in the collection of the data. All authors made

- 501 significant contributions to drafting and revising the manuscript and have approved
- 502 the final version for publication.

503 Conflicts of Interest

504 The authors declare that they have no potential conflicts of interest.

505 Data availability

- 506 Data described in the manuscript, code book, and analytic code will be made available
- 507 by contacting the corresponding author.

508 References

- ECONOMIC UNDF, AFFAIRS. S. World population prospects 2022:
 Summary of results: UN, 2023.
- 511 2. Pot AM, Oliveira D, Hoffman J. Towards healthy ageing in China: shaping
- 512 person-centred long-term care. Lancet. 2022;400(10367):1905-6. doi:
 513 10.1016/s0140-6736(22)02361-3.
- 514 3. Kennedy BK, Berger SL, Brunet A, Campisi J, Cuervo AM, Epel ES, et al.
- 515 Geroscience: linking aging to chronic disease. Cell. 2014;159(4):709-13. doi:
- 516 10.1016/j.cell.2014.10.039.
- Hamczyk MR, Nevado RM, Barettino A, Fuster V, Andrés V. Biological
 Versus Chronological Aging: JACC Focus Seminar. J Am Coll Cardiol.
 2020;75(8):919-30. doi: 10.1016/j.jacc.2019.11.062.
- 520 5. Schaum N, Lehallier B, Hahn O, Pálovics R, Hosseinzadeh S, Lee SE, et al.
- Ageing hallmarks exhibit organ-specific temporal signatures. Nature.
 2020;583(7817):596-602. doi: 10.1038/s41586-020-2499-y.

- 523 6. Jylhävä J, Pedersen NL, Hägg S. Biological Age Predictors. EBioMedicine.
- 524 2017;21:29-36. doi: 10.1016/j.ebiom.2017.03.046.
- 525 7. Li X, Ploner A, Wang Y, Magnusson PK, Reynolds C, Finkel D, Pedersen NL,
- Jylhävä J, Hägg S. Longitudinal trajectories, correlations and mortality
 associations of nine biological ages across 20-years follow-up. Elife. 2020;9.
- 528 doi: 10.7554/eLife.51507.
- Nie C, Li Y, Li R, Yan Y, Zhang D, Li T, et al. Distinct biological ages of
 organs and systems identified from a multi-omics study. Cell Rep.
 2022;38(10):110459. doi: 10.1016/j.celrep.2022.110459.
- 532 9. Tian YE, Cropley V, Maier AB, Lautenschlager NT, Breakspear M, Zalesky A.
 533 Heterogeneous aging across multiple organ systems and prediction of chronic
 534 disease and mortality. Nat Med. 2023;29(5):1221-31. doi: 10.1038/s41591535 023-02296-6.
- 536 10. Bauer SR, Newman JC. Interpreting Geroscience-Guided Biomarker Studies.
- 537 JAMA Intern Med. 2022;182(3):300-2. doi:
 538 10.1001/jamainternmed.2021.7812.
- 539 11. Belsky DW, Caspi A, Houts R, Cohen HJ, Corcoran DL, Danese A, et al.
 540 Quantification of biological aging in young adults. Proc Natl Acad Sci U S A.
 541 2015;112(30):E4104-10. doi: 10.1073/pnas.1506264112.
- 542 12. Elliott ML, Caspi A, Houts RM, Ambler A, Broadbent JM, Hancox RJ, et al.
 543 Disparities in the pace of biological aging among midlife adults of the same
 544 chronological age have implications for future frailty risk and policy. Nat

545 Aging. 2021;1(3):295-308. doi: 10.1038/s43587-021-00044-4.

- 546 13. Fiorito G, Caini S, Palli D, Bendinelli B, Saieva C, Ermini I, et al. DNA
 547 methylation-based biomarkers of aging were slowed down in a two-year diet
 548 and physical activity intervention trial: the DAMA study. Aging Cell.
 549 2021;20(10):e13439. doi: 10.1111/acel.13439.
- Han Y, Hu Y, Yu C, Guo Y, Pei P, Yang L, et al. Lifestyle, cardiometabolic
 disease, and multimorbidity in a prospective Chinese study. Eur Heart J.
 2021;42(34):3374-84. doi: 10.1093/eurheartj/ehab413.
- 553 15. Zhang YB, Chen C, Pan XF, Guo J, Li Y, Franco OH, Liu G, Pan A.
 554 Associations of healthy lifestyle and socioeconomic status with mortality and
 555 incident cardiovascular disease: two prospective cohort studies. Bmj.
 556 2021;373:n604. doi: 10.1136/bmj.n604.
- 16. Henry A, Katsoulis M, Masi S, Fatemifar G, Denaxas S, Acosta D, Garfield V,
 Dale CE. The relationship between sleep duration, cognition and dementia: a
 Mendelian randomization study. Int J Epidemiol. 2019;48(3):849-60. doi:
 10.1093/ije/dyz071.
- 561 17. Klopack ET, Carroll JE, Cole SW, Seeman TE, Crimmins EM. Lifetime
 562 exposure to smoking, epigenetic aging, and morbidity and mortality in older
 563 adults. Clin Epigenetics. 2022;14(1):72. doi: 10.1186/s13148-022-01286-8.
- 564 18. Kresovich JK, Martinez Lopez AM, Garval EL, Xu Z, White AJ, Sandler DP,
 565 Taylor JA. Alcohol Consumption and Methylation-Based Measures of
 566 Biological Age. J Gerontol A Biol Sci Med Sci. 2021;76(12):2107-11. doi:

567 10.1093/gerona/glab149.

- 568 19. Kresovich JK, Park YM, Keller JA, Sandler DP, Taylor JA. Healthy eating
 569 patterns and epigenetic measures of biological age. Am J Clin Nutr.
 570 2022;115(1):171-9. doi: 10.1093/ajcn/nqab307.
- Gao X, Huang N, Guo X, Huang T. Role of sleep quality in the acceleration of 571 20. 572 biological aging and its potential for preventive interaction on air pollution insults: Findings from UK Aging 573 the Biobank cohort. Cell. 2022;21(5):e13610. doi: 10.1111/acel.13610. 574
- 575 21. Wang C, Guan X, Bai Y, Feng Y, Wei W, Li H, et al. A machine learning-based
 576 biological aging prediction and its associations with healthy lifestyles: the
 577 Dongfeng-Tongji cohort. Ann N Y Acad Sci. 2022;1507(1):108-20. doi:
 578 10.1111/nyas.14685.
- 579 22. Liang X, Zhang WS, Jin YL, Jiang CQ, Zhu F, Cheng KK, Lam TH, Xu L.
 580 Association of Multi-Dimensional Factors with Accelerating Age and
 581 Constructing a Healthy Lifestyle Index: Guangzhou Biobank Cohort Study.
 582 Gerontology. 2023;69(6):737-47. doi: 10.1159/000528760.
- 583 23. Thomas A, Belsky DW, Gu Y. Healthy Lifestyle Behaviors and Biological
 584 Aging in the U.S. National Health and Nutrition Examination Surveys 1999585 2018. J Gerontol A Biol Sci Med Sci. 2023;78(9):1535-42. doi:
 586 10.1093/gerona/glad082.
- 587 24. Yang G, Cao X, Li X, Zhang J, Ma C, Zhang N, et al. Association of
 588 Unhealthy Lifestyle and Childhood Adversity With Acceleration of Aging

- 589 Among UK Biobank Participants. JAMA Netw Open. 2022;5(9):e2230690.
- 590 doi: 10.1001/jamanetworkopen.2022.30690.
- 591 25. Zhong X, Lu Y, Gao Q, Nyunt MSZ, Fulop T, Monterola CP, Tong JC, Larbi
 592 A, Ng TP. Estimating Biological Age in the Singapore Longitudinal Aging
 593 Study. J Gerontol A Biol Sci Med Sci. 2020;75(10):1913-20. doi:
 594 10.1093/gerona/glz146.
- 595 26. Zhao X, Hong F, Yin J, Tang W, Zhang G, Liang X, Li J, Cui C, Li X. Cohort
 596 Profile: the China Multi-Ethnic Cohort (CMEC) study. Int J Epidemiol.
- 597 2021;50(3):721-1. doi: 10.1093/ije/dyaa185.
- 598 27. Millwood IY, Im PK, Bennett D, Hariri P, Yang L, Du H, et al. Alcohol intake
 599 and cause-specific mortality: conventional and genetic evidence in a
 600 prospective cohort study of 512 000 adults in China. Lancet Public Health.
 601 2023;8(12):e956-e67. doi: 10.1016/s2468-2667(23)00217-7.
- 602 28. Mazza E, Ferro Y, Pujia R, Mare R, Maurotti S, Montalcini T, Pujia A.
- 603 Mediterranean Diet In Healthy Aging. J Nutr Health Aging. 2021;25(9):1076604 83. doi: 10.1007/s12603-021-1675-6.
- Xiao X, Qin Z, Lv X, Dai Y, Ciren Z, Yangla Y, et al. Dietary patterns and
 cardiometabolic risks in diverse less-developed ethnic minority regions:
 results from the China Multi-Ethnic Cohort (CMEC) Study. Lancet Reg
 Health West Pac. 2021;15:100252. doi: 10.1016/j.lanwpc.2021.100252.
- 609 30. Luo M, Gupta N, Holtermann A, Stamatakis E, Ding D. Revisiting the
 610 'physical activity paradox' in a Chinese context: Occupational physical activity

- and mortality in 142,302 urban working adults from the China Kadoorie
 Biobank study. Lancet Reg Health West Pac. 2022;23:100457. doi:
 10.1016/j.lanwpc.2022.100457.
- 614 31. Chaput JP, Dutil C, Featherstone R, Ross R, Giangregorio L, Saunders TJ, et
- al. Sleep duration and health in adults: an overview of systematic reviews.
- 616 Appl Physiol Nutr Metab. 2020;45(10 (Suppl. 2)):S218-s31. doi:
 617 10.1139/apnm-2020-0034.
- 618 32. Klemera P, Doubal S. A new approach to the concept and computation of
 619 biological age. Mech Ageing Dev. 2006;127(3):240-8. doi:
 620 10.1016/j.mad.2005.10.004.
- 621 33. Chen L, Zhang Y, Yu C, Guo Y, Sun D, Pang Y, et al. Modeling biological age
 622 using blood biomarkers and physical measurements in Chinese adults.
 623 EBioMedicine. 2023;89:104458. doi: 10.1016/j.ebiom.2023.104458.
- 624 34. Xiang Y, Xu H, Chen H, Tang D, Huang Z, Zhang Y, et al. Tea consumption
- and attenuation of biological aging: a longitudinal analysis from two cohort
 studies. Lancet Reg Health West Pac. 2024;42:100955. doi:
 10.1016/j.lanwpc.2023.100955.
- Keil AP, Buckley JP, O'Brien KM, Ferguson KK, Zhao S, White AJ. A
 Quantile-Based g-Computation Approach to Addressing the Effects of
 Exposure Mixtures. Environ Health Perspect. 2020;128(4):47004. doi:
 10.1289/ehp5838.
- 632 36. Sun Q, Yu D, Fan J, Yu C, Guo Y, Pei P, et al. Healthy lifestyle and life

633	expecta	ncy at ag	e 30 years	in the Chinese population:	an obs	servational study.
634	Lancet	Public	Health.	2022;7(12):e994-e1004.	doi:	10.1016/s2468-
635	2667(22	2)00110-4	•			

- 636 37. Lao XQ, Ma WJ, Sobko T, Zhang YH, Xu YJ, Xu XJ, et al. Dramatic
 637 escalation in metabolic syndrome and cardiovascular risk in a Chinese
 638 population experiencing rapid economic development. BMC Public Health.
 639 2014;14:983. doi: 10.1186/1471-2458-14-983.
- 38. Zhou YF, Song XY, Pan XF, Feng L, Luo N, Yuan JM, Pan A, Koh WP.
 Association Between Combined Lifestyle Factors and Healthy Ageing in
 Chinese Adults: The Singapore Chinese Health Study. J Gerontol A Biol Sci
 Med Sci. 2021;76(10):1796-805. doi: 10.1093/gerona/glab033.
- Kim K, Zheng Y, Joyce BT, Jiang H, Greenland P, Jacobs DR, Jr., et al.
 Relative contributions of six lifestyle- and health-related exposures to
 epigenetic aging: the Coronary Artery Risk Development in Young Adults
 (CARDIA) Study. Clin Epigenetics. 2022;14(1):85. doi: 10.1186/s13148-02201304-9.
- Simons RL, Ong ML, Lei MK, Klopach E, Berg M, Zhang Y, Philibert R, 649 40. Gibbons FX, Beach SRH. Shifts in lifestyle and socioeconomic circumstances 650 predict change-for better or worse-in speed of epigenetic aging: A study of 651 Soc 2022;307:115175. 652 middle-aged black women. Sci Med. doi: 10.1016/j.socscimed.2022.115175. 653
- 41. Wang S, Li W, Li S, Tu H, Jia J, Zhao W, et al. Association between plant-

655	based	dietary	pattern	and	biologica	l aging	trajectory	in a	a large	prospe	ctive
656	cohort	t. BMC]	Med. 20	23;21	1(1):310.	doi: 10.1	1186/s1291	6-02	23-0297	74-9.	

- 42. Watts EL, Matthews CE, Freeman JR, Gorzelitz JS, Hong HG, Liao LM,
- 658 McClain KM, Saint-Maurice PF, Shiroma EJ, Moore SC. Association of
- 659 Leisure Time Physical Activity Types and Risks of All-Cause, Cardiovascular,
- 660 and Cancer Mortality Among Older Adults. JAMA Netw Open.
- 661 2022;5(8):e2228510. doi: 10.1001/jamanetworkopen.2022.28510.
- 662 43. Kankaanpää A, Tolvanen A, Bollepalli S, Leskinen T, Kujala UM, Kaprio J,
- Ollikainen M, Sillanpää E. Leisure-Time and Occupational Physical Activity
 Associates Differently with Epigenetic Aging. Med Sci Sports Exerc.
 2021;53(3):487-95. doi: 10.1249/mss.00000000002498.
- 666 44. Galkin F, Kovalchuk O, Koldasbayeva D, Zhavoronkov A, Bischof E. Stress,
- diet, exercise: Common environmental factors and their impact on epigenetic
 age. Ageing Res Rev. 2023;88:101956. doi: 10.1016/j.arr.2023.101956.
- 669 45. Zhang L, Liu Y, Sun Y, Zhang X. Combined Physical Exercise and Diet:
 670 Regulation of Gut Microbiota to Prevent and Treat of Metabolic Disease: A
- 671 Review. Nutrients. 2022;14(22). doi: 10.3390/nu14224774.
- 46. Aleksandrova K, Koelman L, Rodrigues CE. Dietary patterns and biomarkersof oxidative stress and inflammation: A systematic review of observational and
- 674 intervention studies. Redox Biol. 2021;42:101869. doi:
 675 10.1016/j.redox.2021.101869.
- 676 47. El Assar M, Álvarez-Bustos A, Sosa P, Angulo J, Rodríguez-Mañas L. Effect

- 677 of Physical Activity/Exercise on Oxidative Stress and Inflammation in Muscle
- 678 and Vascular Aging. Int J Mol Sci. 2022;23(15). doi: 10.3390/ijms23158713.

679

680

681 Tables

682 Table 1. Characteristics of participants in baseline and repeated surveys, based

683 on the categorized HLI¹

	Baseline H	LI (category)	Follow up HLI (category)		
Chamatanistia	unfavorable	healthy	unfavorable	healthy	
Characteristic	HLI = 0	HLI = 1	HLI = 0	HLI = 1	
	N = 4935	N = 3461	N = 5121	N = 3275	
	52.00 [45.00,	49.00 [43.00,	53.75 [47.01.	50.86 [44.21.	
Age (years)	60.001	57.001	62.24]	59.141	
Female (%)	2511 (50.9)	2666 (77.0)	2685 (52.4)	2492 (76.1)	
Urbanicity (%)	1472 (29.8)	1525 (44.1)	1527 (29.8)	1470 (44.9)	
Majority (%)	2665 (54.0)	2427 (70.1)	2763 (54.0)	2329 (71.1)	
Education (%)	2000 (0 110)	2127 (7011)	2705 (5110)	2323 (111)	
No schooling	1506 (30.5)	653 (18.9)	1554 (30.3)	605 (18 5)	
Primary school	1305(26.4)	721 (20.8)	1377 (26.9)	649 (19.8)	
Middle/high school	1764(35.7)	1525(44.1)	1780(34.8)	1509(461)	
College/university	360(73)	562 (16 2)	1/00 (34.0) /10 (8 0)	512 (15.6)	
$O_{\text{countien}}(9/2)$	300 (7.3)	502 (10.2)	410 (8.0)	512 (15.0)	
Drimony in dustry					
printary industry	1993 (40.4)	820 (23.7)	2093 (40.9)	826 (25.2)	
Secondary industry					
practitioner	335 (6.8)	197 (5.7)	396 (7.7)	174 (5.3)	
Tertiony industry					
nertitioner	1699 (34.4)	1671 (48.3)	1736 (33.9)	1462 (44.6)	
Unamployed	008(184)	272 (22 2)	806 (17 5)	812 (24.8)	
Married (%)	4206 (10.4)	773(22.3)	4505 (80.7)	2066(00.6)	
Wallieu (70)	4390 (09.1)	1750 71	4393 (09.7)	2900 (90.0)	
Total energy intake (kcal/day)	1/90.55	1/39./1 [1404.02	1310.72	1402.29	
	[1394.49,	[1404.92,	[1129.17,	1970 921	
· · ·	2295.57]	2130.38]	1994.04]	18/9.82]	
BMI (kg/m^2)	24.29 [21.94,	24.12 [22.04,	24.61 [22.23,	24.34 [22.31,	
	26.79]	26.47]	27.05]	26.73]	
Depression symptom	289 (5.9)	117 (3.4)	203 (4.0)	86 (2.6)	
(%)	()	× ,			
Anxiety symptom	335 (6.8)	146 (4.2)	212 (4.1)	71 (2.2)	
(%)					
Menopausal status in					
women (%)					
Premenopausal	1042 (41.5)	1336 (50.1)	914 (34.0)	1099 (44.1)	
Perimenopausal	170 (6.8)	189 (7.1)	133 (5.0)	109 (4.4)	
Postmenopausal	1299 (51.7)	1141 (42.8)	1638 (61.0)	1284 (51.5)	
Beverage (%)					
Never	4564 (92.5)	3292 (95.1)	4746 (92.7)	3079 (94.0)	
Former consumer	23 (0.5)	16 (0.5)	27 (0.5)	14 (0.4)	
Currently	3/18(71)	153(4 4)	348 (6.8)	182 (5.6)	
consuming	546 (7.1)	155 (ד.ד)	546 (0.8)	162 (5.0)	
Dietary supplement	711(144)	722(20.0)	774(14.1)	712 (21 7)	
intake (%)	/11 (14.4)	122 (20.9)	/24 (14.1)	/12(21./)	
Major diseases					
Diabetes (%)	245 (5.0)	161 (4.7)	275 (5.4)	159 (4.9)	
CVD (%)	974 (19.7)	628 (18.1)	1054 (20.6)	602 (18.4)	
Cancer (%)	36 (0.7)	31 (0.9)	39 (0.8)	32 (1.0)	
Biological ages²	. /	. /		· /	

Comprehensive BA	52.35 [43.90,	49.15 [40.19,	54.78 [46.48,	51.67 [42.74,
Comprehensive BA	61.07]	58.61]	63.45]	61.31]
Comprehensive BA	0.00 [-2.89,	-0.70 [-3.79,	0 40 [2 66 2 48]	-0.13 [-3.21,
acceleration	3.00]	2.73]	0.40 [-2.00, 5.48]	3.25]
Cardiopulmonary BA	51.86 [45.05,	49.10 [42.48,	53.79 [47.07,	50.82 [43.97,
	60.40]	56.89]	62.14]	59.18]
Cardiopulmonary BA	0.08 [-0.47,	-0.03 [-0.65,	0.04[0.56_0.62]	-0.08 [-0.71,
acceleration	0.59]	0.55]	0.04 [-0.30, 0.02]	0.56]
Matabalia BA	52.87 [43.86,	49.42 [40.53,	55.14 [46.10,	52.29 [42.89,
Metabolic DA	61.78]	59.01]	64.29]	61.77]
Metabolic BA	0.55 [-3.86,	-0.37 [-4.41,	0 46 [3 87 4 04]	0.32 [-3.93,
acceleration	4.85]	3.86]	0.40 [-3.87, 4.94]	4.46]
Liver BA	52.42 [41.08,	49.40 [37.83,	57.61 [46.15,	54.13 [42.31,
LIVEI DA	63.96]	60.43]	68.52]	64.32]
Liver BA	0.15 [-8.16,	-0.88 [-8.50,	2.88 [-5.31,	1.11 [-6.02,
acceleration	8.46]	6.73]	11.58]	9.08]

684 BA, biological age; BMI, body mass index; CVD, cardiovascular disease; HLI,

685 healthy lifestyle indicator.

686 ¹ Data are presented as median (25th, 75th percentile) for continuous variables and

687 count (percentage) for categorical variables.

688 ² Information on each validated BA has been reported. BA acceleration is the

689 difference between each BA and CA in the same survey.

690

691 Table 2 Associations of healthy lifestyle factors and HLI with the BA acceleration

Variables	Comprehensive BA acceleration β (95%CI)	Cardiopulmonary BA acceleration β (95%CI)	Metabolic BA acceleration β (95%CI)	Liver BA acceleration β (95%CI)
HLI (range)				
Per 1-point increase	-0.11 (-0.19, -0.03)	-0.02 (-0.04, -0.01)	-0.12 (-0.22, -0.02)	-0.23 (-0.47, 0.00)
HLI (category)				
To have a healthy lifestyle	-0.19 (-0.34, -0.03)	-0.03 (-0.07, 0.00)	-0.16 (-0.36, 0.03)	-0.19 (-0.66, 0.27)
Healthy lifestyle factors				
∆Smoking	-0.13 (-0.47, 0.21)	-0.03 (-0.10, 0.04)	-0.54 (-0.97, -0.11)	-0.57 (-1.58, 0.44)
∆Alcohol	-0.17 (-0.47, 0.13)	-0.02 (-0.09, 0.04)	-0.18 (-0.56, 0.21)	-0.59 (-1.50, 0.31)
∆Diet	-0.15 (-0.29, -0.00)	-0.01 (-0.04, 0.02)	-0.18 (-0.36, 0.00)	-0.10 (-0.53, 0.33)
∆Exercise	-0.16 (-0.32, 0.00)	-0.03 (-0.07, 0.00)	-0.09 (-0.30, 0.11)	-0.05 (-0.53, 0.44)
∆Sleep	-0.02 (-0.16, 0.12)	-0.02 (-0.05, 0.01)	0.01 (-0.17, 0.19)	-0.36 (-0.78, 0.06)

692 of validated BAs

BA, biological age; HLI, healthy lifestyle indicator.

694 Δ Smoking, change in smoking status between the baseline and repeated survey;

 Δ Alcohol, change in alcohol consumption between the baseline and repeated survey;

696 Δ Diet, change in dietary quality between the baseline and repeated survey; Δ Exercise,

697 change in exercise between the baseline and repeated survey; Δ Sleep, change in sleep

698 between the baseline and repeated survey; CI, confidence interval.

Estimates were obtained using FEMs treating the BA accelerations as the dependent variables and HLI (as either continuous or as categorized) or five individual lifestyle factors as the independent variables. Models were adjusted for age, occupation, marital status, total energy intake, depression symptoms, anxiety symptoms, menopausal status in women, beverage intake, dietary supplement intake, diabetes, cardiovascular disease, cancer, sex, ethnicity, urbanicity, education, and the participants' age at baseline.

706 Figure legends

707 Figure 1. Changes in lifestyle factors from baseline to repeated survey

708 HLI, healthy lifestyle indicator.

The blue bars represent the percentage of individuals who have transitioned to healthier behaviors or improved their HLI score; the gray bars represent the percentage of individuals whose behaviors or HLI score have not changed; the orange bars represent the percentage of individuals who have transitioned to unhealthier behaviors or whose HLI score has decreased.

Figure 2 Relative contributions of five healthy lifestyle components to BAs acceleration

716 Panels: Results of the comprehensive BA acceleration, the cardiopulmonary BA

717 acceleration, the metabolic BA acceleration, and the liver BA acceleration (A–D).

718 Δ Smoking, change in smoking status between the baseline and repeated survey;

719 Δ Alcohol, change in alcohol consumption between the baseline and repeated survey;

720 Δ Diet, change in dietary quality between the baseline and repeated survey; Δ Exercise,

change in exercise between the baseline and repeated survey; Δ Sleep, change in sleep

between the baseline and repeated survey.

Estimates were obtained using QGC, which treated the BA accelerations as the dependent variables and five individual lifestyle factors as the independent variables. Models were adjusted for age, occupation, marital status, total energy intake, depression symptoms, anxiety symptoms, menopausal status in women, beverage intake, dietary supplement intake, diabetes, cardiovascular disease, cancer, sex, ethnicity, urbanicity, education, and the participants' age at baseline.

- 729 The blue bars represent results that are statistically significant in the FEM analysis,
- 730 while the gray bars represent results in the FEM analysis that were not found to be
- 731 statistically significant.

732 Figure 3 Stratified analysis of estimated associations between the HLI and the

733 comprehensive BA acceleration

All models were adjusted for age, occupation, marital status, total energy intake, depression symptoms, anxiety symptoms, menopausal status in women, beverage intake, dietary supplement intake, diabetes, cardiovascular disease, cancer, sex, ethnicity, urbanicity, education, and the participants' age at baseline, with exclusion of the stratified variable as appropriate. The boxes represent point estimations. Horizontal lines represent 95% CI.





HLI (range)

HLI (category)

Category	Ν		β (95%CI)		β (95%CI)
All	8396		-0.11 (-0.19, -0.03)		-0.19 (-0.34, -0.03)
Sex					
Male	3219		-0.02 (-0.12, 0.09)		-0.00 (-0.25, 0.25)
Female	5177		-0.16 (-0.28, -0.05)		-0.23 (-0.43, -0.03)
Heterogeneity test			P = 0.058		P = 0.157
Age (years)					
<60	6325		-0.12 (-0.21, -0.02)		-0.19 (-0.38, -0.01)
≥60	2071		-0.07 (-0.22, 0.08)		-0.09 (-0.40, 0.21)
Heterogeneity test			P = 0.616		P = 0.586
Ethnicity					
Majority	5092		-0.04 (-0.14, 0.06)		-0.10 (-0.30, 0.10)
Minority	3304		-0.22 (-0.35, -0.09)		-0.30 (-0.55, -0.05)
Heterogeneity test			P = 0.028		P = 0.225
Urbanicity					
Rural	5399		-0.15 (-0.25, -0.06)		-0.20 (-0.39, -0.01)
Urban	2997		-0.04 (-0.18, 0.09)		-0.13 (-0.40, 0.14)
Heterogeneity test			P = 0.188		P = 0.685
BA Acceleration (years)					
<0	4384		-0.06 (-0.16, 0.04)		-0.19 (-0.40, 0.01)
≥ 0	4012		-0.13 (-0.24, -0.02)		-0.13 (-0.35, 0.09)
Heterogeneity test			P = 0.332		P = 0.672
Baseline disease status					
No disease	6543		-0.14 (-0.22, -0.05)		-0.21 (-0.38, -0.03)
At least one	1853		-0.04 (-0.21, 0.13)		-0.12 (-0.46, 0.21)
Heterogeneity test			P = 0.324		P = 0.668
		-0.5 0	0.5	-0.5 0 0.5	