

1 **Antibiotic self-medication in Otuke district, northern Uganda: Prevalence and**
2 **associated factors.**

3

4 Denis Diko Adoko ^{1¶} and Rebecca Nakaziba ^{2¶*}

5

6 ¹Department of Midwifery, Lira University, Lira, Uganda

7 ²Department of Pharmacology, Lira University, Lira, Uganda

8 **Corresponding author:* rnakaziba@lirauni.ac.ug/d2rebecca1@gmail.com [RN]

9

10

11 **Abstract**

12 Antibiotic self-medication is a form of irrational drug use that contributes to antimicrobial
13 resistance, which results in increasing health care costs and morbidity and mortality rates in
14 the population. The misuse of antimicrobial agents is highly linked with the growing problem
15 of antimicrobial resistance within the population globally. Unless addressed, antibiotic self-
16 medication will drive the world back to the pre-antibiotic era, with people dying helplessly due
17 to infectious diseases. This study aimed to investigate the prevalence of antibiotic self-
18 medication and its associated factors in the Otuke District, Northern Uganda. A community-
19 based cross-sectional study was conducted in the Otuke Town Council, Otuke district. The data
20 of adults aged 18 years and above were collected using a semi-structured questionnaire, and
21 the data were coded and entered into SPSS software version 26. The data were descriptively
22 analyzed for frequencies and percentages. Bivariant and multivariant analyses were performed
23 to determine associations between the variables. Out of 385 participants, 68% self-medicated
24 with antibiotics in the past 12 months. Freedom from drug use (AOR: 3.071; 95% CI: 1.203,
25 7.876) and unregulated use of antibiotics (AOR at 95% CI: 8.288 (2.815, 24.397)) were more
26 likely to lead to ASM (p value <0.001). Other significant factors included knowledge of
27 antibiotics, previous symptom experience, previous successful treatment, long waiting hours
28 and poor staff attitudes (p value <0.05). The most common self-medicated antibiotics were
29 amoxicillin, Ampiclox and metronidazole.

30 Antibiotic self-medication in the Otuke district is very high due to the availability of medicines
31 and lack of functional drug use regulatory frameworks. The district and government of Uganda
32 should design and implement measures to mitigate this widespread antimicrobial misuse to
33 prevent the development of antimicrobial resistance.

34 **Keywords:** Antibiotic; Self-medication; Prevalence; Factors; Otuke district; Uganda.

35 **Introduction**

36 The World Health Organization (WHO) defines self-medication as “the selection and use of
37 medicines by individuals to treat self-recognized illnesses or symptoms without prescription
38 and medical supervision” (1). It is a common form of healthcare practiced in most parts of the
39 world, with over 50% of antibiotics bought and used over the counter (2). Antimicrobial self-
40 medication (ASM) leads to the irrational use of drugs, contributing to antimicrobial resistance,
41 which results in increasing health care costs coupled with increased mortality and morbidity
42 (3, 4). In Africa, the prevalence of ASM ranges from 12.1% to 93.9%, with a mean prevalence
43 of 55.7%. West Africa, as a sub-region, has the highest prevalence at 70.1%, followed by North
44 Africa at 48.1% (5). In Uganda, the prevalence of ASM is as high as 75.7% (Ocan et al., 2014).
45 The misuse of antimicrobial agents is highly linked to the growing problem of antimicrobial
46 resistance (AMR) within the population (6). According to estimates by the World Health
47 Organization (WHO), AMR could lead to 700,000 deaths annually at a global scale, and it has
48 been estimated that 10 million people could die from drug-resistant bacterial infections
49 worldwide by 2050 if the current trend of increasing AMR is not suppressed (7). Antibiotic
50 resistance is a threat to the achievements of modern medicine. Without effective antibiotics to
51 prevent and treat infections, organ transplants, chemotherapy, and surgeries such as cesarean
52 sections are becoming much more dangerous. (8). In developed countries such as Germany,
53 antibiotics are prescription only drugs (9). Unfortunately in developing countries, medicines
54 use, including antibiotics, is rarely regulated (10). In developing countries, antibiotics can be
55 readily purchased without any control, leading to more cases of antibiotic resistance (ABR)
56 (11). The prevalence of ASM in developed countries ranges from 3% to 19%, while that in
57 developing countries ranges from 24% to 73.9% (12). In Africa alone, the prevalence of ASM
58 ranges from 12.1% to 93.9%, with a mean prevalence of 55.7% (5). For instance, in a studies
59 conducted in Eritrea, Ghana, and Sudan; the prevalence of ASM was 45.1%, 36%, 71.3%

60 respectively (13-15). The main determinants of ASM include education level, age, gender, past
61 successful use, severity of illness and income (16). According to Green, Keenan (17) whereas
62 in low- and middle-income countries, past good experiences and suggestions from friends or
63 relatives promote self-medication (18). In Eritrea, the main reasons for ASM were also
64 previous successful experience and the illness being ‘not serious enough to seek medical care’
65 (13) but in Afghanistan, economic problems, lack of time to visit doctors, and ease of use were
66 cited as the main reasons for practicing ASM (19). In Pakistan, ASM is affected by male
67 gender, age, and education level (20). In a related study performed in Moshi, Kilimanjaro,
68 among university students, shortage of drugs at health facilities, long waiting times at health
69 facilities and long distances to health facilities were the main causes of ASM (12). In a similar
70 study, inability to pay for health care charges, the freedom to choose the preferred drugs, lack
71 of medical professionals, poor quality of healthcare facilities, unregulated distribution of
72 medicines and patients' misconception about physicians were factors for ASM in developing
73 countries (21). According to a study performed in Afghanistan the commonly self-medicated
74 antibiotics were penicillins, metronidazole, and ceftriaxone (19) while in Pakistan,
75 Ciprofloxacin was the most commonly used self-medicated (20). In Western Uganda,
76 Amoxicillin and metronidazole are the most commonly used antibiotics among nursing
77 students (22) whereas in northern Uganda, cotrimoxazole, amoxicillin and metronidazole are
78 the most commonly used (23).

79 Self-medication with antibiotics results in health problems such as treatment failure, an
80 increased incidence of side effects and the development of acquired AMR, which decreases the
81 effectiveness of antibiotics for treating bacterial infections and is now becoming a global health
82 problem (10). The development of antibiotics was one of the great discoveries of modern
83 medicine (24). Antibiotics fight bacteria and can cure life-threatening infectious diseases such
84 as pneumonia, for which there was previously no effective treatment (21). However, the

85 improper use of antibiotics has led to an increasing number of bacteria becoming resistant to
86 these types of drugs (25). Antimicrobial resistance (AMR) is currently increasing. It has been
87 a global public health concern for the past decade (26). The number of antibiotic-resistant
88 microorganisms is on the rise (27), making it difficult to achieve good treatment outcomes
89 during the management of infectious diseases. The costs of AMR to economies and their
90 healthcare systems are significant, affecting the productivity of patients and their caregivers
91 through prolonged hospital stays and the need for more expensive and intensive care (28).
92 Although AMR is a natural phenomenon, resistance develops more rapidly through the misuse
93 and overuse of antimicrobial agents (28, 29). Therefore, the public plays a crucial role in the
94 development or prevention of this disease. The current study aimed to determine the prevalence
95 of antibiotic self-medication in Otuke District, Northern Uganda, and the associated factors to
96 inform the government of the vice so that informed interventions can be designed.

97 **Materials and methods**

98 **Study Area**

99 The study was performed in the Otuke District. Otuke District is located within the Lango
100 subregion in Northern Uganda. Otuke district consists of many government health facilities
101 that are well distributed across all sub-counties and a total of 50 drug shops. The Otuke Town
102 Council alone has five medical clinics (*Source: Records from the National Drug Authority,*
103 *Otuke District*).

104 **Participant recruitment and procedures**

105 The study was a descriptive cross-sectional study employing quantitative methods of data
106 collection. A cross-sectional design is time-saving and inexpensive. The sample size was
107 calculated using the Kish Leslie (1965) formula assuming a proportion of 50% and yielded a

108 sample size of 385. The study was conducted among adult community members (aged 18 years
109 and above) of the Otuke District excluding those who were very sick and could not respond to
110 the questions. The study employed a cluster sampling technique. The clusters were randomly
111 selected from the Adwon-Ibuto, Barodugu, Teogini and Te-Boke cells of the Otuke Town
112 Council. The study participants were conveniently selected from each cluster and the potential
113 participants were accessed by the research team moving into their homes. The study purpose,
114 risks and benefits were clearly explained to potential participants who were required to sign an
115 informed consent form to affirm their acceptance. Those who agreed to participate in the study
116 and signed the consent form were recruited. The data were collected using a pretested
117 questionnaire developed in line with the study objectives. The tool had three parts: Part A
118 (Sociodemographic characteristics such as age, sex, and level of education), Part B (Self-
119 medication practices), and Part C (Factors associated with ASM (Health system factors,
120 individual factors, and interpersonal factors). Following the acquisition of informed consent,
121 data were collected using an interviewer-administered questionnaire. Each section of the
122 questionnaire was well explained to the respondents who then provided answers. The
123 questionnaires were checked for completeness after data collection, and those with incomplete
124 or irrelevant data were excluded from the analysis. Following data collection, the data were
125 coded and double-entered into Statistical Package for Social Sciences (SPSS) software version
126 26 and descriptively analyzed for frequencies and percentages. Bivariate and multivariate
127 analyses were performed to identify associations between the variables.

128 **Ethical issues**

129 The study was presented before the faculty of Nursing and Midwifery, Lira University and to
130 Otuke district administration for approval. Moreover, the study purpose, risks and benefits were
131 clearly explained to potential participants (adults aged 18 years and above) who were required
132 to sign an informed consent form to affirm their acceptance before recruitment into the study.

133 **Results**

134 **Socio-demographic data**

135 A total of 49.4% of the participants were aged between 18 and 28 years, 56.6% were females,
 136 54.3% stopped at the primary level of education, 53.5% were married, 65.5% were self-
 137 employed, 94.8% were members of the Lango tribe, and 44.9% were Anglicans (Table 1).

138 **Table 1: Socio-demographic characteristics of the study participants (n=385)**

Variable	Frequency(f)	Percentage (%)
Age		
18-28	190	49.4
29-40	114	29.6
41-60	64	16.6
61 and above	17	4.4
Gender		
Male	167	43.4
Female	218	56.6
Level of education		
Primary	209	54.3
Secondary	121	31.4
Tertiary	47	12.2
No education	8	2.1
Marital status		
Single	153	39.7
Married	206	53.5
Separated	26	6.8
Occupation		
Self-employed	252	65.5
Government employee	40	10.4
Employed in the private sector	30	7.8
Not employed	63	16.4
Tribe		
Lango	365	94.8
Others	20	5.2
Religion		
Anglican	173	44.9
Seventh-Day Adventist	16	4.2
Islam	8	2.1
Catholic	155	40.3
Pentecostal	31	8.1
Others	2	0.5

139

140 **Prevalence of antibiotic self-medication**

141 The majority of the community members of the Otuke Town Council (68%) practiced self-
142 medication (Fig 1)

143 **Fig. 1: Prevalence of antibiotic self-medication in Otuke district (n=385).**

144 **Commonly self-medicated antibiotics and their sources**

145 The most common self-medicated antibiotic was amoxicillin (51.3%). Most of the respondents
146 (98.5%) obtained antibiotics from drug shops/pharmacy (Table 2).

147 **Table 2: Commonly self-medicated antibiotics and their sources (n=385)**

Variable	Frequency(f)	Percentage (%)
<i>Self- medicated antibiotics</i>		
Amoxicillin	134	51.3
Ampiclox	87	33.3
Cloxacillin	24	9.2
Metronidazole	57	21.9
Erythromycin	28	10.8
Azithromycin	13	5.0
Cotrimoxazole	21	8.1
Ciprofloxacin	42	16.1
Doxycycline	24	9.2
Other antibiotics	35	13.5
<i>Sources of antibiotics for self-medication</i>		
Drug shop/pharmacy	257	98.5
Friends/relatives	3	1.1
Previous remainder	1	0.4
Others	00	0.0

148 **Factors associated with Antibiotic self-medication**

149 **Health system, individual and interpersonal factors**

150 Among the health system factors, individual factors and interpersonal factors; long waiting
151 hours (82.9%), previous successful treatment (55.6%), and suggestions by friends/relatives
152 (24.4%) had the greatest effects respectively (Table 3).

153

154

155 **Table 3: Health system, individual and interpersonal factors associated with ASM**

Variable	Frequency(f)	Percentage (%)
Health system factors		
Poor staff attitude	110	28.6
Health facility too far	55	14.3
Long waiting hours	319	82.9
No drugs at the health facility	268	69.9
Lack of medical professional	35	9.1
Unregulated distribution of antibiotics	89	23.1
Individual factors		
Previous symptom experience	207	53.8
Previous successful treatment	214	55.6
Illness was not serious,	211	54.8
Storage of antibiotics	109	28.3
Lack of time to visit doctors	70	18.2
Misconception about health workers	21	5.5
Inability to pay for the health charges	48	12.5
Freedom to choose the drugs	177	46.1
Knowledge on the antibiotics	157	40.8
Old prescription	113	29.4
Interpersonal factors		
Given by a friend	43	11.2
Suggestion by a relative/friend	94	24.4
Drug brought by health worker	41	10.6

156

157 **Associations between socio-demographic characteristics with ASM**

158 There was no significant association between socio-demographic characteristics and antibiotic
 159 self-medication in the present study (Table 4).

160 **Table 4: Association of sociodemographic characteristics with ASM (at 95% CI)**

Variables	Antibiotic Self-medication: Freq (%)		X ²	df	p value
	Yes	No			
Age					
18-28	127(48.7)	63(50.8)			
29-40	80(30.7)	34(27.4)	1.369	3	0.694
41-60	41(15.7)	23(18.5)			
61 and above	13(5.0)	4(3.2)			
Gender					
Male	114(43.7)	53(42.7)	0.030	1	0.862
Female	147(56.6)	71(57.3)			

Level of education					
Primary	137(52.5)	72(58.1)			
Secondary	78(29.9)	43(34.7)	7.483	3	0.862
Tertiary	39(14.9)	8(6.5)			
No education	7(2.7)	1(0.8)			
Marital status					
Single	107(40.2)	48(38.7)			
Married	140(53.6)	66(53.2)	0.517	2	0.772
Separated	16(6.1)	10(8.1)			
Occupation					
Self-employed	164(62.8)	88(71.0)			
Government employee	30(11.5)	10(8.1)	2.655	3	0.448
Employed in the private sector	21(8.0)	9(7.3)			
Not employed	46(17.6)	17(13.7)			
Tribe					
Lango	246(94.3)	119(96.0)	0.502	1	0.479
Others	15(5.7)	5(4.0)			
Religion					
Anglican	121(69.9)	52(30.1)			
Seventh-Day Adventist	12(75.0)	4(25.0)			
Islam	6(75.0)	2(25.0)	3.511	5	0.622
Catholic	102(65.8)	53(34.2)			
Pentecostal	18(58.1)	13(41.9)			
Others	2(0.5)	0(100)			

161

162 **Associations between health system factors with ASM**

163 Most health system factors were strongly associated with antibiotic self-medication (p value
164 <0.001) (Table 5).

165 **Table 5: Association of health system factors with ASM**

Variables	Antibiotic Self-medication: Freq (%)		df	COR at 95% CI	p value
	Yes	No			
Poor staff attitude			1	12.750(6.208,26.184)	<0.001
No	159(57.8)	116(42.2)			
Yes	102(92.7)	8(7.3)			
Health facility too far			1	1.661(1.330,2.076)	<0.001*
No	206(62.4)	124(37.6)			
Yes	55(100)	0(0.0)			
Long waiting hours			1	15.500(5.639,42.604)	<0.001*

No	62(93.9)	4(6.1)			
Yes	199(62.4)	120(37.6)			
No drugs at the health facility			1	2.079(1.412,3.061)	<0.001*
No	79(67.5)	38(32.5)			
Yes	182(67.9)	86(32.1)			
Lack of medical professional			1		0.998
No	226(64.6)	124(35.4)			
Yes	35(100)	0(0.0)			
Unregulated distribution of antibiotics			1	3.852(2.907,5.106)	<0.001*
No	235(79.4)	61(20.6)			
Yes	28(29.2)	63(70.8)			

166

167 **Associations between interpersonal factors with ASM**

168 There was a very strong association between most interpersonal factors and self-medication (p
169 value<0.05) (Table 6).

170 **Table 6: Association between interpersonal factors with ASM**

Variables	Antibiotic Self-medication: Freq (%)		df	COR at 95% CI	p value
	No	Yes			
Given by a friend			1	1.758(1.410,2.192)	<0.001*
No	124(36.3)	218(63.7)			
Yes	0(0.0)	43(100)			
Suggestion by a relative/friend			1	93.000(12.963,667.189)	<0.001*
No	123(42.3)	168(57.7)			
Yes	1(1.1)	93(98.9)			
Drug brought by health worker			1	0.640(0.591,0.692)	0.997
No	124(36.0)	220(64.0)			
Yes	0(0.0)	41(100)			

171

172 **Associations of individual factors with ASM**

173 As indicated in Table 7 below, there was a strong association between most of the individual
174 factors and ASM (p value <0.001).

175

176 **Table 7: Association of individual factors with ASM**

Variables	Antibiotic Self-medication: Freq (%)		df	COR at 95% CI	p value
	No	Yes			
Previous symptom experience			1	3.684(2.574,5.273)	<0.001*
No	38(21.3)	140(78.7)			
Yes	86(41.5)	121(58.5)			
Previous successful treatment			1	0.462(0.334,0.637)	<0.001*
No	117(68.4)	54(31.6)			
Yes	7(3.3)	207(96.7)			
Illness was not serious			1	0.813(0.603,1.095)	0.173
No	96(55.5)	78(44.8)			
Yes	28(13.3)	183(86.7)			
Storage of antibiotics			1	1.226(0.967,1.554)	0.092
No	124(44.9)	152(55.1)			
Yes	0(0.00)	109(100)			
Lack of time to visit doctors			1	1.540(1.229,1.931)	<0.001*
No	124(39.4)	191(60.6)			
Yes	0(0.00)	70(100)			
Misconception about health workers			1	1.935(1.558,2.404)	<0.001*
No	124(34.1)	240(65.9)			
Yes	0(0.0)	21(100)			
Inability to pay for the health charges			1		0.997
No	124(36.8)	213(63.2)			
Yes	0(0.0)	48(100)			
Freedom to choose the drugs			1	0.580(0.428,0.788)	<0.001*
No	12(5.8)	195(94.2)			
Yes	112(63.3)	65(36.7)			
Knowledge on the antibiotics			1	5.162(3.630,7.340)	<0.001*
No	37(16.2)	191(83.8)			
Yes	87(55.4)	70(44.6)			
Old prescription			1		0.996
No	124(45.6)	148(54.4)			
Yes	0(0.0)	113(100)			

177

178 **Legend:** CI- confidence interval * significant p value

179 **Multivariate analysis of the association of factors with ASM**

180 The results from multivariate analysis of variables showed that freedom to choose drugs (AOR:
 181 3.071; 95% CI (1.203, 7.876)) and unregulated use of antibiotics (AOR at 95% CI: 8.288
 182 (2.815, 24.397)) were more likely to lead to ASM (p value <0.001). Other significant factors

183 included knowledge of antibiotics, previous symptom experience, previous successful
 184 treatment, long waiting hours and poor staff attitudes (p value <0.05) (Table 8).

185 **Table 8: Association between factors and antibiotic self-medication**

Variable	Bivariate p value	df	AOR at 95% CI	p value
Knowledge on the antibiotics No Yes	<0.001	1	1.00 3.079(1.203,7.876)	0.019*
Freedom to choose the drugs No Yes	<0.001	1	1.00 6.085(2.277,16.262)	<0.001*0.124
Misconception about health workers No Yes	<0.001	1	1.00 2.810(0.456,17.319)	0.266
Lack of time to visit doctors No Yes	<0.001	1	1.00 0.312(0.072,1.343)	0.118
Previous symptom experience No Yes	<0.001	1	1.00 8.212(2.688,0.476)	0.008*
Previous successful treatment No Yes	<0.001	1	1.00 0.040(0.013,0.124)	0.002*
Given by a friend No Yes	<0.001	1	1.00 1.785(0.476,6.688)	0.390
Suggestion by a relative/friend No Yes	<0.001	1	1.00 0.305(0.074,1.261)	0.101
Unregulated use of antibiotics No Yes	<0.001	1	1.00 8.288(2.815,24.397)	0.001*
No drugs at the health facility No Yes	<0.001	1	1.00 1.093(0.400,2.986)	0.862
Long waiting hours No Yes	<0.001	1	1.00 21.616(3.776,123.759)	0.006*
Health facility too far No Yes	<0.001	1	1.00 0.633(0.186,2.157)	0.465
Poor staff attitude No Yes	<0.001	1	1.00 0.052(0.004,0.622)	0.020

186 Legend: CI- confidence interval * significant p value

187 **Discussion**

188 ASM is one of the most common causes of AMR worldwide; however, it can be controlled if
189 appropriate measures are taken. The ability to develop preventive measures can only be made
190 possible by tracking the self-medication behaviors of people in the public. Additionally, it
191 reveals the health system's weaker areas. To ascertain the prevalence of ASM in the Otuke
192 District and to gain insight into the contributing factors of ASM, the current cross-sectional
193 study was carried out.

194 Of the 385 participants, 68% (over two-thirds) reported having self-medicated with antibiotics
195 in the past 6 months. This implies that approximately 7 in 10 community members practiced
196 ASM, with 49.4% of ASM being in the age bracket 18-28 years. This could be attributed to the
197 fact that there are no regulations in place to curb the practice, so every member of the
198 community is free to buy antibiotics from the many established drug shops or pharmacies. This
199 could also be the result of the long waiting hours experienced when patients seek health care
200 services in various health facilities. The findings of this study are closely related to those of a
201 similar study in Sudan, where the prevalence of ASM was 71% (15); and to another in northern
202 Uganda, which was 73% (23); and in Afghanistan, at 73.2% (19). Against this background, a
203 way forward is worth seeking to reduce the practice of ASM among the community. On the
204 other hand, a similar study conducted among nursing students in Western Uganda showed a
205 much greater prevalence of ASM, at 85.7% (22). This could be due to the pharmacological
206 knowledge gained by the nursing students. In contrast, related studies in the Middle East and
207 South America revealed lower prevalence rates of ASM, at 36.1% to 45.8% and 29%,
208 respectively (12). This could be due to better drug use regulations that guide antimicrobial use
209 in these countries, whereby antibiotics are prescription only drugs.

210 A variety of factors were analyzed for their association with ASM in this study grouped into
211 health system, individual and interpersonal factors. The findings indicated that freedom to
212 choose drugs (AOR: 3.071; 95% CI (1.203, 7.876)) and unregulated use of antibiotics (AOR
213 at 95% CI: 8.288 (2.815, 24.397)) were more likely to lead to ASM (p value <0.001). This is
214 true because in Uganda, the use of antibiotics is not highly regulated, so members of the public
215 have free access to these antibiotics. This finding is similar to that of Sachdev and others who
216 found that unregulated deliveries were the major driver of ASM (30). Therefore, the enactment
217 of drug regulation laws could be the beginning of an end to ASM. Previous symptom
218 experience was also significantly associated with ASM probably because individuals tend to
219 associate a given symptom with a disease. Therefore, a symptom managed successfully in the
220 past is deemed manageable in the same way in the future. This is in agreement with findings
221 of a research conducted in Pakistan (31). Although many symptoms may be successfully
222 managed by a specific drug, different conditions can manifest in a similar way, leading to the
223 irrational use of antibiotics. Communities should be aware that some diseases mimic each other
224 and that the use of a given antibiotic could be more of a disaster than a blessing. In addition,
225 previous successful treatment was highly significantly associated with ASM. This could have
226 been due to the patient's ability to remember how he/she was managed. This finding is
227 consistent with a study performed in low- and middle-income countries (LMICs) in which the
228 habit of self-medication using previous treatments was identified (18). There is a need for more
229 patient and family sensitization to the dangers of practicing ASM. Furthermore, long waiting
230 hours in health facilities were reported by multiple respondents in the present study were
231 associated with ASM. Many respondents thought it was a waste of time going to a health
232 facility at 8:00 am and returning home late in the afternoon. Therefore, they end up leaving the
233 facility in search of quicker services at drug stores. In Ethiopia, a similar study supported the
234 present findings where long waiting times contributed to self-medication (32). Increased

235 staffing could reduce the time spent at the facility during health seeking and improve patient
236 trust. Moreover, poor staff attitude was another factor that was significantly associated with
237 ASM in this study. Staff attitudes in terms of being rude, asking for money or not willing to
238 help caused the community members to go elsewhere and buy their own drugs without
239 consultation. This finding was consistent with those from studies in Cameroon and LMICs (33,
240 34). Regular staff motivation and appraisal could instill and sustain the drive to serve the
241 medical staff, which could improve medication practices with antibiotics and prevent ASM.

242 Of the 385 participants, 98.5% bought antibiotics from drug shops/pharmacies. This study also
243 explored the common self-medicated antibiotics and discovered Amoxicillin (51.3%),
244 Ampiclox (33.3%), and Metronidazole(21.9%). In western Uganda among nursing students,
245 Amoxicillin and Metronidazole were the most commonly used antibiotics (22) as well as in
246 northern Uganda as reported by Ocan, Bwanga (23). In Sudan, Amoxicillin is the most
247 commonly used antibiotic (15). The increased use of Amoxicillin could be due to its lower cost
248 than other antibiotics and the high prevalence of respiratory tract infections (although not all
249 respiratory infections require antibiotics for treatment) (35). Sensitizing the community to
250 various treatment modalities for respiratory infections could be vital for minimizing ASM.

251 **Conclusion and recommendations**

252 Self-medication with antibiotics is highly practiced in the Otuke District. Factors leading to
253 ASM originate from the health system as well as from individuals predisposing the community
254 to the danger of antimicrobial resistance if not addressed.

255 The health team should provide adequate information regarding the dangers of ASM to the
256 general public and equip community members with knowledge regarding supportive ways of
257 managing common conditions. In addition, the government should enact drug laws that govern
258 the use of antimicrobial agents and improve staff welfare to motivate them to effectively deliver

259 their services to the community. Furthermore, the government should recruit more staff to
260 quicken health service delivery.

261 **Acknowledgments**

262 The community of the Otuke district for participating in this study.

263 **Author contributions**

264 ADD: Conception, methods, data collection; reporting and review; RN: Methods, supervision,
265 manuscript draft and review

266 References

- 267 1. Namususwa S, Nakaziba R. Practice and knowledge of the dangers of self-medication among
268 pregnant mothers attending antenatal care in Iganga District (Uganda). *Nurs Commun.*
269 2022;6:e2022014.
- 270 2. Ocan M, Obuku EA, Bwanga F, Akena D, Richard S, Ogwal-Okeng J, et al. Household
271 antimicrobial self-medication: a systematic review and meta-analysis of the burden, risk factors and
272 outcomes in developing countries. *BMC public health.* 2015;15(1):742.
- 273 3. Haque M, Rahman NAA, McKimm J, Kibria GM, Azim Majumder MA, Haque SZ, et al. Self-
274 medication of antibiotics: investigating practice among university students at the Malaysian National
275 Defence University. *Infect Drug Resist.* 2019;12:1333-51.
- 276 4. Torres NF, Solomon VP, Middleton LE. Patterns of self-medication with antibiotics in Maputo
277 City: a qualitative study. *Antimicrobial Resistance & Infection Control.* 2019;8:1-12.
- 278 5. Yeika EV, Ingelbeen B, Kemah BL, Wirsiy FS, Fomengia JN, van der Sande MAB. Comparative
279 assessment of the prevalence, practices and factors associated with self-medication with antibiotics
280 in Africa. *Tropical medicine & international health : TM & IH.* 2021;26(8):862-81.
- 281 6. Silva A, Costa E, Freitas A, Almeida A. Revisiting the frequency and antimicrobial resistance
282 patterns of bacteria implicated in community urinary tract infections. *Antibiotics.* 2022;11(6):768.
- 283 7. Medina M-j, Legido-Quigley H, Hsu LY. Antimicrobial resistance in one health. *Global Health*
284 *Security: Recognizing Vulnerabilities, Creating Opportunities.* 2020:209-29.
- 285 8. WHO. Antibiotic resistance. 2020.
- 286 9. Alrasheedy AA, Alsalloum MA, Almuqbil FA, Almuzaini MA, Aba Alkhayl BS, Albishri AS, et al.
287 The impact of law enforcement on dispensing antibiotics without prescription: a multi-methods
288 study from Saudi Arabia. *Expert review of anti-infective therapy.* 2020;18(1):87-97.
- 289 10. Nepal G, Bhatta S. Self-medication with antibiotics in WHO Southeast Asian Region: a
290 systematic review. *Cureus.* 2018;10(4).
- 291 11. Ajibola O, Omisakin OA, Eze AA, Omoleke SA. Self-medication with antibiotics, attitude and
292 knowledge of antibiotic resistance among community residents and undergraduate students in
293 Northwest Nigeria. *Diseases.* 2018;6(2):32.
- 294 12. Chuwa BB, Njau LA, Msigwa KI, Shao E. Prevalence and factors associated with self
295 medication with antibiotics among University students in Moshi Kilimanjaro Tanzania. *Afr Health Sci.*
296 2021;21(2):633-9.
- 297 13. Ateshim Y, Bereket B, Major F, Emun Y, Woldai B, Pasha I, et al. Prevalence of self-
298 medication with antibiotics and associated factors in the community of Asmara, Eritrea: a descriptive
299 cross sectional survey. *BMC Public Health.* 2019;19(1):726.
- 300 14. Kretchy J-P, Adase SK, Gyansa-Lutterodt M. The prevalence and risks of antibiotic self-
301 medication in residents of a rural community in Accra, Ghana. *Scientific African.* 2021;14:e01006.
- 302 15. Hussain MA, Mohamed AO, Abdelkarim OA, Yousef BA, Babikir AA, Mirghani MM, et al.
303 Prevalence and Predictors of Antibiotic Self-Medication in Sudan: A Descriptive Cross-Sectional
304 Study. *Antibiotics.* 2023;12(3):612.
- 305 16. Ocan M, Obuku EA, Bwanga F, Akena D, Richard S, Ogwal-Okeng J, et al. Household
306 antimicrobial self-medication: a systematic review and meta-analysis of the burden, risk factors and
307 outcomes in developing countries. *BMC public health.* 2015;15:742.
- 308 17. Green DL, Keenan K, Fredricks KJ, Huque SI, Mushi MF, Kansime C, et al. The role of
309 multidimensional poverty in antibiotic misuse: a mixed-methods study of self-medication and non-
310 adherence in Kenya, Tanzania, and Uganda. *The Lancet Global Health.* 2023;11(1):e59-e68.
- 311 18. Aslam A, Gajdacs M, Zin CS, Ab Rahman NS, Ahmed SI, Zafar MZ, et al. Evidence of the
312 practice of self-medication with antibiotics among the lay public in low-and middle-income
313 countries: a scoping review. *Antibiotics.* 2020;9(9):597.

- 314 19. Roien R, Bhandari D, Hosseini SMR, Mosawi SH, Ataie MA, Ozaki A, et al. Prevalence and
315 determinants of self-medication with antibiotics among general population in Afghanistan. *Expert*
316 *Review of Anti-infective Therapy*. 2022;20(2):315-21.
- 317 20. Aslam A, Zin CS, Jamshed S, Rahman NSA, Ahmed SI, Pallós P, et al. Self-Medication with
318 Antibiotics: Prevalence, Practices and Related Factors among the Pakistani Public. *Antibiotics (Basel,*
319 *Switzerland)*. 2022;11(6).
- 320 21. Chinemerem Nwobodo D, Ugwu MC, Oliseloke Anie C, Al-Ouqaili MT, Chinedu Ikem J, Victor
321 Chigozie U, et al. Antibiotic resistance: The challenges and some emerging strategies for tackling a
322 global menace. *Journal of Clinical Laboratory Analysis*. 2022;36(9):e24655.
- 323 22. Nankya S, Musinguzi P, Arishaba A, Najjuma¹ JN. Prevalence and Practices of Self-medication
324 With Antibiotics Among Nursing Students at a Training Institution in Western Uganda. 2023.
- 325 23. Ocan M, Bwanga F, Bbosa GS, Bagenda D, Waako P, Ogwal-Okeng J, et al. Patterns and
326 predictors of self-medication in northern Uganda. *PloS one*. 2014;9(3):e92323.
- 327 24. Lu C, Tian Y, Tian H, Li B, Peng B, Zheng J, et al. An ultrasound activable metal-phenolic
328 network nano-antibiotics for in vivo on-site infection therapy. *Science China Materials*.
329 2023;66(1):395-406.
- 330 25. Rabaan AA, Alhumaid S, Mutair AA, Garout M, Abulhamayel Y, Halwani MA, et al.
331 Application of Artificial Intelligence in Combating High Antimicrobial Resistance Rates. *Antibiotics*.
332 2022;11(6):784.
- 333 26. Nabaweesi I, Olum R, Sekite AB, Suubi WT, Nakiwunga P, Machali A, et al. Antibiotic
334 practices, perceptions and self-medication among patients at a National Referral Hospital in Uganda.
335 *Infection and drug resistance*. 2021:2155-64.
- 336 27. Naylor NR, Atun R, Zhu N, Kulasabanathan K, Silva S, Chatterjee A, et al. Estimating the
337 burden of antimicrobial resistance: a systematic literature review. *Antimicrobial Resistance &*
338 *Infection Control*. 2018;7(1):58.
- 339 28. WHO. *Antimicrobial Resistance*. 2021.
- 340 29. EClinicalMedicine. *Antimicrobial resistance: a top ten global public health threat*. 41.
341 2021;101221.
- 342 30. Sachdev C, Anjankar A, Agrawal J. Self-medication with antibiotics: an element increasing
343 resistance. *Cureus*. 2022;14(10).
- 344 31. Atif M, Asghar S, Mushtaq I, Malik I, Amin A, Babar Z-U-D, et al. What drives inappropriate
345 use of antibiotics? A mixed methods study from Bahawalpur, Pakistan. *Infection and drug resistance*.
346 2019:687-99.
- 347 32. Dache A, Dona A, Ejeso A. Inappropriate use of antibiotics, its reasons and contributing
348 factors among communities of Yirgalem town, Sidama regional state, Ethiopia: A cross-sectional
349 study. *SAGE Open Medicine*. 2021;9:20503121211042461.
- 350 33. Torres N, Chibi B, Middleton L, Solomon V, Mashamba-Thompson T. Evidence of factors
351 influencing self-medication with antibiotics in low and middle-income countries: a systematic
352 scoping review. *Public health*. 2019;168:92-101.
- 353 34. Elong Ekambi G-A, Okalla Ebongue C, Penda IC, Nnanga Nga E, Mpondo Mpondo E,
354 Eboumbou Moukoko CE. Knowledge, practices and attitudes on antibiotics use in Cameroon: Self-
355 medication and prescription survey among children, adolescents and adults in private pharmacies.
356 *PloS one*. 2019;14(2):e0212875.
- 357 35. Mortazhejri S, Hong PJ, Yu AM, Hong BY, Stacey D, Bhatia RS, et al. Systematic review of
358 patient-oriented interventions to reduce unnecessary use of antibiotics for upper respiratory tract
359 infections. *Systematic reviews*. 2020;9:1-16.

360


361

362

363

364

365



Figure