

ECONOMIC EVALUATION OF A WEARABLE-BASED INTERVENTION TO INCREASE PHYSICAL ACTIVITY AMONG INSUFFICIENTLY ACTIVE MIDDLE-AGED ADULTS

Jack H. Ching

Google; Mountain View, CA
jackching@google.com

Steve Duff

Veritas Health Economics Consulting, Inc.; Kamas, UT
steveduff@veritashec.com

John Hernandez

Google; Mountain View, CA
johnbhernandez@google.com

ABSTRACT

Background: Physical activity levels worldwide have declined over recent decades, with the average number of daily steps decreasing steadily since 1995. Given that physical inactivity is a major modifiable risk factor for chronic disease and mortality, increasing the level of physical activity is a clear opportunity to improve population health on a broad scale. The current study aims to assess the cost-effectiveness and budget impact of a Fitbit-based intervention among healthy, but insufficiently active, adults to quantify the potential clinical and economic value for a commercially insured population in the U.S.

Methods: An economic model was developed to compare physical activity levels, health outcomes, costs, and quality-adjusted life-years (QALYs) associated with usual care and a Fitbit-based intervention that consists of a consumer wearable device alongside goal setting and feedback features provided in a companion software application. Improvement in physical activity was measured in terms of mean daily step count. The effects of increased daily step count were characterized as reduced short-term healthcare costs and decreased incidence of chronic diseases with corresponding improvement in health utility and reduced disease costs. Published literature, standardized costing resources, and data from a National Institutes of Health-funded research program were utilized. Cost-effectiveness and budget impact analyses were performed for a hypothetical cohort of middle-aged adults.

Results: The base case cost-effectiveness results found the Fitbit intervention to be dominant (less costly and more effective) compared to usual care. Discounted 15-year incremental costs and QALYs were -\$1,257 and 0.011, respectively. In probabilistic analyses, the Fitbit intervention was dominant in 93% of simulations and either dominant or cost-effective (defined as less than \$150,000/QALY gained) in 99.4% of simulations. For budget impact analyses conducted from the perspective of a U.S. Commercial payer, the Fitbit intervention was estimated to save approximately \$6.5 million dollars over 2 years and \$8.5 million dollars over 5 years for a cohort of 8,000 participants. Although the economic analysis results were very robust, the short-term healthcare cost savings were the most uncertain in this population and warrant further research.

Conclusions: There is abundant evidence documenting the benefits of wearable activity trackers when used to increase physical activity as measured by daily step counts. Our research provides additional health economic evidence supporting implementation of wearable-based interventions to improve population health, and offers compelling support for payers to consider including wearable-based physical activity interventions as part of a comprehensive portfolio of preventive health offerings for their insured populations.

Introduction

Physical activity (PA) levels worldwide have declined over recent decades, with the average number of daily steps decreasing steadily since 1995.¹ Among those in high-income countries, nearly 37% do not meet recommended levels of physical activity, and the rates of physical inactivity increase with age.² Inadequate physical activity is associated with substantial healthcare costs and higher risk of incident diseases, such as cardiovascular disease, type 2 diabetes, cancer, and mental health conditions. If current prevalence remains unchanged, physical inactivity is projected to result in 500 million new cases of preventable disease globally, which will incur an additional \$300 billion in treatment costs over 10 years.³ In the United States (U.S.), less than 50% of the overall population meet the recommended aerobic physical activity guidelines of at least 150 minutes of moderate-intensity exercise per week. This percentage is even lower among older age groups, with only 18-30% of adults ages 35-64 and 10-15% of adults older than 65 meeting these guidelines in 2020.⁴

Given that physical inactivity is a major modifiable risk factor for chronic disease and mortality, increasing the level of physical activity is a clear opportunity to improve population health on a broad scale. There is overwhelming evidence for the health benefits of increased physical activity, including reductions in all-cause mortality, improved cardiorespiratory, metabolic, and muscular health, and reductions in the risk of developing breast and colon cancer.⁵⁻⁸ Studies also support the role of regular physical activity in improving sleep quality and mental health outcomes such as anxiety and depression.⁹⁻¹² Recent evidence from the All of Us Research Program, the largest longitudinal study with objectively measured physical activity in the U.S., re-affirmed these benefits among participants who used a Fitbit device to track their physical activity. Those who maintained higher daily step counts over a median follow-up of 4 years had 25-46% lower risks of developing multiple chronic conditions, including diabetes, hypertension, obesity, sleep apnea, gastroesophageal reflux disease, and major depressive disorder.^{13,14}

Wearable activity trackers, such as Fitbit devices, offer a promising low-cost approach to help increase physical activity levels, leading to substantial health benefits. These trackers can be used either as the primary component of an intervention (often in conjunction with a software application) to promote self-monitoring and goal setting, or as a part of more comprehensive programs that also incorporate additional counseling, education, and/or structured exercise elements.^{15,16} Across more than a hundred randomized controlled trials (RCTs), these interventions consistently show clinically significant increases in daily step counts, increases in moderate to vigorous physical activity (MVPA), and improvements in body composition, blood pressure, and fitness.^{16,17} Among healthy adults, wearable-based interventions were associated with improvements in physical activity over an average of 21 weeks, and long-term follow-up found that approximately 40% of the short-term improvements were maintained through 3-4 years.^{18,19} These improvements are larger among older adults and those with chronic conditions such as type 2 diabetes and COPD, for whom wearable activity interventions resulted in significant short-term improvements in physical activity and reduced sedentary time.^{20,21} It is also notable that even the provision of a pedometer alone, without additional human-delivered behavioral change interventions, can increase both daily steps and MVPA minutes for up to 12 months.²²

Fitbit devices, in particular, have been effective in promoting healthy lifestyle behaviors and supporting chronic condition management as part of multifaceted behavior change interventions. A systematic review of 37 RCTs found that Fitbit-based interventions increased the number of daily steps, minutes of weekly activity, and led to additional pounds of weight lost.²³ The evidence for Fitbit-based interventions in sub-populations are consistent with those for wearable activity trackers in general. Studies among sedentary adult women found that the Fitbit-based intervention substantially increased steps per day, and the number of bouts and minutes of MVPA per week.^{24,25} Fitbit-based interventions have also been shown to increase physical activity levels in older adults for up to 32 weeks, and to increase various measures of physical fitness in this population.²⁶⁻²⁸

There is also abundant support for the health economic value of greater physical activity over the long-term, and for wearable-based interventions that aim to increase physical activity levels. Adopting or maintaining a physically active lifestyle throughout adulthood is associated with reductions in Medicare costs later in life, compared to those who remained inactive.²⁹ Wearable-based physical activity interventions, ranging from those that use pedometers, exercise prescriptions, individual/group counseling, to mass media promotions, have all been shown to be cost-effective or cost-saving among populations of middle-aged and older adults.³⁰⁻³⁴ In general, studies have found less costly interventions, such as those focused on self management using pedometers, to be more cost-effective or more likely to be cost-saving when compared to higher cost interventions that also include physician visits and/or referral to exercise professionals.

Building on existing health and economic evidence for wearable-based interventions that focus on improving physical activity, the current study aims to assess the cost-effectiveness and budget impact of a Fitbit-based intervention (consisting of a consumer wearable device and app-based goal setting and feedback features) among healthy, but

insufficiently active, adults to quantify the potential clinical and economic value for a commercially insured population in the U.S.

Methods

Modeling Approach

A Markov cohort state-transition model was developed to quantify the clinical, economic, and humanistic outcomes of usual care and a Fitbit intervention aimed at increasing physical activity in adults who are insufficiently active. The modeling approach was informed by previous research^{30–39} and good research practices published by the International Society for Pharmacoeconomics and Outcomes Research (ISPOR).^{40–45} A schematic in **Figure 1** illustrates the modeling framework, disease health states, and outcomes.

The model was designed to estimate the impact of the Fitbit intervention on physical activity in terms of an increase in the mean number of daily steps. In addition to being readily accessible and widely used to measure physical activity, daily step count is a well-supported metric that is inversely and linearly associated with health outcomes including cardiovascular disease events, type 2 diabetes incidence, and all-cause mortality.⁴⁶ In addition, total daily step counts and other physical activity measures have been found to have an effect on both short-term (i.e., 1-2 years) medical resource utilization and costs^{47–52} while the impact on longer-term costs and quality of life has been quantified in numerous modeling studies through decreased incidence of diseases associated with sedentary behavior.^{30–39} The cohort of individuals in the analysis were assumed to be in general good health at baseline (i.e., devoid of any of the chronic diseases of interest) and started in the ‘Well’ health state. During each model cycle (1 year), individuals remain in the ‘Well’ health state or transition to any of the incident disease health states or death, based on background age/sex-based mortality, adjusted according to health state-specific mortality hazard ratios. Transitions into one of the chronic disease states precluded transition back to the ‘Well’ health state; participants only remain in the disease health state or transition to the death absorbing state. Costs, life-years (LYs) and quality-adjusted life-years (QALYs) are accrued in each cycle.

Population

The target population of interest for the intervention in the analysis was insufficiently active adults in the U.S., aged 40–60 (mean age 50) who are insured by Commercial health plans. The age and sex distribution was based on 2022 data from the U.S. Census Bureau.⁵³ The proportion of adults with insufficient physical activity in this population was informed by a recent nationwide survey study of 2,640 participants.⁵⁴ Population details are shown in **Table 1**.

Comparators

As described above, physical activity interventions using wearable activity trackers such as accelerometers and pedometers have resulted in improved physical activity and clinical outcomes. A common alternative assessed in evaluations of physical activity interventions is usual care, which is the comparator we adopted for our economic analysis. Unlike usual care, which does not specifically encourage or increase physical activity and therefore results in no meaningful or lasting change in daily step count, the Fitbit intervention is estimated to increase the average daily step count by 1,287 steps, or approximately 20% from a baseline daily step count of 6,500. These estimates were derived using data from 8 RCTs of Fitbit-based interventions among participants less than 65 years of age (**Supplementary Text**).^{55–62} The durability of the Fitbit intervention was informed by two systematic reviews and meta-analyses of 8 and 57 RCTs (8,480 and 16,355 participants, respectively) that determined step count monitoring interventions to have sustained benefits over at least 4 years.^{19,63} Both analyses demonstrated statistically significant increases in daily step counts in the intervention group over controls, although the magnitude of the increases waned over time. To emulate this in the model, the maximum step count increase for the Fitbit intervention was assumed to peak at 1 year, decline linearly by 25% per year, with average daily step counts returning to baseline (and equivalent to usual care step count levels) beginning at year 5 (**Table 2**).

As is common with comparable wearable-based physical activity interventions, the Fitbit intervention was assumed to consist of a consumer wearable device alongside goal setting and feedback features provided in a companion software application. We further assume that the Fitbit intervention does not include additional human-delivered intervention components, and conducted extensive sensitivity analyses to explore our assumptions of intervention effectiveness.

To model the incidence of chronic diseases related to insufficient physical activity—including obesity (OBES), type 2 diabetes mellitus (T2DM), hypertension (HTN), gastroesophageal reflux disease (GERD), major depressive disorder (MDD), and sleep apnea (SA)—we utilized new analyses of previously published estimates of disease incidence derived

from electronic health records (EHR) in the All of Us Research Program.¹³ This unique ongoing research program is funded by the National Institutes of Health and is currently collecting multiple streams of health-related information from hundreds of thousands of Americans. As of the 2022 publication by Master et al., 6,042 of the 214,206 program participants sharing their EHR data had linked their own Fitbit devices to the data collection effort, thereby providing a rich data source on the relationship between physical activity and the risk of developing chronic disease. Cumulative incidence estimates by year (3, 5, and 7) for six diseases based on a linear function (as opposed to the published estimates based on a cubic spline fit) of average daily step count were obtained from the All of Us researchers for model parameterization to mitigate potential overfitting. Cumulative incidence data were converted to the annual incidence rates as model inputs (**Supplementary Table 1**).

Costs

The costs in the model include those related to the Fitbit intervention (incremental to usual care which is assumed to have no intervention cost), overall short-term healthcare costs in Years 1 and 2, and longer-term health state-related costs (based on ‘Well’ and disease states) in Years 3+. Given the non-invasive nature of the Fitbit intervention, adverse events are uncommon and assumed to have nominal costs; therefore, they were not included in the analysis. Unit costs for the model were derived primarily from published sources.^{52,64–71} In cases where robust evidence was unavailable, conservative assumptions based on clinical expert opinion were used to supplement published estimates. Furthermore, in several instances, published costs required conversion between the Medicare and Commercial perspectives. Inflation and deflation factors of 50% (Medicare to Commercial) and 33% (Commercial to Medicare), respectively, were utilized for this purpose. All unit costs in the model are reported in 2023 U.S. dollars (USD); published costs were inflated using the Medical Care component of the U.S. consumer price index (CPI)⁷² when necessary (**Table 1**).

The cost of the Fitbit intervention is estimated based on the retail price of a Charge 6 device, in addition to a 12-month subscription to Fitbit Premium (which provides goal setting, feedback, and access to a library of workout content and additional detailed metrics in-app), as well as additional intervention deployment costs.

Numerous studies have documented the impact of increased physical activity (using various modalities or interventions) on short-term medical resource use and costs, usually measured over 1-2 years.^{47–52} Given the substantial and clinically meaningful increase in average daily steps observed with Fitbit use,⁶³ we assumed a potential for modest short-term reduction in resource use and costs in our analyses. This model element was informed by an analysis of claims data from over 8,500 Medicare Advantage enrollees participating in the Silver Sneakers® (SS) healthy aging program.⁵² Because the study population and intervention are not an exact match to those in our model, two factors were included in our model to reduce the potential cost savings due to differences in population age (reflecting an assumption of lower overall healthcare costs in a younger population) and intervention intensity (reflecting an assumption that the SS program has a greater impact on resource use/cost). These parameters and conservative assumptions were evaluated extensively in both one-way and probabilistic sensitivity analyses as well as with comparison to published estimates of annual healthcare costs in a comparably aged population.⁶⁸

Finally, longer-term costs associated with the model health states (‘Well’ and incident diseases) were quantified for both comparators. Incremental costs attributable to each disease health state were extracted from the literature and added to the ‘Well’ health state to yield costs for each of the incident disease states.^{52,64–71}

Mortality

The annual probability of death for participants in the ‘Well’ health state was based on age- and sex-adjusted all-cause mortality data from U.S. life tables.⁷³ Published hazard ratios describing the excess mortality associated with each of the chronic diseases of interest were used to adjust background all-cause mortality and the annual probability of death for those disease states.^{74–83}

Utilities

Health-related quality of life utility and disutility weights were used to calculate QALYs for the cost-effectiveness analysis (CEA). Utilities for the ‘Well’ health state (by age strata) were based on a nationally representative catalog of EuroQol-5 Dimension (EQ-5D) utility scores.⁸⁴ For the chronic diseases in the model, we identified published disutilities^{85–90} (or utilities that were subsequently used to derive disutilities using information in Sullivan, et al. 2006⁸⁴) that were applied to the ‘Well’ health state utility to account for the worse quality of life for those conditions.

Validation and Analyses

Consistent with best practices guidance, the model was subjected to extensive validity testing.^{41,42,44} The face validity of the model was evaluated by comparison to other physical activity intervention model structures and through a clinical applicability review (J.H.). Internal validity was assessed by a review of the model programming by a second modeler (J.H.C.) and through a series of quality assurance tests and analyses to confirm proper model functionality. Finally, the external validity of the epidemiologic framework was appraised by comparisons of model outcomes to U.S. life expectancy data and other published studies.^{73,91,92}

Upon conclusion of the validation process, base case cost-effectiveness and budget impact analyses using the default parameter estimates were conducted. Both types of economic analysis took the perspective of a Commercial payer. Therefore, for the CEA, the time horizon was set to 15 years when the mean age in the model reached 65 (i.e., when a Medicare payer perspective would be more relevant). For the budget impact analysis (BIA), two separate time horizons were evaluated. A 2-year horizon was evaluated to focus on the potential short-term clinical and cost benefits while a 5-year horizon was used to also capture longer-term benefits and to be consistent with the modeled durability of the Fitbit intervention. Outcomes were discounted at an annual rate of 3% in the CEA and were not discounted in the BIA as recommended in good research practice guidelines.^{45,93,94} Although guidelines also recommend conduct of analyses from the societal perspective, the simple, non-invasive nature of the intervention is unlikely to result in substantial impacts within the informal health care and non-health care sectors, such that this would not materially change our conclusions.

The impact of uncertainty in parameter estimates and model structure was evaluated with deterministic and probabilistic sensitivity analyses (PSA). Deterministic analyses included one-way sensitivity analyses (OWSA) and scenarios constructed to assess multi-parameter variability. Probabilistic analyses included over 40 parameters varied simultaneously with new estimates drawn at random from their respective probability distributions (e.g., beta distribution for utilities, gamma distribution for costs, etc.) for 5,000 runs. Net monetary benefit (NMB) was calculated at an assumed willingness to pay (WTP) of \$150,000/QALY.^{95,96} Cost-effectiveness acceptability curves and a scatterplot were also constructed based on the results of the PSA.

Finally, an exploratory analysis of an insufficiently active older population (mean starting age 65) was conducted to assess the costs and outcomes of the Fitbit intervention from a Medicare payer perspective. A recent publication by Deidda, et. al.³¹ informed the exploratory analysis. Analysis details and results are presented in the **Supplementary Materials**.

Results

Validation Results

The model was deemed valid based on the evaluations conducted and the external validity check. For the latter, we appraised the epidemiologic framework to ensure that our modeling of incident disease and survival resulted in credible predictions. For both the base case population (middle-aged adults; mean age 50) and exploratory population (older adults; mean age 65), our model estimated mean life expectancy 2.5 – 3 years less than that based on U.S. life table predictions.⁷³ This is consistent with studies from Australia and the U.S. documenting a life expectancy reduction of 1.5 – 4.8 years due to sedentary behavior⁹² or, conversely, an increased life expectancy of 2 years if sedentary behaviors were to be ameliorated.⁹¹

Base Case Results

The results of the base case economic analyses are presented in **Table 3** and **Figure 2**. In the cost-effectiveness analysis, the total 15-year discounted costs of the Fitbit intervention were less than for usual care alone (\$88,284 vs \$89,541) resulting in a net cost savings of \$1,257. The majority of the cost savings (\$1,049) was associated with short-term cost reduction with the remainder due to reductions in incident disease. A nominal increase in life-years was observed for the Fitbit intervention (increment of 0.003 LYs) related to lower incident disease mortality. Similarly, because Fitbit use results in a greater proportion of time spent in the ‘Well’ health state, QALYs also are higher than for usual care (9.042 vs 9.031). As the Fitbit intervention is less costly and more effective than usual care, an ICER is not calculated and the Fitbit intervention is considered to “dominate” usual care.

Examining the economics from the perspective of a Commercial healthcare payer, in a 1-million member plan, it was estimated that there would be approximately 160,000 members in the 40-60 year old age group that would be classified as having insufficient physical activity. Of those, we assumed that 8,000 would be eligible and choose to adopt a Fitbit intervention and be included in the budget impact analysis.

Total costs for a cohort adopting the Fitbit intervention were lower than if they received usual care and remained insufficiently active (**Table 3**). Two-year and five-year cost savings were \$6,514,000 and \$8,548,644, respectively which equate to -\$0.27 per member per month (PMPM) and -\$0.14 PMPM. Given the shorter time horizon in the budget impact analysis, the contribution of incident disease cost savings is even lower than in the cost-effectiveness analysis.

Sensitivity and Scenario Analysis Results

One-way deterministic sensitivity analyses were conducted by varying key parameter values within credible ranges. Results of the sensitivity analyses are illustrated in a tornado diagram (**Supplementary Figure 1**); however, rather than displaying the results in terms of the impact of parameter changes on the ICER results, the tornado diagram shows the change in net monetary benefit at a willingness to pay threshold of \$150,000/QALY since all changes in individual parameters do not result in the calculation of an ICER (i.e., the Fitbit intervention remains dominant). These sensitivity analyses illustrate the importance of the short-term cost parameters on the results with 3 of the top 5 most influential parameters relating to this model element. However, even when these key parameter estimates are varied, the net monetary benefit is positive and the Fitbit intervention remains cost saving (not shown in a Figure). Furthermore, these analyses demonstrate that, even with step count improvements as low as 750 steps per day or if the intervention were only effective for 1 year (i.e., step count levels return to baseline at Year 2), the Fitbit intervention remains dominant.

While the deterministic results are informative regarding individual model parameters, the results of the PSA, in which the uncertainty in numerous parameters is evaluated simultaneously, are much more informative about the robustness of the cost-effectiveness results. The PSA results are presented in the scatterplot in **Figure 3**. Each of the 5,000 model runs is shown as an incremental QALY and cost pair (Fitbit intervention relative to usual care). To facilitate interpretation, lines representing the ICER thresholds of \$50,000, \$100,000, and \$150,000/QALY gained are superimposed on the graph. The vast majority (93.0%) of the simulation results fell into the southeast quadrant of the graph indicating that the Fitbit intervention is dominant (less costly and more effective). Over 99% of the model runs resulted in an ICER less than \$150,000/QALY gained or Fitbit intervention dominating usual care. These results provide very high confidence that, even when accounting for parameter uncertainty (including in the key short-term cost parameters), a Fitbit intervention would provide favorable value or cost savings compared to usual care.

Good research practices in modeling call for an examination of structural uncertainty in the model.^{41,42,44} We attempted to address this issue in two ways; first, we evaluated an alternative approach for characterizing intervention effectiveness in terms of daily step counts (a ‘tiered steps approach’ that allows for varying levels of step count increases among population subgroups vs the ‘mean steps approach’ [the model default]) and second, we varied how health state costs were integrated into the analysis (inclusion of these costs in the first 2 years vs. exclusion [the model default]). Modifications to each structural element resulted in a nominal change to the model outcomes, none of which materially changed the interpretation or conclusions of the analysis (**Supplementary Table 3**). More information on the analyses is available in the **Supplementary Materials**.

In addition to the various types of sensitivity analyses described above, we also performed several ad-hoc scenario analyses in which one or more parameters were varied. These included:

- Use of the ‘post-intervention’ steps estimate rather than the ‘difference in difference’ approach described in the **Supplementary Materials**;
- An increase and decrease of the proportion of eligible members adopting a Fitbit intervention (budget impact analysis only);
- Use of the upper and lower bounds of disease incidence estimated by the linear function of daily steps and incidence as obtained from the All of Us researchers;
- Extension of the cost-effectiveness analysis time horizon from 15 years to a lifetime horizon; and
- Substitution of health state costs (‘Well’ and incident diseases) in Years 1 and 2 for the general short-term healthcare cost component.

As presented in **Supplementary Table 3**, modification of the mean daily steps estimate, intervention adoption rates, the cumulative incidence of chronic disease, or the time horizon had a nominal impact on the results with the Fitbit intervention remaining dominant and cost saving compared to usual care. The results changed substantially only when health state costs were the only type of cost considered in the analysis. The Fitbit intervention was dominant and cost saving compared to usual care across the range of scenario analyses with the 15-year per-person cost savings ranging from -\$1,257 (when the short-term cost savings were included) to -\$318 (when the only source of cost savings was attributable to reduction in the onset of chronic disease). While complete exclusion of the literature-supported overall short-term healthcare costs is an extreme assumption, it illustrates the need for further research to ascertain the potential

reduction in short-term healthcare resource use associated with a physical activity intervention in the middle-aged population.

The exploratory analysis of a Fitbit-based intervention in an older population insured by Medicare demonstrated greater cost savings associated with the intervention than in the base case analysis of a middle-aged population (**Supplementary Tables 4 and 5, Supplementary Figure 2**). This is primarily due to the magnitude of short-term healthcare cost savings, for which the evidence is more robust in an older population, which is 2- to 4-times greater in the exploratory analysis than in the base case population. As a result, the cost savings associated with incident disease reduction in the exploratory analysis make up a much smaller proportion of the overall cost savings, partially because the inclusion of fewer chronic diseases in the exploratory analysis limits the magnitude of achievable cost savings. In addition, over the lifetime horizon in the exploratory analysis, only a fraction of disease occurrence can be completely prevented by an intervention that, on average, has effects over a 5-year period. Additional analysis details and results are presented in the **Supplementary Materials**.

Discussion

In this economic analysis, we quantified both the cost effectiveness and budget impact of a Fitbit intervention relative to usual care in an insufficiently active, middle-aged population of U.S. adults. Over a 15-year time horizon, the CEA demonstrated that the Fitbit intervention would be cost saving and result in more QALYs than if participants only received usual care. When assessed over a shorter time horizon from a hypothetical Commercial health plan perspective, the BIA results showed the Fitbit intervention would save \$6-8 million dollars over a 2- to 5-year period for a cohort of plan participants. The results were most sensitive to assumptions about short-term healthcare cost impacts but the cost-saving conclusions remained consistent across all scenario analyses that explored model and parameter uncertainty.

The overall findings are consistent with prior health economic models of interventions that aim to increase physical activity through the use of consumer wearable activity trackers, particularly those that leverage pedometer-based interventions. A systematic review and modeling analysis by Gc et. al. reported that brief interventions leveraging pedometers as a motivational tool (with or without additional exercise counseling) were likely to be cost-saving at a population level.^{39,97} Specifically, a program encouraging the use of pedometers was more effective and less costly when compared to usual care in the general Australian population; pairing a pedometer with a step-based exercise prescription goal was similarly dominant when compared to exercise prescription alone for those older than age 65 in New Zealand.^{32,98} In Belgium, a multi-strategy health promotion campaign that incorporated pedometer use was also dominant when compared to no intervention among a general adult population.⁹⁹

Our findings of short-term cost reductions from a fully remotely-delivered intervention have also been demonstrated among adults receiving primary care in the U.K. Anokye et. al.³⁰ found that postal delivery of a pedometer intervention had a 50% chance of being cost-effective after 1 year of the intervention compared to control, and is dominant in the long-term when compared to both usual care and a version of the pedometer intervention that included nurse consultations. In particular, short-term cost savings in the postal pedometer group in the first year were due to lower health services use.³⁰ While the modeled Fitbit intervention differs slightly in its mode of implementation (i.e. not delivered within primary care practice), Fitbit's robust tracking of physical activity metrics, including step counts and active minutes, as well as goal setting and feedback features compares favorably to pedometer interventions that may be delivered during a brief healthcare encounter.

The strengths of our research include its robust model design and conservative approach to model parameters and assumptions. The model concept and structure were informed by, and are generally consistent with, numerous other economic analyses of physical activity interventions.³⁰⁻³⁹ One substantial enhancement over older analyses was the use of mean daily step count improvements from RCTs of Fitbit-based interventions and longitudinal health outcomes from over 6,000 Fitbit users with 4 years of median follow-up to estimate rates of incident disease. When calculating step count improvement, we used the most conservative approach to bias results against the Fitbit intervention. Finally, our PSA was extensive, including over 40 key model parameters, strengthening the confidence and robustness of our findings.

Despite its strengths, our study has important limitations. First, although there is abundant literature documenting short-term cost savings associated with improved physical activity,⁴⁷⁻⁵² most of those studies were conducted in older patients which limits the generalizability of those results to our middle-aged cohort. We attempted to minimize the effect of this data issue by making conservative assumptions, thus reducing the cost offset of the Fitbit intervention due to potential differences in population age and intervention efficacy. In addition, our PSA allowed for short-term costs to be greater in the Fitbit intervention than control. Finally, even without short-term cost savings, our analysis suggests that a Fitbit intervention is likely to still be cost-saving compared to usual care.

Next, although the Fitbit intervention has many important elements other than simply providing daily step count information (e.g., app-based goal setting and personalized feedback that may substitute for similar elements that would have incurred additional costs if provided by clinical staff, physical therapist, etc.), our analysis does not use *de novo* data in the population described. Rather, the analysis relies on RCTs that may not exactly replicate the intervention as described or the modeled population, nor have sufficient follow-up to inform the longer-term impact on physical activity levels beyond the duration of the intervention. This issue was addressed with sensitivity analyses that varied the intervention efficacy (in terms of daily steps) and durability (time to return to baseline step counts).

Finally, we acknowledge limitations in how our model structure captures the benefits of increased levels of physical activity. Estimates of disease incidence stratified by daily step counts in our model were based on reported associations rather than causal estimates. While an RCT would be the optimal approach to prove causality; in the absence of such data, we leveraged detailed disease incidence estimates from a robust longitudinal cohort study that provided multiple years of objectively tracked levels of physical activity, as measured by Fitbit devices. These estimates are likely a better reflection of physical activity levels under real-world conditions as opposed to prior estimates that relied on self-reports or short-term accelerometry measurements. We additionally varied these disease incidence estimates broadly in multi-way sensitivity analyses. Across these analyses, the Fitbit intervention remained very likely to be cost-saving compared to usual care. In addition, since the present analysis characterized the benefits of physical activity only through reduced disease incidence among the ‘Well’ population, it is possible that the overall magnitude of benefits could be even larger if improvements in chronic condition management were also included.

Our research strengthens support for implementing wearable-based interventions to improve population health through increasing physical activity among insufficiently active adults. We illustrate the health economic value of deploying a relatively low-cost wearable-based intervention, as it can be potentially cost-saving through lower healthcare utilization and disease-related costs, even if the resulting improvements in clinical outcomes are modest. Taken together, this offers compelling support for healthcare payers to consider including wearable-based physical activity interventions as part of a comprehensive portfolio of preventive health offerings for their insured populations.

Funding Statement

This study was funded by Google. SD is a shareholder in Veritas Health Economics Consulting Inc, which was contracted by Google to conduct this study.

Competing Interests

JHC and JH are employees of Google and own Alphabet stock as part of the standard compensation package.

Data Availability

All data produced in the present study are available upon reasonable request to the authors.

Acknowledgements

We gratefully acknowledge Amy McDonough and the Strategic Health Solutions team for supporting this study, and Jeffrey Annis (Vanderbilt Institute for Clinical and Translational Research) for his assistance with parameter estimates used in our modeling analyses.

References

1. Conger SA, Toth LP, Cretsinger C, et al. Time Trends in Physical Activity Using Wearable Devices: A Systematic Review and Meta-analysis of Studies from 1995 to 2017. *Med. Sci. Sports Exerc.* 2022;54:288–98.
2. Guthold R, Stevens GA, Riley LM, and Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million participants. *Lancet Glob Health* 2018;6:e1077–e1086.
3. World Health Organization. Global status report on physical activity 2022. <https://www.who.int/teams/health-promotion/physical-activity/global-status-report-on-physical-activity-2022>. Accessed: 2023-12-1. 2022.
4. Elgaddal N, Kramarow EA, and Reuben C. Physical Activity Among Adults Aged 18 and Over: United States, 2020. NCHS Data Brief 2022:1–8.
5. Ekelund U, Tarp J, Steene-Johannessen J, et al. Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ* 2019;366:14570.
6. Saint-Maurice PF, Troiano RP, Bassett Jr DR, et al. Association of Daily Step Count and Step Intensity With Mortality Among US Adults. *JAMA* 2020;323:1151–60.
7. Kyu HH, Bachman VF, Alexander LT, et al. Physical activity and risk of breast cancer, colon cancer, diabetes, ischemic heart disease, and ischemic stroke events: systematic review and dose-response meta-analysis for the Global Burden of Disease Study 2013. *BMJ* 2016;354:i3857.
8. Del Pozo Cruz B, Ahmadi MN, Lee IM, and Stamatakis E. Prospective Associations of Daily Step Counts and Intensity With Cancer and Cardiovascular Disease Incidence and Mortality and All-Cause Mortality. *JAMA Intern. Med.* 2022;182:1139–48.
9. Kredlow MA, Capozzoli MC, Hearon BA, Calkins AW, and Otto MW. The effects of physical activity on sleep: a meta-analytic review. *J. Behav. Med.* 2015;38:427–49.
10. Herring MP, O'Connor PJ, and Dishman RK. The effect of exercise training on anxiety symptoms among patients: a systematic review. *Arch. Intern. Med.* 2010;170:321–31.
11. Schuch FB, Vancampfort D, Firth J, et al. Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. *Am. J. Psychiatry* 2018;175:631–48.
12. Noetel M, Sanders T, Gallardo-Gómez D, et al. Effect of exercise for depression: systematic review and network meta-analysis of randomised controlled trials. *BMJ* 2024;384:e075847.
13. Master H, Annis J, Huang S, et al. Association of step counts over time with the risk of chronic disease in the All of Us Research Program. *Nat. Med.* 2022;28:2301–8.
14. Perry AS, Annis JS, Master H, et al. Association of Longitudinal Activity Measures and Diabetes Risk: An Analysis From the National Institutes of Health All of Us Research Program. *J. Clin. Endocrinol. Metab.* 2023;108:1101–9.
15. Lyons EJ, Lewis ZH, Mayrsohn BG, and Rowland JL. Behavior change techniques implemented in electronic lifestyle activity monitors: a systematic content analysis. *J. Med. Internet Res.* 2014;16:e192.
16. Brickwood KJ, Watson G, O'Brien J, and Williams AD. Consumer-Based Wearable Activity Trackers Increase Physical Activity Participation: Systematic Review and Meta-Analysis. *JMIR Mhealth Uhealth* 2019;7:e11819.
17. Ferguson T, Olds T, Curtis R, et al. Effectiveness of wearable activity trackers to increase physical activity and improve health: a systematic review of systematic reviews and meta-analyses. *Lancet Digit Health* 2022;4:e615–e626.
18. Tang MSS, Moore K, McGavigan A, Clark RA, and Ganesan AN. Effectiveness of Wearable Trackers on Physical Activity in Healthy Adults: Systematic Review and Meta-Analysis of Randomized Controlled Trials. *JMIR Mhealth Uhealth* 2020;8:e15576.
19. Chaudhry UAR, Wahlich C, Fortescue R, Cook DG, Knightly R, and Harris T. The effects of step-count monitoring interventions on physical activity: systematic review and meta-analysis of community-based randomised controlled trials in adults. *Int. J. Behav. Nutr. Phys. Act.* 2020;17:129.
20. Wu S, Li G, Du L, Chen S, Zhang X, and He Q. The effectiveness of wearable activity trackers for increasing physical activity and reducing sedentary time in older adults: A systematic review and meta-analysis. *Digit Health* 2023;9:20552076231176705.
21. Li C, Chen X, and Bi X. Wearable activity trackers for promoting physical activity: A systematic meta-analytic review. *Int. J. Med. Inform.* 2021;152:104487.
22. Harris T, Kerry S, Victor C, et al. A pedometer-based walking intervention in 45- to 75-year-olds, with and without practice nurse support: the PACE-UP three-arm cluster RCT. *Health Technol. Assess.* 2018;22:1–274.

23. Ringeval M, Wagner G, Denford J, Paré G, and Kitsiou S. Fitbit-Based Interventions for Healthy Lifestyle Outcomes: Systematic Review and Meta-Analysis. *J. Med. Internet Res.* 2020;22:e23954.
24. Oppezzo MA, Tremmel JA, Kapphahn K, et al. Feasibility, preliminary efficacy, and accessibility of a twitter-based social support group vs Fitbit only to decrease sedentary behavior in women. *Internet Interv* 2021;25:100426.
25. Butryn ML, Arigo D, Raggio GA, Colasanti M, and Forman EM. Enhancing physical activity promotion in midlife women with technology-based self-monitoring and social connectivity: A pilot study. *J. Health Psychol.* 2016;21:1548–55.
26. McMahan SK, Lewis B, Oakes JM, Wyman JF, Guan W, and Rothman AJ. Assessing the Effects of Interpersonal and Intrapersonal Behavior Change Strategies on Physical Activity in Older Adults: a Factorial Experiment. *Ann. Behav. Med.* 2017;51:376–90.
27. Aguiñaga S, Marques IG, Kitsiou S, et al. BAILAMOS With mHealth Technology! Improving Physical Activity and Well-Being in Middle-Aged and Older Latinxs: A Pre-Post Feasibility Study. *Health Educ. Behav.* 2021;48:575–83.
28. Halloway S, Wilbur J, Braun LT, Schoeny ME, and Volgman AS. The Feasibility of a Combined Lifestyle Physical Activity and Cognitive Training Intervention to Prevent Cognitive Impairment in Older Women With Cardiovascular Disease. *J. Phys. Act. Health* 2021;18:70–5.
29. Coughlan D, Saint-Maurice PF, Carlson SA, Fulton J, and Matthews CE. Leisure time physical activity throughout adulthood is associated with lower medicare costs: evidence from the linked NIH-AARP diet and health study cohort. *BMJ Open Sport Exerc Med* 2021;7:e001038.
30. Anokye N, Fox-Rushby J, Sanghera S, et al. Short-term and long-term cost-effectiveness of a pedometer-based exercise intervention in primary care: a within-trial analysis and beyond-trial modelling. *BMJ Open* 2018;8:e021978.
31. Deidda M, Coll-Planas L, Tully MA, et al. Cost-effectiveness of a programme to address sedentary behaviour in older adults: results from the SITLESS RCT. *Eur. J. Public Health* 2022;32:415–21.
32. Cobiac LJ, Vos T, and Barendregt JJ. Cost-effectiveness of interventions to promote physical activity: a modelling study. *PLoS Med.* 2009;6:e1000110.
33. Roux L, Pratt M, Tengs TO, et al. Cost effectiveness of community-based physical activity interventions. *Am. J. Prev. Med.* 2008;35:578–88.
34. Over EA, Wendel-Vos GW, Berg M van den, et al. Cost-effectiveness of counseling and pedometer use to increase physical activity in the Netherlands: a modeling study. *Cost Eff. Resour. Alloc.* 2012;10:13.
35. Anokye NK, Trueman P, Green C, Pavey TG, Hillsdon M, and Taylor RS. The cost-effectiveness of exercise referral schemes. *BMC Public Health* 2011;11:954.
36. Pavey TG, Anokye N, Taylor AH, et al. The clinical effectiveness and cost-effectiveness of exercise referral schemes: a systematic review and economic evaluation. *Health Technol. Assess.* 2011;15:i–xii, 1–254.
37. Coyle D, Coyle K, Kenny GP, et al. Cost-effectiveness of exercise programs in type 2 diabetes. *Int. J. Technol. Assess. Health Care* 2012;28:228–34.
38. Campbell F, Holmes M, Everson-Hock E, et al. A systematic review and economic evaluation of exercise referral schemes in primary care: a short report. *Health Technol. Assess.* 2015;19:1–110.
39. Gc VS, Suhrcke M, Hardeman W, Sutton S, Wilson ECF, and Very Brief Interventions Programme Team. Cost-Effectiveness and Value of Information Analysis of Brief Interventions to Promote Physical Activity in Primary Care. *Value Health* 2018;21:18–26.
40. Briggs AH, Weinstein MC, Fenwick EAL, et al. Model parameter estimation and uncertainty: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force–6. *Value Health* 2012;15:835–42.
41. Caro JJ, Briggs AH, Siebert U, Kuntz KM, and ISPOR-SMDM Modeling Good Research Practices Task Force. Modeling good research practices—overview: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force–1. *Value Health* 2012;15:796–803.
42. Eddy DM, Hollingworth W, Caro JJ, et al. Model transparency and validation: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force–7. *Value Health* 2012;15:843–50.
43. Roberts M, Russell LB, Paltiel AD, et al. Conceptualizing a model: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force–2. *Value Health* 2012;15:804–11.
44. Siebert U, Alagoz O, Bayoumi AM, et al. State-transition modeling: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force–3. *Value Health* 2012;15:812–20.
45. Sullivan SD, Mauskopf JA, Augustovski F, et al. Budget impact analysis-principles of good practice: report of the ISPOR 2012 Budget Impact Analysis Good Practice II Task Force. *Value Health* 2014;17:5–14.

46. Kraus WE, Janz KF, Powell KE, et al. Daily Step Counts for Measuring Physical Activity Exposure and Its Relation to Health. *Med. Sci. Sports Exerc.* 2019;51:1206–12.
47. Ackermann RT, Cheadle A, Sandhu N, Madsen L, Wagner EH, and LoGerfo JP. Community exercise program use and changes in healthcare costs for older adults. *Am. J. Prev. Med.* 2003;25:232–7.
48. Ackermann RT, Williams B, Nguyen HQ, Berke EM, Maciejewski ML, and LoGerfo JP. Healthcare cost differences with participation in a community-based group physical activity benefit for medicare managed care health plan members. *J. Am. Geriatr. Soc.* 2008;56:1459–65.
49. Martinson BC, Crain AL, Pronk NP, O'Connor PJ, and Maciosek MV. Changes in physical activity and short-term changes in health care charges: a prospective cohort study of older adults. *Prev. Med.* 2003;37:319–26.
50. Nguyen HQ, Ackermann RT, Maciejewski M, et al. Managed-Medicare health club benefit and reduced health care costs among older adults. *Prev. Chronic Dis.* 2008;5:A14.
51. Nguyen HQ, Maciejewski ML, Gao S, Lin E, Williams B, and Logerfo JP. Health care use and costs associated with use of a health club membership benefit in older adults with diabetes. *Diabetes Care* 2008;31:1562–7.
52. Teigland C, Bilder S, and Slay D. Evaluating a healthy aging program's impact on health outcomes in Medicare Advantage beneficiaries. <https://www.inovalon.com/resource/evaluating-a-healthy-aging-programs-impact-on-health-outcomes-in-medicare-advantage-beneficiaries/>. 2022.
53. US Census Bureau Population Division. Annual estimates of the resident population by single year of and sex for the United States: April 1, 2020 to July 1, 2022 (NC-EST2022-SYASEXN). <https://www.census.gov/data/tables/time-series/demo/popest/2020s-national-detail.html>. Accessed: 2023-12-10. 2023.
54. Matthews CE, Carlson SA, Saint-Maurice PF, et al. Sedentary Behavior in U.S. Adults: Fall 2019. *Med. Sci. Sports Exerc.* 2021;53:2512–9.
55. Amorim AB, Pappas E, Simic M, et al. Integrating Mobile-health, health coaching, and physical activity to reduce the burden of chronic low back pain trial (IMPACT): a pilot randomised controlled trial. *BMC Musculoskelet. Disord.* 2019;20:71.
56. Ashe MC, Winters M, Hoppmann CA, et al. “Not just another walking program”: Everyday Activity Supports You (EASY) model-a randomized pilot study for a parallel randomized controlled trial. *Pilot Feasibility Stud* 2015;1:4.
57. Cadmus-Bertram L, Tevaarwerk AJ, Sesto ME, Gangnon R, Van Remortel B, and Date P. Building a physical activity intervention into clinical care for breast and colorectal cancer survivors in Wisconsin: a randomized controlled pilot trial. *J. Cancer Surviv.* 2019;13:593–602.
58. Duscha BD, Piner LW, Patel MP, et al. Effects of a 12-week mHealth program on peak VO₂ and physical activity patterns after completing cardiac rehabilitation: A randomized controlled trial. *Am. Heart J.* 2018;199:105–14.
59. Katz P, Margaretten M, Gregorich S, and Trupin L. Physical Activity to Reduce Fatigue in Rheumatoid Arthritis: A Randomized Controlled Trial. *Arthritis Care Res.* 2018;70:1–10.
60. Li LC, Sayre EC, Xie H, et al. Efficacy of a Community-Based Technology-Enabled Physical Activity Counseling Program for People With Knee Osteoarthritis: Proof-of-Concept Study. *J. Med. Internet Res.* 2018;20:e159.
61. Paxton RJ, Forster JE, Miller MJ, Geron KL, Stevens-Lapsley JE, and Christiansen CL. A Feasibility Study for Improved Physical Activity After Total Knee Arthroplasty. *J. Aging Phys. Act.* 2018;26:7–13.
62. Van Blarigan EL, Chan H, Van Loon K, et al. Self-monitoring and reminder text messages to increase physical activity in colorectal cancer survivors (Smart Pace): a pilot randomized controlled trial. *BMC Cancer* 2019;19:218.
63. Gasana J, O'Keeffe T, Withers TM, and Greaves CJ. A systematic review and meta-analysis of the long-term effects of physical activity interventions on objectively measured outcomes. *BMC Public Health* 2023;23:1697.
64. American Diabetes Association. Economic Costs of Diabetes in the U.S. in 2017. *Diabetes Care* 2018;41:917–28.
65. Greenberg PE, Fournier AA, Sisitsky T, et al. The Economic Burden of Adults with Major Depressive Disorder in the United States (2010 and 2018). *Pharmacoeconomics* 2021;39:653–65.
66. Hong YD, Onukwugha E, and Slejko JF. The Economic Burden of Comorbid Obstructive Sleep Apnea Among Patients with Chronic Obstructive Pulmonary Disease. *J Manag Care Spec Pharm* 2020;26:1353–62.
67. Kirkland EB, Heincelman M, Bishu KG, et al. Trends in Healthcare Expenditures Among US Adults With Hypertension: National Estimates, 2003-2014. *J. Am. Heart Assoc.* 2018;7.
68. Leung MYM, Carlsson NP, Colditz GA, and Chang SH. The Burden of Obesity on Diabetes in the United States: Medical Expenditure Panel Survey, 2008 to 2012. *Value Health* 2017;20:77–84.
69. Sharma P, Falk GW, Bhor M, et al. Healthcare Resource Utilization and Costs Among Patients With Gastroesophageal Reflux Disease, Barrett's Esophagus, and Barrett's Esophagus-Related Neoplasia in the United States. *J Health Econ Res* 2023;10:51–8.

Economic Evaluation of a Wearable-based Intervention

70. Wang G, Zhou X, Zhuo X, and Zhang P. Annual Total Medical Expenditures Associated with Hypertension by Diabetes Status in U.S. Adults. *Am. J. Prev. Med.* 2017;53:S182–S189.
71. Chhatre S, Chang YHA, Gooneratne NS, Kuna S, Strollo P, and Jayadevappa R. Association between adherence to continuous positive airway pressure treatment and cost among medicare enrollees. *Sleep* 2020;43.
72. US Bureau of Labor Statistics. Consumer price index for medical care in the U.S. city average, all urban consumers, seasonally adjusted. <https://www.bls.gov/cpi/data.htm>. Accessed: 2023-12-12. 2023.
73. Arias E, Xu J, and Kochanek K. United States Life Tables, 2021. *Natl. Vital Stat. Rep.* 2023;72:1–64.
74. Aune D, Huang W, Nie J, and Wang Y. Hypertension and the Risk of All-Cause and Cause-Specific Mortality: An Outcome-Wide Association Study of 67 Causes of Death in the National Health Interview Survey. *Biomed Res. Int.* 2021;2021:9376134.
75. Ford ES. Trends in mortality from all causes and cardiovascular disease among hypertensive and nonhypertensive adults in the United States. *Circulation* 2011;123:1737–44.
76. Gilman SE, Sucha E, Kingsbury M, Horton NJ, Murphy JM, and Colman I. Depression and mortality in a longitudinal study: 1952-2011. *CMAJ* 2017;189:E1304–E1310.
77. Li S, Wang J, Zhang B, Li X, and Liu Y. Diabetes Mellitus and Cause-Specific Mortality: A Population-Based Study. *Diabetes Metab. J.* 2019;43:319–41.
78. Lin Y, Wu Y, Lin Q, et al. Objective Sleep Duration and All-Cause Mortality Among People With Obstructive Sleep Apnea. *JAMA Netw Open* 2023;6:e2346085.
79. Ness-Jensen E, Santoni G, Gottlieb-Vedi E, Lindam A, Pedersen N, and Lagergren J. Mortality in gastro-oesophageal reflux disease in a population-based nationwide cohort study of Swedish twins. *BMJ Open* 2020;10:e037456.
80. Pratt LA, Druss BG, Manderscheid RW, and Walker ER. Excess mortality due to depression and anxiety in the United States: results from a nationally representative survey. *Gen. Hosp. Psychiatry* 2016;39:39–45.
81. Preis SR, Hwang SJ, Coady S, et al. Trends in all-cause and cardiovascular disease mortality among women and men with and without diabetes mellitus in the Framingham Heart Study, 1950 to 2005. *Circulation* 2009;119:1728–35.
82. Richter JE and Rubenstein JH. Presentation and Epidemiology of Gastroesophageal Reflux Disease. *Gastroenterology* 2018;154:267–76.
83. Ward ZJ, Willett WC, Hu FB, Pacheco LS, Long MW, and Gortmaker SL. Excess mortality associated with elevated body weight in the USA by state and demographic subgroup: A modelling study. *EClinicalMedicine* 2022;48:101429.
84. Sullivan PW and Ghushchyan V. Preference-Based EQ-5D index scores for chronic conditions in the United States. *Med. Decis. Making* 2006;26:410–20.
85. Hanmer J, Cherepanov D, Palta M, Kaplan RM, Feeny D, and Fryback DG. Health Condition Impacts in a Nationally Representative Cross-Sectional Survey Vary Substantially by Preference-Based Health Index. *Med. Decis. Making* 2016;36:264–74.
86. Jain S, Gupta S, Li VW, Suthoff E, and Arnaud A. Humanistic and economic burden associated with depression in the United States: a cross-sectional survey analysis. *BMC Psychiatry* 2022;22:542.
87. Janicak PG, Dunner DL, Aaronson ST, et al. Transcranial magnetic stimulation (TMS) for major depression: a multisite, naturalistic, observational study of quality of life outcome measures in clinical practice. *CNS Spectr.* 2013;18:322–32.
88. McCarty TR, Jirapinyo P, James LP, Gupta S, Chan WW, and Thompson CC. Transoral incisionless fundoplication is cost-effective for treatment of gastroesophageal reflux disease. *Endosc Int Open* 2022;10:E923–E932.
89. Shah CH and Brown JD. Reliability and Validity of the Short-Form 12 Item Version 2 (SF-12v2) Health-Related Quality of Life Survey and Disutilities Associated with Relevant Conditions in the U.S. Older Adult Population. *J. Clin. Med. Res.* 2020;9.
90. Walia HK, Thompson NR, Katzan I, Foldvary-Schaefer N, Moul DE, and Mehra R. Impact of Sleep-Disordered Breathing Treatment on Quality of Life Measures in a Large Clinic-Based Cohort. *J. Clin. Sleep Med.* 2017;13:1255–63.
91. Katzmarzyk PT and Lee IM. Sedentary behaviour and life expectancy in the USA: a cause-deleted life table analysis. *BMJ Open* 2012;2.
92. Veerman JL, Healy GN, Cobiac LJ, et al. Television viewing time and reduced life expectancy: a life table analysis. *Br. J. Sports Med.* 2012;46:927–30.
93. Lipscomb J, Weinstein MC, and Torrance GW. Time preference. In: *Cost-Effectiveness in Health and Medicine*. Ed. by Gold MR, Siegel JE, Russell LB, and Weinstein MC. New York: Oxford University Press, 1996:214–35.

Economic Evaluation of a Wearable-based Intervention

94. Weinstein MC, Siegel JE, Gold MR, Kamlet MS, and Russell LB. Recommendations of the Panel on Cost-effectiveness in Health and Medicine. *JAMA* 1996;276:1253–8.
95. Neumann PJ, Cohen JT, and Weinstein MC. Updating cost-effectiveness—the curious resilience of the \$50,000-per-QALY threshold. *N. Engl. J. Med.* 2014;371:796–7.
96. Neumann PJ and Kim DD. Cost-effectiveness Thresholds Used by Study Authors, 1990-2021. *JAMA* 2023;329:1312–4.
97. Gc V, Wilson ECF, Suhrcke M, Hardeman W, Sutton S, and VBI Programme Team. Are brief interventions to increase physical activity cost-effective? A systematic review. *Br. J. Sports Med.* 2016;50:408–17.
98. Leung W, Ashton T, Kolt GS, et al. Cost-effectiveness of pedometer-based versus time-based Green Prescriptions: the Healthy Steps Study. *Aust. J. Prim. Health* 2012;18:204–11.
99. De Smedt D, De Cocker K, Annemans L, De Bourdeaudhuij I, and Cardon G. A cost-effectiveness study of the community-based intervention '10 000 Steps Ghent'. *Public Health Nutr.* 2012;15:442–51.
100. Harvey JA, Chastin SFM, and Skelton DA. Prevalence of sedentary behavior in older adults: a systematic review. *Int. J. Environ. Res. Public Health* 2013;10:6645–61.
101. Alzheimers Association. 2023 Alzheimer's disease facts and figures. *Alzheimers. Dement.* 2023;19:1598–695.
102. Birger M, Kaldjian AS, Roth GA, Moran AE, Dieleman JL, and Bellows BK. Spending on Cardiovascular Disease and Cardiovascular Risk Factors in the United States: 1996 to 2016. *Circulation* 2021;144:271–82.
103. Burns ER, Stevens JA, and Lee R. The direct costs of fatal and non-fatal falls among older adults - United States. *J. Safety Res.* 2016;58:99–103.
104. Lancôt KL, Hviid Hahn-Pedersen J, Eichinger CS, et al. Burden of Illness in People with Alzheimer's Disease: A Systematic Review of Epidemiology, Comorbidities and Mortality. *J Prev Alzheimers Dis* 2024;11:97–107.
105. Shaked O, Cohen G, Goshen A, Shimony T, Shohat T, and Gerber Y. Physical Activity and Long-Term Mortality Risk in Older Adults with and without Cardiovascular Disease: A Nationwide Cohort Study. *Gerontology* 2022;68:529–37.
106. Bjerk M, Brovold T, Davis JC, and Bergland A. Evaluating a falls prevention intervention in older home care recipients: a comparison of SF-6D and EQ-5D. *Qual. Life Res.* 2019;28:3187–95.
107. Lacey EA and Walters SJ. Continuing inequality: gender and social class influences on self perceived health after a heart attack. *J. Epidemiol. Community Health* 2003;57:622–7.
108. Vandepitte S, Putman K, Van Den Noortgate N, Verhaeghe N, and Annemans L. Cost-effectiveness of an in-home respite care program to support informal caregivers of persons with dementia: A model-based analysis. *Int. J. Geriatr. Psychiatry* 2020;35:601–9.
109. InterAct Consortium, Ekelund U, Palla L, et al. Physical activity reduces the risk of incident type 2 diabetes in general and in abdominally lean and obese men and women: the EPIC-InterAct Study. *Diabetologia* 2012;55:1944–52.
110. Abbott RD, White LR, Ross GW, Masaki KH, Curb JD, and Petrovitch H. Walking and dementia in physically capable elderly men. *JAMA* 2004;292:1447–53.
111. Podewils LJ, Guallar E, Kuller LH, et al. Physical activity, APOE genotype, and dementia risk: findings from the Cardiovascular Health Cognition Study. *Am. J. Epidemiol.* 2005;161:639–51.
112. Larsson SC and Wolk A. The Role of Lifestyle Factors and Sleep Duration for Late-Onset Dementia: A Cohort Study. *J. Alzheimers. Dis.* 2018;66:579–86.
113. Jefferis BJ, Parsons TJ, Sartini C, et al. Does total volume of physical activity matter more than pattern for onset of CVD? A prospective cohort study of older British men. *Int. J. Cardiol.* 2019;278:267–72.
114. Lacey B, Golledge J, Yeap BB, et al. Physical activity and vascular disease in a prospective cohort study of older men: The Health In Men Study (HIMS). *BMC Geriatr.* 2015;15:164.
115. Patel A, Berdunov V, Quayyum Z, King D, Knapp M, and Wittenberg R. Estimated societal costs of stroke in the UK based on a discrete event simulation. *Age Ageing* 2020;49:270–6.
116. Buchner DM, Rillamas-Sun E, Di C, et al. Accelerometer-Measured Moderate to Vigorous Physical Activity and Incidence Rates of Falls in Older Women. *J. Am. Geriatr. Soc.* 2017;65:2480–7.

TABLES & FIGURES

Table 1. Base Case Model Parameter Estimates and Sources

Parameter	Value	Sources
Population Details		
Health plan size (# members) ^a	1,000,000	Assumption
% of plan aged 40-60 ^a	32%	US Census Bureau 2023 ⁵³
% of cohort that are insufficiently active ^a	50%	Matthews 2021 ⁵⁴
% eligible and willing to adopt Fitbit intervention ^a	5%	Assumption
Cohort mean starting age (years)	50	Assumption
% female	50%	US Census Bureau 2023 ⁵³
Intervention Details		
Usual care cost	\$0	Assumption
Fitbit intervention cost	\$250	Retail Fitbit pricing ^b ; assumptions
Baseline number of daily steps	6,500	Amorim 2019; ⁵⁵ Ashe 2015; ⁵⁶ Cadmus-Bertram 2019; ⁵⁷ Duscha 2018; ⁵⁸ Katz 2018; ⁵⁹ Li 2018; ⁶⁰ Paxton 2018; ⁶¹ Van Blarigan 2019 ⁶²
Mean peak change in daily steps for Fitbit intervention	1,287	Amorim 2019; ⁵⁵ Ashe 2015; ⁵⁶ Cadmus-Bertram 2019; ⁵⁷ Duscha 2018; ⁵⁸ Katz 2018; ⁵⁹ Li 2018; ⁶⁰ Paxton 2018; ⁶¹ Van Blarigan 2019 ⁶²
Durability of Fitbit intervention (years) ^c	5	Chaudhry 2020; ¹⁹ Gasana 2023; ⁶³ assumptions
Clinical Details		
% of diabetes with comorbid obesity	50%	Leung 2017 ⁶⁸
% of diabetes with comorbid hypertension	50%	Wang 2017 ⁷⁰
Background all-cause mortality	Age-based	Arias 2023 ⁷³
Excess all-cause mortality hazard ratios		
Obesity (age 35-49)	2.40	Ward 2022 ⁸³
Obesity (age 50-69)	1.90	Ward 2022 ⁸³
Obesity (age 70+)	1.50	Ward 2022 ⁸³
Type 2 diabetes	1.95	Preis 2009; ⁸¹ Li 2019 ⁷⁷
Hypertension	1.41	Ford 2011; ⁷⁵ Aune 2021 ⁷⁴
GERD	1.00	Richter 2018; ⁸² Ness-Jensen 2020 ⁷⁹
Major depressive disorder	1.70	Pratt 2016; ⁸⁰ Gilman 2017 ⁷⁶
Sleep apnea	1.40	Lin 2023 ⁷⁸
Healthcare Cost Details^d		
Annual short-term healthcare costs (Years 1 and 2) ^e		
Usual care (per participant)	\$4,344	Teigland 2022; ⁵² Leung 2017; ⁶⁸ assumptions

Parameter	Value	Sources
Fitbit intervention (per participant)	\$3,811	Teigland 2022; ⁵² Leung 2017; ⁶⁸ assumptions
Annual health state costs^f		
Age 40-64		
Well	\$4,720	Teigland 2022; ⁵² Leung 2017 ⁶⁸
Obesity	\$1,093	Leung 2017 ⁶⁸
Type 2 diabetes	\$9,673	ADA 2018; ⁶⁴ Leung 2017; ⁶⁸ Wang 2017 ⁷⁰
Hypertension	\$3,001	Kirkland 2018; ⁶⁷ Wang 2017 ⁷⁰
Diabetes + obesity	\$14,179	Leung 2017 ⁶⁸
Diabetes + hypertension	\$12,072	Wang 2017 ⁷⁰
GERD	\$7,341	Sharma 2023 ⁶⁹
Major depressive disorder	\$6,886	Greenberg 2021 ⁶⁵
Sleep apnea	\$9,222	Hong 2020 ⁶⁶
Age ≥65		
Well	\$6,554	Leung 2017; ⁶⁸ Wang 2017 ⁷⁰
Obesity	\$1,327	Leung 2017 ⁶⁸
Type 2 diabetes	\$9,882	ADA 2018; ⁶⁴ Leung 2017; ⁶⁸ Wang 2017 ⁷⁰
Hypertension	\$2,853	Kirkland 2018; ⁶⁷ Wang 2017 ⁷⁰
Diabetes + obesity	\$11,561	Leung 2017 ⁶⁸
Diabetes + hypertension	\$9,713	Wang 2017 ⁷⁰
GERD ^e	\$4,918	Sharma 2023 ⁶⁹
Major depressive disorder ^g	\$4,614	Greenberg 2021 ⁶⁵
Sleep apnea	\$11,538	Chhatre 2020 ⁷¹
Utility/Disutility Details^h		
Well		
Age 40-49	0.871	Sullivan 2006 ⁸⁴
Age 50-59	0.842	Sullivan 2006 ⁸⁴
Age 60-69 ⁱ	0.823	Sullivan 2006 ⁸⁴
Age 70-79 ⁱ	0.790	Sullivan 2006 ⁸⁴
Age ≥80 ⁱ	0.736	Sullivan 2006 ⁸⁴
Obesity		
Type 2 diabetes	-0.127	Sullivan 2006 ⁸⁴
Hypertension	-0.071	Shah 2020; ⁸⁹ Sullivan 2006 ⁸⁴
Diabetes + obesity	-0.057	Shah 2020; ⁸⁹ Sullivan 2006 ⁸⁴
Diabetes + hypertension	-0.127	Minimum of diabetes, obesity
GERD	-0.071	Minimum of diabetes, hypertension
Major depressive disorder	-0.058	McCarty 2022; ⁸⁸ Sullivan 2006 ⁸⁴
Sleep apnea	-0.156	Hanmer 2016; ⁸⁵ Jain 2022; ⁸⁶ Janicak 2013; ⁸⁷ Sullivan 2006 ⁸⁴
Miscellaneous Analysis Details		
Budget impact time horizon (years)	2 and 5	Assumption
Cost-effectiveness time horizon (years)	15	Assumption; until age 65
Annual discount rate (costs, QALYs) ^j	3%	Lipscomb 1996; ⁹³ Weinstein 1996 ⁹⁴

Abbreviations: GERD, gastroesophageal reflux disease; QALY, quality-adjusted life-year

^aBudget impact analysis only; ^bThe cost of the Fitbit intervention is estimated based on the retail price of a Charge 6 device, in addition to a 12-month subscription to Fitbit Premium (which provides goal setting, feedback, and access to a library of workout content and additional detailed metrics in-app), as well as additional intervention deployment costs; ^cAfter peaking at 1 year, the mean daily step count is assumed to decrease linearly until returning to baseline at 5 years; ^dCosts from published literature not reported in 2023 USD were inflated using the Medical Care consumer price index;⁷² ^eThe displayed cost estimates were calculated from an economic analysis of the Silver Sneakers program (Teigland 2022⁵²) with adjustments to better match our modeled cohort in terms of age (50% age-based reduction factor) and intervention efficacy (25% efficacy reduction factor); these adjusted cost estimates were then validated by comparison to published annual healthcare costs of a ‘well’ population (Leung 2017); ^fIncident disease costs are incremental to the ‘Well’ health state cost; Commercial costs were used for participants aged 40-64 and Medicare costs were used for participants aged ≥ 65 (only in lifetime sensitivity analyses); ^gMedicare cost was estimated by reducing the published Commercial cost by 33%; ^hDisutilities were used for incident diseases to avoid the counterintuitive result in which the utility of an incident disease health state is higher (more favorable) than the ‘Well’ state which could occur at older ages; ⁱAges ≥ 65 are only relevant in lifetime sensitivity analyses; ^jCost-effectiveness analysis only

Table 2. Intervention Efficacy (Mean Daily Steps per Day)^a

Mean Number of Daily Steps	Year 0	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7^b
Usual Care Cohort	6,500	6,500	6,500	6,500	6,500	6,500	6,500	6,500
Fitbit Cohort	6,500	7,787	7,465	7,144	6,822	Equivalent to Usual Care		
Incremental Steps	0	1,287	965	644	322	0	0	0

Sources: STEPS—Amorim 2019;⁵⁵ Ashe 2015;⁵⁶ Cadmus-Bertram 2019;⁵⁷ Duscha 2018;⁵⁸ Katz 2018;⁵⁹ Li 2018;⁶⁰ Paxton 2018;⁶¹ Van Blarigan 2019;⁶² DURABILITY—Chaudhry 2020;¹⁹ Gasana 2023;⁶³ assumptions

^aSensitivity analyses allowed the mean number of daily steps in the Fitbit cohort to return to baseline as early as Year 2 or as late as Year 6; ^bYear 7 daily steps are carried forward in subsequent years

Table 3. Base Case Economic Analysis Results

	Usual Care	Fitbit	Incremental
Cost Effectiveness^a			
Total costs	\$89,541	\$88,284	-\$1,257
Total LYs	11.490	11.493	0.003
Total QALYs	9.031	9.042	0.011
ICER (cost/LY gained)			Fitbit dominates UC ^c
ICER (cost/QALY gained)			Fitbit dominates UC ^c
Budget Impact^b			
2-year total costs	\$69,496,000	\$62,982,000	-\$6,514,000
2-year incremental budget impact (PMPM)			-\$0.27
5-year total costs	\$220,166,612	\$211,617,969	-\$8,548,644
5-year incremental budget impact (PMPM)			-\$0.14

Abbreviations: ICER, incremental cost-effectiveness ratio; LY, life-year; PMPM, per-member per-month; QALY, quality-adjusted life-year; UC, usual care

^aExpected costs and outcomes per participant quantified over a 15-year time horizon, discounted at 3% annually; ^bCosts for a cohort of 8,000 participants over both a 2-year and 5-year time horizon (undiscounted); ^c“Dominates” indicates that Fitbit is less costly and more effective than UC





Figure 1. A simplified schematic of the physical activity intervention model. During and after a physical activity intervention (Fitbit or Usual Care), short-term (years 1 and 2) and long-term (post-2 years) healthcare resource use and costs are quantified. The long-term impact of the interventions on 6 conditions associated with insufficient physical activity also is considered in terms of survival and quality-adjusted survival. Abbreviations: GERD, gastroesophageal reflux disease; HTN, hypertension; LY, life-year; MDD, major depressive disorder; OBES, obesity; QALY, quality-adjusted life-year; SA, sleep apnea; T2DM, type 2 diabetes mellitus

A)



B)




Figure 2. Stratified cost results of the A) cost-effectiveness and B) budget impact analyses. The time horizons of the analyses are 15 years and 5 years, respectively. The majority of the cost savings is associated with reductions in healthcare resource use in the first two years after initiation of the interventions.




Figure 3. Scatterplot of the probabilistic sensitivity analysis (5,000 model runs). Incremental costs and QALYs of the Fitbit intervention compared to Usual Care are demonstrated. The orange triangle denotes the base case ICER. ICER lines representing \$50,000, \$100,000, and \$150,000/QALY thresholds are displayed as is the 95% confidence interval ellipse. Abbreviations: CI, confidence interval; ICER, incremental cost-effectiveness ratio; QALY, quality-adjusted life-year




Figure 4. Cost-effectiveness acceptability curve. At all willingness to pay thresholds (\$/QALY gained), a physical activity intervention including a Fitbit is the preferred treatment strategy over Usual Care. Abbreviation: QALY, quality-adjusted life-year

SUPPLEMENTARY MATERIALS

Supplementary Text

Calculation of step improvement for Fitbit

The increase in daily step count associated with the Fitbit intervention was estimated from 8 RCTs that compared a Fitbit-based intervention to a control group without the device.⁵⁵⁻⁶² The mean step count difference between the two comparators was calculated using two approaches: 1) a “mean change” or difference in difference approach that accounts for disparities in the baseline step counts for the Fitbit and control groups; and 2) a “post-intervention” approach that assumes baseline step counts are not significantly different between the two groups and only considers the difference in the final step count estimates. In their meta-analysis of Fitbit RCTs, Ringeval and colleagues (2020)²³ used a mix of these two approaches in their calculations. We preferred consistency and chose the “mean change” approach for our estimation as it yielded a more conservative result (mean difference of +1,287 steps) compared to the “post-intervention” approach (+1,677 steps).

Evaluation of model structural uncertainty

As recommended in good research practice guidelines,^{41,42,44} model structural uncertainty should be evaluated. The first method employed to meet this recommendation was to substitute a “tiered steps approach” for the “mean steps approach” used as the model default. The latter uses the simple weighted average daily step improvement quantified from RCTs using a Fitbit device.⁵⁵⁻⁶² The tiered steps approach assumes a more complex (and realistic) structure in which the mean daily steps are stratified into 3 tiers at baseline: 1) 5,500 daily steps (25% of cohort); 2) 6,500 daily steps (50%); and 3) 7,500 daily steps (25%). As with the base case, for usual care, there is no improvement in mean daily steps. For the Fitbit intervention for the 3 groups, there is improvement of 0 steps in tier 1, 1,200 daily steps in tier 2, and 2,700 steps in tier 3. The distribution and daily step count improvement are arbitrary but net out to similar baseline and maximum improvement step counts as with the base case—6,500 and 1,275, respectively. This attempt to instill heterogeneity into the population and daily step count model structure appears to have minimal impact on the results.

A second approach for examining structural uncertainty pertains to the types of resource use and costs that might be considered in the initial 2-year period. In the base case, short-term costs consisted solely of adjusted estimates from a U.S. economic analysis of the Silver Sneakers program.⁵² It was uncertain whether inclusion of health state costs ('Well' and incident diseases) in the initial 2-year period would represent a more thorough accounting of health costs or the degree to which some costs would be double-counted. Therefore, a modification to the model programming was made to allow for health state cost inclusion in all model cycles. As expected, given that the Fitbit intervention leads to reductions in incident disease, inclusion of those costs in the first two years led to a modest increase in cost savings. While the conclusions of the analysis remained unchanged with this assessment of model uncertainty, we maintain that the conservative method (exclusion of health state costs in the initial 2-year period) remains the desired base case approach.

Exploratory analysis of an older, insufficiently active population

The base case model and analysis was designed to estimate the impact of the Fitbit intervention on physical activity in a middle-aged cohort (starting age = 50) in terms of an increase in the number of daily steps and reductions in medical resource use, costs, and incident disease. Few economic analyses have evaluated physical activity interventions in older adults (age ≥ 65) in terms of daily steps and none, to our knowledge, have linked steps to a robust source of incident disease data which remains a gap warranting future research. However, a recent economic analysis by Deidda and colleagues (2022)³¹ in older adults did provide sufficient data to inform an exploratory analysis of the cost effectiveness of a Fitbit intervention in a Medicare population.

The Deidda study examined the cost-effectiveness of physical activity interventions (an exercise referral scheme [ERS] with or without self-management strategies [SMS]) using a long-term economic model extending the analysis of the SITLESS randomized controlled trial in older adults. The model used two physical activity health states ('active' and 'inactive'), corresponding to maintaining (or not) at least 150 minutes of moderate to vigorous aerobic physical activity, and seven health states representing conditions associated with an inactive lifestyle in an older population (e.g., Type 2 diabetes, Alzheimer's disease, falls, etc.). This

approach differs from our model concept which uses daily step counts and chronic diseases more relevant to a middle-aged population. Therefore, some modification to both model structure and parameter estimates was required for our exploratory analysis. Furthermore, given that Medicare is the predominant U.S. payer for a population aged 65+, the time horizon for the cost-effectiveness analysis was increased from 15 years to a lifetime horizon.

Population

The target population of interest for the exploratory analysis was older adults with insufficient physical activity, aged ≥ 65 (starting age=65), insured by Medicare. The age and sex distribution was based on 2022 data from the U.S. Census Bureau.⁵³ The proportion with insufficient physical activity in this population was informed by a systematic review of 23 published reports of the prevalence of sedentary behavior in older adults.¹⁰⁰ Population details are shown in **Supplementary Table 4**.

Comparators

Unlike our base case analysis, Deidda compared Usual Care to physical activity interventions that included an exercise referral scheme. Such schemes have a mixed history with some economic analyses suggesting unfavorable cost effectiveness,^{32,38} favorable cost effectiveness,³⁵ or as Deidda reported, potential cost savings. In their analysis, Deidda and colleagues used data from the SITLESS RCT to generate probabilities of moving between the two physical activity health states for each of the modeled comparators. A different set of transition probabilities was used for the first year in the analysis than all subsequent years. To estimate the incidence of acute (i.e., falls) and chronic conditions (e.g., Alzheimer's disease) related to insufficient physical activity in the older population over time, the transition probabilities and published incidence data (stratified by activity level) were utilized. Unit costs, utilities, and mortality rates were then applied to the acute/chronic health states to calculate total costs and QALYs (**Supplementary Table 4**).

Our model structure was modified to better accommodate the approach used by Deidda and colleagues. First, a review of U.S. costs, utilities, disutilities, and mortality for the health state

conditions in the Deidda model was conducted. Based on the findings, the health states included in the exploratory analysis were: 1) Alzheimer's disease/dementia; 2) Type 2 diabetes; 3) cardiovascular disease; and 4) falls. Second, the ERS transition probabilities were adjusted so that the annual proportion of patients in the active and inactive health states for Fitbit were equivalent to Usual Care at 5 years and beyond (a similar durability assumption to our base case) rather than maintaining an advantage in physical activity relative to Usual Care as was the case for ERS in the Deidda study. Third, the health state distributions were used in conjunction with the disease incidence data to calculate a weighted average incidence for each condition for each comparator. These incidence estimates were then applied recursively over time to the proportion alive in each annual model cycle to calculate the proportion in each of the health states. Finally, unit costs, utilities/disutilities, and mortality rates were applied to each health state to calculate total costs and QALYs.

Costs

As in the base case analysis, the costs in the exploratory model include those related to the Fitbit intervention (incremental to usual care which is assumed to have no cost), short-term healthcare costs (2 years), and health state costs ('Well' and incident conditions incremental to the 'Well' health state). Unit costs for the model were derived primarily from published sources.^{52,64,68,70,101-103} When applicable, published costs were inflated to 2023 USD using the Medical Care component of the U.S. CPI (**Supplementary Table 4**).⁷²

In the exploratory analyses, we assumed a greater potential for some short-term cost reduction given the similarity in our population to the Medicare Advantage enrollees participating in the Silver Sneakers® (SS) healthy aging program.⁵² Because the SS intervention is not an exact match to those in our model, we maintained the efficacy reduction factor from the base case analysis (assuming the SS program has a greater impact on resource use/cost). Longer-term costs associated with the model health states ('Well' and incident acute and chronic conditions) were quantified for both comparators in a similar fashion to the base case analysis.

Mortality

The annual probability of death for participants in the ‘Well’ health state was based on age- and sex-adjusted all-cause mortality data from U.S. life tables.⁷³ Published hazard ratios^{77,81,104,105} describing the excess mortality associated with each of the health conditions of interest were used to adjust background all-cause mortality and the annual probability of death for those diseases. Falls were assumed to be an acute event that could occur in any cycle with a conditional probability of death for any fall event (**Supplementary Table 4**).¹⁰³

Utilities

Utilities for the ‘Well’ health state (by age strata) were based on a nationally representative catalog of EuroQol-5 Dimension (EQ-5D) utility scores.⁸⁴ For the acute/chronic conditions in the model, we identified published disutilities¹⁰⁶⁻¹⁰⁸ (or utilities that were subsequently used to derive disutilities using information in Sullivan, et al. 2006⁸⁴) that were applied to the ‘Well’ health state utility to account for the worse quality of life for those conditions.

Supplementary Table 1. Annual Incidence Rates for Chronic Diseases in the Model^a

Chronic Disease	Year 0	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7	Year 8	Year 9	Year 10	Year 11
Usual Care												
Obesity	0.00%	1.01%	1.02%	1.03%	2.10%	2.14%	2.85%	2.93%	3.23%	3.55%	3.90%	4.29%
Type 2 diabetes	0.00%	0.26%	0.26%	0.26%	0.51%	0.52%	0.64%	0.65%	0.71%	0.78%	0.86%	0.95%
Hypertension	0.00%	1.34%	1.36%	1.38%	3.28%	3.39%	4.02%	4.19%	4.61%	5.07%	5.58%	6.14%
GERD	0.00%	1.75%	1.78%	1.81%	3.16%	3.27%	4.50%	4.72%	5.19%	5.71%	6.28%	6.91%
MDD	0.00%	1.40%	1.42%	1.44%	2.87%	2.95%	3.84%	3.99%	4.39%	4.83%	5.31%	5.84%
Sleep apnea	0.00%	0.92%	0.93%	0.94%	1.77%	1.80%	1.97%	2.01%	2.21%	2.43%	2.68%	2.95%
Fitbit Intervention												
Obesity	0.00%	0.90%	0.93%	0.96%	2.05%							
Type 2 diabetes	0.00%	0.24%	0.24%	0.25%	0.50%							
Hypertension	0.00%	1.23%	1.26%	1.31%	3.19%							
GERD	0.00%	1.58%	1.65%	1.73%	3.12%							
MDD	0.00%	1.25%	1.29%	1.36%	2.76%							
Sleep apnea	0.00%	0.79%	0.82%	0.87%	1.69%							

Equivalent to Usual Care

Sources: STEPS—Amorim 2019;⁵⁵ Ashe 2015;⁵⁶ Cadmus-Bertram 2019;⁵⁷ Duscha 2018;⁵⁸ Katz 2018;⁵⁹ Li 2018;⁶⁰ Paxton 2018;⁶¹ Van Blarigan 2019;⁶² INCIDENCE AT STEP COUNT—Master 2022¹³

Abbreviations: GERD, gastroesophageal reflux disease; MDD, major depressive disorder

^aIncidence estimates from the All of Us study concluded at Year 7; therefore an annual growth rate factor of 10% was applied in Years 8-11 to simulate the increasing incidence over time observed in the All of Us Study; starting at Year 11, incidence rates were held constant for the remainder of the analysis


Supplementary Table 2. Parameters Used in Probabilistic Sensitivity Analyses (Cost Effectiveness Only)

Parameter	Value	SE	Distribution	Source Informing SE^b
Intervention Details				
Baseline number of daily steps	6,500	125	Normal	Ringeval 2020 ²³
Mean peak change in daily steps for Fitbit intervention	1,287	125	Normal	Ringeval 2020 ²³
Durability of Fitbit intervention (years) ^a	5	0.05	Normal	Chaudhry 2020; ¹⁹ Gasana 2023, ⁶³ assumption
Clinical Details				
% of diabetes with comorbid obesity ^a	50%	10%	Beta	Assumption
% of diabetes with comorbid hypertension ^a	50%	10%	Beta	Assumption
Excess all-cause mortality hazard ratios				
Obesity (age 35-49)	2.40	0.08	Lognormal	Ward 2022 ⁸³
Obesity (age 50-69)	1.90	0.05	Lognormal	Ward 2022 ⁸³
Obesity (age 70+)	1.50	0.03	Lognormal	Ward 2022 ⁸³
Type 2 diabetes	1.95	0.16	Lognormal	Preis 2009 ⁸¹
Hypertension	1.41	0.03	Lognormal	Aune 2021 ⁷⁴
Major depressive disorder	1.70	0.13	Lognormal	Pratt 2016; ⁸⁰ Gilman 2017 ⁷⁶
Sleep apnea	1.40	0.20	Lognormal	Lin 2023 ⁷⁸
Annual incidence growth rate factor after 7 years	10%	1%	Beta	Assumption
Healthcare Cost Details^c				
Annual short-term healthcare costs (Years 1 and 2)				
Usual care (per participant)	\$8,687	\$869	Gamma	Gc 2018; ³⁹ assumption
Fitbit intervention (per participant)	\$7,268	\$727	Gamma	Gc 2018; ³⁹ assumption
Age-based adjustment factor	50%	10.8%	Beta	Assumption
Efficacy-based adjustment factor	25%	10.8%	Beta	Assumption
Annual health state costs				
Age 40-64				
Well	\$4,720	\$472	Gamma	Gc 2018; ³⁹ assumption
Obesity	\$1,093	\$109	Gamma	Gc 2018; ³⁹ assumption
Type 2 diabetes	\$9,673	\$967	Gamma	Gc 2018; ³⁹ assumption
Hypertension	\$3,001	\$300	Gamma	Gc 2018; ³⁹ assumption
Diabetes + obesity	\$14,179	\$1,418	Gamma	Gc 2018; ³⁹ assumption

Parameter	Value	SE	Distribution	Source Informing SE ^b
Diabetes + hypertension	\$12,072	\$1,207	Gamma	Gc 2018; ³⁹ assumption
GERD	\$7,341	\$734	Gamma	Gc 2018; ³⁹ assumption
Major depressive disorder	\$6,886	\$689	Gamma	Gc 2018; ³⁹ assumption
Sleep apnea	\$9,222	\$922	Gamma	Gc 2018; ³⁹ assumption
Age ≥65				
Well	\$6,554	\$655	Gamma	Gc 2018; ³⁹ assumption
Obesity	\$1,327	\$133	Gamma	Gc 2018; ³⁹ assumption
Type 2 diabetes	\$9,882	\$988	Gamma	Gc 2018; ³⁹ assumption
Hypertension	\$2,853	\$285	Gamma	Gc 2018; ³⁹ assumption
Diabetes + obesity	\$11,561	\$1,156	Gamma	Gc 2018; ³⁹ assumption
Diabetes + hypertension	\$9,713	\$971	Gamma	Gc 2018; ³⁹ assumption
GERD ^c	\$4,918	\$492	Gamma	Gc 2018; ³⁹ assumption
Major depressive disorder	\$4,614	\$461	Gamma	Gc 2018; ³⁹ assumption
Sleep apnea	\$11,538	\$1,154	Gamma	Gc 2018; ³⁹ assumption
Disutility Details^d				
Obesity	-0.127	-0.0009	Beta	Sullivan 2006 ⁸⁴
Type 2 diabetes	-0.071	-0.0004	Beta	Sullivan 2006 ⁸⁴
Hypertension	-0.057	-0.0005	Beta	Sullivan 2006 ⁸⁴
GERD	-0.058	-0.0005	Beta	Sullivan 2006 ⁸⁴
Major depressive disorder	-0.156	-0.0036	Beta	Hanmer 2016; ⁸⁵ Sullivan 2006 ⁸⁴
Sleep apnea	-0.075	-0.0126	Beta	Hanmer 2016; ⁸⁵ Walia 2017 ⁹⁰

Abbreviations: GERD, gastroesophageal reflux disease; SE, standard error

^aSome parameters were truncated to ensure realistic values—Fitbit durability forced to be in the 2-6 year range; diabetes comorbidity percentages ≤50%; ^bWhen assumptions were required for a standard error estimate, 95% confidence intervals were approximated to ensure that ranges over which the parameter would vary were reasonable; ^cIn most cases, published costs were not reported with a measure of uncertainty; therefore, a standard error of 10% was assumed as was the approach in Gc 2018³⁹ (an economic analysis of physical activity interventions); ^dDisutilities may be expressed as negative numbers but a beta distribution is limited to positive numbers; therefore, the absolute values of the disutility and standard errors were used and then the sampled estimate used in the model was converted to a negative number



Supplementary Figure 1. One-way sensitivity analysis tornado diagram of net monetary benefit (at a willingness to pay threshold of \$150,000/QALY gained) in the cost-effectiveness analysis. Lower and upper bound parameter estimates for each variable are listed within the diagram. Abbreviations: GERD, gastroesophageal reflux disease; HR, hazard ratio; yrs, years

Supplementary Table 3. Additional Scenario Analysis Results

	Costs			QALYs			ICER or PMPM
	Usual Care	Fitbit	Incremental	Usual Care	Fitbit	Incremental	
Base Case							
Cost effectiveness ^a	\$89,541	\$88,284	-\$1,257	9.031	9.042	0.011	Fitbit dominant ^b
Budget impact ^a	\$220,166,612	\$211,617,969	-\$8,548,644	NA	NA	NA	-\$0.14
Structural Uncertainty							
Tiered steps approach^c							
Cost effectiveness	\$89,539	\$88,242	-\$1,297	9.032	9.041	0.009	Fitbit dominant ^b
Budget impact	\$220,169,841	\$211,713,811	-\$8,456,030	NA	NA	NA	-\$0.14
Inclusion of health state costs in Years 1 & 2							
Cost effectiveness	\$99,888	\$98,522	-\$1,367	9.031	9.042	0.011	Fitbit dominant ^b
Budget impact	\$304,209,839	\$294,766,789	-\$9,443,050	NA	NA	NA	-\$0.16
Scenarios							
Use of post-intervention steps estimate							
Cost effectiveness	\$89,541	\$88,173	-\$1,368	9.031	9.045	0.014	Fitbit dominant ^b
Budget impact	\$220,166,612	\$211,061,720	-\$9,104,892	NA	NA	NA	-\$0.15
Reduced population eligibility and adoption rate^d							
Budget impact	\$110,083,306	\$105,808,984	-\$4,274,322	NA	NA	NA	-\$0.07
Increased population eligibility and adoption rate^d							
Budget impact	\$330,249,918	\$317,426,953	-\$12,822,965	NA	NA	NA	-\$0.21
Reduced cumulative incidence of chronic disease^e							

	Costs			QALYs			ICER or PMPM
	Usual Care	Fitbit	Incremental	Usual Care	Fitbit	Incremental	
Cost effectiveness	\$85,736	\$84,540	-\$1,196	9.105	9.114	0.009	Fitbit dominant ^b
Budget impact	\$212,916,291	\$204,778,351	-\$8,137,940	NA	NA	NA	-\$0.14
Increased cumulative incidence of chronic disease^e							
Cost effectiveness	\$92,582	\$91,280	-\$1,302	8.972	8.984	0.012	Fitbit dominant ^b
Budget impact	\$227,009,258	\$218,097,985	-\$8,911,272	NA	NA	NA	-\$0.15
Lifetime horizon							
Cost effectiveness	\$162,005	\$160,710	-\$1,295	13.463	13.476	0.013	Fitbit dominant ^b
Health state costs only^f							
Cost effectiveness	\$91,328	\$91,010	-\$318	9.031	9.042	0.011	Fitbit dominant ^b
Budget impact	\$234,713,839	\$233,784,789	-\$929,050	NA	NA	NA	-\$0.02

Abbreviations: ICER, incremental cost-effectiveness ratio; PMPM, per-member per-month; QALY, quality-adjusted life-year

^a15-year time horizon for CEA, 5 year horizon for BIA; ^b“Dominant” indicates that Fitbit is less costly and more effective than Usual Care; ^cThe tiered approach uses the following 3 strata (“tiers”) of step counts for Usual Care (25% with 5,500 mean daily steps; 50% with 6,500 mean daily steps; and 25% with 7,500 mean daily steps) none of whom have increased mean daily step counts over time; the 3 strata for the Fitbit intervention are the same at baseline but have maximum improvements of 0 steps (strata 1), 1,200 steps (strata 2), and 2,700 steps (strata 3); ^dThe base case adoption rate estimate (5%) was decreased and increased to 2.5% and 7.5%, respectively; ^eChronic disease cumulative incidence estimates were set to the lower and upper limits of the linear function obtained from the *All of Us* researchers; ^fShort-term healthcare costs in Years 1 and 2 were eliminated and health state costs—‘Well’ and incident diseases—added to the first 2 years

Supplementary Table 4. Exploratory Analysis Model Parameter Estimates and Sources

Parameter	Value	Sources
Population Details		
Health plan size (# members) ^a	1,000,000	Assumption
% of plan aged 65+ ^a	100%	US Census Bureau 2023 ⁵³
% of cohort that is insufficiently active ^a	67%	Harvey 2013 ¹⁰⁰
% eligible and willing to adopt Fitbit intervention ^a	5%	Assumption
Cohort mean starting age (years)	65	Assumption
% female	55%	US Census Bureau 2023 ⁵³
Intervention Details		
Usual care cost	\$0	Assumption
Fitbit intervention cost	\$250	Retail Fitbit pricing ^b ; assumptions
Durability of Fitbit intervention (years) ^c	5	Deidda 2022, ³¹ assumptions
Clinical Details		
Background all-cause mortality	Age-based	Arias 2023 ⁷³
Annual acute condition/chronic disease incidence		
Type 2 diabetes (active)	0.0034	InterAct Consortium 2012 ¹⁰⁹ in Deidda 2022 ³¹
Type 2 diabetes (inactive)	0.0049	InterAct Consortium 2012 ¹⁰⁹ in Deidda 2022 ³¹
Alzheimer’s disease/dementia (active)	0.0125	Abbott 2004; ¹¹⁰ Podewils 2005; ¹¹¹ Larsson 2018 ¹¹² in Deidda 2022 ³¹
Alzheimer’s disease/dementia (inactive)	0.0167	Abbott 2004; ¹¹⁰ Podewils 2005; ¹¹¹ Larsson 2018 ¹¹² in Deidda 2022 ³¹
Cardiovascular disease (active)	0.0255	Jefferis 2019; ¹¹³ Lacey 2015, ¹¹⁴ Patel 2020 ¹¹⁵ in Deidda 2022 ³¹
Cardiovascular disease (inactive)	0.0410	Jefferis 2019; ¹¹³ Lacey 2015, ¹¹⁴ Patel 2020 ¹¹⁵ in Deidda 2022 ³¹
Falls (active)	0.0776	Buchner 2017 ¹¹⁶ in Deidda 2022 ³¹
Falls (inactive)	0.1085	Buchner 2017 ¹¹⁶ in Deidda 2022 ³¹
Excess all-cause mortality hazard ratios or % ^c		
Type 2 diabetes	1.95	Preis 2009; ⁸¹ Li 2019 ⁷⁷
Alzheimer’s disease/dementia	2.00	Lanctot 2024 ¹⁰⁴
Cardiovascular disease	1.40	Shaked 2022 ¹⁰⁵
Falls	0.75%	Burns 2016 ¹⁰³
Healthcare Cost Details^d		
Annual short-term healthcare costs (Years 1 and 2)		
Usual care (per participant)	\$8,687	Teigland 2022 ⁵²
Fitbit intervention (per participant)	\$7,268	Teigland 2022; ⁵² assumption
Efficacy-based adjustment factor	25%	Assumption
Annual health state costs ^e		
Well	\$6,554	Leung 2017; ⁶⁸ Wang 2017 ⁷⁰
Type 2 diabetes	\$9,882	ADA 2018; ⁶⁴ Leung 2017; ⁶⁸ Wang 2017 ⁷⁰

Parameter	Value	Sources
Alzheimer's disease/dementia	\$31,193	Alzheimer's Association 2023 ¹⁰¹
Cardiovascular disease	\$5,066	Birger 2021 ¹⁰²
Falls	\$12,650	Burns 2016 ¹⁰³
Utility/Disutility Details^f		
Well		
Age 60-69	0.823	Sullivan 2006 ⁸⁴
Age 70-79	0.790	Sullivan 2006 ⁸⁴
Age ≥80	0.736	Sullivan 2006 ⁸⁴
Type 2 diabetes	-0.071	Sullivan 2006 ⁸⁴
Alzheimer's disease/dementia	-0.180	Vandepitte 2020 ¹⁰⁸ in Deidda 2022 ³¹
Cardiovascular disease	-0.060	Lacey 2003 ¹⁰⁷ in Deidda 2022 ³¹
Falls	-0.160	Bjerk 2019 ¹⁰⁶ in Deidda 2022 ³¹
Miscellaneous Analysis Details		
Budget impact time horizon (years)	2 and 5	Assumption
Cost-effectiveness time horizon (years)	Lifetime	Assumption
Annual discount rate (costs, QALYs) ^g	3%	Lipscomb 1996; ⁹³ Weinstein 1996 ⁹⁴

Abbreviations: QALY, quality-adjusted life-year

^aBudget impact analysis only; ^bThe cost of the Fitbit intervention is estimated based on the retail price of a Charge 6 device, in addition to a 12-month subscription to Fitbit Premium (which provides goal setting, feedback, and access to a library of workout content and additional detailed metrics in-app), as well as additional intervention deployment costs; ^cUnlike the base case analysis, the exploratory analysis does not use daily step count as an efficacy measure but rather uses the transition probabilities in Deidda 2022³¹ as an efficacy measure and for calculation of health condition incidence; using adjusted transition probabilities and incidence estimates from the Deidda study, annual incidence rates converge between the Usual Care and Fitbit groups by 5 years; ^dCosts from published literature not reported in 2023 USD were inflated using the Medical Care consumer price index; ^eIncident disease costs are incremental to the 'Well' health state cost; ^fDisutilities were used for incident diseases to avoid the counterintuitive result in which the utility of an incident disease health state is higher (more favorable) than the 'Well' state which could occur at older ages; ^gCost-effectiveness analysis only


Supplementary Table 5. Exploratory Economic Analysis Results

	Usual Care	Fitbit	Incremental
Cost Effectiveness^a			
Total costs	\$148,776	\$146,798	-\$1,977
Total LYs	12.131	12.135	0.004
Total QALYs	9.921	9.923	0.002
ICER (cost/LY gained)			Fitbit dominates UC ^c
ICER (cost/QALY gained)			Fitbit dominates UC ^c
Budget Impact^b			
2-year total costs	\$582,029,000	\$510,724,250	-\$62,929,750
2-year incremental budget impact (PMPM)			-\$2.62
5-year total costs	\$1,575,338,659	\$1,509,713,110	-\$65,625,550
5-year incremental budget impact (PMPM)			-\$1.09


Abbreviations: ICER, incremental cost-effectiveness ratio; LY, life-year; PMPM, per-member per-month; QALY, quality-adjusted life-year; UC, usual care

^aExpected costs and outcomes per participant quantified over a lifetime horizon, discounted at 3% annually; ^bCosts for a cohort of 33,500 participants over both a 2-year and 5-year time horizon (undiscounted); ^c“Dominates” indicates that Fitbit is less costly and more effective than UC

A)



B)



Supplementary Figure 2. Stratified cost results of the exploratory A) cost-effectiveness and B) budget impact analyses. The time horizons of the analyses are lifetime and 5 years, respectively. The majority of the cost savings is associated with reductions in healthcare resource use in the first two years after initiation of the interventions.