

# **Full title: Alterations in Aortic Elasticity Indices among Type 2 Diabetes Patients in a Low and Middle Income Country Using M-Mode Echocardiography: A Cross-Sectional Comparative Study**

## **Short title: Alterations in Aortic Elasticity Indices among Type 2 Diabetes Patients**

- 1 Hai Nguyen Ngoc Dang<sup>1¶</sup>, Thang Viet Luong<sup>2¶</sup>, Quan Nguyen Khoi<sup>3,4</sup>, Uyen Ngoc
- 2 Phuong Nguyen<sup>3</sup>, Nguyen Nguyen Khoi Pham<sup>2</sup>, Hieu Thi Nguyen Tran<sup>2</sup>, Hung
- 3 Khanh Tran<sup>2</sup>, Mai Thi Thu Cao<sup>2</sup>, Binh Anh Ho<sup>5</sup>, Thang Chi Doan<sup>5</sup>, Hung Minh
- 4 Nguyen, Tien Anh Hoang<sup>2</sup>, Minh Van Huynh<sup>2</sup>.
  
- 5 <sup>1</sup>The Faculty of Medicine, Duy Tan University, Da Nang, Vietnam
  
- 6 <sup>2</sup>University of Medicine and Pharmacy, Hue University, Hue, Vietnam
  
- 7 <sup>3</sup>College of Health Sciences, VinUniversity, Hanoi, Vietnam
  
- 8 <sup>4</sup>Johns Hopkins University Bloomberg School of Public Health, Baltimore, MD
  
- 9 <sup>5</sup>Cardiovascular Center, Hue Central Hospital, Hue, Vietnam
  
- 10 <sup>6</sup>Bach Mai Hospital, Ha Noi, Vietnam

11 \*Corresponding author:

12 Email: [hatien@huemed-univ.edu.vn](mailto:hatien@huemed-univ.edu.vn) or [hatien@hueuni.edu.vn](mailto:hatien@hueuni.edu.vn)

13 ¶Hai Nguyen Ngoc Dang and Thang Viet Luong are co-first authors and contributed  
14 equally to this work.

15 Author contributions:

16 Conceptualization: Hai Nguyen Ngoc Dang, Thang Viet Luong, Quan Nguyen  
17 Khoi, Nguyen Nguyen Khoi Pham.

18 Data curation: Hai Nguyen Ngoc Dang, Thang Viet Luong.

19 Formal analysis: Hai Nguyen Ngoc Dang, Thang Viet Luong.

20 Investigation: Hai Nguyen Ngoc Dang, Thang Viet Luong.

21 Methodology: Hai Nguyen Ngoc Dang, Thang Viet Luong.

22 Supervision: Hai Nguyen Ngoc Dang, Thang Viet Luong.

23 Writing – original draft: Hai Nguyen Ngoc Dang, Thang Viet Luong, Quan Nguyen  
24 Khoi, Uyen Ngoc Phuong Nguyen, Nguyen Nguyen Khoi Pham, Hieu Thi Nguyen  
25 Tran, Hung Khanh Tran, Mai Thi Thu Cao, Binh Anh Ho, Thang Chi Doan, Hung  
26 Nguyen Minh, Tien Anh Hoang, Minh Van Huynh.

27 Writing – review & editing: Hai Nguyen Ngoc Dang, Thang Viet Luong, Quan  
28 Nguyen Khoi, Uyen Ngoc Phuong Nguyen, Nguyen Nguyen Khoi Pham, Hieu Thi  
29 Nguyen Tran, Hung Khanh Tran, Mai Thi Thu Cao, Binh Anh Ho, Thang Doan  
30 Chi, Hung Nguyen Minh, Tien Anh Hoang, Minh Van Huynh.

31 All authors have read and agreed to the published version of the manuscript.

## 32 **Abstract**

33 **Background:** Diabetes is one of the leading causes of noncommunicable  
34 diseases worldwide. It is known to induce cardiovascular remodeling, which can  
35 result in a variety of complications, including a considerable increase in aortic  
36 stiffness. While studies in Western populations have explored these effects, data on  
37 Asians, particularly Vietnamese, are limited. This study aimed to assess aortic  
38 elasticity in type 2 diabetes mellitus (T2DM) patients compared to healthy  
39 individuals.

40 **Methods:** Aortic elasticity indices were measured using M-mode  
41 echocardiography. A comparison between the healthy group and the T2DM group  
42 revealed substantial differences in aortic elasticity indices.

43 **Results:** The aortic stiffness index (ASI) was significantly greater in the  
44 T2DM group than in the control group, with median values of 6.10 (3.64-12.47)  
45 and 3.79 (2.40-8.50), respectively ( $p = 0.003$ ). Aortic strain (AS) was substantially

46 lower in the T2DM group than in the control group, with median values of 8.21  
47 (4.24 - 13.07) and 10.66 (6.01 - 18.23), respectively ( $p = 0.039$ ). Furthermore, the  
48 median aortic compliance (AC) and aortic distensibility (AD) in individuals with  
49 T2DM were 4.07 (2.28-7.44) and 3.08 (1.57-5.26), respectively, lower than those  
50 in the control group, with median values of 6.40 (3.08-10.75) and 5.33 (2.80-9.79),  
51 respectively. A longer diabetes duration was linked to a greater ASI ( $r = 0.43$ ,  $p <$   
52  $0.05$ ), while the AS decreased ( $r = -0.37$ ,  $p < 0.05$ ).

53 **Conclusions:** Substantial variations in aorta elasticity indices were found  
54 using M-mode echocardiography. More research is needed to investigate the  
55 consequences of these discrepancies and their significance for clinical purposes.

56 **Graphical Abstract**

## 57 **Introduction**

58           Type 2 diabetes mellitus (T2DM) is currently an emerging  
59 noncommunicable illness worldwide. According to the International Diabetes  
60 Federation (IDF), the number of individuals with T2DM is expected to increase  
61 globally, with 10.5% of people aged 20 to 75 years being diagnosed with diabetes  
62 by 2021. By 2045, this figure will increase by 46%, with an estimated 783 million  
63 people living with diabetes. Diabetes caused 2.3 million fatalities in the Western  
64 Pacific region in 2021, the highest among all IDF regions. Among these, Vietnam  
65 has almost 57 thousand diabetes-related fatalities in adults aged 20 to 79 years [1].

66           Many studies have shown that T2DM patients experience changes in both  
67 cardiac morphology and function over time after being diagnosed with the condition  
68 [2,3]. In addition to damaging the heart, vascular remodeling has recently been  
69 shown to occur in patients with diabetes patients. Numerous studies on diabetes  
70 mellitus have revealed a strong link between vascular remodeling and disease,  
71 which is affected by a variety of variables [4–6]. The most important factors are  
72 insulin resistance and dyslipidemia [7]. Furthermore, because diabetes mellitus  
73 involves the risk of persistent hyperglycemia, glucose creates covalent connections  
74 with proteins, nucleic acids, and lipids by nonenzymatic glycosylation, also known  
75 as glycation. This irreversible process results in advanced glycation end products  
76 (AGEs). These chemicals accumulate in the vascular wall and, with time, impair

77 the flexibility of blood vessels [8]. As diabetes progresses, its complications may  
78 worsen. The duration of the condition (since first diagnosis) can be seen as the  
79 natural progression of vascular complications that are also associated with an  
80 increased risk of death from coronary artery disease or stroke due to structural  
81 abnormalities in the cardiovascular system. There are several invasive and  
82 noninvasive procedures for measuring vascular elasticity. These include pulse  
83 pressure, invasive blood pressure, and monitoring variations in blood vessel size.  
84 Imaging techniques such as magnetic resonance imaging, computed tomography,  
85 and ultrasound provide noninvasive but effective methods for measuring arterial  
86 elasticity. Among these approaches, ultrasound stands out for its low cost,  
87 ubiquitous availability, and broad applicability, making it the recommended tool  
88 for measuring arterial elasticity.

89         Aortic elasticity, a measure of arterial stiffness, has been demonstrated to be  
90 a strong independent predictor of atherosclerosis and other cardiovascular events.  
91 This shows that aortic flexibility could be a useful technique for detecting vascular  
92 issues in their early stages [9]. However, most previous studies have concentrated  
93 on Western or Caucasian populations. There is a considerable knowledge gap about  
94 how aortic elasticity may change in people with T2DM in populations of low- and  
95 middle-income countries in general and specifically in the population of Vietnam.  
96 This study was conducted to investigate the indications of aortic elasticity in T2DM

97 patients and compare the aorta elasticity indices of T2DM patients and healthy  
98 control participants to determine patterns of change. By analyzing these parameters,  
99 this study aimed to gain valuable insights into how T2DM affects cardiovascular  
100 health, especially aortic elasticity, and contributes to an effort to detect and reduce  
101 disease burdens globally early.

## 102 **Methods**

### 103 **Study population**

104 This is a quantitative, comparative cross-sectional study with a comparison  
105 group. We integrated components from the STROBE declaration to enhance the  
106 quality of reporting observational studies [10]. This study included 109 subjects  
107 who underwent echocardiography at the Cardiac Ultrasound Unit of the Hue  
108 University of Medicine and Pharmacy from 15/04/2022 to 01/06/2023. The study  
109 received approval from The Institutional Ethics Committee of Hue University of  
110 Medicine and Pharmacy (Approval number: H2022/025) and was conducted under  
111 the principles outlined in the Declaration of Helsinki 2013. Prior to participation,  
112 written informed consent was obtained from each patient or their legal  
113 representative.

114 Among the 109 participants, 47 were healthy subjects who underwent regular  
115 health check-ups, had no history of diabetes or cardiac pathology, and were

116 currently free from cardiac or diabetic conditions. Another 62 individuals were  
117 diagnosed with T2DM and had no known comorbidities according to the diagnostic  
118 criteria stipulated by the American Diabetes Association in 2022 [11].

119 Patients who refused to participate in the study, as well as those with known  
120 comorbidities such as hypertension, were excluded. Patients with any other  
121 cardiovascular disease (severe aortic stenosis, moderate to severe aortic  
122 regurgitation, severe mitral valve stenosis, congenital heart disease,  
123 cardiomyopathy, myocardial ischemia, reduced ejection fraction, arrhythmia, or  
124 chronic atrial fibrillation) were also excluded. We also excluded patients who had  
125 pacemakers inserted, pregnant or breastfeeding women, people who suffered from  
126 acute life-threatening medical illnesses, and those who had hazy or inadequate  
127 ultrasound images.

## 128 **Data collection**

129 The data were collected using an ultrasound machine, paper medical records,  
130 and data collection forms to extract information from the study subjects. This  
131 process involved the following steps:

## 132 **Clinical examination and blood tests**

133           After consenting to participate in this study, the patient's medical history was  
134   obtained, with an emphasis on characteristics such as the time of T2DM diagnosis  
135   and blood sugar management at each visit.

136           Body mass index (BMI) was computed using the following formula:  $BMI =$   
137   weight (kg)/[height (m)  $\times$  height (m)]. Body surface area (BSA) was calculated  
138   using the formula of Du Bois [12]:  $BSA = 0.007184 \times (\text{weight})^{0.425} \times (\text{height})^{0.725}$ .  
139   Blood pressure was measured following the guidelines of the American Heart  
140   Association in 2019 [13,14].

141           Blood samples for biochemical analysis were collected from fasting venous  
142   blood after waking. Biochemical tests were performed on a Roche Cobas E601  
143   automated clinical chemistry analyzer at the Central Testing Unit of Hue University  
144   of Medicine and Pharmacy Hospital.

## 145   **Transthoracic echocardiography**

146           Our study utilized a specialized Affiniti 70 echocardiography machine from  
147   Philips, the Netherlands, with the ability to perform time-motion mode (M-mode),  
148   two-dimensional mode, continuous-wave Doppler, color Doppler, and tissue  
149   Doppler imaging. The ultrasound machine simultaneously recorded the  
150   electrocardiography data along with the echocardiographic images. Transthoracic

151 echocardiography was performed following the American Society of  
152 Echocardiography guidelines for adults [15].

### 153 **Arterial stiffness indices**

154 M-mode echocardiography recorded the diameter of the ascending aorta 3  
155 cm above the aorta valve in a parasternal long-axis image.

156 **Fig 1. The aorta was visualized in M-mode.** AoD: end-diastolic diameter of the  
157 ascending aorta; AoS: end-systolic diameter of the ascending aorta.

158 The end-diastolic diameter of the ascending aorta (AoD) was measured at the  
159 peak of the R wave on the ECG, and the end-systolic diameter of the ascending  
160 aorta (AoS) was measured at the maximum anterior motion of the aorta (see **Fig 1**).  
161 The diameter of the ascending aorta was measured in centimeters (cm). Arterial  
162 stiffness was assessed using five indices: aortic strain (AS) % =  $100 \times (\text{AoS} - \text{AoD})/\text{AoD}$ ,  
163 aortic stiffness index (ASI) =  $\ln(\text{SBP}/\text{DBP})/[(\text{AoS} - \text{AoD})/\text{AoD}]$ ,  
164 aortic compliance (AC) mm/mmHg =  $[10 \times (\text{AoS} - \text{AoD})]/(\text{SBP} - \text{DBP})$ , aortic  
165 distensibility (AD) %mmHg<sup>-1</sup> =  $[1000 \times 2 \times (\text{AoS} - \text{AoD})]/[\text{AoD} \times (\text{SBP} - \text{DBP})]$ ,  
166 and aortic Peterson's elastic modulus (AEM) mmHg =  $[(\text{SBP} - \text{DBP}) \times \text{AoD}]/(\text{AoS} - \text{AoD})$ . Note: AoS refers to the aortic end-systolic diameter; AoD denotes the  
167 aortic end-diastolic diameter; SBP represents systolic blood pressure; DBP stands  
168 for diastolic blood pressure [16–18].  
169

## 170 **Statistical analysis**

171           The statistical analysis was carried out using SPSS software version 26 (IBM,  
172 Armonk, NY, USA). The Kolmogorov–Smirnov test was used to determine the  
173 normality of the data. Normally distributed continuous variables are reported as the  
174 means  $\pm$  standard deviations. Variables that were not normally distributed are  
175 represented as medians with interquartile ranges (IQRs = 25<sup>th</sup> – 75<sup>th</sup> percentiles).  
176 Categorical variables are represented as frequencies and percentages. Missing data  
177 were omitted from the analyses. Individuals with and without T2DM were  
178 compared based on their baseline characteristics. Continuous variables were  
179 analyzed using independent t tests, and the results are reported as the means  $\pm$   
180 standard deviations. The Mann–Whitney nonparametric test was applied for the  
181 analysis of variables that were not normally distributed. Categorical variables are  
182 presented as n (%) and were compared using the chi-squared test.

183           To compare arterial elastic parameters between the T2DM and non-T2DM  
184 groups, the Mann–Whitney test was used for nonnormally distributed continuous  
185 variables. Spearman’s correlation coefficient (r) and its related P value elucidated  
186 the correlation between continuous variables. All probability values were two-  
187 sided, and P values less than 0.05 were considered to indicate statistical  
188 significance. We followed the SAMPL guidelines in our statistical analysis to  
189 prevent avoidable errors or omissions in reporting statistical data [19].

## 190 **Results**

### 191 **Clinical and echocardiographic features**

192 This study included 109 subjects, including 62 T2DM patients and 47 healthy  
193 individuals who met the diagnostic criteria and composed the comparison group.

194 Our research revealed the following results.

195 There were no significant differences in age, sex, height, weight, BMI, or  
196 BSA between the groups ( $p > 0.05$ ). The differences in the left atrial diameter  
197 (LAd), left ventricular mass index (LVMI), and ejection fraction (EF) between the  
198 T2DM group and control group were not statistically significant ( $p > 0.05$ ). Detailed  
199 information regarding these indices is presented in **Table 1**.

200 **Table 1. Anthropometric and echocardiographic characteristics of**  
201 **participants by group**

<b>Characteristics</b>	<b>T2DM group (n = 62)</b>	<b>Comparison group (n = 47)</b>	<b>P value</b>
<b>Age (years)</b>	61.63 ± 10.46	60.40 ± 7.08	0.491

<b>Female (%)</b>	40 (64.5%)	26 (55.3%)	0.561
<b>BMI (kg/m<sup>2</sup>)</b>	22.26 ± 3.06	22.49 ± 2.35	0.657
<b>BSA (m<sup>2</sup>)</b>	1.53 ± 0.15	1.56 ± 0.13	0.300
<b>HbA1C (%)</b>	10.92 ± 3.02	5.03 ± 0.38	<0.001
<b>Duration of diabetes (years)</b>	6.00 [4.00 – 14.25]	NA	
<b>SBP (mmHg)</b>	127.63 ± 8.85	115.1 ± 12.19	<0.001
<b>DBP (mmHg)</b>	72.63 ± 7.02	72.32 ± 7.59	0.824
<b>LAd (cm)</b>	3.15 ± 0.42	3.19 ± 0.45	0.645
<b>LVMI (g/m<sup>2</sup>)</b>	107.31 ± 30.39	103.58 ± 23.08	0.488
<b>LVEF (%)</b>	67.92 ± 7.53	70.23 ± 5.27	0.075

<b>Total cholesterol (mmol/L)</b>	4.97 [3.63 – 6.49]	4.98 [4.15 – 5.00]	0.503
<b>Triglyceride (mmol/L)</b>	1.85 [1.06 – 2.95]	1.20 [1.05 – 1.25]	0.001
<b>HDL-C (mmol/L)</b>	0.96 [0.81 – 1.22]	1.25 [1.22 – 1.30]	< 0.001
<b>LDL-C (mmol/L)</b>	3.13 [1.96 - 3.67]	3.45 [2.73 – 3.55]	0.264

202 The values are presented as the means  $\pm$  standard deviations, medians [Q25 - Q75]  
203 or numbers (%). T2DM: type 2 diabetes mellitus; BSA: body surface area; BMI:  
204 body mass index; SBP: systolic blood pressure; DBP: diastolic blood pressure;  
205 LAd: left atrial diameter; LVMI: left ventricular mass index; LVEF: left ventricular  
206 ejection fraction; HbA1c: glycated hemoglobin; HDL-C: high-density lipoprotein  
207 cholesterol; LDL-C: low-density lipoprotein cholesterol.

## 208 **Aortic elasticity indices in the T2DM and control groups**

209 The T2DM group had significantly lower AC, AD, and AS than did the  
210 control group. The T2DM patients also had significantly greater ASI and AEM than  
211 did the controls ( $p < 0.05$ ). Additionally, the control group had a greater AS of 10.66

212 (6.01 - 18.23) than did the T2DM group at 8.21 (4.24 - 13.07). Detailed information  
 213 is provided in **Table 2** and **Fig 2**.

214 **Table 2. Computed aortic elasticity parameters of participants by group**

<b>Arterial elastic parameters</b>	<b>T2DM group (n = 62)</b>	<b>Comparison group (n = 47)</b>	<b>P values</b>
<b>AoS (mm)</b>	31.20 ± 4.95	27.08 ± 4.67	< 0.001
<b>AoD (mm)</b>	28.40 ± 4.32	24.04 ± 4.97	< 0.001
<b>AC (%mm/mmHg)</b>	4.07 (2.28 - 7.44)	6.40 (3.08 - 10.75)	0.008
<b>AD (%mmHg<sup>-1</sup>)</b>	3.08 (1.57 - 5.26)	5.33 (2.80 - 9.79)	< 0.001
<b>AS (%)</b>	8.21 (4.24 - 13.07)	10.66 (6.01 - 18.23)	0.039
<b>ASI</b>	6.10 (3.64 - 12.47)	3.79 (2.40 - 8.50)	0.003
<b>AEM (Kpa)</b>	86.68 (50.80 - 169.73)	50.00 (27.24 - 95.33)	< 0.001

215 The values are presented as the means  $\pm$  standard deviations or medians [Q25 -  
216 Q75]. T2DM: Type 2 diabetes mellitus; AoS: aortic end-systolic diameter; AoD:  
217 aortic end-diastolic diameter; AC: aortic compliance; AD: aortic distensibility; AS:  
218 aortic strain; ASI: aortic stiffness index; AEM: Aortic Peterson's elastic modulus

219 **Fig 2. Comparison of aortic elasticity indices.** (A) Comparison of aortic  
220 distensibility between the diabetes group and control group. (B) Comparison of  
221 aortic compliance between the diabetes group and the control group. (C)  
222 Comparison of strains between the diabetes group and control group. (D)  
223 Comparison of the aortic stiffness indices between the diabetes group and the  
224 control group. (E) Comparison of Peterson's elastic modulus between the diabetes  
225 group and control group. T2DM: diabetes mellitus.

## 226 **Correlations between indices of aortic elasticity and relevant** 227 **variables**

228 An analysis was performed to explore the relationships between  
229 anthropometric and echocardiographic features and computed aortic indices. **Fig 3**  
230 visually illustrates the results in a heatmap. The results revealed that in the T2DM  
231 group, the duration of diabetes was associated with arterial elasticity indicators. A  
232 longer duration of diabetes was positively associated with greater ASI and AEM  
233 (Spearman's correlation coefficients of 0.43 and 0.43, respectively). In contrast,

234 AC, AD, and AS tended to decrease as diabetes duration increased (Spearman's  
235 correlation coefficients of -0.39, -0.43, and -0.37, respectively).

236 **Fig 3. Correlations between anthropometric and echocardiographic features**  
237 **and computed aortic indices.** BSA: body surface area; BMI: body mass index;  
238 HbA1c: glycated hemoglobin; LAd: left atrial diameter; RWT: relative wall  
239 thickness; LVEF: left ventricular ejection fraction; LVMI: left ventricular mass  
240 index; AC: aortic compliance; AD: aortic distensibility; AS: aortic strain; ASI:  
241 aortic stiffness index; AEM: Aortic Peterson's elastic modulus.

## 242 **Discussion**

243 There is growing interest in studying the association between aortic  
244 flexibility and cardiovascular disease. In contrast, the stage of myocardial  
245 performance in diabetes and prediabetes patients remains overlooked [20]. Among  
246 the wide range of choices for echocardiography, M-mode echocardiography is  
247 still effective at capturing the motions and functions of the heart and surrounding  
248 structures. Despite the better temporal resolution of M-mode echocardiography, it  
249 is still being disregarded in international guidelines [21].  
250 Substantial efforts have been made to investigate the relationships between insulin  
251 resistance (and T2DM) and aortic elastic characteristics. This study was performed  
252 in Vietnam with a group of T2DM patients from low- and middle-income countries

253 to assess the feasibility and cost-effectiveness of M-mode echocardiography in  
254 daily practice.

## 255 **Comparison of aortic elasticity indices between patients with T2DM** 256 **and healthy controls**

257 It is generally known that vascular elasticity often begins to decline in middle  
258 age and worsens over time [17]. Throughout the aging process, the elasticity of the  
259 endothelium layer gradually decreases, which has a significant impact on the  
260 structure of the arteries due to reduced elastic fiber activity, particularly in major  
261 arteries such as the aorta [22]. The deterioration of arterial elasticity is caused by a  
262 variety of mechanisms, including the impairment of vasodilation mechanisms via  
263 nitric oxide (NO) and the absence of ETB receptors, which increase the plasma  
264 endothelin-1 concentration because these receptors contribute to endothelin-1  
265 destruction [23,24].

266 In this study, we measured the elasticity of the aorta using five indices: AC,  
267 AD, ASI, AS, and AEM. The T2DM group had considerably lower AC, AD, and  
268 AS indices than did the comparison group but had higher ASI and AEM. This study  
269 successfully reproduced the findings previously reported by Malgorzata Dec-  
270 Gilowska et al [17]. Furthermore, our findings are consistent with those of Liu et  
271 al., who conducted a study on individuals with diabetes patients and concluded that

272 these patients are more likely to experience arterial stiffness than healthy  
273 individuals are [25]. According to Zapolski et al., patients with diabetes have  
274 greater ASI readings than healthy people, independent of whether their disease is  
275 accompanied by hypertension [26]. Another study by Aslan also confirmed that  
276 patients with prediabetes have a greater ASI than healthy people according to  
277 echocardiography [20].

278 An increased ASI is an independent risk factor for cardiovascular disease and  
279 mortality [27]. Our findings are consistent with those of Dec-Gilowska et al., who  
280 reported that diabetes patients tend to have a significantly decreased AC. In parallel,  
281 our investigation revealed a rise in ASI akin to the observations made by Dec-  
282 Gilowska et al [28]. We found that both the ASI and the AEM nearly doubled in  
283 patients with T2DM compared to those in the control group. It is well known that  
284 the ASI is strongly correlated with the mean arterial pressure, and increased ASI  
285 may account for the development of hypertension. However, this study is unable to  
286 shed light on the question of whether either ASI or hypertension precedes the other  
287 in this progression among T2DM patients.

288 Despite major breakthroughs in T2DM treatment, the risk of cardiovascular  
289 morbidity and mortality continues to increase [29]. Recent research has revealed a  
290 complex link between cardiovascular mortality risk and T2DM, which is thought to  
291 be caused in part by increased arterial stiffness [30]. The literature has shown that

292 people with T2DM have a much greater incidence of increased arterial stiffness  
293 than individuals without diabetes [31]. This can be due to insulin resistance,  
294 hyperglycemia, and persistent hyperinsulinemia. These drugs activate the renin-  
295 angiotensin-aldosterone system and enhance the expression of type 1 angiotensin  
296 receptors in vascular tissue, inducing arterial wall hypertrophy and fibrosis [32–34].  
297 Insulin resistance inhibits PI3-kinase-dependent enzyme signaling, which is  
298 involved in insulin metabolism, cell growth, proliferation, differentiation, motility,  
299 survival, and intracellular transport, resulting in hyperinsulinemia. Furthermore,  
300 decreased glucose tolerance increases glycation and changes the mechanical  
301 characteristics of the artery wall's interstitial tissue via a variety of complex  
302 mechanisms [35,36]. Increased arterial stiffness causes irreversible pathogenic  
303 loads, particularly in patients with diabetes. A constant increase in arterial stiffness  
304 results in increased transfer of pulse pressure into the microcirculation, which is  
305 directly associated with a greater risk of disorders such as hypertension, chronic  
306 renal disease, coronary artery disease, heart failure, and stroke [37–40]. Our study  
307 revealed substantial differences in AC, AD, AS, ASI, and AEM between  
308 individuals with T2DM and individuals without diabetes, which suggests further  
309 screenings and interventions for these patients are needed for better outcomes.

### 310 **Correlations between aortic elasticity indices and diabetic conditions**

311 Our study did not identify any correlation between the HbA1c level and  
312 aortic elasticity indices (**Fig 3**).

313 The association between HbA1c, an indicator of long-term blood glucose  
314 regulation, and aortic stiffness seems to be constrained in certain studies [41–43].  
315 This constraint can be explained by the fact that HbA1c measurements reflect only  
316 a snapshot of glucose levels attached to hemoglobin over the preceding three  
317 months rather than the comprehensive control of blood sugar over an extended  
318 period. Consequently, it may not accurately correlate with the development of  
319 arterial stiffness, which occurs gradually over time [44].

320 Emerging evidence suggests that blood glucose variability could exert a more  
321 significant impact on aortic stiffness. Individuals with diabetes, particularly those  
322 with poorly managed blood glucose levels and significant variability, face an  
323 elevated risk of aortic stiffness and related cardiovascular complications [45,46].  
324 Therefore, emphasizing the control of blood glucose fluctuations through  
325 comprehensive diabetes management strategies is pivotal for mitigating the risk of  
326 aortic stiffness and enhancing cardiovascular health outcomes among individuals  
327 with diabetes.

328 Nevertheless, we did find significant correlations between the duration of  
329 diabetes and arterial elasticity indices as shown in the heatmap (**Fig 3**). A longer

330 duration of diabetes is linked to greater stiffness, whereas elasticity decreases  
331 accordingly.

332 In diabetes, the duration of T2DM is intricately linked to the progression of  
333 disease complications [47]. In a study led by Agnoletti et al., a significant  
334 correlation emerged between the duration of diabetes and the worsening of arterial  
335 stiffness. Their meticulous findings revealed that the duration of diabetes correlates  
336 with microvascular complications, regardless of renal function, and with  
337 macrovascular complications due to the presence of increased aortic stiffness [48].  
338 By recognizing the pivotal role of diabetes duration in the progression of arterial  
339 stiffness, early diagnosis, and proactive treatment become paramount in optimizing  
340 patient care and outcomes.

### 341 **Limitations of the study**

342 The single-center design and limited sample size constrain our study. The  
343 small sample size is particularly restrictive, potentially introducing bias and  
344 diminishing statistical power. Therefore, there is a need for multicenter research  
345 with a larger sample size to obtain more vital clinical evidence. Additionally, our  
346 study is only a cross-sectional investigation that can demonstrate associations but  
347 not causal relationships. Hence, longitudinal studies are required to address this  
348 issue. Furthermore, arterial stiffness parameters were measured in the central aorta,  
349 whereas blood pressure was measured in peripheral arteries. However, due to

350 ethical considerations, our study solely relied on peripheral blood pressure  
351 measurements instead of invasive arterial pressure measurements. Additionally,  
352 given the circumstances at our research site, where alterations in medication  
353 regimens are common, our study did not assess the impact of different drug  
354 treatments on arterial stiffness. Thus, more significant and longer-term studies are  
355 needed to evaluate this aspect.

## 356 **Conclusions**

357 A notable difference in aortic elasticity indices was detected between the  
358 T2DM patient group and the control group via M-mode echocardiography.

## 359 **References**

- 360 1. Magliano D, Boyko EJ. IDF diabetes atlas. 10th edition. Brussels: International  
361 Diabetes Federation; 2021.
- 362 2. Dang HNN, Thang VL, Binh AH. Evaluation of the relationship between left  
363 atrial stiffness, left ventricular stiffness, and left atrioventricular coupling index  
364 in type 2 diabetes patients: a speckle tracking echocardiography study. *Front*  
365 *Cardiovasc Med.* 2024;11: 1372181. doi:10.3389/fcvm.2024.1372181
- 366 3. Dang HNN, Luong TV, Tran TT. Evaluating left atrial function changes by  
367 speckle tracking echocardiography in type 2 diabetes patients in Central

- 368 Vietnam: a cross-sectional comparative study. *Egypt Heart J.* 2024;76: 38.  
369 doi:10.1186/s43044-024-00470-w
- 370 4. McCallinhart PE, Sunyecz IL, Trask AJ. Coronary Microvascular Remodeling  
371 in Type 2 Diabetes: Synonymous With Early Aging? *Front Physiol.* 2018;9:  
372 1463. doi:10.3389/fphys.2018.01463
- 373 5. Li Y, Liu Y, Liu S, Gao M, Wang W, Chen K, et al. Diabetic vascular diseases:  
374 molecular mechanisms and therapeutic strategies. *Sig Transduct Target Ther.*  
375 2023;8: 152. doi:10.1038/s41392-023-01400-z
- 376 6. Ahmadizar F, Wang K, Roos M, Bos M, Mattace- Raso F, Kavousi M.  
377 Association between arterial stiffness/remodeling and new-onset type 2  
378 diabetes mellitus in general population. *Diabetes Research and Clinical  
379 Practice.* 2023;196: 110237. doi:10.1016/j.diabres.2023.110237
- 380 7. Bornfeldt KE, Tabas I. Insulin Resistance, Hyperglycemia, and  
381 Atherosclerosis. *Cell Metabolism.* 2011;14: 575–585.  
382 doi:10.1016/j.cmet.2011.07.015
- 383 8. Prasad C, Davis KE, Imrhan V, Juma S, Vijayagopal P. Advanced Glycation  
384 End Products and Risks for Chronic Diseases: Intervening Through Lifestyle  
385 Modification. *American Journal of Lifestyle Medicine.* 2019;13: 384–404.  
386 doi:10.1177/1559827617708991

- 387 9. Cavalcante JL, Lima JAC, Redheuil A, Al-Mallah MH. Aortic Stiffness.  
388 Journal of the American College of Cardiology. 2011;57: 1511–1522.  
389 doi:10.1016/j.jacc.2010.12.017
- 390 10. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke  
391 JP, et al. The Strengthening the Reporting of Observational Studies in  
392 Epidemiology (STROBE) statement: guidelines for reporting observational  
393 studies. Lancet. 2007;370: 1453–1457. doi:10.1016/S0140-6736(07)61602-X
- 394 11. American Diabetes Association. Standards of Medical Care in Diabetes—2022  
395 Abridged for Primary Care Providers. Clinical Diabetes. 2022;40: 10–38.  
396 doi:10.2337/cd22-as01
- 397 12. Du Bois D, Du Bois EF. A formula to estimate the approximate surface area if  
398 height and weight be known. 1916. Nutrition. 1989;5: 303–311; discussion  
399 312-313.
- 400 13. Casey DE, Thomas RJ, Bhalla V, Commodore-Mensah Y, Heidenreich PA,  
401 Kolte D, et al. 2019 AHA/ACC Clinical Performance and Quality Measures  
402 for Adults With High Blood Pressure. Journal of the American College of  
403 Cardiology. 2019;74: 2661–2706. doi:10.1016/j.jacc.2019.10.001
- 404 14. Muntner P, Shimbo D, Carey RM, Charleston JB, Gaillard T, Misra S, et al.  
405 Measurement of Blood Pressure in Humans: A Scientific Statement From the

- 406 American Heart Association. Hypertension. 2019;73.  
407 doi:10.1161/HYP.0000000000000087
- 408 15. Mitchell C, Rahko PS, Blauwet LA, Canaday B, Finstuen JA, Foster MC, et al.  
409 Guidelines for Performing a Comprehensive Transthoracic Echocardiographic  
410 Examination in Adults: Recommendations from the American Society of  
411 Echocardiography. Journal of the American Society of Echocardiography.  
412 2019;32: 1–64. doi:10.1016/j.echo.2018.06.004
- 413 16. Nemes A, Geleijnse ML, Forster T, Soliman OII, Ten Cate FJ, Csanády M.  
414 Echocardiographic Evaluation and Clinical Implications of Aortic Stiffness  
415 and Coronary Flow Reserve and their Relation. Clin Cardiol. 2008;31: 304–  
416 309. doi:10.1002/clc.20252
- 417 17. Song X, Fan L, Yan Z, Rui Y. Echocardiographic evaluation of the elasticity  
418 of the ascending aorta in patients with essential hypertension. J of Clinical  
419 Ultrasound. 2021;49: 351–357. doi:10.1002/jcu.22925
- 420 18. Stefanadis C, Stratos C, Boudoulas H, Kourouklis C, Toutouzas P.  
421 Distensibility of the ascending aorta: comparison of invasive and non-invasive  
422 techniques in healthy men and in men with coronary artery disease. European  
423 Heart Journal. 1990;11: 990–996.  
424 doi:10.1093/oxfordjournals.eurheartj.a059639

- 425 19. Lang TA, Altman DG. Basic statistical reporting for articles published in  
426 Biomedical Journals: The “Statistical Analyses and Methods in the Published  
427 Literature” or the SAMPL Guidelines. *International Journal of Nursing  
428 Studies*. 2015;52: 5–9. doi:10.1016/j.ijnurstu.2014.09.006
- 429 20. Aslan AN, Ayhan H, Çiçek ÖF, Akçay M, Durmaz T, Keles T, et al.  
430 Relationship between Aortic Stiffness and the Left Ventricular Function in  
431 Patients with Prediabetes. *Intern Med*. 2014;53: 1477–1484.  
432 doi:10.2169/internalmedicine.53.1720
- 433 21. Feigenbaum H. Role of M-mode Technique in Today’s Echocardiography.  
434 *Journal of the American Society of Echocardiography*. 2010;23: 240–257.  
435 doi:10.1016/j.echo.2010.01.015
- 436 22. Petrovic DJ. The Influence of Aging-Related Decrease in Endothelial Function  
437 on the Stiffness of Peripheral Conducting Arteries. *Journal for Vascular  
438 Ultrasound*. 2023;47: 184–192. doi:10.1177/15443167231193164
- 439 23. Wilkinson IB, Webb DJ, Cockcroft JR. Nitric Oxide and the Regulation of  
440 Arterial Elasticity: Right Idea, Wrong Vascular Bed? *Hypertension*. 2002;39.  
441 doi:10.1161/01.HYP.0000014617.84543.CA
- 442 24. Kostov K. The Causal Relationship between Endothelin-1 and Hypertension:  
443 Focusing on Endothelial Dysfunction, Arterial Stiffness, Vascular

- 444 Remodeling, and Blood Pressure Regulation. *Life*. 2021;11: 986.  
445 doi:10.3390/life11090986
- 446 25. Liu J-J, Sum CF, Tavintharan S, Yeoh LY, Ng XW, Moh AMC, et al. Obesity  
447 is a determinant of arterial stiffness independent of traditional risk factors in  
448 Asians with young-onset type 2 diabetes. *Atherosclerosis*. 2014;236: 286–291.  
449 doi:10.1016/j.atherosclerosis.2014.07.017
- 450 26. Song X-T, Rui Y-F, Fan L, Yan Z-N. Echocardiographic Association of  
451 Epicardial Adipose Tissue with Ascending Aorta Elasticity in Patients with  
452 Type 2 Diabetes Mellitus. *Angiology*. 2023;74: 325–332.  
453 doi:10.1177/00033197221098298
- 454 27. Laurent S, Boutouyrie P, Asmar R, Gautier I, Laloux B, Guize L, et al. Aortic  
455 Stiffness Is an Independent Predictor of All-Cause and Cardiovascular  
456 Mortality in Hypertensive Patients. *Hypertension*. 2001;37: 1236–1241.  
457 doi:10.1161/01.HYP.37.5.1236
- 458 28. Dec-Gilowska M, Trojnar M, Makaruk B, Mosiewicz-Madejska B, Dzida G,  
459 Mosiewicz J. Local elasticity of the aorta in patients with type 2 diabetes  
460 mellitus. *J Int Med Res*. 2020;48: 030006052095264.  
461 doi:10.1177/0300060520952643

- 462 29. Tancredi M, Rosengren A, Svensson A-M, Kosiborod M, Pivodic A,  
463 Gudbjörnsdottir S, et al. Excess Mortality among Persons with Type 2  
464 Diabetes. *N Engl J Med*. 2015;373: 1720–1732. doi:10.1056/NEJMoa1504347
- 465 30. Stehouwer CDA, Henry RMA, Ferreira I. Arterial stiffness in diabetes and the  
466 metabolic syndrome: a pathway to cardiovascular disease. *Diabetologia*.  
467 2008;51: 527–539. doi:10.1007/s00125-007-0918-3
- 468 31. De Oliveira Alvim R, Santos PCJL, Musso MM, De Sá Cunha R, Krieger JE,  
469 Mill JG, et al. Impact of diabetes mellitus on arterial stiffness in a  
470 representative sample of an urban Brazilian population. *Diabetol Metab Syndr*.  
471 2013;5: 45. doi:10.1186/1758-5996-5-45
- 472 32. Salomaa V, Riley W, Kark JD, Nardo C, Folsom AR. Non–Insulin-Dependent  
473 Diabetes Mellitus and Fasting Glucose and Insulin Concentrations Are  
474 Associated With Arterial Stiffness Indexes: The ARIC Study. *Circulation*.  
475 1995;91: 1432–1443. doi:10.1161/01.CIR.91.5.1432
- 476 33. Scuteri A, Najjar SS, Muller DC, Andres R, Hougaku H, Metter EJ, et al.  
477 Metabolic syndrome amplifies the age-associated increases in vascular  
478 thickness and stiffness. *Journal of the American College of Cardiology*.  
479 2004;43: 1388–1395. doi:10.1016/j.jacc.2003.10.061

- 480 34. Sutton-Tyrrell K, Newman A, Simonsick EM, Havlik R, Pahor M, Lakatta E,  
481 et al. Aortic Stiffness Is Associated With Visceral Adiposity in Older Adults  
482 Enrolled in the Study of Health, Aging, and Body Composition. *Hypertension*.  
483 2001;38: 429–433. doi:10.1161/01.HYP.38.3.429
- 484 35. Jesmin S, Sakuma I, Hattori Y, Kitabatake A. Role of Angiotensin II in Altered  
485 Expression of Molecules Responsible for Coronary Matrix Remodeling in  
486 Insulin-Resistant Diabetic Rats. *ATVB*. 2003;23: 2021–2026.  
487 doi:10.1161/01.ATV.0000094235.78783.D1
- 488 36. Rizzoni D, Porteri E, Guelfi D, Muiesan ML, Valentini U, Cimino A, et al.  
489 Structural Alterations in Subcutaneous Small Arteries of Normotensive and  
490 Hypertensive Patients With Non–Insulin-Dependent Diabetes Mellitus.  
491 *Circulation*. 2001;103: 1238–1244. doi:10.1161/01.CIR.103.9.1238
- 492 37. Kaess BM, Rong J, Larson MG, Hamburg NM, Vita JA, Levy D, et al. Aortic  
493 Stiffness, Blood Pressure Progression, and Incident Hypertension. *JAMA*.  
494 2012;308: 875. doi:10.1001/2012.jama.10503
- 495 38. Sedaghat S, Mattace-Raso FUS, Hoorn EJ, Uitterlinden AG, Hofman A, Ikram  
496 MA, et al. Arterial Stiffness and Decline in Kidney Function. *Clinical Journal*  
497 *of the American Society of Nephrology*. 2015;10: 2190–2197.  
498 doi:10.2215/CJN.03000315

- 499 39. Van Bortel LM, Laurent S, Boutouyrie P, Chowienczyk P, Cruickshank JK,  
500 De Backer T, et al. Expert consensus document on the measurement of aortic  
501 stiffness in daily practice using carotid-femoral pulse wave velocity. *Journal of*  
502 *Hypertension*. 2012;30: 445–448. doi:10.1097/HJH.0b013e32834fa8b0
- 503 40. Zhong Q, Hu M-J, Cui Y-J, Liang L, Zhou M-M, Yang Y-W, et al. Carotid–  
504 Femoral Pulse Wave Velocity in the Prediction of Cardiovascular Events and  
505 Mortality: An Updated Systematic Review and Meta-Analysis. *Angiology*.  
506 2018;69: 617–629. doi:10.1177/0003319717742544
- 507 41. Chen Y, Huang Y, Li X, Xu M, Bi Y, Zhang Y, et al. Association of arterial  
508 stiffness with HbA1c in 1,000 type 2 diabetic patients with or without  
509 hypertension. *Endocr*. 2009;36: 262–267. doi:10.1007/s12020-009-9221-z
- 510 42. Martagón AJ, Fermín-Martínez CA, Antonio-Villa NE, Mehta R, Almeda-  
511 Valdés P, Vargas-Vázquez A, et al. Arterial Stiffness and HbA1c: Association  
512 Mediated by Insulin Resistance in Hispanic Adults. *IJERPH*. 2022;19: 11017.  
513 doi:10.3390/ijerph191711017
- 514 43. Gilstrap LG, Chernew ME, Nguyen CA, Alam S, Bai B, McWilliams JM, et  
515 al. Association Between Clinical Practice Group Adherence to Quality  
516 Measures and Adverse Outcomes Among Adult Patients With Diabetes. *JAMA*  
517 *Netw Open*. 2019;2: e199139. doi:10.1001/jamanetworkopen.2019.9139

- 518 44. Sun R, Duan Y, Zhang Y, Feng L, Ding B, Yan R, et al. Time in Range  
519 Estimation in Patients with Type 2 Diabetes is Improved by Incorporating  
520 Fasting and Postprandial Glucose Levels. *Diabetes Ther.* 2023;14: 1373–1386.  
521 doi:10.1007/s13300-023-01432-2
- 522 45. Fang Q, Shi J, Zhang J, Peng Y, Liu C, Wei X, et al. Visit-to-visit HbA1c  
523 variability is associated with aortic stiffness progression in participants with  
524 type 2 diabetes. *Cardiovasc Diabetol.* 2023;22: 167. doi:10.1186/s12933-023-  
525 01884-7
- 526 46. Ceriello A, Lucisano G, Prattichizzo F, La Grotta R, Franzén S, Svensson A-  
527 M, et al. HbA1c variability predicts cardiovascular complications in type 2  
528 diabetes regardless of being at glycemic target. *Cardiovasc Diabetol.* 2022;21:  
529 13. doi:10.1186/s12933-022-01445-4
- 530 47. De Jong M, Woodward M, Peters SAE. Duration of diabetes and the risk of  
531 major cardiovascular events in women and men: A prospective cohort study of  
532 UK Biobank participants. *Diabetes Research and Clinical Practice.* 2022;188:  
533 109899. doi:10.1016/j.diabres.2022.109899
- 534 48. Agnoletti D, Mansour AS, Zhang Y, Protogerou AD, Ouerdane S, Blacher J,  
535 et al. Clinical interaction between diabetes duration and aortic stiffness in type

536 2 diabetes mellitus. J Hum Hypertens. 2017;31: 189–194.

537 doi:10.1038/jhh.2016.58




Fig1




Fig2




Fig3