

Prevalence and Risk Factors of Hypertension Among Young Adults (26-35 Years Old) in Indonesia: Analysis of Secondary Data from Riskesdas 2018

Laluna Rachma Putri¹, Arulita Ika Fibriana¹, Mahalul Azam^{1,*}

¹Prodi Kesehatan Masyarakat, Fakultas Kedokteran, Univeritas Negeri Semarang

*) correspondent author: mahalul.azam@mail.unnes.ac.id

Abstract

Background: Worldwide, there are 54% adults diagnosed with hypertension. The prevalence of hypertension increased, based on the basic health survey (Riset Kesehatan Dasar; RISKESDAS) in 2013, the prevalence of hypertension was 25.8%, while the on the RISKESDAS 2018 data increased to 34.1%. This research aimed to find out the risk factors of hypertension among young adults in Indonesia using RISKESDAS 2018 data.

Method: The research design approach observational quantitative by cross-sectional design. This research used secondary data from RISKESDAS 2018 with a total of 140,073 subjects aged 26-35 years. All subjects who fulfil the inclusion criteria will be analyzed (n=78,000). Data analysed by chi-square test and binary logistic regression.

Results: The result of research exposed gender ($p<0.001$), marital status ($p<0.001$), education levels ($p<0.001$), working status ($p<0.001$), BMI ($p<0.001$), diabetes mellitus ($p<0.001$), emotional mental disorder ($p<0.001$), and instant food consumption ($p<0.001$) has a significant consequence of hypertension.

Conclusions: The results of research can be used as a consideration for creating programs and policies to control risk factors or hypertension among young adults in Indonesia. Promotive and preventive policies programs related to hypertension must reach the community to reduce hypertension morbidity and mortality rates.

Keywords: *Hypertension, risk factors, young adults, RISKESDAS 2018, cross-sectional*

Introduction

Non-communicable diseases (NCDs) are not caused by viruses, germs, or bacteria but by daily lifestyle. One of the NCDs that threatens human health is hypertension (Minasari et al., 2022). Hypertension is a circulatory system disorder that increases blood pressure (Azinar et al., 2019). The number of hypertension patients continues to increase every year. In 2025, there will be 1.5 billion people with hypertension. Every year, it is estimated that there will be 9.4 million deaths due to hypertension and its complications (World Health Organization, 2023). Worldwide, there are 54% of adults diagnosed with hypertension (Global Health Observatory, 2023). An estimated 46% of adults with hypertension are unaware of their condition (World Health Organization, 2023). The prevalence of hypertension increased, based on the basic health survey (Riset Kesehatan Dasar; RISKESDAS). In 2013, the prevalence of hypertension was 25.8%, while the RISKESDAS 2018 data increased to 34.1%. There are a plenty of unreported cases; merely 1/3 of hypertension cases in Indonesia are diagnosed (Rokom, 2021).

Hypertension affected economic losses, productivity decline, and the emergence of complications. Noteworthy, hypertension can cause complications in diabetes mellitus, making it a considerable risk factor for 39 cardiovascular diseases (Azam et al., 2023). Hypertension patients have a 2.6 times greater risk of developing coronary heart disease compared to non-hypertensive individuals (Amisi et al., 2018). Hypertension patients lose productivity without controlling for health risks and chronic conditions (MacLeod et al., 2022).

Hypertension medication ranks first in terms of the highest cost in Indonesia, particularly IDR 12.1 trillion (Social Security Agency of Health, 2017).

The majority of young adults with hypertension are ignorant of abnormally high blood pressure and acknowledge the impact when they get older. There is a long-term impact of early-life risk factors of hypertension in young adults (Gooding et al., 2014). Hypertension patients aged below 40 years have low awareness, low diagnosis, and poor blood pressure control (Johnson et al., 2014). The prevalence of undiagnosed hypertension in men aged 25 to 34 years is 55% and 44% in women aged 25 to 34 years. The prevalence of undiagnosed hypertension in young adults is higher than in people aged 75 years or over (17% in men and 21% in women) (Lacobucci, 2023).

We must undertake preventive efforts against hypertension among young adults. It is essential to identify and maintain modifiable risk factors for hypertension. This study aimed to provide information on the risk factors for hypertension among young adults based on the results of basic health research in 2018 in Indonesia. The results of the analysis are expected to impact suitable hypertension prevention in Indonesia. Efforts against hypertension prevention positively impact the health of each individual and community welfare, reducing costs for hypertension medicine and reducing the disadvantage of disease caused by hypertension.

Methods

Design and Study Population

This research is an observational quantitative approach cross-sectional design. This study utilized secondary data from the basic health survey (Riset Kesehatan Dasar; RISKESDAS) 2018. RISKESDAS is a population health study conducted by the Agency for Health Policies Development, Ministry of Health of the Republic of Indonesia. RISKESDAS aims to monitor indicators of public health and indicators of public health services. RISKESDAS provides information on key health indicators such as health status, nutritional status, environmental health, health behaviors, and various aspects of health care. RISKESDAS data provides health data at district/city and national levels. This study population included subjects aged 26–35 who were interviewed by the RISKESDAS 2018 enumerator, with a total of 140,073 subjects aged 26–35. Of the 140,073 subjects, 78,300 met the criteria to be analyzed (Figure 1).




Figure 1. Sampling flow

Data Collection

This study met the ethical criteria by Universitas Negeri Semarang of No.169/KEPK/FK/KLE/2024. The Agency for Health Policies Development, Ministry of Health of the Republic of Indonesia, has approved the use of RISKESDAS data for study purposes. This study has received permission to utilize, exploit, and process data in scientific studies. The subjects used in this research were subjects who already measured their weight and height; there were also no missing data.

This study used certain parameters from RISKESDAS 2018, which were analyzed in this research. These include sociodemographic data, hypertension status, diabetes mellitus status, emotional mental disorder status, smoking status, physical activity status, alcohol consumption, body mass index, salt food consumption, fat food consumption, and instant food consumption.

Statistical Analysis

The characteristics of the subjects were categorical data, so data was presented in a proportion table. This study performed a chi-square test to find out the risk factors associated with hypertension among young adults in Indonesia. The p-values <0.05 reflected as statistically significant. This study involved multivariate analysis using binary logistic regression for all parameters with p-value <0.25 in the chi-square test. The statistical analysis was accomplished in this study using IBM SPSS Statistics software (for Windows, Version 22.0, Armonk, NY: IBM Corp.).

Result

Subjects' Characteristics

Of the 140,073 subjects aged 26-35, 73,800 fulfilled the criteria and were included in further analysis. Among these, 3,225 (4.1%) young adults in Indonesia suffered from hypertension, whereas 75,075 (95.9%) did not. Table 1 explains the characteristics of these subjects.

Table 1. The characteristics of study subjects

Variables	Frequencies (N)	Percentages (%)
Gender		
Male	27227	34.8
Female	51073	65.2
Marital Status		
Unmarried	8398	10.7
Married	67895	86.7
Widowed	2007	2.6
Residence Area		
Urban	34367	43.9
Rural	43933	55.1
Education Level		
Low	37482	47.9
High	40818	52.1
Working Status		
No	26342	33.6
Yes	51958	66.4

Body Mass Index		
BMI \geq 25	31430	40.1
BMI<25	46870	59.9
Diabetes Mellitus		
Yes	274	0.3
No	78026	99.7
Emotional Mental Disorder		
Yes	7705	9.8
No	70595	90.2
Smoking		
Yes	22,45	28.4
No	56055	71.6
Physical Activity		
No	7067	9.0
Yes	71233	91.0
Alcohol Consumption		
Yes	3540	4.5
No	74760	95.5
Salt Food Consumption		
Frequent (\geq 3-6 times/week)	32153	41.1
Rarely (<3-6 times/week)	48147	58.9
Fat Food Consumption		
Frequent (\geq 3-6 times/week)	47346	60.5
Rarely (<3-6 times/week)	30954	39.5
Instant Food Consumption		
Frequent (\geq 3-6 times/week)	21864	27.9
Rarely (<3-6 times/week)	56436	72.1
Hypertension		
Yes	3225	4.1
No	75075	95.9

Risk Factors Associated to Hypertension among Young Adults (26-35 years) in Indonesia

Based on the result of the chi-square test (Tabel 2), some parameters show p-value <0.005; it refers to evidence of influence hypertension among young adults, i.e. gender (PR=0.856; 95% CI: 0.794-0.924), education levels (PR=1.355; 95% CI: 1.262-1.454), marital status (PR=0.612; 95% CI: 0.481-0.777), working status (PR=1.358; 95% CI: 1.264-1.460), body mass index (PR=2.424; 95% CI: 2.255-2.606), diabetes mellitus (PR=4.257; 95% CI: 3.059-5.924), emotional mental disorder (PR=2.105; 95% CI: 1.919-2.311), smoking (PR=0.876; 95% CI: 0.808-0.949), and instant food consumption (PR=1.159; 95% CI: 1.074-1.251). Variables with a p-value <0.25 will undergo continued analysis by binary logistic regression. Table 3 shows the final regression model results.

Table 2. The characteristics of study subjects based on hypertension status

Variables	Hypertension				p-value	PR (95% CI)
	Yes		No			
	N	%	N	%		
Gender						
Male	1015	3.7	26212	96.3	<0.001*	0.856 (0.794-0.924) 1 (Reference)
Female	2210	4.3	48863	95.7		

Marital Status						
Unmarried	253	3.0	8145	97.0	<0.001*	0.612 (0.481-0.777)
Married	2875	4.2	65020	95.8	0.190	0.871 (0.708-1.071)
Widowed	97	4.8	1910	95.2		1 (Reference)
Residence Area						
Urban	1425	4.1	32942	95.9	0.731	1.013 (0.943-1.087)
Rural	1800	4.1	42133	95.9		1 (Reference)
Education Level						
Low	1778	4.7	35704	95.3	<0.001*	1.355 (1.262-1.454)
High	1447	3.5	39371	96.5		1 (Reference)
Working Status						
No	1305	5.0	25037	95.0	<0.001*	1.358 (1.264-1.460)
Yes	1920	3.7	50038	96.3		1 (Reference)
Body Mass Index						
BMI \geq 25	1968	6.3	29462	93.7	<0.001*	2.424 (2.255-2.606)
BMI<25	1257	2.7	45613	97.3		1 (Reference)
Diabetes Mellitus						
Yes	42	15.3	232	84.7	<0.001*	4.257 (3.059-5.924)
No	3183	4.1	74843	95.9		1 (Reference)
Emotional Mental Disorder						
Yes	583	7.6	7122	92.4	<0.001*	2.105 (1.919-2.311)
No	2642	3.7	67953	96.3		1 (Reference)
Smoking						
Yes	835	3.8	21410	96.2	0.001*	0.876 (0.808-0.949)
No	2390	4.3	53665	95.7		1 (Reference)
Physical Activity						
No	298	4.2	6769	95.8	0.663	1.027 (0.910-1.160)
Yes	2927	4.1	68306	95.9		1 (Reference)
Alcohol Consumption						
Yes	159	4.5	3381	95.5	0.260	1.100 (0.934-1.294)
No	3066	4.1	71694	95.9		1 (Reference)
Salt Consumption						
\geq 3-6 times/week	1309	4.1	30844	95.9	0.583	0.980 (0.912-1.053)
<3-6 times/week	1916	4.2	44231	95.8		1 (Reference)
Fat Consumption						
\geq 3-6 times/week	1954	4.1	45392	95.9	0.897	1.005 (0.935-1.081)
<3-6 times/week	1271	4.1	29683	95.9		1 (Reference)
Instant Food Consumption						
\geq 3-6 times/week	995	4.6	20869	95.4	<0.001*	1.159 (1.074-1.251)
<3-6 times/week	2230	4.0	54206	96.0		1 (Reference)

* = *p*-value <0.05

Through binary logistic regression analysis, this study found that male (PR=1.225; 95% CI: 1.119-1.341), unmarried subject (PR=0.711; 95% CI: 0.556-0.908), low education level (PR=1.286; 95% CI: 1.195-1.384), unemployed subject (PR=1.267; 95% CI: 1.166-1.377), BMI above average (PR=2.472; 95% CI: 2.295-2.662), diabetes mellitus subject (PR=3.688; 95% CI: 2.630-5.173), emotional mental disorder subject (PR=2.041; 95% CI: 1.856-2.244), frequent consume instant food (PR=1.108; 95% CI: 1.025-1.197), all together were linked to hypertension among young adults.

Table 3. Binary logistic regression of hypertension risk factors among young adults subject

Variables	B	S.E	Wald	p-value	PR (95% CI)
Gender	0.203	0.046	19.368	<0.001*	1.225 (1.119-1.341)
Marital Status	-0.341	0.125	7.434	0.006*	0.711 (0.556-0.908)

Education Levels	0.251	0.037	45.177	<0.001*	1.286 (1.195-1.384)
Working Status	0.237	0.043	30.930	<0.001*	1.267 (1.166-1.377)
BMI	0.905	0.038	573.050	<0.001*	2.472 (2.295-2.662)
Diabetes Mellitus	1.305	0.173	57.182	<0.001*	3.688 (2.630-5.173)
Emotional Mental Disorder	0.713	0.048	216.699	<0.001*	2.041 (1.856-2.244)
Instant Food Consumption	0.102	0.040	6.722	0.010*	1.108 (1.025-1.197)
Constant	-3.844	0.113	1165.952	<0.001	0.021

* = *p-value* <0.05

Discussion

Most young adults are unaware of their abnormally high blood pressure and only realize its effects when they age (Gooding et al., 2014). Young adults are less sensitive to hypertension than older people because young adults visit doctors less often than older people (Everett & Zajacova, 2015). There is a long-term impact of early-life risk factors of hypertension in young adults. Hypertension patients aged below 40 years have low awareness, low diagnosis, and poor blood pressure control (Johnson et al., 2014). This study provided information regarding the risk factors related to hypertension.

Among 73,800 subjects, the prevalence of hypertension among young adults in this study is 4.1% based on RISKESDAS 2018 data. The prevalence of hypertension among young adults in this population is lower than in a study in Zimbabwe, i.e., 7.4%, based on survey conducted in Harare, Bulawayo, and Mashonaland East (Sabapathy et al., 2023). Similarly, a study conducted in Lahore, Pakistan showed that the prevalence of hypertension among young adults was 15.7% (Zaib Zahid et al., 2023). The study conducted in India has a higher prevalence of hypertension among young adults, i.e., 13.8%, based on The National Family Health Survey-4 data (Krishnamoorthy et al., 2023).

Males are more likely to suffer from hypertension than females (Ahammed et al., 2021; Liu et al., 2017; Singh et al., 2017; Wyszynska et al., 2023). Low education levels are more likely to suffer from high blood pressure (Mohammed Nawi et al., 2021; Singh et al., 2017; Wyszynska et al., 2023). Marital status affects the occurrence of hypertension, i.e., an unmarried subject has a greater risk of hypertension (Ramezankhani et al., 2019; Singh et al., 2017; Thawornchaisit et al., 2013). Unemployed subjects risk developing hypertensive disease more than employed people (Kato et al., 2023; Mohammed Nawi et al., 2021; Rismadi et al., 2021). Having exceeded body mass index is more at risk of developing hypertension (Ondimu et al., 2019; Patil et al., 2017; Thawornchaisit et al., 2013; Wyszynska et al., 2023; Zaki et al., 2021). People with diabetes mellitus are more at risk of developing hypertension (Singh et al., 2017; Thawornchaisit et al., 2013; Tsimihodimos et al., 2018; Zaki et al., 2021). Emotional mental disorder is one of the conditions that promotes the development of hypertension (Alwani et al., 2023; Hu et al., 2015; Munakata, 2018; Ojike et al., 2016). Smoking habits have a significant influence on hypertension (Jang, 2021; Lusno et al., 2020; Maharjan, 2018; Meher et al., 2023; Saladini et al., 2016; Wang et al., 2020).

Frequently eating instant sodium-rich foods also affects hypertension (Alsabieh et al., 2019; De Deus Mendonça et al., 2017; Shi et al., 2019).

Males are more at risk than females for hypertension. The results of this study are in line with the study in Albania regarding risk factors for hypertension among young adults, which states that males are more at risk of hypertension (Ahammed et al., 2021), and a study in Poland also showed similar results (Wyszyńska et al., 2023). A cross-sectional study conducted in Varanasi stated that males are at 1.97 times greater risk of hypertension (Singh et al., 2017). A study of young adults in Zimbabwe also found that males were 1.53 times more likely to develop hypertension than females (Sabapathy et al., 2023). Generally, males have less favorable habits compared to females, such as excess consumption of instant food, smoking, and alcohol consumption, leading to hypertension (Ahammed et al., 2021; Thawornchaisit et al., 2013). The low number of male visits to health services is also the reason for the higher rate of hypertension in males. Females are more likely to visit health services for consultation regarding birth control and regular gynecological services, where every visit will check their blood pressure so that females' blood pressure is more controlled compared to males (Everett & Zajacova, 2015).

Unmarried subjects are more likely to suffer hypertension. The result was in line with a study conducted in Iran that showed unmarried subjects had a 1.55 higher risk of hypertension (Ramezankhani et al., 2019). The study conducted in Poland showed that married men had 1.58 more significant risk of hypertension than unmarried men (Lipowicz & Lopuszanska, 2005). Being married may be a protective factor against hypertension. Marriage is associated with a greater likelihood of lower nighttime systolic blood pressure (SBP) in men. Marriage might have some advantages, i.e., better health, good nutritional, and maintained psychiatric wellness (Causland et al., 2014). The treatment for hypertension among young adults was better in married men. Married improved men's compliance against medical treatment (Kim & Kim, 2020).

Education affects the incidence of hypertension. According to a study conducted in China, education limited to elementary school or below has a greater risk of hypertension compared to someone who graduated from middle school or above (Sun et al., 2022). A low education level has an impact on low knowledge, and this affects a person's behaviour. Education influences a person's behaviour in receiving and implementing information. There is a relationship between education and the occurrence of hypertension. Highly educated individuals have easier access to health information. A person's education level influences the quality of behaviour in receiving and processing information, which will impact their health status (Pebrisiana et al., 2022; Singh et al., 2017).

This study found working status has a significant influence on hypertension, whereas unemployed have a greater risk of hypertension. This study's results align with a systematic review and meta-analysis study in Southeast Asian Countries, which states that the unemployed have a greater risk of hypertension (Mohammed Nawi et al., 2021). Unemployed people are 1.39 more at risk of hypertension compared to employed people (Blok et al., 2022). The results of a nationwide study in China state that unemployment conditions are a high risk of hypertension because not working will affect economic capacity. Thus unemployment will indirectly make someone less aware of hypertension treatment and control (Zheng et al., 2020). Unemployment is more at risk of hypertension because of low income; a person will have a limited variety of food, and health services received are more limited (Rismadi et al., 2021).

This study shows that body mass index has a significant effect on hypertension. Having a BMI \geq 25 has a 2.47 times greater risk of hypertension compared to having a

BMI<25. Some studies show similar results, including a study in Poland which states that having a BMI above average is a greater risk of hypertension (Wyszyńska et al., 2023), a study conducted in Nagpur (Patil et al., 2017), and the study conducted on South Asian population-based from nationally representative surveys (Shahimi et al., 2022). A cohort study conducted in Thailand over four years showed that BMI was significantly associated with an increased risk of hypertension (Thawornchaisit et al., 2013). A study conducted among young adults in Tenwek Mission Hospital, Bomet County, Kenya, in 2018 stated that having a BMI \geq 25 had a 3.05 more significant risk of hypertension (Ondimu et al., 2019). A study conducted in Malaysia stated that obese people have a 4.29 times risk of developing hypertension (Zaki et al., 2021). Likened to people with normal BMI, people with BMI \geq 25 are more likely to have metabolic disorders, such as fat metabolism disorder and insulin antibody disorder. Metabolic disorders can trigger cardiovascular disease, one of which is hypertension (Tang et al., 2022). Apart from other risk factors, BMI directly affects blood pressure. Nevertheless, the mechanism of BMI's influence on blood pressure is not definite (Landi et al., 2018). Several studies show that BMI is a critical and sensitive indicator for predicting high blood pressure (H. A. Lee & Park, 2018; Wu et al., 2019)

Comorbidities, such as diabetes mellitus, result in increased blood pressure. This study showed that diabetes mellitus has a significant impact on hypertension. A study conducted by (Petrie et al., 2018) indicates that the presence of comorbid diabetes mellitus puts a person at twice the risk of hypertension. A prospective cohort study conducted over four years in a national Thai Cohort Study showed similar results, where diabetes mellitus was strongly associated with hypertension (Thawornchaisit et al., 2013). Diabetes mellitus is also a variable that affects hypertension, as shown by a study conducted in Malaysia (Zaki et al., 2021). Research conducted in South Korea through The Fourth Korea National Health and Nutrition Examination Survey (KNHANES IV) states that hypertension is more prevalent in adults with diabetes mellitus in the general population and for both genders (H. S. Lee et al., 2013). Hypertension occurs due to complex metabolic disorders caused by diabetes mellitus. Increased circulating fluid volume and peripheral vascular resistance in patients with diabetes mellitus occur due to insulin resistance, which increases body fluid volume. This condition is closely related to increased blood pressure (Ohishi, 2018).

Psychological conditions also influence cardiovascular diseases, including hypertension. Psychologically, distress that occurs causes an increased risk of hypertension, 1.53 greater than a person without a mental disorder (Ojike et al., 2016). Stress was significantly related to hypertension in a cross-sectional study in China, where stressful conditions put a person 1.2 times more at risk of hypertension (Hu et al., 2015), and the results of a study in Jakarta, Indonesia showed that mental-emotional disorders made the risk of hypertension 4.095 times greater (Ekaningrum, 2021). Emotional mental disorders, e.g., stress, will trigger endothelial dysfunction that allows for an increase in blood pressure (Munakata, 2018). Emotional mental disorders may trigger habits that can cause an increase in blood pressure, such as physical inactivity due to a lack of enthusiasm for daily activities, smoking, consuming alcohol, and eating instant foods that contain high amounts of sodium.

This study found that instant food consumption leads to hypertension condition. A prospective cohort study in Spanish stated that high consumption of ultra-processed food creates a greater risk of hypertension (De Deus Mendonça et al., 2017). An 8-year cohort study in Thailand showed that dietary patterns are associated with hypertension. An unhealthy diet includes consuming processed foods, including instant foods that are low fiber, high in carbohydrates, saturated fat, and high in sodium (Shi et al., 2019). Generally, the combinations contained in instant food are carbohydrates, fat, and salt (Thawornchaisit

et al., 2013). Frequent consumption of instant food will trigger cardiometabolic, which causes an increase in blood pressure (Shin et al., 2014). People often consume high-sodium instant foods. If high amounts of sodium are consumed continuously, it will lead to hypertension (Ekaningrum, 2021).

Conclusion

The prevalence of hypertension among young adults in this study was 4.1% based on RISKESDAS 2018 data. Gender, education levels, working status, body mass index, diabetes mellitus, emotional mental disorder, and instant food consumption were associated with hypertension among young adults in Indonesia. The results of this study can be used to predict risk factors for hypertension among young adults in Indonesia. Preventive efforts against hypertension in Indonesia should be performed by educating the public regarding behaviors that trigger hypertension so that modifiable risk factors will be more controlled. Early detection of health among young adults also needs to be improved, considering the low awareness and willingness to go to health services. This study has limitations, so the recommendation for future researchers is to examine more variables that may become a risk for hypertension.

References

- Ahamed, B., Maniruzzaman, M., Talukder, A., & Ferdausi, F. (2021). Prevalence and Risk Factors of Hypertension Among Young Adults in Albania. *High Blood Pressure and Cardiovascular Prevention*, 28(1), 35–48. <https://doi.org/10.1007/s40292-020-00419-5>
- Alsabieh, M., Alqahtani, M., Altamimi, A., Albasha, A., Alsulaiman, A., Alkhamshi, A., Habib, S. S., & Bashir, S. (2019). Fast Food Consumption and its Associations with Heart Rate, Blood Pressure, Cognitive Function and Quality of Life. Pilot Study. *Heliyon*, 5(5), e01566. <https://doi.org/10.1016/j.heliyon.2019.e01566>
- Alwani, A. A., Singh, U., Sankhyan, S., Chandra, A., Rai, S. K., & Nongkynrih, B. (2023). Hypertension-related distress and its associated factors: findings from an urban primary health centre of South Delhi, India. *Journal of Family Medicine and Primary Care*, 2(9), 1885–1895. <https://doi.org/10.4103/jfmpc.jfmpc>
- Amisi, W. G., Nelwan, J. E., & Kolibu, F. K. (2018). Hubungan antara Hipertensi dengan Kejadian Penyakit Jantung Koroner pada Pasien yang Berobat di Rumah Sakit Umum Pusat Prof. Dr. R. D. Kandou Manado. *Kemas*, 7(4), 1–7.
- Azam, M., Hidayati, F. N., Fibriana, A. I., Bahrudin, U., & Aljunid, S. M. (2023). Isolated Systolic Hypertension among Diabetes Mellitus Subjects; a national cross-sectional study in Indonesia. *Kemas*, 19(1), 53–64. <https://doi.org/10.15294/kemas.v19i1.40548>
- Azinar, M., Azam, M., & Sugiharto, S. (2019). Peer Support Group as a Model for Improving Behavior of Hypertension Risk and Complication Control. *Journal of Health Education*, 4(1), 17–21. <https://doi.org/https://doi.org/10.15294/jhe.v4i1.24192>
- Blok, S., Haggengburg, S., Collard, D., Van Der Linden, E. L., Galenkamp, H., Moll Van Charante, E. P., Agyemang, C., & Van Den Born, B. J. H. (2022). The association between socioeconomic status and prevalence, awareness, treatment and control of hypertension in different ethnic groups: the Healthy Life in an Urban Setting study. *Journal of Hypertension*, 40(5), 897–907. <https://doi.org/10.1097/HJH.0000000000003092>
- Causland, F. R. M., Sacks, F. M., & Forman, J. P. (2014). Marital status, dipping and nocturnal blood pressure: Results from the Dietary Approaches to Stop Hypertension trial. *Journal of Hypertension*, 32(4), 756–761. <https://doi.org/10.1097/HJH.000000000000107>
- De Deus Mendonça, R., Souza Lopes, A. C., Pimenta, A. M., Gea, A., Martinez-Gonzalez, M. A., & Bes-Rastrollo, M. (2017). Ultra-processed food consumption and the incidence

- of hypertension in a mediterranean cohort: The seguimiento universidad de navarra project. *American Journal of Hypertension*, 30(4), 358–366.
<https://doi.org/10.1093/ajh/hpw137>
- Ekaningrum, A. Y. (2021). Hubungan Asupan Natrium, Lemak, Gangguan Mental Emosional, Dan Gaya Hidup Dengan Hipertensi Pada Dewasa Di Dki Jakarta. *Journal of Nutrition College*, 10(2), 82–92. <https://doi.org/10.14710/jnc.v10i2.30435>
- Everett, B., & Zajacova, A. (2015). Gender differences in hypertension and hypertension awareness among young adults. *Biodemography and Social Biology*, 61(1), 1–17. <https://doi.org/10.1080/19485565.2014.929488>
- Global Health Observatory. (2023). *Noncommunicable disease: risk factors*. Global Health Observatory.
- Gooding, H. C., McGinty, S., Richmond, T. K., Gillman, M. W., & Field, A. E. (2014). Hypertension awareness and control among young adults in the National Longitudinal Study of Adolescent Health. *Journal of General Internal Medicine*, 29(8), 1098–1104. <https://doi.org/10.1007/s11606-014-2809-x>
- Hu, B., Liu, X., Yin, S., Fan, H., Feng, F., & Yuan, J. (2015). Effects of psychological stress on hypertension in middle-aged Chinese: A cross-sectional study. *PLoS ONE*, 10(6), 1–13. <https://doi.org/10.1371/journal.pone.0129163>
- Jang, I. (2021). Pre-hypertension and Its Determinants in Healthy Young Adults: Analysis of Data from the Korean National Health and Nutrition Examination Survey VII. *International Journal of Environmental Research and Public Health*, 18(17). <https://doi.org/10.3390/ijerph18179144>
- Johnson, H. M., Thorpe, C. T., Bartels, C. M., Schumacher, J. R., Palta, M., Pandhi, N., Sheehy, A. M., & Smith, M. A. (2014). Undiagnosed hypertension among young adults with regular primary care use. *Journal of Hypertension*, 32(1), 65–74. <https://doi.org/10.1097/HJH.0000000000000008>
- Kato, A. M., Kibone, W., Okot, J., Baruch Baluku, J., & Bongomin, F. (2023). Self-Reported Hypertension and Associated Factors Among Adults in Butambala District, Central Uganda: A Community-Based Prevalence Study. *Integrated Blood Pressure Control*, 16(November), 71–80. <https://doi.org/10.2147/IBPC.S434230>
- Kim, J. S., & Kim, C. G. (2020). Gender differences in hypertension treatment and control in young adults. *Journal of Nursing Research*, 28(3), 1–14. <https://doi.org/10.1097/JNR.0000000000000359>
- Krishnamoorthy, Y., Rajaa, S., Sulgante, S., Chinnakali, P., Jaswal, N., & Goel, S. (2023). Prevalence of Hypertension and Determinants of Treatment-Seeking Behaviour among Adolescents and Young Adults in India: An Analysis of NFHS-4. *Journal of Public Health (Oxford, England)*, 45(1), e48–e56. <https://doi.org/10.1093/pubmed/fdac006>
- Lacobucci, G. (2023). Adults at lowest risk are most likely to have undiagnosed hypertension, data show. *BMJ*, 381, page 976.
- Landi, F., Calvani, R., Picca, A., Tosato, M., Martone, A. M., Ortolani, E., Sisto, A., D'angelo, E., Serafini, E., Desideri, G., Fuga, M. T., & Marzetti, E. (2018). Body mass index is strongly associated with hypertension: Results from the longevity check-up 7+ study. *Nutrients*, 10(12), 1–12. <https://doi.org/10.3390/nu10121976>
- Lee, H. A., & Park, H. (2018). Diet-related risk factors for incident hypertension during an 11-year follow-up: The Korean genome epidemiology study. *Nutrients*, 10(8). <https://doi.org/10.3390/nu10081077>
- Lee, H. S., Lee, S. S., Hwang, I. Y., Park, Y. J., Yoon, S. H., Han, K., Son, J. W., Ko, S. H., Park, Y. G., Yim, H. W., Lee, W. C., & Park, Y. M. (2013). Prevalence, awareness, treatment and control of hypertension in adults with diagnosed diabetes: The Fourth Korea National Health and Nutrition Examination Survey (KNHANES IV). *Journal of Human Hypertension*, 27(6), 381–387. <https://doi.org/10.1038/jhh.2012.56>
- Lipowicz, A., & Lopuszanska, M. (2005). Marital Differences in Blood Pressure and the Risk of Hypertension among Polish Men. *European Journal of Epidemiology*, 20(5), 421–427. <https://doi.org/10.1007/s10654-005-1752-x>
- Liu, X., Xiang, Z., Shi, X., Schenck, H., Yi, X., Ni, R., & Liu, C. (2017). The Risk Factors of

- High Blood Pressure among Young Adults in the Tujia-Nationality Settlement of China. *BioMed Research International*, 2017. <https://doi.org/10.1155/2017/8315603>
- Lusno, M. F. D., Haksama, S., Wulandari, A., & Sriram, S. (2020). Association between smoking and hypertension as a disease burden in Sidoarjo: a case-control study. *International Journal of Applied Biology*, 4(2), 9–16. <https://journal.unhas.ac.id/index.php/ijoab/article/view/10951>
- MacLeod, K. E., Ye, Z., Donald, B., & Wang, G. (2022). A Literature Review of Productivity Loss Associated with Hypertension in the United States. *Population Health Management*, 25(3), 297–308. <https://doi.org/10.1089/pop.2021.0201>
- Maharjan, B. (2018). Prevalence and Awareness of Hypertension among Adults and its Related Risk Factors. *Journal of Nepal Health Research Council*, 15(3), 242–246. <https://doi.org/10.3126/jnhrc.v15i3.18848>
- Meher, M., Pradhan, S., & Pradhan, S. R. (2023). Risk Factors Associated With Hypertension in Young Adults: A Systematic Review. *Cureus*, 15(4). <https://doi.org/10.7759/cureus.37467>
- Minasari, S., Azam, M., & Rahayu, S. R. (2022). Pengembangan Aplikasi Monitoring Penyakit Hipertensi dan Diabetes Mellitus Terintegrasi. *Jurnal Manajemen Kesehatan Indonesia*, 10(1), 28–42. <https://doi.org/10.14710/jmki.10.1.2022.28-42>
- Mohammed Nawi, A., Mohammad, Z., Jetly, K., Abd Razak, M. A., Ramli, N. S., Wan Ibadullah, W. A. H., & Ahmad, N. (2021). The Prevalence and Risk Factors of Hypertension among the Urban Population in Southeast Asian Countries: A Systematic Review and Meta-Analysis. *International Journal of Hypertension*, 2021. <https://doi.org/10.1155/2021/6657003>
- Munakata, M. (2018). Clinical significance of stress-related increase in blood pressure: current evidence in office and out-of-office settings. *Hypertension Research*, 41(8), 553–569. <https://doi.org/10.1038/s41440-018-0053-1>
- Ohishi, M. (2018). Hypertension with diabetes mellitus: Physiology and pathology review-article. *Hypertension Research*, 41(6), 389–393. <https://doi.org/10.1038/s41440-018-0034-4>
- Ojike, N., Sowers, J. R., Seixas, A., Ravenell, J., Rodriguez-Figueroa, G., Awadallah, M., Zizi, F., Jean-Louis, G., Ogedegbe, O., & McFarlane, S. I. (2016). Psychological distress and hypertension: results from the national health interview survey for 2004–2013. *CardioRenal Medicine*, 6(3), 198–208. <https://doi.org/10.1159/000443933>
- Ondimu, D. O., Kikuvi, G. M., & Otieno, W. N. (2019). Risk factors for hypertension among young adults (18-35) years attending in tenwek mission hospital, Bomet county, Kenya in 2018. *Pan African Medical Journal*, 33, 1–8. <https://doi.org/10.11604/pamj.2019.33.210.18407>
- Patil, C. R., Sahoo, D. P., Dhoble, M., Kherde, A., & Inamdar, A. (2017). Prevalence of hypertension and its associated risk factors in young adults attending a tertiary care institute of Nagpur: a cross sectional study. *International Journal Of Community Medicine And Public Health*, 4(10), 3630. <https://doi.org/10.18203/2394-6040.ijcmph20174223>
- Pebrisia, P., Tambunan, L. N., & Baringbing, E. P. (2022). Hubungan Karakteristik dengan Kejadian Hipertensi pada Pasien Rawat Jalan di RSUD Dr. Doris Sylvanus Provinsi Kalimantan Tengah. *Jurnal Surya Medika*, 8(3), 176–186. <https://doi.org/10.33084/jsm.v8i3.4511>
- Petrie, J. R., Guzik, T. J., & Touyz, R. M. (2018). Diabetes, Hypertension, and Cardiovascular Disease: Clinical Insights and Vascular Mechanisms. *Canadian Journal of Cardiology*, 34(5), 575–584. <https://doi.org/10.1016/j.cjca.2017.12.005>
- Ramezankhani, A., Azizi, F., & Hadaegh, F. (2019). Associations of marital status with diabetes, hypertension, cardiovascular disease and all-cause mortality: A long term follow-up study. *PLoS ONE*, 14(4), 1–15. <https://doi.org/10.1371/journal.pone.0215593>
- Rismadi, K., Siagian, A., & Siregar, F. A. (2021). Pengaruh Penghasilan dan Gaya Hidup Terhadap Kejadian Hipertensi pada Nelayan di Kota Medan. *Journal Health Sains*, 2(3), 1–13.

- Rokom. (2021). *Hipertensi Penyebab Utama Penyakit Jantung, Gagal Ginjal, dan Stroke*. Ministry of Health of the Republic of Indonesia. <https://sehatnegeriku.kemkes.go.id/baca/rilis-media/20210506/3137700/hipertensi-penyebab-utama-penyakit-jantung-gagal-ginjal-dan-stroke/>
- Sabapathy, K., Mwita, F. C., Dauya, E., Bandason, T., Simms, V., Dziva Chikwari, C., Doyle, A. M., Ross, D., Shah, A., Hayes, R. J., Schutte, A. E., Kranzer, K., & Ferrand, R. A. (2023). Prevalence of Hypertension and High-normal Blood Pressure among Young Adults in Zimbabwe: Findings from a Large, Cross-sectional Population-based Survey. *The Lancet Child & Adolescent Health*, 8(2), 101–111. [https://doi.org/10.1016/s2352-4642\(23\)00287-0](https://doi.org/10.1016/s2352-4642(23)00287-0)
- Saladini, F., Benetti, E., Fania, C., Mos, L., Casiglia, E., & Palatini, P. (2016). Effects of smoking on central blood pressure and pressure amplification in hypertension of the young. *Vascular Medicine*, 21(5), 422–428. <https://doi.org/10.1177/1358863X16647509>
- Shahimi, N. H., Lim, R., Mat, S., Goh, C. H., Tan, M. P., & Lim, E. (2022). Association between mental illness and blood pressure variability: a systematic review. *BioMedical Engineering Online*, 21(1), 1–22. <https://doi.org/10.1186/s12938-022-00985-w>
- Shi, Z., Papier, K., Yiengprugsawan, V., Kelly, M., Seubsman, S. A., & Sleigh, A. C. (2019). Dietary patterns associated with hypertension risk among adults in Thailand: 8-year findings from the Thai Cohort Study. *Public Health Nutrition*, 22(2), 307–313. <https://doi.org/10.1017/S1368980018002203>
- Shin, H. J., Cho, E., Lee, H. J., Fung, T. T., Rimm, E., Rosner, B., Manson, J. A. E., Wheelan, K., & Hu, F. B. (2014). Instant noodle intake and dietary patterns are associated with distinct cardiometabolic risk factors in Korea. *Journal of Nutrition*, 144(8), 1247–1255. <https://doi.org/10.3945/jn.113.188441>
- Singh, S., Shankar, R., & Singh, G. P. (2017). Prevalence and Associated Risk Factors of Hypertension: A Cross-Sectional Study in Urban Varanasi. *International Journal of Hypertension*, 2017. <https://doi.org/10.1155/2017/5491838>
- Social Security Agency of Health. (2017). *Tangkis Risiko Kardiometabolik dengan Optimalisasi PROLANIS*.
- Sun, K., Lin, D., Li, M., Mu, Y., Zhao, J., Liu, C., Bi, Y., Chen, L., Shi, L., Li, Q., Yang, T., Wan, Q., Wu, S., Wang, G., Luo, Z., Qin, Y., Tang, X., Chen, G., Huo, Y., ... Yan, L. (2022). Association of education levels with the risk of hypertension and hypertension control: a nationwide cohort study in Chinese adults. *Journal of Epidemiology and Community Health*, 76(5), 451–457. <https://doi.org/10.1136/jech-2021-217006>
- Tang, N., Ma, J., Tao, R., Chen, Z., Yang, Y., He, Q., Lv, Y., Lan, Z., & Zhou, J. (2022). The effects of the interaction between BMI and dyslipidemia on hypertension in adults. *Scientific Reports*, 12(1), 1–7. <https://doi.org/10.1038/s41598-022-04968-8>
- Thawornchaisit, P., De Looze, F., Reid, C. M., Seubsman, S. A., Sleigh, A. C., Chokhanapitak, J., Hounthasarn, S., Khamman, S., Pandee, D., Pangsap, S., Prapamontol, T., Puengson, J., Somboonsook, B., Sripaiboonkij, N., Somsamai, P., Vilainerun, D., Wimonwattanaphan, W., Pachanee, C. A., Tangmunkongvorakul, A., ... McClure, R. (2013). Health risk factors and the incidence of hypertension: 4-Year prospective findings from a national cohort of 60 569 Thai Open University students. *BMJ Open*, 3(6), 1–10. <https://doi.org/10.1136/bmjopen-2013-002826>
- Tsimihodimos, V., Gonzalez-Villalpando, C., Meigs, J. B., & Ferrannini, E. (2018). Hypertension and Diabetes Mellitus Coprediction and Time Trajectories. *Hypertension*, 71(3), 422–428. <https://doi.org/10.1161/HYPERTENSIONAHA.117.10546>
- Wang, M., Li, W., Zhou, R., Wang, S., Zheng, H., Jiang, J., Wang, S., Yu, C., Gao, W., Lv, J., Wu, T., Cao, W., Hu, Y., Li, L., & Ji, J. S. (2020). The paradox association between smoking and blood pressure among half million chinese people. *International Journal of Environmental Research and Public Health*, 17(8), 7–17. <https://doi.org/10.3390/ijerph17082824>
- World Health Organization. (2023). *Hypertension*. <https://www.who.int/news-room/fact-sheets/detail/hypertension>
- Wu, X., Li, B., Lin, W. Q., Huang, L. L., Wang, X. X., Fu, L. Y., Li, B. B., & Wang, P. X.

- (2019). The association between obesity indices and hypertension: Which index is the most notable indicator of hypertension in different age groups stratified by sex? *Clinical and Experimental Hypertension*, 41(4), 373–380.
<https://doi.org/10.1080/10641963.2018.1489546>
- Wyszyńska, J., Łuszczki, E., Sobek, G., Mazur, A., & Dereń, K. (2023). Association and Risk Factors for Hypertension and Dyslipidemia in Young Adults from Poland. *International Journal of Environmental Research and Public Health*, 20(2).
<https://doi.org/10.3390/ijerph20020982>
- Zaib Zahid, Muhammad Abdullah Shah, Haider Tanvir, Awais Amjad, Noor e Sehar Butt, & Fahad Asim. (2023). Prevalence and Associated Factors of Hypertension in Young Adults: A Cross-sectional Study. *The Professional Medical Journal*, 31(01), 44–48.
<https://doi.org/10.29309/tpmj/2024.31.01.7839>
- Zaki, N. A. M., Ambak, R., Othman, F., Wong, N. I., Man, C. S., Morad, M. F. A., He, F. J., MacGregor, G., Palaniveloo, L., & Baharudin, A. (2021). The prevalence of hypertension among Malaysian adults and its associated risk factors: data from Malaysian Community Salt Study (MyCoSS). *Journal of Health, Population and Nutrition*, 40(Suppl 1), 4–9. <https://doi.org/10.1186/s41043-021-00237-y>
- Zheng, C., Wang, Z., Wang, X., Chen, Z., Zhang, L., Kang, Y., Yang, Y., Jiang, L., & Gao, R. (2020). Social determinants status and hypertension: A Nationwide Cross-sectional Study in China. *Journal of Clinical Hypertension*, 22(11), 2128–2136.
<https://doi.org/10.1111/jch.14023>