- 1 Title
- 2 Geospatial analysis of malaria burden in Kagera region, North-western Tanzania using
- 3 health facility and community survey data
- 4 Authors
- 5 Daniel A. Petro, Nyimvua Shaban, Sijenunu Aaron, Frank Chacky, Samwel Lazaro,
- 6 Maciej F. Boni, and Deus S. Ishengoma
- 7 Corresponding author: Daniel Anderson Petro
- 8 Phone number: +255762001943
- 9 Email address: <u>petrodaniel945@gmail.com</u>
- 10 Author affiliations: University of Dar es Salaam, Dar es Salaam, Tanzania (D.A. Petro, N.
- 11 Shaban); National Malaria Control Programme, Dodoma, Tanzania (S. Aaron, F. Chacky, S.
- 12 Lazaro); Temple University, Philadelphia, USA (M.F. Boni); National Institute for Medical
- 13 Research; Kampala International University, Dar es Salaam, Tanzania (D.S. Ishengoma)
- 14 Running title: Geospatial burden of malaria in Kagera region, TZ
- 15 Keywords: Malaria, malaria hotspots, artemisinin partial resistance, spatial autocorrelation,
- 16 Kagera, Tanzania

#### 17 Abstract

- 18 Generating evidence of malaria burden in areas with confirmed artemisinin partial resistance
- 19 (ART-R) is the first step before developing a response strategy to the resistance. In this study,
- we assessed the burden of malaria in Kagera region (with recently confirmed ART-R) by
- 21 geospatial analysis using data from health facilities (HFs) and community surveys from 2017 to

2023. In 2983717/8124363 patients from HFs, rapid diagnostic test positivity rate was 36.7%
(range: 0-80%) and was similar in patients aged under-fives (33.1%, range: 0-79%) and ≥5
years (33.7%, range: 3-80%). The prevalence of malaria was 10.0% (range: 0-40.5%,
n=84999/ 853761) in pregnant women but it was 26.1% (3409/13065) in school children
(range: 0-78.4%). We identified hotspots and coldspots, and persistently high burden in 69/192
wards, school children, and patients aged ≥5 years, providing evidence for planning and
developing an ART-R response for the region.

Tanzania is one of the priority countries defined by the World Health Organization (WHO) as a 29 30 high-burden high-impact country; it accounted for an estimated 3.2% of all malaria cases and 4.4% of deaths reported worldwide in 2022 (1). In Tanzania, about 96% of all cases are 31 caused by Plasmodium falciparum, and the remaining are due to other species (P. malariae 32 and P. ovale) individually or in mixed infections (2-4). The National Malaria Control 33 Programme's (NMCP) strategy is to reduce the malaria burden and progress to substantially 34 control and eliminate using different effective interventions. The main strategies include 35 integrated malaria vector control (mainly with insecticide-treated nets), case management, 36 preventive therapies, and other supportive strategies (5). However, progress to malaria 37 38 elimination is under key biological threats such as the emergence and spread of insecticide (6) and antimalarial drug resistance (7), emergence of parasites with histidine-rich protein 2/339 (hrp2/3) gene deletions (8), and invasive vector species (Anopheles stephensi) in some 40 41 malaria-endemic countries in Africa (1).

42 Malaria transmission in Tanzania is heterogeneous with 93% of the population living in 43 areas where transmission occurs (*5*). Regions with very low and low transmission risk are 44 located in the central, northeastern, and southwestern parts surrounded on both sides by

moderate transmission regions and high transmission in the northwestern, southern, and
western parts (*9*). Various factors including the types of mosquito vectors, environmental
conditions, and other human-related factors are responsible for the current pattern of malaria
transmission in Tanzania (*10–12*).

Over the past four decades, malaria-endemic countries in sub-Saharan Africa have 49 reported resistance of *P. falciparum* to all major antimalarial drugs, including chloroquine, 50 sulfadoxine-pyrimethamine (SP) (7), and artemisinin derivatives and their currently used 51 partner drugs (13,14). Nevertheless, artemisinin-based combination therapies (ACTs) are still 52 53 highly effective against parasites of African origin (7). However, artemisinin partial resistance (ART-R) has been confirmed in Rwanda (15,16), Uganda (17), Tanzania (18) and Eritrea (19), 54 and one particular genotype has only been detected in the Horn of Africa, in Ethiopia (20), 55 Somalia (21) and Sudan (22,23). In addition, the mutation that originated in Rwanda has been 56 also reported in Kenya (24). In Tanzania, little was known about ART-R until in 2021 when a 57 country-wide survey through the project on molecular surveillance of malaria in Tanzania 58 59 (MSMT) reported a focus in Kagera region only, with high frequency (0.077) of parasites with mutations in Kelch13 gene (K13) and as high as 0.22 in Karagwe District Council (DC) which 60 are linked to ART-R in Africa (25). Based on WHO's criteria (7), a follow-up therapeutic 61 efficacy study conducted in 2022 confirmed the presence of ART-R in the region although the 62 tested drugs still had high efficacy exceeding 98.0% (18). 63

As per WHO's strategy to respond to ART-R in Africa, when resistance is confirmed, new strategies to contain, prevent, and mitigate the spread and consequences of ART-R must be considered urgently (*26*). As the first step towards the development of a response strategy, WHO recommends mapping and providing a detailed profile of the disease burden and its

spread in different areas (27). Therefore, this study aimed to investigate the temporal trends and spatial patterns of malaria burden in Kagera region and identify areas at the highest or lowest risk of malaria transmission as an initial stage towards the development of a response to ART-R in Tanzania. The findings from this study provide the government and other stakeholders with evidence to support the designing, deployment, and implementation of interventions to monitor, prevent, and contain the spread of ART-R within the region and to other areas of Tanzania.

## 75 Material and Methods

## 76 Study design and site

This study utilized aggregated secondary data collected retrospectively from health facilities
(HFs) and community surveys in Kagera region of north-western Tanzania. The region
consists of 8 councils, 192 wards, and 734 villages covering an area of approximately 35868
square kilometers. A description of the region's geographical boundaries and the source of its
economy is provided in Appendix A.





Figure 1. A map showing the 26 regions of Mainland Tanzania (white) and Kagera region (red), Lake Victoria and
Indian Ocean (lightblue), and neighboring countries (gray).

### 85 Data collation, management, and analysis

The data for this were downloaded from the District Health Information System – version 2 (DHIS2), a web-based software platform for reporting, analyzing, storing, and disseminating health data. This (DHIS2) is an electronic format for the paper-based Health Management Information System of the Tanzania Ministry of Health which is used by HFs to collect and report their routine data (*5*). The description of how the information is entered and maintained as well as quality control of the data is given elsewhere (*9,28*). The data used in this study

included monthly malaria testing of pregnant women attending first antenatal care (ANC)
clinics, symptomatic malaria patients seeking care at HFs (aged under-fives and ≥5 years)
both from 2017 to 2023 and school children through school malaria parasitological surveys
(SMPS) of 2015, 2017, 2019, and 2021 from the 37 selected wards in all councils of Kagera
region.

Using Excel (Office 2013) and R software version 4.3.2: (www.r-project.org), data 97 cleaning and preprocessing were done to ensure accuracy and consistency (Appendix B). 98 Monthly data of total malaria tests performed and positive tests (by RDTs) reported by all HFs 99 100 in the region were aggregated to provide annualized estimates. Malaria test-positivity rate (TPR) and prevalence were defined as the proportion of positive malaria tests among all 101 malaria tests in patients aged under-fives and  $\geq 5$  years, and pregnant women, respectively. 102 103 For school children, the average prevalence for each council and ward was used. Using R software, data analysis was done for overall, per year and study group, council-wise, and at 104 ward level, and the results presented in tables, figures or text. Python plotting tools (pandas 105 106 v2.2.0, matplotlib v3.8.2, numpy v1.26.3) (Python.org) were used to show temporal trends of TPR and prevalence in the councils. 107

Kruskal-Wallis test was used to assess for any significant differences in malaria burden among councils, wards, and across years. Data on administrative boundaries of Tanzania at lower levels and its neighboring countries were downloaded from various sources including the Tanzanian <u>National Bureau of Statistics</u> (NBS), gadm.org, and naturalearthdata.com accessed via the R package *rnaturalearth* v1.0.1. These were used to create the study site and used for choropleth mapping, spatial autocorrelation (SA), and hotspot analysis. The spatial distribution of malaria burden in the region was mapped using R packages *ggplot2* v3.5.1 and *sf* v1.0.15.

115 To investigate the spatial and temporal patterns of malaria burden in the region, the Global Moran's Index was calculated for each group and year. The value of this index ranges from -1 116 (negative SA) to +1 (positive SA) (29.30). For hotspot analysis, local spatial statistical methods 117 118 were applied through a spdep package v1.3.3 found in R software. The description of wards 119 and HFs included in the analysis is found in Appendix B and Figure 1, while the cut-offs applied to characterize malaria prevalence were; <1% (very low), 1 - <5% (low), 5 -<30% 120 (moderate), and  $\geq$ 30% (high) as previously described (31.32). For malaria TPR, the cut-offs 121 were <5% (very low), 5-<15% (low), 15-<30% (moderate), and  $\geq$ 30% (high) (Appendix C and 122 123 Table 1). The formulae and illustration of how SA and hotspot analysis were conducted can be

124 found elsewhere (29,30) and in Appendix D.

### 125 **Results**

### 126 **Reported malaria cases from Health Facilities (HFs)**

127 In the HFs in Kagera region, 36.7% of patients (range: 0-80%; n = 2983717/8124363) with 128 malaria-like symptoms seeking care from 2017 to 2023, had RDT-positive results. The TPRs 129 were similar in patients aged under-fives (33.1%, range: 0-79%) and  $\geq$ 5 years (33.7%, range: 130 3-80%), with clear geographic and temporal variations (p<0.001). Among under-fives, most of 131 the wards in the high stratum (TPR ≥30%) were from rural areas while wards in the very low 132 stratum (TPR <5%) came from urban areas (Appendix Table 2). Furthermore, five councils had 133 >10% of the wards with consistently high TPR (≥30%) throughout the years. These included 134 6/17 (35.3%) wards in Biharamulo DC, 12/43 (27.9%) in Muleba DC, 5/22 (22.7%) in Ngara DC, 4/23 (17.4%) in Karagwe DC and 4/29 (13.8%) in Bukoba DC. In Kyerwa DC, only 2/24 135 (8.3%) wards had consistently higher TPRs while only one ward (Kashai in Bukoba MC) 136 137 recorded the lowest TPR (<5%) (Appendix Table 4).

#### 138

Among patients aged  $\geq$ 5 years, the pattern was similar to that of under-fives, with a large number of wards observed in the high stratum throughout the years (Appendix Table 3). Similar to under-fives, the same five councils had >10% of the wards with consistently high TPRs ( $\geq$ 30%) over the years and these included, 7/17 (41.2%) wards in Biharamulo DC, 7/23 (30.4%) in Karagwe DC, 11/43 (25.6%) in Muleba DC, 5/22 (22.7%) in Ngara DC, and 4/29 (13.8%) wards in Bukoba DC. Only 1/24 (4.2%) ward in Kyerwa DC had consistently higher TPR ( $\geq$ 30%) (Appendix Table 4) (**Figure 2B**).



146

Figure 2. Spatial distribution of malaria test positivity rates (TPRs) in (A) under-fives (B) patients aged ≥5 years at
 ward level in Kagera region from 2017 to 2023. White areas are wards excluded from the analysis

# 149 Monthly variations of test positivity rates (TPRs) at health facility level

150 Monthly malaria TPR was higher (≥30%) in patients aged ≥5 years compared to under-fives in

all councils (in Muleba DC, Ngara DC, Biharamulo DC, Bukoba DC, and Karagwe DC) and the

pattern was similar in all years, except in Bukoba MC which recorded the lowest TPR (<5%) in

both age groups. In all councils except Bukoba MC and Missenyi DC, TPRs less than 20%



were rarely observed (**Figure 3**).



158

155

### 160 Malaria prevalence among pregnant women

- 161 From 2017 to 2023, the prevalence of malaria in pregnant women was 10.0% (range: 0-40.5%,
- 162 n=84999/ 853761), with significant differences among wards and across years (p<0.001). The
- 163 prevalence was consistently higher in Biharamulo DC, Karagwe DC, Muleba DC, and Kyerwa
- 164 DC, and low in Bukoba MC (<1%) throughout the years (**Figure 4**).



Figure 4. Trends of malaria prevalence in pregnant women by council, in Kagera region from 2017 to 2023. The
 pink-shaded regions show the Clopper-Pearson 95% confidence bands.

168	The largest number of wards (76.8%, n=116/151) were located in the moderate stratum
169	while the lowest (0.6%, n=1/159) was in the high-risk stratum of prevalence (Appendix Table
170	5). In addition, only 3/8 (37.5%) councils had ≥5% of the wards with higher prevalence (≥30%)
171	(including 3/17 (17.6%) wards in Biharamulo DC, 4/43 (9.3%) in Muleba DC and 2/23 (8.7%)
172	wards in Karagwe DC) while Kyerwa DC (n=1/24, 4.2%) and Bukoba DC $(n=1/29, 3.4\%)$ had
173	<5% of the wards with high prevalence. At least 5% of the wards from Bukoba DC (n= $2/29$ ,
174	6.9%) and Missenyi DC (n = $1/20$ , 5%) consistently recorded the lowest prevalence (Appendix

175 Table 6) (**Figure 5)**.



- Figure 5. Spatial distribution of malaria prevalence in pregnant women at ward level in Kagera region from
   2017 to 2023. White areas are wards excluded from the analysis.
- 1,0
- 179
- 180

## 181 Malaria prevalence in school children

- 182 The prevalence of malaria in school children was 26.1% (range: 0-78.4%; n=3409/1306)
- (Appendix E). A large number of wards (19/34, 55.9%) were located in the high stratum
- 184 (≥30%) and few wards (1/34, 2.9%) came from areas with very low stratum (<1%) (Appendix
- 185 Table 7). Furthermore, two wards from Ngara DC and two from Bukoba DC had persistently
- higher prevalence (≥30%) throughout the study period, while no ward recorded the lowest
- prevalence (<1%) for all years (Appendix Table 8) (**Figure 6**).



Figure 6. Spatial distribution of malaria prevalence in school children at ward level in Kagera region for 2015,
 2017, 2019, and 2021 surveys. White areas are wards not selected for the survey or excluded from the
 analysis.

### 192 Overall malaria test positivity rates at ward level

Among patients aged between under-fives and ≥5 years, a declining trend was observed in

under-fives and started to shift to older patients at the same time (Appendix Table 9). In both

age groups, an increasing trend was observed in 2023 (Appendix Figure 2).

### **196 Overall malaria prevalence at ward level**

- 197 Malaria prevalence was higher in school children than in pregnant women, a trend that was
- similar across years (Appendix Figure 3). Furthermore, prevalence in pregnant women showed
- a decreasing trend with minor variability before 2022 but increased by approximately 15% in
- 200 2023 (Appendix Table 9).

## 201 Global spatial autocorrelation analysis of malaria burden

- <sup>202</sup> Malaria burden in the region was clustered in under-fives, patients aged ≥5 years, and
- 203 pregnant women. Councils/wards with high/low burden were surrounded by other areas
- with high/low values as well for all years and the observed patterns were statistically
- significant (p<0.05) in all groups except school children (**Tables 1** and **2**).
- **Table 1**. Results of global spatial autocorrelation of malaria test positivity rates in patients aged under-fives
- 207 and  $\geq$  5 years in Kagera region

Group	Year	Global moran's	z score	p value	Pattern
		index			
	2017	0.7281	12.621	<0.0001	Clustered
	2018	0.7522	13.263	<0.0001	Clustered
	2019	0.7815	13.771	<0.0001	Clustered
Under-fives	2020	0.7391	13.528	<0.0001	Clustered
	2021	0.6745	12.331	<0.0001	Clustered
	2022	0.6724	12.084	<0.0001	Clustered
	2023	0.5229	9.440	<0.0001	Clustered
	2017	0.6561	11.392	<0.0001	Clustered
	2018	0.6852	12.091	<0.0001	Clustered
	2019	0.6806	11.995	<0.0001	Clustered
≥ 5 years	2020	0.6104	11.197	<0.0001	Clustered
	2021	0.6324	11.572	<0.0001	Clustered
	2022	0.5779	10.413	<0.0001	Clustered
	2023	0.4784	8.709	<0.0001	Clustered

- 208
- 209
- 210
- 211

Group	Year	Global Moran's	z-score	p-value	Pattern
		Index			
	2017	0.3556	6.2041	<0.0001	Clustered
	2018	0.5267	9.2397	<0.0001	Clustered
	2019	0.5518	9.663	<0.0001	Clustered
ANC*	2020	0.5383	9.990	<0.0001	Clustered
	2021	0.5706	10.294	<0.0001	Clustered
	2022	0.4961	9.275	<0.0001	Clustered
	2023	0.3487	6.589	<0.0001	Clustered
	2015	0.1688	0.7905	0.4292	Clustered
	2017	0.0465	0.4611	0.6448	Clustered
$SMPS^\dagger$	2019	-0.0445	0.1648	0.8691	Dispersed
	2021	-0.0313	0.1626	0.8709	Dispersed

#### 212 **Table 2**. Year-wise results of global spatial autocorrelation of malaria prevalence in Kagera region

213 \*ANC, Antenatal Care,

<sup>†</sup>SMPS, School malaria parasitological survey

#### 215 Hotspot analysis of malaria burden

Malaria hotspots were concentrated in the southern, central, and northwestern parts of Kagera region. Four councils (Ngara DC, Biharamulo DC, Muleba DC, and Bukoba DC) had >17% of the wards classified as malaria hotspots in patients aged under-fives and ≥5 years, and pregnant women while Karagwe DC and Kyerwa DC had <17% of wards as malaria hotspots in the same groups. The number of wards classified as hotspots of malaria in each group at 95% and 99% confidence levels (CLs) is presented in **Table 3** (Appendix Table 10).

223

#### Table 3. Number of wards per council classified as malaria hotspots in patients aged under-fives

		Hotspot wards	Hotspot wards
Group	Council	at 95% CL <sup>†</sup> (n/N, %)	at 99% CL (n/N, %)
	Ngara DC*	(19/22 ,86.4)	(17/22, 77.3)
	Biharamulo DC	(11/17, 64.7)	(8/17, 47.1)
Under-fives	Muleba DC	(15/43 ,34.9)	(12/43, 27.9)
	Bukoba DC	(9/29, 31.0)	(6/29, 20.7)
	Karagwe DC	(2/23, 8.7)	(1/23, 4.3)
	Kyerwa DC	(1/24, 4.2)	-
	Ngara DC	(20/22, 90.9)	(16/22, 72.7)
	Biharamulo DC	(9/17, 52.9)	(7/17, (41.2)
	Muleba DC	(16/43, 37.2)	(13/43, 30.2)
≥5 years	Bukoba DC	(8/29, 27.6)	(5/29, 17.2)
	Kyerwa DC	(3/24, 12.5)	(1/24, 4.2)
	Karagwe DC	(1/23, 4.3)	(1/23, 4.3)
	Biharamulo DC	(13/17, 76.5)	(11/17, 64.7)
	Ngara DC	(7/22 ,31.8)	(4/22, 18.2)
	Muleba DC	(13/43 ,30.2)	(10/43, 23.3)
	Bukoba DC	(7/29, 24.1)	(7/29, 24.1)
Pregnant women	Karagwe DC	(4/23, 17.4)	(3/23, 13.0)
	Kyerwa DC	(3/24 ,12.5)	(1/24, 4.2)
	Missenyi DC	(2/20, 10)	-

and ≥5 years, and pregnant women in Kagera region at 95 and 99% confidence levels

\*DC, district council, <sup>†</sup>CL, confidence level. N, number of wards within a council,

228 n, number of wards identified as hotspots

229 On the other hand, malaria coldspots were situated in the northeast and few in the

remaining parts of the region. Three councils (Bukoba MC, Missenyi DC, and Bukoba DC) had

231 >20% of the wards classified as malaria coldspots in patients aged under-fives, ≥5 years, and

- pregnant women while Kyerwa DC and Muleba DC had <20% of the wards. The number of
- wards classified as malaria coldspots in each group at 95% and 99% CLs is presented in
- Table 4 (Appendix Table 11).
- 235 Table 4. Number of wards per council classified as malaria coldspots in patients aged under-fives
- and ≥5 years, and pregnant women

		Coldspot wards	Coldspot wards
Group	Council	at 95% CL (n/N, %)	at 99% CL (n/N, %)
	Bukoba MC*	(13/14, 92.9)	(12/14, 85.7)
	Missenyi DC $^{\dagger}$	(17/20, 85)	(14/20, 70)
Under-fives	Bukoba DC	(15/29, 51.7)	(12/29, 41.4)
	Karagwe DC	(1/23, 4.3)	-
	Kyerwa DC	(1/24, 4.2)	-
	Muleba DC	(1/43, 2.3)	-
	Bukoba MC	(12/14, 85.7)	(11/14, 78.6)
	Missenyi DC	(17/20, 85)	(14/20, 70)
≥5 years	Bukoba DC	(14/29, 48.3)	(11/29, 37.9)
	Kyerwa DC	(1/24, 4.2)	-
	Muleba DC	(1/43, 2.3)	(1/43, 2.3)
	Bukoba MC	(10/14, 71.4)	(3/14, 21.4)
	Bukoba DC	(11/29, 37.9)	(9/29 31.0)
Pregnant women	Missenyi DC	(8/20, 40)	(2/20, 10)
	Kyerwa DC	(4/24, 16.7)	-
	Ngara DC	(1/22, 4.5)	-

237 \*DC, district council

<sup>†</sup>MC, municipal council,

<sup>239</sup> <sup>‡</sup>CL, confidence level. N, number of wards within a council, n, number of wards identified as coldspot

The locations of malaria hotspots and coldspots at 95% and 99% CLs in under-fives, ≥5

years, and pregnant women respectively are indicated in Appendix Figures S4 to S6.

## 242 **Discussion**

This study was conducted as an initial stage to respond to ART-R as recommended by WHO 243 (26) and aimed to undertake an assessment to generate, and provide evidence of the current 244 245 burden of malaria in Kagera region where ART-R was recently confirmed (18). We used spatial statistical and disease mapping techniques to assess the local burden of malaria and 246 identify potential malaria hotspots in Kagera region to support NMCP's efforts of developing a 247 response strategy against ART-R. Using aggregated secondary data from multiple sources, 248 we showed that over 35% (69/192) of all wards in Kagera region, mostly from rural areas, had 249 250 a high malaria burden, with these wards being clustered across years. The TPR was higher in 251 patients aged  $\geq$ 5 years than under-fives and this is comparable to previous studies conducted in Tanzania (11,12). The findings reported here provide important evidence to support the 252 253 designing and implementation of mitigation and response strategies to recently confirmed 254 ART-R in Kagera region and Tanzania in general.

255 Consistent with previous studies (12), this study reported high temporal and spatial 256 heterogeneity of malaria burden at council and ward levels within the region. Most of the wards, especially from rural areas, recorded a high malaria burden for most of the years; while 257 258 some wards mainly in urban and mixed/rural areas recorded the lowest burden (see Appendix 259 Tables 4,6, and 8). This finding was consistent across years and study groups possibly due to conditions supporting malaria transmission in rural compared to urban areas as previously 260 261 reported (33–35). The higher burden of malaria in rural areas is normally attributed to several factors including the presence of favorable conditions for malaria transmission in rural areas 262 such as stagnant water bodies that tend to create desirable breeding sites for mosquito 263 264 vectors (35). In addition, a large proportion of poor people tend to reside in rural areas and

265 normally live in poorly constructed houses which support higher mosquito biting rates and transmission of malaria (12,36,37). To reduce the burden in these areas, various interventions 266 such as distribution of insecticide-treated nets (ITNs), house improvement strategies, bio-267 268 larviciding in areas that are surrounded by water bodies, and behavioral change and 269 communication strategies need to be implemented, as these have been reported to 270 significantly reduce malaria transmission and the burden (38,39). On the other hand, the low burden of malaria in urban areas observed in this study is consistent with previous studies 271 (33,35). This is possibly due to the high socio-economic status of the urban population and, 272 273 the availability and access to malaria treatment and control measures when compared to rural areas (35). 274

A significant clustered spatial pattern of malaria burden reported in patients aged under-275 276 fives and  $\geq 5$  years, and pregnant women within the region aligns with other studies conducted elsewhere (29,30). The reports of areas with persistently high malaria burden in the southern, 277 central, and north-western parts of the region as reported in this study may be attributed to the 278 279 presence of similar risk factors such as environmental conditions, socio-economic status, healthcare infrastructures, and the presence of suitable mosquito breeding sites that tend to 280 281 create foci of malaria transmission (10,12,40). Hence, future interventions in the region to 282 reduce the local disease burden need to be designed based on these results to target the areas that have been identified to have a persistently high burden of malaria. 283 284 Malaria prevalence in Kagera region was high in school children compared to pregnant women, reaching as high as 78.4% in some wards. In previous studies, it was reported that 285 286 school children spend most of their time in the evening playing outside and less often use bed 287 nets, which tends to increase their chances of mosquito bites and hence increase the risk of

288 malaria infections (41–43). To reduce malaria transmission in school children, various initiatives have been undertaken by NMCP and other stakeholders to protect this risky group 289 including the distribution of ITNs through different channels like the school net program (5.44). 290 291 However, the prevalence of malaria infections in this asymptomatic group is still high (12,45). 292 Future studies are needed to explore more reasons and factors for this recurring observation in the region and the whole country and whether this will support the spread of the reported ART-293 R. On the other hand, pregnant women in Tanzania are normally given two or more doses of 294 SP for intermittent preventive treatment as well as ITNs during their first ANC visits in HFs, 295 296 which might have significantly contributed to reducing the risk of malaria infections in this group (28). However, the increasing trend of malaria prevalence in pregnant women observed 297 in 2023 needs further follow-up to uncover all possible reasons for this change. 298

299 The overall malaria TPR at the ward level was high  $(\geq 30\%)$  in both under-fives and patients aged  $\geq$ 5 years with minor differences. These findings are comparable to what has 300 been reported by previous studies in Tanzania and elsewhere (11, 12). The year-wise 301 302 comparison of TPR revealed a shift of TPR from under-fives to ≥5 years, similar to previous studies (46). This may be attributed to various efforts that have been undertaken by NMCP to 303 implement effective malaria interventions in vulnerable groups like under-fives while these 304 interventions may not have been widely considered in older age groups (11,28). This might 305 have led to the shift of malaria infections to individuals aged  $\geq 5$  years, with the peak of malaria 306 307 prevalence reported in children aged between 5 and 15 years (11). In addition, malaria TPR in under-fives and  $\geq$ 5 years showed an increasing trend in 2023. An in-depth analysis to uncover 308 possible reasons for these epidemiological changes in the region is required so that 309 appropriate measures can be planned, now that ART-R has been confirmed. 310

311 Malaria transmission in Kagera region was observed throughout the year with some of the months recording high values of TPR compared to others. This indicates that malaria 312 transmission in the region was not seasonal; an observation that can signify the presence of 313 314 other factors like population movements, changes in malaria control, or unexpected increases/decreases in breeding sites or mosquito populations (12,47,48). Causal effects for 315 316 these factors are in general difficult to establish, future studies are needed to explore and uncover the key drivers of persistent malaria transmission in this malaria-endemic region in the 317 country. 318

The findings presented here had some limitations that may be addressed in future 319 studies. Since this study included HFs with a reporting rate of 50% or more, future studies may 320 consider including all HFs during analysis and use different geostatistical methods to account 321 for monthly missing data, differences in utilization of HFs, treatment-seeking behaviors, and 322 distance to HFs. In addition, the HF data did include personal information such as age and 323 sex, and this limited additional analysis. For instance, previous studies showed a high burden 324 325 of malaria in school children (aged 5 - 15 years) but this could not be teased out because all patients aged ≥5 years were put in one group. Future studies may also include populations at 326 risk and use incidence to track changes in the disease burden within the region. 327

Overall, malaria infections in the region were higher in most of the wards from rural areas compared to urban areas, among school children compared to pregnant women, and among patients aged  $\geq$ 5 years than in under-fives. The observed heterogeneity of malaria burden underscores prioritization of Kagera region for additional malaria control mainly due to the currently circulating *P. falciparum* parasites with *K13*561H genotypes in the region (*18*). As early and preemptive action is preferred to late response (*49*), identifying specific

- resistance containment strategies (50) for 2024 and 2025 in addition to general malaria
- 335 control strategies will be critical for slowing down the spread of artemisinin-resistant parasites
- 336 from Kagera region to other parts of Tanzania.
- 337
- 338

#### 339 Biographical Sketch

- 340 Mr. Petro is a doctoral student at the University of Dar es Salaam, Dar es Salaam. His
- 341 research interests focus on malaria epidemiology.

### 342 Acknowledgment

- The authors would like to thank the MoH through NMCP, and all members of the Department
- of Mathematics, University of Dar es Salaam, and MSMT project team for their participation
- and invaluable support to make this work complete.
- 346 This study was partially funded by the National Institute for Medical Research (NIMR)
- under the MSMT project. The MSMT project is funded in whole, by the Bill & Melinda Gates
- Foundation [grant number INV 02202]. Under the grant conditions of the Foundation, a
- Creative Commons Attribution 4.0 Generic License has already been assigned to the author of
   the accepted Manuscript version that might arise from this submission.
- 351 PAP and DSI designed the study, and SA, FC and SL supported the acquisition and
- 352 cleaning of the data. DAP conducted the initial analysis under the supervision of DSI and NS.
- 353 DSI, NS, and MFB were involved in the interpretation of the results. DAP and MFB performed
- a final analysis of the data under the guidance of DSI, and DAP drafted the initial manuscript.
- 355 All co-authors contributed to the revision of the initial draft and subsequent versions of the

manuscript. All authors read and approved the final manuscript, and declared no competing
 interest.

#### 358 **References**

- 1. World Health Organization. World Malaria Report 2023 [Internet]. World Health
- 360 Organization; 2023. Available from:
- 361 https://www.who.int/publications/i/item/9789240086173
- 2. Popkin Hall ZR, Seth MD, Madebe RA, Budodo R, Bakari C, Francis F, et al. Malaria
- 363 species positivity rates among symptomatic individuals across regions of differing
- transmission intensities in Mainland Tanzania. J Infect Dis [Internet]. 2023 Nov 22;

365 Available from: http://dx.doi.org/10.1093/infdis/jiad522

- 366 3. Popkin-Hall ZR, Seth MD, Madebe RA, Budodo R, Bakari C, Francis F, et al. Prevalence
- of non-falciparum malaria infections among asymptomatic individuals in four regions of
- Mainland Tanzania. Parasit Vectors [Internet]. 2024;17(1):153. Available from:
- 369 https://doi.org/10.1186/s13071-024-06242-4
- 4. Sendor R, Mitchell CL, Chacky F, Mohamed A, Mhamilawa LE, Molteni F, et al. Similar
- 371 Prevalence of Plasmodium falciparum and Non-P. falciparum Malaria Infections among
- 372 Schoolchildren, Tanzania1. Emerg Infect Dis [Internet]. 2023 Jun;29(6):1143–53.
- 373 Available from: http://dx.doi.org/10.3201/eid2906.221016
- 5. National Malaria Control Programme (NMCP). "National Malaria Strategic Plan 2021-
- 2025: Transitioning to Malaria Elimination Phase", Tanzania Ministry of Health,
- 376 Community Development, Gender, Elderly and Children. Dodoma; 2020 p. 1–179.
- 377 6. World Health Organization. Global-report-on-insecticide-resistance-in-malaria-vectors-
- 378 2010-2016 [Internet]. 2018. Available from: https://fundacionio.com/wp-

- content/uploads/2020/11/Global-report-on-insecticide-resistance-in-malaria-vectors-2010 2016.pdf
- World Health Organization. Report on antimalarial drug efficacy, resistance and response:
  10 years of surveillance (2010-2019). World Health Organization; 2020.
- 8. Watson OJ, Tran TNA, Zupko RJ, Symons T, Thomson R, Visser T, et al. Global risk of
   selection and spread of Plasmodium falciparum histidine-rich protein 2 and 3 gene

385 deletions. medRxiv. 2023;

- 386 9. Thawer SG, Golumbeanu M, Munisi K, Aaron S, Chacky F, Lazaro S, et al. The use of
- routine health facility data for micro-stratification of malaria risk in mainland Tanzania.
- Malar J [Internet]. 2022 Nov;21(1):345. Available from: http://dx.doi.org/10.1186/s12936 022-04364-7
- 10. Rumisha SF, Shayo EH, Mboera LEG. Spatio-temporal prevalence of malaria and

anaemia in relation to agro-ecosystems in Mvomero district, Tanzania. Malar J.

**2019;18(1):1–14**.

- 11. Ishengoma DS, Mmbando BP, Mandara CI, Chiduo MG, Francis F, Timiza W, et al.
- 394 Trends of Plasmodium falciparum prevalence in two communities of Muheza district

395 North-eastern Tanzania: correlation between parasite prevalence, malaria interventions

- and rainfall in the context of re-emergence of malaria after two decades of progressive.
- <sup>397</sup> Malar J [Internet]. 2018 Jul;17(1):252. Available from: http://dx.doi.org/10.1186/s12936-

398 **018-2395-1** 

Mandai SS, Francis F, Challe DP, Seth MD, Madebe RA, Petro DA, et al. High prevalence
 and risk of malaria among asymptomatic individuals from villages with high rates of
 artemisinin partial resistance in Kyerwa district, North-western Tanzania. medRxiv.

402 **2023;2010–23**.

- 13. Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, et al. Artemisinin resistance in
   Plasmodium falciparum malaria. N Engl J Med. 2009;361(5):455–67.
- 14. Noedl H, Se Y, Schaecher K, Smith BL, Socheat D, Fukuda MM. Evidence of artemisinin resistant malaria in western Cambodia. N Engl J Med. 2008;359(24):2619–20.
- 15. Uwimana A, Legrand E, Stokes BH, Ndikumana JLM, Warsame M, Umulisa N, et al.
- 408 Emergence and clonal expansion of in vitro artemisinin-resistant Plasmodium falciparum
- kelch13 R561H mutant parasites in Rwanda. Nat Med. 2020;26(10):1602–8.
- 16. Uwimana A, Umulisa N, Venkatesan M, Svigel SS, Zhou Z, Munyaneza T, et al.
- 411 Association of Plasmodium falciparum kelch13 R561H genotypes with delayed parasite
- 412 clearance in Rwanda: an open-label, single-arm, multicentre, therapeutic efficacy study.
- 413 Lancet Infect Dis. 2021;21(8):1120–8.
- 414 17. Balikagala B, Fukuda N, Ikeda M, Katuro OT, Tachibana SI, Yamauchi M, et al. Evidence
  415 of artemisinin-resistant malaria in Africa. N Engl J Med. 2021;385(13):1163–71.
- 18. Ishengoma DS, Mandara CI, Bakari C, Fola AA, Madebe RA, Seth MD, et al. Evidence of
- 417 artemisinin partial resistance in North-western Tanzania: clinical and drug resistance

418 markers study. medRxiv [Internet]. 2024 Feb 1; Available from:

- 419 http://dx.doi.org/10.1101/2024.01.31.24301954
- 19. Mihreteab S, Platon L, Berhane A, Stokes BH, Warsame M, Campagne P, et al.
- 421 Increasing Prevalence of Artemisinin-Resistant HRP2-Negative Malaria in Eritrea. N Engl
- 422 J Med [Internet]. 2023 Sep 28;389(13):1191–202. Available from:
- 423 http://dx.doi.org/10.1056/NEJMoa2210956
- 424 20. Fola AA, Feleke SM, Mohammed H, Brhane BG, Hennelly CM, Assefa A, et al.

- 425 Plasmodium falciparum resistant to artemisinin and diagnostics have emerged in Ethiopia.
- 426 Nat Microbiol [Internet]. 2023 Oct;8(10):1911–9. Available from:
- 427 http://dx.doi.org/10.1038/s41564-023-01461-4
- 428 21. Jalei AA, Na-Bangchang K, Muhamad P, Chaijaroenkul W. Monitoring antimalarial drug-
- resistance markers in Somalia. Parasites Hosts Dis [Internet]. 2023 Feb;61(1):78–83.
- 430 Available from: http://dx.doi.org/10.3347/PHD.22140
- 431 22. Mohamed AO, Hussien M, Mohamed A, Suliman A, Elkando NS, Abdelbagi H, et al.
- 432 Assessment of Plasmodium falciparum drug resistance molecular markers from the Blue
- 433 Nile State, Southeast Sudan. Malar J [Internet]. 2020 Feb 18;19(1):78. Available from:
- 434 http://dx.doi.org/10.1186/s12936-020-03165-0
- 435 23. Molina-de la Fuente I, Sagrado Benito MJ, Ousley J, Gisbert F de B, García L, González
- V, et al. Screening for K13-Propeller Mutations Associated with Artemisinin Resistance in
- 437 Plasmodium falciparum in Yambio County (Western Equatoria State, South Sudan). Am J
- 438 Trop Med Hyg [Internet]. 2023 Nov 1;109(5):1072–6. Available from:
- 439 http://dx.doi.org/10.4269/ajtmh.23-0382
- 440 24. Owoloye A, Olufemi M, Idowu ET, Oyebola KM. Prevalence of potential mediators of
- 441 artemisinin resistance in African isolates of Plasmodium falciparum. Malar J [Internet].
- 442 2021 Dec 2;20(1):451. Available from: http://dx.doi.org/10.1186/s12936-021-03987-6
- 443 25. Juliano JJ, Giesbrecht DJ, Simkin A, Fola AA, Lyimo BM, Pereus D, et al. Country wide
- surveillance reveals prevalent artemisinin partial resistance mutations with evidence for
- 445 multiple origins and expansion of high level sulfadoxine-pyrimethamine resistance
- 446 mutations in northwest Tanzania. medRxiv [Internet]. 2023 Nov 30; Available from:
- 447 http://dx.doi.org/10.1101/2023.11.07.23298207

- 448 26. World Health Organization. Strategy to respond to antimalarial drug resistance in Africa
- 449 2022 [Internet]. Vol. 04. Geneva, Switzerland: World Health Organization; 2022. Available
- 450 from:
- 451 https://www.who.int/publications/i/item/9789240060265%0Ahttps://cdn.who.int/media/docs
- 452 /default-source/malaria/who-antimalarial-drug-resistance-strategy-for-
- 453 consultation.pdf?sfvrsn=9d4eaa0\_6
- 454 27. Zupko RJ, Nguyen TD, Ngabonziza JCS, Kabera M, Li H, Tran TNA, et al. Modeling policy
- 455 interventions for slowing the spread of artemisinin-resistant pfkelch R561H mutations in
- 456 Rwanda. Nat Med [Internet]. 2023 Nov;29(11):2775–84. Available from:
- 457 http://dx.doi.org/10.1038/s41591-023-02551-w
- 458 28. National Malaria Control Programme (NMCP). Ministry Of Health, Community
- 459 Development, Gender, Elderly and Children, "National Guidelines For Malaria Diagnosis,
- 460 Treatment And Preventive Therapies." Dodoma; 2020.
- 29. Tewara MA, Mbah-Fongkimeh PN, Dayimu A, Kang F, Xue F. Small-area spatial statistical
- 462 analysis of malaria clusters and hotspots in Cameroon;2000–2015. BMC Infect Dis
- 463 [Internet]. 2018 Dec 7;18(1):636. Available from: https://doi.org/10.1186/s12879-018-
- 464 **3534-6**
- 465 30. Saxena R, Nagpal BN, Das MK, Srivastava A, Gupta SK, Kumar A, et al. A spatial
- statistical approach to analyze malaria situation at micro level for priority control in Ranchi
- district, Jharkhand. Indian J Med Res [Internet]. 2012 Nov;136(5):776–82. Available from:
- 468 https://www.ncbi.nlm.nih.gov/pubmed/23287124
- 31. Ministry of Health Community Development Gender Elderly & Children, National Malaria
- 470 Control Programme. Supplementary malaria midterm strategic plan 2018–2020. 2019.

471	32.	Ministry of Health, Community Development, Gender, Elderly and Children, National
472		Malaria Control Programme. Consultative Malaria Expert Meeting Report 2018. 2018.
473	33.	Nundu SS, Culleton R, Simpson SV, Arima H, Muyembe JJ, Mita T, et al. Malaria parasite
474		species composition of Plasmodium infections among asymptomatic and symptomatic
475		school-age children in rural and urban areas of Kinshasa, Democratic Republic of Congo.
476		Malar J. 2021;20:1–13.
477	34.	Sultana M, Sheikh N, Mahumud RA, Jahir T, Islam Z, Sarker AR. Prevalence and
478		associated determinants of malaria parasites among Kenyan children. Trop Med Health.
479		2017;45:1–9.
480	35.	Mbah CE, Ambe LA, Ngwewondo A, Kidzeru EB, Akwah L, Mountchissi C, et al. A
481		Comparative Study of Asymptomatic Malaria in a Forest Rural and Depleted Forest Urban
482		Setting during a Low Malaria Transmission and COVID-19 Pandemic Period. Biomed Res
483		Int. 2022;2022.
484	36.	Bofu RM, Santos EM, Msugupakulya BJ, Kahamba NF, Swilla JD, Njalambaha R, et al.
485		The needs and opportunities for housing improvement for malaria control in southern
486		Tanzania. Malar J. 2023;22(1):69.
487	37.	Tusting LS, Rek J, Arinaitwe E, Staedke SG, Kamya MR, Cano J, et al. Why is malaria
488		associated with poverty? Findings from a cohort study in rural Uganda. Infectious
489		Diseases of Poverty [Internet]. 2016 Aug;5(1):78. Available from:
490		http://dx.doi.org/10.1186/s40249-016-0164-3
491	38.	McCann RS, van den Berg H, Diggle PJ, van Vugt M, Terlouw DJ, Phiri KS, et al.
492		Assessment of the effect of larval source management and house improvement on
493		malaria transmission when added to standard malaria control strategies in southern

494		Malawi: study protocol for a cluster-randomized controlled trial. BMC Infect Dis [Internet].
495		2017 Sep 22;17(1):639. Available from: http://dx.doi.org/10.1186/s12879-017-2749-2
496	39.	Orkoh E, Efobi U. Effects of Behaviour Change Communication on Knowledge and
497		Prevention of Malaria Among Women in Ghana. Eval Rev [Internet]. 2023 Aug
498		11;193841X231194565. Available from: http://dx.doi.org/10.1177/0193841X231194565
499	40.	Dabaro D, Birhanu Z, Negash A, Hawaria D, Yewhalaw D. Effects of rainfall, temperature,
500		and topography on malaria incidence in elimination targeted district of Ethiopia. Malar J.
501		2021;20(1):1–10.
502	41.	Finda MF, Moshi IR, Monroe A, Limwagu AJ, Nyoni AP, Swai JK, et al. Linking human
503		behaviors and malaria vector biting risk in south-eastern Tanzania. PLoS One.
504		2019;14(6):e0217414.
505	42.	Kihwele F, Gavana T, Makungu C, Msuya HM, Mlacha YP, Govella NJ, et al. Exploring
506		activities and behaviors potentially increases school-age children's vulnerability to malaria
507		infections in south-eastern Tanzania. Malar J. 2023;22(1):293.
508	43.	Kihwele F, Gavana T, Makungu C, Msuya HM, Govella NJ, Mlacha YP, et al. Outdoor
509		activities and behaviors increase the vulnerability of school-age children to malaria
510		infections in south-eastern Tanzania [Internet]. Research Square. 2023. Available from:
511		https://www.researchsquare.com/article/rs-2832075/v1
512	44.	Yukich J, Stuck L, Scates S, Wisniewski J, Chacky F, Festo C, et al. Sustaining LLIN
513		coverage with continuous distribution: the school net programme in Tanzania. Malar J
514		[Internet]. 2020;19:158. Available from: http://dx.doi.org/10.1186/s12936-020-03222-8
515	45.	Chacky F, Runge M, Rumisha SF, Machafuko P, Chaki P, Massaga JJ, et al. Nationwide
516		school malaria parasitaemia survey in public primary schools, the United Republic of

- 517 Tanzania [Internet]. Vol. 17, Malar J. 2023. Available from:
- 518 http://dx.doi.org/10.1186/s12936-018-2601-1
- 46. Kigozi SP, Kigozi RN, Epstein A, Mpimbaza A, Sserwanga A, Yeka A, et al. Rapid shifts in
- 520 the age-specific burden of malaria following successful control interventions in four
- regions of Uganda. Malar J [Internet]. 2020 Mar 30;19(1):128. Available from:
- 522 http://dx.doi.org/10.1186/s12936-020-03196-7
- 47. Dida GO, Anyona DN, Abuom PO, Akoko D, Adoka SO, Matano AS, et al. Spatial
- distribution and habitat characterization of mosquito species during the dry season along
- 525 the Mara River and its tributaries, in Kenya and Tanzania. Infectious diseases of poverty.

# 526 **2018;7(1):1–16**.

- 48. Fornace KM, Diaz AV, Lines J, Drakeley CJ. Achieving global malaria eradication in
  changing landscapes. Malar J. 2021;20(1):1–14.
- 49. Boni MF, White NJ, Baird JK. The Community As the Patient in Malaria-Endemic Areas:
- 530 Preempting Drug Resistance with Multiple First-Line Therapies. PLoS Med [Internet]. 2016
- 531 Mar;13(3):e1001984. Available from: http://dx.doi.org/10.1371/journal.pmed.1001984
- 532 50. Boni MF. Breaking the cycle of malaria treatment failure. Front Epidemiol [Internet]. 2022
- 533 Dec 14;2:1041896. Available from: http://dx.doi.org/10.3389/fepid.2022.1041896

# 534 Appendices

- 535 Appendix A. Description of the study site
- 536 **Appendix B.** Administrative units and health facilities in Kagera region that were included in the analysis
- 537 Appendix C: Stratification of malaria burden in Kagera region
- 538 Appendix D: Description of spatial autocorrelation analysis
- 539 Appendix E: Description of wards and schools selected for school malaria parasitological surveys
- 540 **Appendix Figure 1**. Descriptive summary of the wards and health facilities included in the analysis

- 541 Appendix Figure 2 Comparison of malaria test positivity rates in Kagera region from 2017 to 2023 based on
- 542 the data from health facilities
- 543 **Appendix Figure 3.** Comparison of malaria prevalence in Kagera region from 2017 to 2023 based on the
- 544 data from pregnant women and school children
- 545 Appendix Figure 4 Locations of hotspots and coldspots of malaria in under-fives at ward level in Kagera
- 546 region from 2017 to 2023
- 547 Appendix Figure 5 Locations of hotspots and coldspots of malaria in patients aged ≥5 years at ward level in
- 548 Kagera region from 2017 to 2023
- 549 **Appendix Figure 6.** Locations of hotspots and coldspots of malaria in pregnant women at ward level in
- 550 Kagera region from 2017 to 2023
- 551 Appendix Table 1. Cut-offs adopted to categorize councils/wards in Kagera region into different risk strata
- 552 Appendix Table 2. Number of wards by year in each stratum of malaria test positivity rates in under-fives in
- 553 Kagera region from 2017 to 2023
- 554 **Appendix Table 3.** Number of wards by year in each stratum of malaria test positivity rates in patients aged  $\geq 5$
- 555 years in Kagera region from 2017 to 2023
- 556 Appendix Table 4. Ward names by council that had a high, moderate, low, and very low malaria test positivity
- 557 rates in under-fives and ≥5 years from 2017 to 2023, Kagera region
- 558 Appendix Table 5. Number of wards by year in each stratum of malaria prevalence in pregnant women in Kagera
- 559 region from 2017 to 2023
- 560 Appendix Table 6. Ward names by council that had a high, moderate, low, and very low malaria prevalence in
- 561 pregnant women from 2017 to 2023, Kagera region
- 562 Appendix Table 7. Number of wards by year in each stratum of malaria prevalence in school children in Kagera
- 563 region from 2017 to 2023
- 564 Appendix Table 8. Ward names by council that had a high, moderate, low, and very low malaria prevalence in
- school children from 2017 to 2023, Kagera region
- 566 Appendix Table 9. Overall malaria burden at ward level in Kagera region by years and study groups from
- 567 2015 to 2023

- 568 **Appendix Table 10.** Names of hotspots of malaria burden at ward level in Kagera region at 95% and 99%
- 569 confidence levels from 2017 to 2023
- 570 Appendix Table 11. Names of coldspots of malaria burden at ward level in Kagera region at 95% and 99%
- 571 confidence levels from 2017 to 2023