

## The Cystic Fibrosis Transmembrane Regulator Controls Tolerogenic Responses to Food Allergens in Mice and Humans

Marc Emmenegger<sup>1,2,#</sup>, Chryssa Zografou<sup>1</sup>, Yile Dai<sup>3</sup>, Laura R. Hoyt<sup>4</sup>, Ravindra Gudneppanavar<sup>5</sup>, Andra Chincisan<sup>1</sup>, Hubert Rehrauer<sup>6</sup>, Falko J. Noé<sup>6</sup>, Natalia Zajac<sup>6</sup>, Georg Meisl<sup>7</sup>, Matthias M. Schneider<sup>7</sup>, Hang Nguyen<sup>5</sup>, Katja Höpker<sup>8</sup>, Tuomas P. J. Knowles<sup>7</sup>, Mireia Sospedra<sup>9</sup>, Roland Martin<sup>9</sup>, Aaron M. Ring<sup>3,10</sup>, Stephanie Leeds<sup>5</sup>, Stephanie C. Eisenbarth<sup>4,11</sup>, Marie E. Egan<sup>5,12</sup>, Emanuela M. Bruscia<sup>5</sup>, and Adriano Aguzzi<sup>1,#</sup>

<sup>1</sup>Institute of Neuropathology, University of Zurich, CH-8091 Zurich, Switzerland.

<sup>2</sup>Division of Medical Immunology, Laboratory Medicine, University Hospital Basel, CH-4031 Basel, Switzerland.

<sup>3</sup>Department of Immunology, Yale School of Medicine, Yale University, New Haven, Connecticut, USA.

<sup>4</sup>Department of Immunobiology, Yale University School of Medicine, New Haven, Connecticut, USA.

<sup>5</sup>Department of Paediatrics, Yale School of Medicine, New Haven, Connecticut, USA.

<sup>6</sup>Functional Genomics Center Zurich, ETH Zurich and University of Zurich, Zurich Switzerland.

<sup>7</sup>Centre for Misfolding Diseases, Yusuf Hamied Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, United Kingdom.

<sup>8</sup>University of Cologne, Faculty of Medicine and University Hospital Cologne, Clinic III for Internal Medicine, Cologne, Germany.

<sup>9</sup>Neuroimmunology and Multiple Sclerosis Research, Neurology Clinic, Frauenklinikstrasse 26, 8091 Zurich, University Hospital Zurich, University Zurich, Switzerland.

<sup>10</sup>Translational Science and Therapeutics Division, Fred Hutchinson Cancer Center, Seattle, Washington, 98102, USA.

<sup>11</sup>Department of Medicine, Division of Allergy and Immunology, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA.

<sup>12</sup>Department of Cellular and Molecular Physiology, Yale University School of Medicine, New Haven, Connecticut, USA.

#corresponding authors: [marc.emmenegger@usz.ch](mailto:marc.emmenegger@usz.ch) and [adriano.aguzzi@usz.ch](mailto:adriano.aguzzi@usz.ch)

**NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.**

## Highlights

- We investigated serum IgG against Ara-h2 in 24,536 patients and identified 133 seropositives.
- Seropositivity was associated with cystic fibrosis, and these patients had IgG4 but not IgE.
- 30 • We reproduced these results in a paediatric validation cohort and in a *Cftr*<sup>-/-</sup> mouse model.
- A compromised epithelial barrier and the cytokine milieu might explain this phenotype.

## Abstract

IgE antibodies against the allergen Ara-h2 can cause life-threatening anaphylaxis upon exposure to peanuts. Desensitization strategies aim at inducing IgG responses against Ara-h2 which may compete with anaphylactogenic IgE. Here we assessed anti-Ara-h2 titers in an unselected cohort of 24,536 adult patients admitted to a general hospital for disparate medical reasons. Surprisingly, adult (n=177) and pediatric (n=76) patients with cystic fibrosis (pwCF) had IgG4, but not IgE, against several peanut and soybean allergens, yet did not suffer from peanut allergy. Antibody repertoires were not globally perturbed in pwCF, and heterozygous Cystic Fibrosis Transmembrane Regulator (*CFTR*) mutation carriers had the same prevalence of food allergies as pwCF. Peanut sensitisation of *Cftr*<sup>-/-</sup> mice failed to induce IgE and was associated with elevated IFN- $\gamma$ . We conclude that *CFTR* is a key regulator of anaphylactogenic and tolerogenic responses to food allergens. *CFTR*-controlled cytokine responses including IFN- $\gamma$ , in combination with a compromised epithelial barrier, may trigger a preferential IgG4 response resulting in tolerance to food allergens.

35  
40  
45

## Introduction

Considerable quantities of food-derived antigens are absorbed through the intestinal epithelia every day. This requires oral immunological tolerance to these intrinsically innocuous food constituents. Oral tolerance permits selective responsiveness to food antigens or commensals, a phenomenon conventionally  
50 termed 'systemic unresponsiveness' (Moran & Burks, 2015; Pabst & Mowat, 2012; Tordesillas & Berin, 2018). Mucosal immunoglobulins (Ig) A directed against food constituents are common in non-allergic individuals (Berin, 2012; Frossard et al., 2004; E. G. Liu et al., 2022); serum IgG to some food allergens is occasionally detected in those without food allergy (Husby et al., 1985; Leviatan et al., 2022), and antigen-  
55 experienced CD4<sup>+</sup> T cells are required for the induction of tolerance (Lockhart et al., 2023). Conversely, malfunctioning oral tolerance is associated with IgE-mediated hypersensitivity reactions to food allergens whose prevalence in industrialized countries has sharply risen over the past thirty years (Branum & Lukacs, 2009; Perkin et al., 2016), as has the prevalence of other chronic diseases (Agache et al., 2022; C. A. Akdis, 2021; C. A. Akdis & Nadeau, 2022). Food allergies entail a high level of morbidity and contribute to mor-  
60 tality worldwide. While different therapeutic interventions have been tested and applied (Durham & Shamji, 2022), many food-related allergic diseases remain incurable (Wood et al., 2024).

Passive immunotherapy with monoclonal IgG antibodies that bind the allergen and prevent IgE engagement are promising candidates to alleviate the disease burden (Orengo et al., 2018; Paolucci et al., 2023; Shamji et al., 2021) and may become an important instrument in treating anaphylactogenic allergies, like  
65 peanut allergy (PA). Although naturally occurring mucosal antigen-specific IgA may not predict tolerance (E. G. Liu et al., 2022), systemic blood-based antibody signatures, assessed in sizeable populations, can reveal immunological mechanisms of natural tolerance and highlight the prevalence of antigen-specific antibodies in certain diseases. Conversely, they can point to their association with absence of disease. We have investigated the antibody repertoires of large collections of patients for the occurrence of autoanti-  
70 bodies against the prion protein (Senatore et al., 2020), the tau microtubule-binding domain (Magalhães et al., 2021), and apoptosis-associated speck-like protein containing a CARD (Losa et al., 2023), and have studied seroepidemiological correlations of immune reactivity to SARS-CoV-2 proteins in 72,250 plasma samples of 54,153 individuals (Emmenegger, De Cecco, et al., 2023).

Here, we have interrogated a large unselected hospital cohort for the presence of IgG against the *Arachis hypogaea* 2 (Ara-h2) protein, the major peanut allergen (Fig. **1A**, overview). We identified 127 seropositive  
75 among 24,536 patients, of which about 0.2% (n=45) suffered from documented peanut allergy. An exploratory data-driven analysis indicated a strong association with cystic fibrosis (CF), a life-limiting autosomal recessive monogenic disorder prevalent in Europe, North America, and Australia (Elborn, 2016; Rowe et al., 2005). We propose that the combination of gut epithelial dysregulation and with the inflammatory


80 milieu typical of persons with CF (pwCF) may trigger enhanced antigen penetration, here Ara-h2, and may elicit an IgG4 dominated B cell response.

## Results

### 85 **High-throughput antibody profiling identifies individuals with IgG against Ara-h2.**

Using an automated high-throughput screening (HTS) platform (**Fig. 1A**), we assayed the presence of IgG antibodies directed against “natural” (i.e. purified from delipidated extract by multi-step chromatography) Ara-h2 (nAra-h2) in 24,536 patients admitted to the University Hospital Zurich (USZ) for a broad variety of reasons. The plasma samples (n=27,239) had been collected for routine diagnostics and patients had con-  
90 sented to their use for biomedical research. Our cohort comprised 48% female (n=11,944) and 52% male (n=12,592) patients with a median age of 54 (IQR: 49-68) years (**Fig. 1B** and **Table 1**). No selection was imposed on this cohort other than being treated as inpatients or outpatients in any of 39 clinical divisions of the University Hospital of Zurich.

The largest collections of samples originated from the Departments of Cardiology, followed by Rheuma-  
95 tology, Internal Medicine, Gastroenterology and Hepatology, Dermatology, and Neurology (**Fig. 1C**). Accordingly, the spectrum of diseases was highly heterogeneous and encompassed 1,532 individual ICD-10



three-letter codes. The five most frequent ICD-10 codes observed were I10 ('Essential (primary) hypertension', n=6,235), Z92 ('Personal history of medical treatment', n=3,437), I25 ('Chronic ischaemic heart disease', n=3,282), Y84 ('Other medical procedures as the cause of abnormal reaction of the patient, or of  
100 later complication, without mention of misadventure at the time of the procedure', n=3,237), and N18 ('Chronic kidney disease', n=3,095), as annotated in **Fig. 1D**.

Approximately 25% of individuals (n=6,280) had some indication of having self-anamnestic, suspected, confirmed, or non-confirmed allergy, and 0.2% (n=45) reported peanut allergy, which reflects the approximately expected frequency of clinically diagnosed peanut allergy in adult European populations (Kotz et  
105 al., 2011; Nwaru et al., 2014; Spolidoro et al., 2023). Other common allergies and food intolerances had the following roughly estimated prevalence in the dataset: nut allergy without further specification (0.4%, n=103), lactose intolerance (0.7%, n=163), fish allergy (0.2%, n=44), pollen allergy (1.5%, n=377), house dust mite allergy (0.6%, n=155), cat hair allergy (0.4%, n=103), dog hair allergy (0.1%, n=19), penicillin allergy (4.9%, n=1197), paracetamol allergy (0.5%, n=126), and nickel allergy (0.8%, n=190). The most frequently prescribed medications (from a total n=5,854 medications listed) were paracetamol (n=11,409),  
110 metamizole (n=8,992), esomeprazole (n=6,338), pantoprazole (n=5,874), and cholecalciferol (n=5,254). The sampling for the screen occurred during a period from August 2016 to December 2019 and the primary HTS was conducted over the same time span with a break between March 2018 and October 2019.

For serological analyses, plasma samples were serially diluted using a twofold geometric progression  
115 (range: 1:50-1:6,000) in high-protein binding 1536-well plates using contactless acoustic dispensing. Optical densities (ODs) were recorded as reported (Emmenegger, De Cecco, et al., 2023) and logistic regression curves were derived. Antibody titers were defined as the inverse decadic logarithm of the dilution at the inflection point of the sigmoid ( $-\log_{10}(EC_{50})$ , abbreviated as  $p(EC_{50})$ ) (**Fig 1E**).  $p(EC_{50})$  values ( $\geq 1.7$ ) aligned well with the averaged ODs at the three highest concentrations (Pearson correlation coefficient  $R = 0.90$ ,  
120 **Fig. S1A**). We visualized individual data in half-violin plots (**Fig 1F**). Seropositive samples were defined as having  $p(EC_{50})$  values  $\geq 2$ , corresponding to half-maximal binding at a 1:100 plasma dilution with mean squared residual error of the logistic regression  $< 20\%$  of the  $p(EC_{50})$  value (black dots in **Fig. 1F**) based on cut-offs previously defined (Senatore et al., 2020). Values  $< 2$  were regarded as seronegative. Thus, we detected 133 (0.49%) seropositive samples from 127 seropositive individuals.


### 125 **Association between antibodies against nAra-h2 and Cystic Fibrosis.**

Seropositive samples showed a balanced proportion of females (n=53) and males (n=74), similarly to seronegative samples ( $\chi^2$ ,  $p = 0.238$ , **Fig. 2A**). The median age of seropositives (45, IQR: 32-58.5 years) was significantly lower than that of seronegatives (54, IQR: 49-68 years; Mood's median test for age:  $p < 0.00001$ , **Fig. 2B**). The largest fraction of the seropositive patients came from Pulmonology/Pneu-

130 monology, Dermatology, or Gastroenterology and Hepatology, with a five-fold increase seen for Pulmo-  
 nology ( $p < 0.0001$ , Fisher's exact test, **Fig. 2C**). None of the other clinical divisions displayed a statistically  
 significant enrichment of seropositive patients.

We then asked whether the seropositive patients possessed distinguishing features on a global scale. We  
 employed Uniform Manifold Approximation and Projection (UMAP) and Principal Component Analysis  
 135 (PCA) to project the maximum variability contained in the dataset, containing 1,532 ICD-10 codes and sex  
 information from 23,035 unique individuals and 25,597 samples. Neither UMAP (**Fig. 2D**), using a cosine  
 metric to account for binary data, nor PCA (**Fig. 2E**) revealed any markedly distinct clusters between sero-  
 positive and seronegative samples.

To better understand the features characteristic of the seropositive samples, we collected ICD-10 codes  
 140 occurring at least 5 times (i.e. in  $> 3.75\%$ ) in the seropositive group and conducted  $\chi^2$  statistical testing ( $p =$   
 $0.01$ ) applying Bonferroni's correction for multiple comparisons. We obtained a table of 131 ICD-10 codes,  
 36 of which displayed statistical significance (**Supplemental File 01**). The most enriched ICD-10 code, E84  
 (Bonferroni-corrected  $p < 10^{-10}$ ), encodes CF, the most common autosomal recessive monogenic disorder



**Fig. 2. Association between IgG to Ara-h2, CF, and CF comorbidities.** **A.** Sex distribution in Ara-h2 seropositive and seronegative individuals. *Ns*:  $\chi^2$  non-significant. **B.** Age of Ara-h2 seropositive and seronegative females and males. Mood's median test for age was significant ( $p < 0.00001$ ). **C.** Contributions of clinical departments to seropositive individuals (%). Enrichment of seropositives for pneumonology ( $p < 0.0001$ , Fisher's exact test). **D-E.** UMAP (**D**) and PCA (**E**) analysis of ICD-10 codes and sex showing similar clusters of seropositive and negative fractions. **F.** Top 20 enriched ICD-10 codes, all statistically significant ( $\chi^2 p < 0.0001$  adjusted for multiple comparisons). **G.** Multiple logistic regression analysis using Bayesian priors (U: unregularized. L: LASSO. RH: Regularised horseshoe) showing the ten top positive and top negative odds ratios for ICD-10 codes, age, and sex. **H.** Top 20 significantly enriched medications ( $\chi^2, p < 0.00001$  adjusted for multiple comparisons). **I.** Multiple logistic regression analysis using Bayesian priors (U: unregularized. L: LASSO. RH: Regularised horseshoe) showing the ten top positive and negative odds ratios for medication, age, and sex. **J.** UMAP analysis using all ICD-10 codes, medication and sex for all seropositive individuals. **K.** PCA of the data in (**J**) with annotation of important features.

145 in the Caucasian population. CF was followed by ‘diseases of the gallbladder, biliary tract, and of the pancreas, with diseases elsewhere classified’ (K87, Bonferroni-corrected  $p < 10^{-10}$ ), ‘osteoporosis, with disease elsewhere classified’ (M82, Bonferroni-corrected  $p < 10^{-10}$ ), ‘other diseases of the pancreas’ (K86, Bonferroni-corrected  $p < 10^{-10}$ ), ‘other form of diabetes mellitus’ (K77, Bonferroni-corrected  $p < 10^{-10}$ ), ‘chronic sinusitis’ (J32, Bonferroni-corrected  $p < 10^{-10}$ ), and ‘acute infection of the lower respiratory tract’ (J22, Bonferroni-corrected  $p < 10^{-10}$ ), among others (**Fig. 2F**). As many of these codes appeared to be related to CF, we removed patients with ICD-10 code E84 from the dataset and repeated the analysis. Although some ICD-10 codes fulfilled the criteria of being shared by  $\geq 5$  seropositive individuals, only one of the ICD-10 codes reached statistical significance (L70, ‘Acne’, see **Supplemental File 02**) whereas all previously significantly enriched codes disappeared. This confirms that ICD code E84 dominated the dataset, with multiple other ICD-10 codes co-segregating with CF. Indeed, diabetes (Gibson-Corley et al., 2016; Kayani et al., 2018), osteoporosis (Gibbens et al., 1988; Hecker & Aris, 2004), chronic sinusitis (Davidson et al., 1995; Okafor et al., 2020; Ramsey & Richardson, 1992) and infections (Cosgriff et al., 2020; Lyczak et al., 2002) are known highly prevalent comorbidities of CF (Bell et al., 2020; Cutting, 2015; Elborn, 2016) expected to be enriched within this patient group.

160 The threshold of  $p(\text{EC}_{50}) \geq 2$  is arbitrary and results in binarization of the dataset (seropositive and seronegative patients). We therefore repeated the analysis on a continuous scale, using the entire spectrum of  $p(\text{EC}_{50})$  values (**Fig. 1F** and **Fig. S1B** for qq-plot). We conducted a regression on logit-transformed  $\text{EC}_{50}$  values (denominated as  $-\text{logit}(\text{EC}_{50})$ ) in a Bayesian framework using an uninformative (normal(0,10) and two shrinkage priors: LASSO (Gelman et al., 2013) and a regularized horseshoe (Piiironen & Vehtari, 2017). We then plotted the odds ratio and the 95% credible intervals of the 10 top-most positively and negatively associated features (**Fig. 2G**). The ICD-10 code for CF (E84) was the only parameter unambiguously correlating with high  $-\text{logit}(\text{EC}_{50})$  values, corroborating results presented above (odds ratio with horseshoe prior: 2.66, 95% credible interval: 2.18-3.14). E44 (“Protein-energy malnutrition of moderate and mild degree”) displayed a slightly increased odds ratio, followed by ICD-10 codes H26 (“Other cataract”), and A49 (“Bacterial infection of unspecified site”). Among the ICD-10 codes that were found to be most negatively associated are HIV-related codes B20 (“Human immunodeficiency virus [HIV] disease resulting in infectious and parasitic diseases”) and U60 (“Clinical category of HIV disease”). Female sex and age were slightly negatively associated with  $-\text{logit}(\text{EC}_{50})$  values, but the decrease in odds ratio was minimal (with horseshoe prior: 0.99, 95% credible interval: 0.97-1.00) and the associations are most likely spurious. Odds ratios for all ICD-10 codes are listed in **Supplemental File 03**.

175 Next, we used the same  $\chi^2$ -based approach to correlate the active principles of medication prescribed to the patients. Of 114 active ingredients appearing at least 5 times in the seropositive fraction, 45 were statistically significant after Bonferroni correction (**Supplemental File 04**). The following medications were




associated with Bonferroni-corrected  $p < 10^{-10}$ : Colistin (last-resort antibiotic medication for multidrug-resistant Gram-negative bacteria), Tazobactam, Piperacillin, Meropenem and Teicoplanin (against Gram-positive bacteria including methicillin-resistant *Staphylococcus aureus*), antifungal compounds like Itraconazole and Amphotericin, digestive enzymes, and Tacrolimus (immunosuppressant,  $p < 10^{-6}$ ) (**Fig. 2H**). All these drugs are typically used for the treatment of CF, and indeed the removal from the dataset of pwCF led again to the disappearance of any significantly enriched medication. The application of afore Bayesian multiple regression model identified candidates similar to those found with  $\chi^2$  statistics (**Fig. 2I**) including Colistin (odds ratio with horseshoe prior: 2.19, 95% credible interval: 1.63-2.87), phosphates, Carvedilol and Piperacillin. The HIV medication Emtricitabine was among the most negatively associated features, in line with ICD-10 codes U60 and B20 (**Fig. 2G, Supplemental File 05**). As a limitation, the regression model – unlike the  $\chi^2$  analysis – was restricted to active ingredients assigned to at least 100 patients in the data set to increase the computational robustness.

The seropositive collective ( $n=127$ ) with  $p(EC_{50}) \geq 2$  consisted of 31 pwCF and 89 with other diseases but no record of CF. Seven patients for whom no information was available were excluded from the analysis. Using ICD-10 codes, active ingredients, and sex as features, we clustered the seropositives, with ICD-10 code E84 as classifier. UMAP and PCA (**Fig. 2J-K**) revealed a distinct cluster formed by pwCF with characteristic ICD-10 codes (e.g. D90, E44, E84, J15, Z94) and medications (e.g. Meropenem, Amphotericin, Domperidone, Piperacillin, Colistin) aligning along the axis of the first principal component (**Fig. 2K**).

In summary, we found an association between IgG antibodies against the major peanut allergen nAra-h2 and CF, but not with any other disease (**Table 2**). Similar studies performed with the cellular prion protein, PrP<sup>c</sup> (Senatore et al., 2020) and with calcitonin-gene related peptide (CGRP), in part with a largely overlapping collection of samples, did not indicate any association with CF. The seropositives did not overlap with PrP<sup>c</sup> or CGRP reactivity (**Fig. S1C**) except for one individual seropositive for both nAra-h2 and PrP<sup>c</sup>. Prior reports do not suggest a higher prevalence of peanut allergy or other food allergies in pwCF (Lucarelli et al., 1994), and additional data on allergies indicated that only two of the seropositive pwCF had an anamnestically reported peanut allergy (1.6%) whereas penicillin allergy was reported in twelve seropositive individuals (9.5%).

205 **Post hoc clinical profiling does not distinguish between seropositive and seronegative pwCF.**

First, we investigated whether ICD-10 codes of pwCF with antibodies against nAra-h2 (n=31) differ from those without (n=88). Neither hierarchical clustering (**Fig. 3A**), PCA (**Fig. 3B**), or UMAP (**Fig. 3C**) detected differences in the assigned codes. Also age (p: 0.8217, Fisher's exact test) and sex (p: 0.7187, Wilcoxon rank sum test) were equally distributed in pwCF with and without anti-Ara-h2 antibodies (**Fig. 3D-E** and 210 **Table 1**). The seropositive and seronegative pwCF did not significantly differ in CRP as well as leukocyte and lymphocyte counts (Wilcoxon rank sum test with Benjamini-Hochberg correction, **Fig. 3F**) suggesting that disease exacerbation was no decisive factor. Because forced expiratory volumes in the first second FEV1 scores – an important feature of lung function – were not available in our dataset, we could therefore not classify lung disease severity. We therefore investigated selected features of CF pathology (Bruscia & 215 Bonfield, 2016, 2022; De Lisle & Borowitz, 2013; Dorsey & Gonska, 2017; Elborn, 2016; Garg & Ooi, 2017; Nichols & Chmiel, 2015; Ooi & Durie, 2016; Roesch et al., 2018). The distributions of Ara-h2 antibody-positive and negative pwCF did not differ regarding the presence of a lung transplant (p = 0.6185, Fisher's exact test), infection with *Pseudomonas aeruginosa* (p = 0.412, Fisher's exact test), pancreatic insufficiency (p = 0.5727, Fisher's exact test), or aspergillosis (p = 0.7994, Fisher's exact test, **Fig. 3G**).



220 For 19 pwCF, we could measure two blood samples taken at different time points at least 104 days apart. The resulting  $p(EC_{50})$  values between these two samples were highly replicable (Pearson correlation coefficient  $R = 0.93$ , **Fig. S1D**), suggesting that the presence of such IgG antibodies is a stable phenomenon. We then plotted the temporal evolution of  $p(EC_{50})$  reactivity in these matched samples (**Fig. 3H**). Although the number of samples available was limited, the anti-Arah h 2 antibody seropositive individuals with CF stayed  
225 mostly seropositive, and the seronegatives stayed seronegative over the period investigated (up to 3 years). Hence, using the parameters at hand, we were unable to cluster seropositive and seronegative pwCF based on salient features of CF pathology.


### ***Antibody isotypes, subclasses, and affinity of antibodies against nAra-h2.***

Immune responses to food allergens in allergic patients are dominated by Th2-dependent priming of B  
230 cells into IgE-secreting plasma cells (Rogosch et al., 2010) which disrupts the tolerogenic local IgA-driven response (Berin, 2012; Tordesillas et al., 2017). However, patients with peanut allergy can develop both IgE and IgG antibodies against peanut allergens (El-Khouly et al., 2007; McKendry et al., 2021; Tay et al., 2007). We therefore characterized the antibody response to nAra-h2 in selected seropositive pwCF of whom we were able to obtain a large blood sample by means of patient recontacting. Peanut-allergic pa-  
235 tients and healthy donors served as controls.

First, we conducted an IgG ELISA, with nAra-h2 coated to microplates, to re-evaluate the results obtained in the HTS. While pwCF displayed high  $p(EC_{50})$  values against nAra-h2, peanut-allergic patients were found to only have moderate IgG titers, and no reactivity was found in healthy controls (**Fig. 4A**). One patient with CF (CF5), whose previous sample showed a high titer against Ara-h2, did not show anti-Ara-h2 reac-  
240 tivity above threshold. Serum IgA against nAra-h2 was detectable in only two pwCF and in one patient with peanut allergy (**Fig. 4A**). To determine nAra-h2-specific serum IgE antibodies, we used the commercial ImmunoCAP technology, an automated sandwich fluorescence immunoassay, with whole peanut extract as antigen (**Fig. 4B**). The peanut-allergic patients, but none of the pwCF or patient controls, displayed high levels of IgE antibodies against peanut extract, with some of them exceeding  $100 \text{ kU}_A/\text{l}$  and all falling into  
245 CAP class 3 or higher, indicating at least moderate and mostly high allergen sensitisation. We then investigated IgG subclasses using a panel of previously validated secondary antibodies (Emmenegger et al., 2022) in the same patients. Both IgG1 and IgG2, but not IgG3, showed minor contributions to the overall IgG response in pwCF, whereas the strength of the response was governed by the IgG4 subclass (**Fig. 4C-D**), which is remarkable as this phenotype is usually observed after allergic desensitization (M. Akdis &  
250 Akdis, 2014; Couroux et al., 2019).

Next, we characterized samples by Western blotting. Strips containing a mix of Ara-h2 and BSA were incubated with plasma samples at a dilution of 1:100, optionally pre-incubated with soluble nAra-h2 for com-

petition. PwCF, but no controls, showed the expected signal for nAra-h2 at 17-19 kDa in the non-preincubated condition. The signal almost entirely disappeared when the sample was pre-incubated with soluble nAra-h2, except for CF patient #2 (Fig. 4E), suggesting high-affinity interactions at low antibody concentrations.




**Fig. 4. Analysis of the antibody response against Ara-h2 in PwCF and in controls.** A. Analysis of the IgG and IgA response against Ara-h2 in six pwCF, three control patients (CT), and five peanut allergy (PA) patients. B. Analysis of the IgE response against peanut extract using ImmunoCAP. C. Analysis of IgG subclasses. D. Representation of data shown in (C) as boxplot with individual dots. E. Competitive Western Blot analysis. In each lane, an electrophoresis with nAra-h2 and BSA was conducted.

To characterize the affinity of antibodies against nAra-h2, we used microfluidic diffusional sizing (MDS) (Emmenegger, De Cecco, et al., 2021; Emmenegger, Worth, et al., 2023; Schneider et al., 2022, 2023) on purified IgG fractions of plasma samples. The affinity and concentration profiles were determined for all pwCF except CF5 which did not display an increase in hydrodynamic radius even after concentrating IgG. For the three samples for which a maximum likelihood value for the affinity was measurable, the measured affinity ranges were 0.75-3.93 nM, and the concentrations in undiluted serum were 0.02-0.1  $\mu$ M (Fig. 4F). One sample (CF2) did not give a maximum likelihood value for the affinity, but the K<sub>d</sub> was below 7.69 nM. We conclude that the anti-Ara-h2 IgG antibodies contained in the plasma of pwCF neutralize Ara-h2 with high affinity *in vitro*.

***pwCF develop IgG against certain food allergens but not pollen allergens.***

Next, we investigated whether pwCF harbour antibodies against multiple food allergens beyond Ara-h2. We selected 177 pwCF samples (median age 34 (IQR: 26-42) years, female:male ratio = 51:49) from our biobank and 76 samples from paediatric pwCF (median age 13 (IQR: 9-17) years, female:male ratio = 45:55) from a geographically distant locale. Thus, we introduced a validation cohort. For controls, we randomly selected 319 samples from hospital patients (median age 54 (IQR: 39-67) years, female:male ratio = 48:52),



**Fig. 5. pwCF have IgG against food allergens but not pollen allergens. A-B.** Heatmap (A) and percent positives (B) of IgG responses against select allergens and control proteins in adult (CF) and paediatric (PCF) pwCF, non-CF patients (PC) and healthy blood donors Fisher's exact test. Detailed statistics in **Supplemental File 06**. **C.** Two-dimensional representation of data from (A) after PCA. Right side: violin plot displaying the distributional differences of the second PCA dimension according to patient groups. **D.** Circularized unsupervised hierarchical clustering suggests that the dataset (from A) can be classified into two main groups, shown in green (predominantly adult or paediatric pwCF) and red (mostly non-CF controls). **E.** IgG4 (top) and IgE (bottom) reactivity in large allergen panel in subset of pwCF and controls, assessed using the ISAC technology.

and from 358 healthy donors (median age 44 (IQR: 31-55) years, female:male ratio = 60:40) from the blood donation services of the Canton of Zurich (**Table 1**). These samples were tested for the presence of IgG antibodies against some of the common food allergens, here nAra-h1, nAra-h2, nAra-h3, nAra-h6, *Corylus avellana* (Cor-a) 9, *Glycine max* (Gly-m) 5, Gly-m 6, *Daucus carota* (Dau-c) 1, *Malus domestica* (Mal-d) 1, *Bos domesticus* (Bos-d) 11, *Betula verrucosa* (Bet-v) 1, and *Lolium perenne* (Lol-p) 1 using our HTS platform. As further controls, we used LAG3 extracellular domain containing a his-tag (Emmenegger, De Cecco, et al., 2021) and an uncoated condition to account for spurious binding to the plastic of the plate. The respective p(EC<sub>50</sub>) values were plotted as heatmap (**Fig. 5A**). The IgG reactivity of pwCF, but not of controls, against nAra-h2 was confirmed. Multiple additional epitopes displayed binding, including Ara h proteins, nGly-m5, nGly-m6, and nCor-a9. Using a cut-off p(EC<sub>50</sub>) value of 2, we classified individuals as seropositive or negative, and plotted the positive samples by condition (**Fig. 5B**). Among those showing consistently significant changes in adult and pediatric pwCF (**Fig. 5B, Supplemental File 06**), 4.5-10.5% of pwCF but only 0.6-1.4% of patient controls or healthy donors had antibodies against nAra-h1 (p: < 0.0365, Fisher's exact test), 19.7-19.8% of pwCF but only 0-0.3% of controls had antibodies against nAra-h2 (p: < 0.0001, Fisher's exact test), 26-31.6% of pwCF and only 1.3-2.8% of controls had antibodies against nAra-h3 (p: < 0.0001, Fisher's exact test), 21.1-23.2% of pwCF and 0% of controls had antibodies against nAra-h6 (p: < 0.0001, Fisher's exact test), 11.8-16.9% of pwCF and 0.6-1.1% of controls had antibodies against nGly-m6 (p: < 0.0001, Fisher's exact test). Importantly, IgG reactivity against the two pollen allergens was almost entirely absent in adult or pediatric pwCF (p: > 0.999 for nBet-v1) or lower (adult) or equal (pediatric) to controls (for nLol-p1), indicating that allergens are not generically targeted by IgG antibodies in CF.

The IgG profiles of pediatric and adult pwCF were similar (**Fig. 5A-B**) with slight differences for nCor-a9, nGly-m5, and nLol-p1. A PCA-based cluster analysis performed on the p(EC<sub>50</sub>) values on all patients and antigens indicated that Ara-h2, Ara-h3, and Ara-h6 built a cluster mainly driven by adult (purple triangles) and pediatric (lila squares) pwCF (**Fig. 5C**). The patient controls (blue crosses) and the healthy donors (brown dots) formed two overlapping yet partially distinct clouds. We performed circularised hierarchical clustering on the same data set and colored the two main branches of the dendrogram (red and green, respectively), aiming to find whether pwCF and controls fall into either of these two main branches (**Fig. 5D**). The distribution of pwCF was shifted towards cluster 1 (green, 61% adults and 37% paediatric pwCF) much more than the distribution of the controls (3% patient controls and 9% healthy controls), corroborating the PCA-based visualization.

We then investigated the allergen profiles with the ImmunoCAP ISAC technology, incorporating an array of 112 allergens from 48 sources, including food (oral), pollen (airway), venom, and contact allergens (Jakob et al., 2015; Matricardi et al., 2016). IgG4 and IgE antibodies were measured in a subset of pwCF and controls. Using a standardized cutoff of 0.3 international standard units (ISU), we observed that pwCF

310 carry IgG4, but not IgE, antibodies against Ara-h2 and Ara-h6 (**Fig. 5E**), as shown with ELISA (**Fig. 5A-B**). In the same patients, there was no reactivity with allergens entering through the airways, an observation we substantiated with raw data where no cutoff has been applied (**Fig. S2** and **Supplemental File 07**). This is opposed to allergic patients displaying IgE reactivity against a multitude of airway, contact, food, or venom allergens. While the cutoff-based representation draws a clear picture due to its stringency (**Fig. 5E**), the raw value map indicates that more food allergens may be targeted by IgG4 in CF (**Fig. S2**), confirming our ELISA data.

315 We conclude that a subset of pwCF develops selective immunity to food proteins, in particular peanut allergens Ara-h2 and Ara-h6, at much higher levels than controls. Importantly, our observation that healthy controls harbor serum IgG against multiple food proteins expands the finding that certain food antigens elicited systemic IgG in up to 50% of individuals (Leviatan et al., 2022).

***Lack of evidence for cross-reactive antibodies and absence of broad reactivity against endogenous antigens in pwCF.***


320 We addressed the possibility that reactivity is governed by cross-reactive antibodies primarily matured to neutralize infectious pathogens. A protein blast using BLOSUM-62 matrix suggested various homologies of Ara-h2 with seed storage proteins of related plant species (e.g. with Ara d 2 of *Arachis duranensis*, Ara h 6 of *Arachis hypogaea*, with related conglutinins of *Lupinus albus* and *angustifolius*, or with albumin of *Glycine max* or *Punica granatum*, see **Supplemental File 08**), and previous structural investigations have not revealed similarities beyond the kingdom of plants (Mueller et al., 2011).

325 We further investigated the immune profile in pwCF by Rapid Extracellular Antigen Profiling (Wang, Dai, et al., 2021). Using samples from adult and pediatric pwCF and of a range of controls, we subjected yeast cells transfected with an array of 6,184 plasmids encoding transmembrane and extracellular proteins (**Supplemental File 09**) to the purified IgG fraction of the above patients and calculated the fold-change in detection, suggestive for antibody binding. We then selected proteins with a log<sub>2</sub>-fold increase of  $\geq 2$  in at least one sample ( $n = 293$ , **Fig. 6A**) and excluded proteins upregulated (presumably nonspecifically) in all conditions ( $n = 53$ , see **Supplemental File 09**). No striking patterns among the different patient conditions emerged and no significant differences in the number of autoreactive antibodies in pwCF and controls was observed (one-way ANOVA, **Fig. S3**). A PCA analysis of the above dataset showed no clusters in adult and paediatric pwCF (green) as well as controls (yellow) (**Fig. 6B**), although some adult and paediatric pwCF showed a tendency towards higher PC1 and negative PC2 values. A separate feature cluster was formed by a group of alpha interferons (e.g. IFNA8, IFNA4) that appeared as positive in two control patients but not in pwCF (**Fig. 6A-B**), with IFNAs being important contributors to the second PC (blue dots) as measured by the squared cosine ( $\cos^2$ ) function (Abdi & Williams, 2010). Conversely, the dataset is suggestive of an

330

335

enrichment of autoantibodies targeting the type III interferons IFNL2/IL28A and IL28B, important contributors to the first PC, in some pwCF. To analyse these patterns using an alternative approach, we mapped all data (n = 6,184), including the nonspecifically upregulated ones, in a group-wise fashion, displaying the difference of the log<sub>2</sub>-fold change of pwCF and controls as a function of the average log<sub>2</sub>-fold-change in all individuals. All but five proteins are within 2 log-fold changes (from -1 to +1), suggesting no aberrant autoreactivity in pwCF compared to controls (**Fig. 6C**). IL28A, IL28B, and IL29 appeared to be enriched in pwCF, indicating that autoreactivity against type III interferons might be increased. This data suggests a normal autoimmune profile in CF, judged by the protein antigens that were employed with this technology, as opposed to previous investigations on systemic lupus erythematosus (Wang, Dai, et al., 2021) or COVID-19 (Wang, Mao, et al., 2021). On the other hand, CF has been associated with elevated IgA autoantibodies





350 against double-stranded DNA (Yadav et al., 2020) and with anti-PAD4 autoantibodies (Linnemann et al., 2022).

***No marked changes in B cell receptor repertoires of pwCF, peanut allergic, and healthy donors.***

To investigate the entire antibody repertoire in an unbiased manner, we performed adaptive immune receptor repertoire (AIRR) sequencing (Miho et al., 2018; Rubelt et al., 2017) in anti-Ara-h2 seropositive pwCF (n = 6), patients with peanut allergy (n = 3), and in healthy donors (n = 3). Sequencing of the variable heavy (VH) and variable light (VL) BCR repertoire from each subject produced >10<sup>6</sup> raw reads for most samples (Fig. S4A). After quality control and processing, a high-fidelity data set consisting of unique error-corrected IgM, IgD, IgG, IgA, and IgE sequences and unique error-corrected IgL and IgK sequences was generated (Fig. S4B-C). The pwCF showed the highest percentages of IgA sequences among the total VH repertoire (44.5%, with IgG contribution of 28.1%), whereas the VH repertoire of the peanut allergy group was composed of predominantly IgG (38.6%), IgA (29.1%) and to a lesser extent, IgD (5.4%). The peanut allergy group isotype repertoire contained some IgE sequences (0.23% on average), while it was almost completely absent in pwCF (0.023% on average). IgM was the largest component of the healthy VH repertoire (41.0%) while IgG (29.3%) and IgA (19.8%) followed. None of the distributions showed significant differences between patient groups (pairwise t-test). In line with comparisons among isotypes, the heavy and light chain distributions were not significantly different among the three patient groups (Fig. S4C, pairwise t-test). As expected, VH3 family usage was more pronounced for all the Ig isotypes and patient groups, followed by VH4, VH1, with hardly any contribution by VH2 and VH5. Statistical comparisons between patient groups did not reveal any significant difference (Fig. S4D).

370 We analyzed the abundance of antibody variable domains to test whether the different isotype compartments in CF or peanut allergy resulted from clonal expansion. Sequences that differ only by isotype were clustered into groups consisting of expanded clones, where the size of each clone is determined by the number of sequence members (Vander Heiden et al., 2017). The largest clone size in the CF group reached 6.18% (95% CI: 1.61-10.7%) for IGHG (Fig. 7A), suggestive of an active immune response (Laserson et al., 2014). Clone size in the IGHA, IGHD, and IGHM compartments of all patient groups did not exceed 5% of the B cell repertoire. IGHE clones were expanded in peanut-allergic patients (but not in any other group), exceeding 20% of the total BCR repertoire.

By calculating the variance of the abundance of the clones we also estimated the evenness of the CF, peanut allergy, and healthy donor groups. Evenness serves as a measure of clonal dominance; samples with low evenness would have a less uniform distribution and would therefore be characterized by fewer but highly expressed clonotypes, whereas samples with similarly abundant clones would have higher evenness. The evenness was inferred from the Hill diversity index (Hill, 1973) calculated over a range of diversity orders (q) ranging from 0 to 4, in equally spaced increments, for each subset (Fig. S5A). The evenness score


tended to be slightly higher in CF than in the two other groups for IGHG, indicating that expansion was  
385 more diverse and not bound to few clones (**Fig. 7B**). However, none of the group-wise comparisons  
reached statistical significance, suggesting that there were no profound differences in the BCR repertoire  
between the three groups.

To determine whether somatic hypermutation (SHM) is perturbed in CF, we analyzed somatic mutation  
patterns of IGHV gene sequences. SHM is known to be biased toward DNA motifs prone to mutations  
390 (hotspots). Mutation frequency for each sequence was calculated as the number of base changes from  
germline in the V region. Mutational frequency displayed isotype-specific differences. As expected, the  
mutational frequency was high for the class-switched IGHA and IGHG but comparatively low for the non-  
class-switched IGHM and IGHD. IGHE appeared at medium mutational frequency; however, the number  
of sequences were low (**Fig. 7C** and **Fig. S5B**). Group-wise comparisons among the IGH isotypes did not  
395 reveal any significant difference, suggesting that pwCF do not have a broadly altered rate of SHM.

In summary, AIRR sequencing revealed that the antibody repertoire in pwCF does not substantially deviate  
from those observed in healthy controls or patients with peanut allergies with regard to isotype composi-  
tion, clonal expansion, clonal dominance, and SHM. This finding further highlights the specificity of the IgG  
response against peanut allergens identified in pwCF in the absence of global alterations of the BCR rep-  
400 ertoire. In the light of the many chronic conditions connected to CF, including the pwCF investigated in this  
study, e.g. chronic infections and immune suppression, the scarcity of difference in the BCR repertoire is  
surprising.

**Absence of CDR3 convergence between sequences obtained via BCR repertoire sequencing of pwCF and publicly available clones known to bind Ara-h2.**

405 Since peanut allergy and pwCF both have antibodies targeting peanut allergens, we hypothesized that there might be common clones with identical or highly similar CDR3 regions that diverged from the germline sequence. Common clonal groups have been found among unrelated peanut-allergic patients that have been treated with oral immunotherapy (OIT) (Kiyotani et al., 2018; Patil et al., 2015). Additionally, common clones of the IgE isotype were also detected in untreated peanut-allergic patients with identical gene rearrangements within VH and VL chains (Croote et al., 2018). Thus, we aligned the heavy and light chain CDR3 regions of our cohorts to publicly available sequences known to recognize Ara-h2 retrieved from several studies (Aguzzi et al., 2020; Croote et al., 2018; Hoh et al., 2016). We identified a small number of convergent antibody responses across all disease and control groups that did not exceed 0.2% of the heavy chain CDR3 sequences (Fig. S6). As expected, light chain convergence was higher reaching 415 15% of the CDR3 kappa chain and no more than 10% of the CDR3 lambda chain across all groups since light chain heterogeneity is low in the general population (Collins & Watson, 2018). There was no evidence for convergent antibody responses in the BCR repertoire in pwCF and peanut allergy.



### ***T cells are activated with Ara-h2 peptides.***

Several studies did not show a clear association between peanut allergy and HLA (Dreskin et al., 2010; Shreffler et al., 2006) whereas others showed a higher frequency of DR8 (Boehncke et al., 1998; Howell et al., 1998). In addition, the HLA-DQ gene region, especially DQ6, was linked to peanut allergy in two comprehensive studies (Hong et al., 2015; Madore et al., 2013). HLA studies performed on pwCF revealed increased frequencies of DR4 and DR7 alleles compared to controls (Aron et al., 2012), however, findings were not reproduced in a large study (Adriaanse et al., 2014). Here, we characterized the human leukocyte antigen (HLA) class II locus of pwCF with IgG reactivity to peanut allergens. This approach was governed by the assumption that the high-affinity B cell responses observed in this collective (**Fig. 4**) require T cell help. The identification of T cell epitopes may provide further explanation for the mechanism by which some pwCF develop an adaptive immune response to certain food allergens.


We genotyped HLA-class II alleles DRA\*, DRB1\*, DRB3\*, DRB4\*, DRB5\*, DQA1\* and DQB1\*, DPA1\*, and DPB1\* in pwCF 1-6 (for summary of results, see **Supplemental File 10**). Certain HLA-DRB1\* and -DRB3\* alleles (DRB1\*03:01:01G in 50%; DRB1\*11:01:01G in 50%; DRB3\*01:01:02G in 67% and DRB3\*02:02:01G) and even more certain DQA\* (DQA1\*01:02:01G in 67% and DQA1\*05:01:01G in 100%) and DQB1\* (DQB1\*02:01:01G in 50%, DQB1\*03:01:01G in 50% and DQB1\*06:02:01G in 50%) and DPA1\* (DPA1\*01:03:01G in 100%; 4/6 patients homozygote for this allele) and DPB1\* alleles (DPB1\*04:02:01G in 50%) were found at much higher frequencies than in the normal European population. Particularly the frequency of HLA-DQA1\*05:01 was elevated 2-6-fold compared to control populations of similar ethnicity (Gonzalez-Galarza et al., 2020)). Moreover, the comparison of the amino acid sequences of HLA-DQA1\*05:01:01 with the DQA1\*01:01:01 allele and other DQA1\* alleles of the six individuals revealed changes in size, charge and hydrophobicity at multiple positions that participate in forming the HLA-DQ binding groove, but also outside (see **Supplemental File 11**), and HLA-DQA1\*05:01:01 differs most from the other alleles (Barker et al., 2023).

Finally, we asked if T-cell reactivity against overlapping peptides of the main peanut allergen Ara-h2, against which the antibodies are directed, provides further insight. We used Fluorospot testing of whole PBMCs with cytokines IL-4 and TNF- $\alpha$  as readout and included PBS as a negative and anti-CD3/CD28 antibodies as well CEFII, a mixture of peptides derived from cytomegalovirus, Epstein Barr Virus and influenza A virus, as positive controls. Two seropositive pwCF, for whom sufficient PBMCs were available, responded to multiple Ara-h2 peptides, with respective peaks at peptides corresponding to amino acids 1-20, 61-80, and 121-140 (**Fig. 7D**). When performing in silico peptide binding predictions of Ara-h2 to the HLA-DQA/B combinations DQA1\*05:02/DQB1\*02:01:01 and DQA1\*05:02/DQB1\*03:01:01 (NetMHCII-

450 pan-4.0; see (Jensen et al., 2018)), no strongly or weakly binding peptide was identified in the Ara-h2 sequences considered important in the cellular assay (see **Supplemental File 12**), and therefore, strong HLA-DQ binding of this region is unlikely.

**CFTR heterozygosity is not associated with peanut allergy or food allergy in general.**

455 Mutations in the CFTR affect not only epithelial cells but also macrophages and lymphocytes (Bruscia & Bonfield, 2016, 2022; Hartl et al., 2012) of heterozygous CFTR mutation carriers (Miller et al., 2020; Polgreen et al., 2018). We wondered if CFTR heterozygous carrier states may be associated with altered antibody responses against peanut and other food allergens. We examined an anonymized collective of confirmed peanut-allergic patients (n=90) for the presence of mutations in the CFTR gene and compared it with the expected frequency in the non-Finnish European population. We restricted the analysis to variants with available information contained in gnomad3.1.1 and additionally displayed data from the CFTR2 database (**Fig. 8A**). The cumulative allele frequency, which correlates with heterozygosity (**Fig. 8B**), was



**Fig. 8. CFTR sequencing in patients with peanut allergy and health survey.** A. Repositories used to analyze and compare the data. B. Cumulative allele frequency correlates with heterozygosity. C. Allele frequencies for all CFTR variants in CFTR-sequenced PA cohort (n=90), and in controls. D. Health surveys assessing the frequency of food allergies in pwCF compared with heterozygous controls in Europa and in North America. E-F. Frequency of mutations (E) and health conditions (F) as indicated by study participants, for mutation carriers and pwCF (logistic regression analysis).

4.0% for peanut-allergic patients, almost identical with the population (4.1%, according to gnomad3.1.1) and similar to the non-Finnish Europeans (3.3%), while the allele frequency of pwCF, based on data from CFTR2, was lower (1.2%). We then displayed the allele frequencies for all variants (**Fig. 8C**). We observed  
465 two significant alterations for variants c.743+40A>G ( $p=0.005$ , t-test) and for c.1585-1G>A ( $p<0.0001$ , t-test). The c.1521\_1523del (p.Phe508del) mutation, which is the most common cause of CF (69%), has an allele frequency of only 1.4% (non-Finnish European population) and 1.1% (peanut-allergic patients).

We then invited individuals with known CFTR status (wildtype, mutation carrier, or pwCF) to participate to an anonymized survey on medical symptoms including allergies (**Fig. 8D**). The survey was distributed online  
470 by the German CF association "[Mukoviszidose e.V.](https://www.mukoviszidose.de/)" and was broadly accessible without restriction. It queried basic demographic information and a panel of comorbidities including allergy (**Supplemental File 13**). 153 individuals participated between July 2022 and January 2023, of which two did not include any information and were excluded. Among the 151 individuals, four (2.6%) indicated to not carry any CFTR mutation. 91 individuals (60.3%) were heterozygous mutation carriers, and 56 individuals (37.1%) reported to  
475 have CF. The median age was 36 (IQR:16.5-43) years, with 62.3% of participants being females (**Table 3**). The majority of carriers (43%) and pwCF (55%) reported a deletion of Phe<sup>508</sup>; some (31%) did not know the details of their sequence while being parent of a child with CF, and 23% of pwCF were trans-heterozygous (**Fig. 8E**).

As the number of non-mutated individuals was too low for a statistical comparison, we directly compared  
480 the prevalence of medical conditions between mutation carriers and pwCF using multivariable logistic regression adjusted for age and sex. There was no significant difference regarding the prevalence of allergy in general (46.2% mutation carriers and 23.2% pwCF self-reported an allergy), or of subgroups thereof (the most frequent allergy type was pollen allergy, and 13.2% mutation carriers and 7.1% pwCF indicated they have some food allergy), between these groups (**Fig. 8F**), consolidating the result from the sequencing-  
485 based approach. The only two conditions significantly different between the two groups were CF (odds ratio: 132.6 (CI95%: 38.1-607.0),  $p < 0.0001$ ) and diabetes mellitus (odds ratio: 32.6 (CI95%: 5.1-645.0),  $p=0.0018$ ). While this survey was conducted primarily in Germany, a registry-based investigation conducted at the Yale New Haven Hospital center in New Haven, USA, resulted in the same conclusions. In  
490 115 individuals (median age 5.0 (IQR: 2.0-15.0), 48.7% of female sex) whereof 56 individuals were CFTR mutation carriers, and 59 individuals were pwCF, the prevalence of food allergy was non-significantly different ( $p=0.345$ , multivariable logistic regression adjusted for age and sex), with 7.1% mutation carriers and 8.5% of pwCF having a clinically diagnosed food allergy (see **Table 4**).


Given the sum of evidence collected, we refute our initial hypothesis that peanut allergy or other food allergies are associated with heterozygosity in CFTR.

495 ***Peanut sensitization in *Cftr*<sup>-/-</sup> mouse model.***

We next assessed whether our finding could be recapitulated in a well-characterized murine model of CF (Snouwaert et al., 1992) which includes intestinal pathology. We speculated that intestinal dysfunction in CF (Collaco et al., 2010; Ooi & Durie, 2016), which includes aberrant mucus release (Garcia et al., 2009; J. Liu et al., 2015) and goblet cell hyperplasia with mucus accumulation in the crypt lumen and along the villous surfaces (Gorrieri et al., 2016; J. Liu et al., 2015; Walker et al., 2022), and increased gut permeability (De Lisle, 2014) combined with a state of chronic infections, may alter the immune reactions triggered by peanut allergens and prevent the development of allergy. Oral anaphylaxis to peanut has previously been shown to be associated with enhanced gut permeability (Gertie et al., 2022). *Cftr*<sup>-/-</sup> and wildtype (WT) mice (both C57BL/6) were orally immunized with ground peanut (5 mg/immunisation, with or without cholera toxin as adjuvant) for six weeks, allowed to recover for ten days, and subjected to an intraperitoneal challenge with 100 µg ground peanut (**Fig. 9A** for overview).

At week 2, modest IgG and IgE responses against crude peanut extract became visible in *Cftr*<sup>-/-</sup> mice and WT control mice (**Fig. 9B**). Antibody levels in cholera toxin-treated *Cftr*<sup>-/-</sup> mice were higher than in WT mice, likely due to an increased antigenic permeability of the intestinal barrier in connection with goblet cell hyperplasia. At week 5 and post challenge, we observed a marked IgE response in WT but not in *Cftr*<sup>-/-</sup> mice ( $p = 0.0019$  and  $0.074$ , respectively, Wilcoxon rank sum test with Holm's correction for multiple comparisons), while the concomitant IgG response was equally strong in both groups (see **Fig. S7A** for statistics). Clinically, mice of both groups reacted with an initial small increase in body temperature post challenge (a possible stress reaction measured also in unchallenged control mice) followed by a sudden drop in body temperature in mice challenged and sensitized using cholera toxin as adjuvant (**Fig. 9C**). The drop in body temperature was delayed, less pronounced, and recovered faster in *Cftr*<sup>-/-</sup> mice. Unchallenged mice or mice whose sensitization was without cholera toxin did not display signs of anaphylaxis (**Fig. 9D**), in contrast to peanut-sensitized and cholera toxin-treated *Cftr*<sup>-/-</sup> mice and WT mice. In contrast, *Cftr*<sup>-/-</sup> displayed a delayed onset of anaphylactic reactions, a reduced peak, and a more pronounced recovery. Thus, *Cftr*<sup>-/-</sup> mice were partially protected from severe allergic reactions. Although the kinetic analyses showed a clear trend, the overall distributions between cholera toxin-induced *Cftr*<sup>-/-</sup> and WT mice did not significantly differ (**Figs. S7B** and **S7C**,  $p = 0.415$  and  $0.0983$ , respectively, Wilcoxon rank sum test).

We then asked whether the cytokine profiles differed between the groups and whether they changed after challenge with peanuts. We speculated that a Th1 environment may skew toward productions of IgG, rather than IgE, which could alter the severity and course of an anaphylactic response. We chose three cytokines, IL-4, IL-10, and IFN- $\gamma$ ; the former a typical Th2 marker, the latter associated with a Th1-based inflammatory response. Naïve *Cftr*<sup>-/-</sup> and WT mice had no detectable IL-4 and IFN- $\gamma$  levels in serum (with one exception) and median IL-10 levels of approximately 3.75 ng/ml (**Fig. 9E**). While serum IL-10 levels at



week 5 and after challenge were close to baseline for both *Cftr*<sup>-/-</sup> and WT mice in the absence of cholera toxin treatment, cholera toxin-treated *Cftr*<sup>-/-</sup> and WT mice had IL-10 concentrations below detection. At week 5 pre-challenge, IL-4 and IFN- $\gamma$  were slightly elevated in cholera toxin-treated *Cftr*<sup>-/-</sup> and WT mice compared with sensitized but non-cholera toxin-treated or non-peanut sensitized mice. Two hours post challenge, the IL-4 concentrations rose approximately 40-fold in peanut-sensitized cholera toxin-treated *Cftr*<sup>-/-</sup> and WT mice, whereas that of IFN- $\gamma$  rose approximately 20-fold in peanut-sensitized cholera toxin-treated *Cftr*<sup>-/-</sup>, but not in WT mice.

In a small human cohort, plasma IFN- $\gamma$  concentrations did not significantly differ between anti-Ara h2-seropositive individuals with CF (n=8), a collection of multimorbid hospital patients (including systemic lupus erythematosus, chronic kidney disease, and COVID-19, named 'mixed', n=26), and six healthy controls (**Fig. S7D**). However, plasma concentrations may only partially reflect local mucosal cytokine levels.



540 Together, there is a distinct cytokine signature in cholera toxin-treated *Cftr*<sup>-/-</sup> mice, visible post intraperi-  
toneal (IP) challenge with peanut extract and characterised by high IFN- $\gamma$ , low IL-10, and high IL-4. While  
IL-10 and IL-4 concentrations matched those in WT mice, high IFN- $\gamma$  was a distinguishing feature at 5 weeks  
immunisation (p=0.0813, Wilcoxon rank sum test) and, most notably, after IP challenge (p=0.00117). This  
545 particular IFN- $\gamma$  signature is remarkable as it has been shown to potently inhibit Th2 and Th17 cytokines  
(Sun et al., 2023), and its tolerogenic role has been described on few occasions (Ca, 2001; Noh & Lee,  
2009).

## Discussion

We conducted a serological investigation to identify IgG against Ara-h2 in a large cohort of hospital patients and consulted the clinical records of individuals showing high titers to identify medical correlates of anti-peanut immunoreactivity. This approach relies on the availability and utilization of surplus samples collected for routine diagnostics, and of data associated with it. The study design is pragmatic (Ioannidis, 2016), it partially circumvents the usual enrolment bias (Grimes & Schulz, 2002), and enables the unconstrained generation of pathophysiological hypotheses. Our screen provided a strong indication that IgG against Ara-h2 are associated with CF. The most parsimonious explanation for this association would have been a higher prevalence of peanut allergy in pwCF, associated with IgE responses (Chen et al., 2021). However, this was not the case. We have therefore set to explore the immunological underpinnings of this association.

We identified high-affinity IgG antibodies against Ara-h2, but also against other peanut allergens and other food constituents, in the absence of IgE and without further indications that the repertoires of autoantibodies or alloantibodies in CF were globally perturbed. We found that such antibodies are affinity-matured due to food constituents rather than cross-reactivity with bacterial or viral proteins. We have additionally found that *CFTR* mutation carriers have the same prevalence of food allergies as pwCF, and these results are in agreement with observations in the population (Branum & Lukacs, 2009; Cianferoni & Spergel, 2009; Sicherer & Sampson, 2018; Soller et al., 2012; Tordesillas et al., 2017). Importantly, many of our findings have been reproduced in a cohort of paediatric pwCF from a different continent, bolstering our confidence in these observations. Lastly, we speculated that CF-associated gut pathology, in conjunction with a permissive cytokine environment, may be causative of the association; we therefore employed a *Cftr*<sup>-/-</sup> mouse model and found, upon allergic sensitisation with peanuts, a signature with low IgE and high IgG levels as well as an altered IFN- $\gamma$  profile that associates with a milder anaphylactic reaction to peanut IP challenge when compared to WT mice.

Oral tolerance provides selective immunity to molecules ingested via the orogastric route in a microbially abundant environment. The IgG response to Ara-h2 was associated with pwCF, but with no other disease entity. While CF-related intestinal pathology is among the most burdensome aspects of the disorder, many other non-CF conditions also affect the gastrointestinal system. None of them displayed an association with IgG against Ara-h2, apart from what is mediated through CF. So how might this link with CF arise? CF is caused by the absence, deficiency or instability of the CFTR protein, an ion channel that maintains the balance of salt and water on many surfaces in the body. pwCF have a multiorgan manifestations leading to progressive lung disease, pancreatic insufficiency, CF-related diabetes, intestinal dysbiosis and inflammation, meconium ileus, intestinal obstruction, biliary cirrhosis and bronchiectasis (Filkins & O'Toole,

2015; Ooi & Durie, 2016). Moreover, there is strong evidence for an altered immune system in CF (Bojanowski et al., 2021; Bruscia & Bonfield, 2016, 2022; Hartl et al., 2012). In this regard, the absence of globally noteworthy aberrances at the level of the immunoglobulin repertoire in CF, shown in our study, might be surprising. We have further shown that CF is not associated with a higher prevalence of food allergies, both compared with *CFTR* mutation carriers and the general population. Consequently, there is an absence of a concomitant IgE response against Ara-h2 or other food ingredients, but an IgG response. The IgG response is dominated by IgG4, reminiscent of the profile observed in allergy patients after desensitization, i.e. following prolonged exposition to an antigen (Emmenegger et al., 2022; Irrgang et al., 2023; Pillai, 2023). Here, CD23<sup>+</sup>IgG1<sup>+</sup> memory B cells poised to switch to IgE, upon activation with IL-4/IL-13, in food allergy (Jiménez-Saiz et al., 2017, 2018; Ota et al., 2023) appear to class-switch from IgG1 to IgG4, as in OIT (Jiménez-Saiz et al., 2018) and thereby lead to desensitization. Thus, one potential extrapolation from our observations could be a lower prevalence of peanut allergy in pwCF.

Exaggeration of defensive mechanisms to prevent ingestion of toxins can result in pathological food allergy (Florsheim et al., 2021). Therefore, to be able to elicit allergic sensitization, dietary proteins must be associated with noxious cues present in the food, as is shown e.g. with cholera toxin experiments performed in our study. A compromised epithelial barrier has been suggested to be involved in causing allergies (C. A. Akdis, 2021; Mattila et al., 2011). Because of lack of *CFTR*, epithelial integrity is compromised in the CF gut (De Lisle, 2014). Additionally, goblet cell hyperplasia is well-documented in CF (Garcia et al., 2009; Gustafsson et al., 2012; J. Liu et al., 2015) and goblet cells assume a pivotal role in supporting epithelial layer integrity and microbiota regulation (Gustafsson & Johansson, 2022; McDole et al., 2012; Weiner et al., 2011), with implications for oral tolerance (Gertie et al., 2022; Mazzini et al., 2014). In this context, pwCF may be more susceptible to epithelial malfunctioning, resulting in an overstimulation of the immune system by oral allergens.

It is extremely intriguing and unexpected that *CFTR* deficiency is associated with tolerogenic rather than anaphylactogenic responses to food allergens. Our peanut allergy sensitisation experiments conducted in wildtype and *Cftr*<sup>-/-</sup> mice indicated an altered serum cytokine profile, with an increase in IFN- $\gamma$  concentration in CF. IFN- $\gamma$ , unlike other IFN types, is selectively produced by T cells to inhibit Th2 T cell responses (Sun et al., 2023). Moreover, CF has also been associated with a predominant proinflammatory type 17 immunity (Price & O'Toole, 2021; Sheikh et al., 2023) and with increased IFN- $\gamma$  concentrations in bronchoalveolar lavage fluid (Tiringer et al., 2013), while a Th2 T helper cell environment is predisposing to allergy (Abbas et al., 2018; Lloyd & Snelgrove, 2018; Tordesillas et al., 2017). However, the systemic cytokine signature does not necessarily reflect local mucosal milieu (Ferreira et al., 2009; Maina et al., 2022; Smith et al., 2021), and differences between the gut and the lung microenvironment are conceivable.

Based on our observations, the combination of a compromised gut epithelium, dysbiosis, and a proinflammatory cytokine milieu may (1) prime the immune system with antigens, in this case with a partially protease-resistant and heat-stable allergen, Ara-h2, in the proximity of noxious substances, and may then (2) poise the B cell response towards IgG4, a tolerance signal. Importantly, while colonization with *Pseudomonas aeruginosa*, which is common in pwCF, has shown to manipulate the host towards a type 2 immune response, including mucus production, to promote nutrient availability (Agaronyan et al., 2022), this immune bias may not generally apply to pwCF but perhaps only to a subset of them (Albon et al., 2023).

### **Limitations of the study**

This study analyzed a heterogeneous unselected cohort, a true excerpt of the population requiring medical aid in a university hospital setting. Its retrospective nature poses challenges owing to incompleteness of records and occasionally lack of relevant medical details. We have only restricted knowledge of whether certain pwCF had received mutation-specific treatment with potentiators like Ivacaftor (Swissmedic approval: 2012) or combinations of Ivacaftor with correctors like Lumacaftor (Swissmedic approval: 2016) or Elaxacaftor and Tezacaftor (Swissmedic approval: 2020), although the time of sampling (75% until end of 2017, 21% in 2018, less than 4% in 2019) suggests that this treatment had not been employed widely. Moreover, the specific mutation was not always available for all pwCF. Similar limitations pertain to the availability of detailed information on patients' allergies in the medical records. Allergies are often self-reported, without being serologically and/or clinically confirmed, and were interpreted with due caution. To offset these limitations, we have utilized complementary methodologies, we have examined multiple cohorts to validate findings, and experiments were conducted at multiple sites. Finally, we have translated an observation on humans onto a model system, a *Cftr*<sup>-/-</sup> mouse. A particularly intriguing prediction derived from the above findings is that transient CFTR inhibition, e.g. in antigen-presenting cells like macrophages or B lymphocytes expressing CFTR (Bruscia & Bonfield, 2022), may quench allergogenic immune responses and might become useful in the context of desensitisation regimens.

## STAR★Methods

### Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
<b>Antibodies</b>		
Goat anti-human IgG (1:4,000 ELISA, 1:5,000 WB)	Jackson	109-035-098; RRID: AB_2337586
Goat anti-human IgA (1:750 ELISA)	Thermo Fisher Scientific	31417; RRID: AB_228253
Mouse anti-human IgG1 (1:3,000, ELISA)	SouthernBiotech	9054-05; RRID: AB_2796627
Mouse anti-human IgG2 (1:3,000, ELISA)	SouthernBiotech	9060-05; RRID: AB_2796633
Mouse anti-human IgG3 (1:3,000, ELISA)	SouthernBiotech	9210-05; RRID: AB_2796699
Mouse anti-human IgG4 (1:3,000, ELISA)	SouthernBiotech	9200-05; RRID: AB_2796691
Anti-mouse IgG	SouthernBiotech	Cat. No.: 1110-05
Anti-mouse IgE	SouthernBiotech	Cat. No.: 1036-05
T cell activation kit (anti-CD3, anti-CD28, anti-CD2 beads)	Miltenyi Biotec	N/A
<b>Biological samples</b>		
University Hospital Zurich unselected patient cohort	N/A	N/A
Paediatric CF patient cohort	N/A	N/A
Healthy blood donors	N/A	N/A
<b>Chemicals, Peptides, and Recombinant Proteins</b>		
Natural Ara-h2	Indoor Biotechnologies	NA-AH2-1
Natural Ara-h1	Indoor Biotechnologies	NA-AH1-1
Natural Ara-h3	Indoor Biotechnologies	NA-AH3-1
Natural Ara-h6	Indoor Biotechnologies	NA-AH6-1
Natural Bos-d11	Indoor Biotechnologies	NA-BD11-1
Natural Cor-a9	Indoor Biotechnologies	NA-CA9-1
Natural Gly-m5	Indoor Biotechnologies	NA-GM5-1
Natural Gly-m6	Indoor Biotechnologies	NA-GM6-1
Natural Bet-v1	Indoor Biotechnologies	NA-BV1-1
Natural Lol-p1	Indoor Biotechnologies	NA-LP1-1
Recombinant Dau-c1.0103	Biomay	Dau c 1
Recombinant Mal-d1.0108	Biomay	Mal d 1
Recombinant human LAG3 <sub>23-450</sub>	Acro Biosystems	LA3-H5222
Recombinant cellular prion protein	In-house	N/A
Calcitonin-gene related peptide (CGRP)	Schafer	Custom-designed
CEFI peptide (T cell activation assay)	N/A	N/A
<b>Critical commercial assays</b>		

ImmunoCAP (IgE)	Thermo Fisher Scientific	N/A
ImmunoCAP ISAC Test (IgG4, IgE)	Thermo Fisher Scientific	N/A
Pierce™ BCA Protein Assay Kit	Thermo Fisher Scientific	23225
Advanta CFTR NGS Library Preparation	FLUIDIGM	PN 101-627- C2
Murine cytokine	Meso Scale Discovery	U-PLEX Personalized multiplexing
IFN gamma Human ELISA Kit, High Sensitivity	Invitrogen	BMS228-2HS
<b>Experimental models: Organisms/strains</b>		
C57BL/6 control mice		
C57BL6-CFTR-KO <sup>Cftr<sup>tm1Unc</sup></sup> ( <i>Cftr<sup>+/+</sup></i> or <i>Cftr<sup>-/-</sup></i> )	The Jackson Laboratory	Strain #002196 backcrossed with C57BL6 mice in house
<b>Software and algorithms</b>		
Python 3	Python Software Foundation	N/A
R version 4.3.2	R Core Team	N/A
R Studio 2024.04.1 Build 748	R Studio, PBC	N/A
PRISM 9	GraphPad Software	N/A

## 640 **Resource Availability**

### *Lead contact*

Further information and requests for resources should be directed to and will be fulfilled by the lead contact, Marc Emmenegger ([marc.emmenegger@usz.ch](mailto:marc.emmenegger@usz.ch)).

### *Materials availability*

645 Small amounts of the biological samples can be shared if available, upon reasonable request, and if approval by an ethics committee as well as an MTA is in place. This study did not generate new unique reagents.

### *Data and code availability*

- Specific data sets can be shared upon reasonable request and if approval by an ethics committee as well as a data transfer agreement is in place.
  - Code employed in this study will be publicly available on Zenodo in the peer-reviewed version of this manuscript after publication.
  - Any additional information required to reanalyse the data reported in this paper is available from the lead contact upon request.
- 650

## 655 **Experimental model and subject details**

### *Ethics statement*

All experiments and analyses involving samples from human donors were conducted with the approval of the local ethics committee (*Ethikkommission Kanton Zürich*, KEK-ZH-Nr. 2015-0561, BASEC-Nr. 2018-01042, and BASEC-Nr. 2020-01731, Yale IRB #0906005332) and were approved by the University of Zurich, 660 the University Hospital Zurich, and the Yale University Medical School Human Investigation Committee, in accordance with the provisions of the Declaration of Helsinki and the Good Clinical Practice guidelines of the International Conference on Harmonisation. Part of the study conducted on anonymized data was performed after clarification of responsibility where a declaration of no-objection was obtained (*Ethikkommission Kanton Zürich*, Req-2019-00572 and Req-2022-00638). Whenever required, written general or 665 informed consent was obtained from study participants.

### *Study design and sampling*

To study the prevalence of IgG antibodies against nAra-h2, we made use of surplus plasma samples (n=27,239) from inpatients and outpatients admitted to the University Hospital of Zurich (USZ, n=24,536) collected daily (Monday-Friday) and used for population-wide interrogations of the antibody repertoire 670 (*Emmenegger, De Cecco, et al., 2023; Magalhães et al., 2021; Senatore et al., 2020*). The criteria for our study to include a sample into the analysis were: (1) The patients' blood was sent to the Institute of Clinical Chemistry (at USZ), (2) there was enough residual heparin plasma (150 µl) for the automated generation of a research aliquot, (3) no aliquot from the same patient was already provided within the same month, (4) additional information (age, sex, clinical ward to which patient was admitted) was available, as described recently (*Emmenegger, De Cecco, et al., 2023*). The demographic characteristics of this patient 675 collective is described in **Table 1**. Briefly, our unselected patient cohort comprised 48% female (n=11,944) and 52% male (n=12,592) patients, with a median age of 54 (IQR: 49-68) years.

For secondary screens and as controls, we additionally investigated samples from blood donors of the Blood Donation Service of the Canton of Zurich who consented to further use of their samples for research. 680 This collective was characterized by 40% male and 60% sex participants and a median age of 44 (IQR: 31-55) years. The criteria to be admitted for blood donation are in line with international standards of blood donation services, see (*Blutspendedienst, 2021a*) and are as outlined before (*Emmenegger, De Cecco, et al., 2023*). The detailed inclusion and exclusion criteria are enumerated here (*Blutspendedienst, 2021b*). Moreover, blood samples from children with CF were obtained during routine clinic visits at the Yale Pae- 685 diatric CF Center with informed consent in accordance with relevant laws and institutional guidelines. This cohort consisted of 45% female and 55% male participants with a median age of 13 (9-17) years, as outlined in **Table 1**.

151 individuals participated in an online health survey. Their median age was 36 (IQR:16.5-43) years. 62.3% of participants were female and 37.7% male (see **Table 3**). Additionally, a registry-based study was conducted in 115 individuals (median age 5.0 (IQR: 2.0-15.0), 48.7% of female sex) whereof 56 individuals were *CFTR* mutation carriers, and 59 individuals were pwCF (see **Table 4**).

## **Method Details**

### *Exploratory high-throughput screening*

The profiling of antibodies in an unselected university hospital patient cohort using a custom antibody discovery platform was conducted as recently shown (Emmenegger, De Cecco, et al., 2023; Emmenegger, Kumar, et al., 2021; Senatore et al., 2020), with minor modifications. Heparin plasma samples (hospital patients) or EDTA plasma samples (healthy blood donors) were investigated for the presence of antibodies directed against the major peanut allergen, natural *Arachis hypogaea* 2 (nAra-h2) purified from peanut extract. As control, some samples were tested for the presence of antibodies against calcitonin-gene related peptide (CGRP). We first coated high-binding 1536-well assay plates (Perkin Elmer SpectraPlates) with 0.5 µg/ml antigen, at a volume of three µl/well (end volume in all subsequent dispensing steps, except for blocking) on the Fritz Gyger CertusFlex or the Thermo Fisher Multidrop Combinl nanodispenser. Plates were incubated for one hour at 37 °C in an automated Thermo Fisher plate incubator and subsequently washed three times with PBS 0.1% Tween-20 (wash buffer) on the Biotek EI406 washer/dispenser. Blocking with 10 µl/well 5% milk in PBS 0.1% Tween-20 (blocking buffer) was performed on Biotek MultifloFX using the peristaltic dispensing unit. Following the incubation at RT for one hour and the subsequent aspiration of blocking buffer using Biotek EI406, plates were sealed with Agilent plate sealer and stored at -30 or -80 °C until the day of experiment. Frozen plates were thawed and peeled with Brooks plate peeler. 1536-well assay plates were pre-filled with 1% milk in PBS 0.1% Tween-20 (sample buffer) according to a pre-defined plate layout using Fritz Gyger CertusFlex or Thermo Fisher Multidrop Combinl, and plates entered the ECHO555 acoustic dispenser (Labcyte) where pre-diluted plasma samples were dispensed at different volumes (from 1200-7.5 nl) from a source plate using ultrasound. Reaching a final well volume of three µl, the samples were thus diluted between 0.02 and  $1.67 \times 10^{-4}$ . Plates were incubated for two hours at RT and washed five times with wash buffer. Secondary antibody (Peroxidase AffiniPure Goat Anti-Human IgG, Fcγ Fragment Specific, Jackson, 109-035-098, at 1:4,000 dilution in sample buffer) was dispensed on the MultifloFX peristaltic pump, incubated for one hour at RT, and washed three times. Lastly, TMB was added, incubated for three minutes at RT, and the chromogenic reaction was stopped with 0.5 M H<sub>2</sub>SO<sub>4</sub>. The plates were centrifuged after all dispensing steps, except for the addition of TMB. The absorbance at 450 nm was measured in a plate reader (Perkin Elmer, EnVision). Optical densities were fitted using logistic regression as recently introduced (Emmenegger, De Cecco, et al., 2023). To filter out potential false-positive results from the HTS due to outliers, the quality of the regression was assessed by means of the mean squared



residual error. Thus, only plasma samples with a  $p(EC_{50}) \geq 2$  and the mean squared residual error of the logistic regression  $< 20\%$  of the actual  $p(EC_{50})$  value were considered as hits, an approach we have applied before (Senatore et al., 2020).

#### 725 *Isolation of peripheral blood mononuclear cells and plasma from patients*

Peripheral blood mononuclear cells (PBMCs) were isolated from patients as described (Emmenegger, De Cecco, et al., 2021). Briefly, patients were re-contacted via the Clinical Trial Center at University Hospital Zurich and consented for a blood donation. The blood was first diluted 1:2 with sterile PBS, separated via Ficoll gradient, purified using 10% B cell medium (IMDM (21980-065, Gibco), 10% FBS, 1x P/S (15140-122, 730 200x stock, Gibco), 1x MEM non-essential amino acids (11140-035, 100x stock, Gibco), 1x MEM sodium pyruvate (11360-039, 100x stock, Gibco), and  $\beta$ -mercaptoethanol (1:1,000, 31350-010, Gibco)) diluted in PBS, and aliquots of 5-10 million PBMCs/cryo vial were frozen in FBS with 10% DMSO. The 1:2 diluted plasma was harvested and used for experiments.

#### *B cell receptor repertoire analysis*

735 High quality RNA (RIN $>7$ ) was extracted from total PBMCs and reverse-transcribed into cDNA with an oligo dT primer. A unique molecular identifier (UMI) and an adaptor sequence were added to the cDNA prior to purification with streptavidin magnetic beads. VDJ amplification of IGHA, IGHD, IGHE, IGHG, IGHM genes and index barcode incorporation was done with a primary PCR reaction (PCR1). PCR products were purified with AMPure XP beads and subjected to a secondary PCR reaction (PCR2) whereby dual-indexed primers 740 (i7 and i5) were added along with the Illumina P5 adaptor. A qPCR reaction, prior to PCR2, was used to determine the optimal number of cycles to use to avoid the plateau phase. PCR2 products were purified and measured on an Agilent TapeStation. Samples were pooled equimolarly and 20% PhiX sequences were added to the pool. The pool was sequenced on the Illumina MiSeq platform in 2x300 bp paired-end, sequencing mode. For the analysis of the read data we relied on the Immcantation suite Version 4.0.0 745 (<https://immcantation.readthedocs.io>). The analysis steps followed the workflow described in (Vander Heiden et al., 2014) and relied on the tools presto 0.6.0, changeo 1.0.0, alakazam 1.0.1, shazam 1.0.0 and tigger 1.0.0.

We applied the manual pRESTO pipeline to perform quality control, filter and process the reads, as described (Vander Heiden et al., 2014). As an output we generated consensus sequences and the assembled 750 receptor sequences. In a modification of the workflow, we performed deduplication of the UMIs with the package UMI-Tools (<https://github.com/CGATOxford/UMI-tools>) and allowed two mismatches when clustering UMIs.

For the subsequent postprocessing and analysis of the clone sequences, we used the pipeline templates provided in the Immcantation tools and custom R scripts to perform the following steps: performs V(D)J 755 alignment using IgBLAST and post-processes the output into the Change-O data standard (changeo-igblast

pipeline), infer V segment genotypes using TlgGER (tigger-genotype pipeline), automated detection of the clonal assignment threshold (shazam-threshold pipeline), assign Ig sequences into clonally related lineages and build full germline sequences (changeo-clone pipeline). All pipelines were run with default settings. With the corresponding R packages (tigger and shazam), we estimated the abundances of the BCR clones, and we computed the Hill diversity curves (alakazaam package). Somatic hypermutation and mutational frequencies were assessed with the shazam package.

#### *Analysis of convergence between sequences obtained through BCR repertoire sequencing and publicly available repertoires*

Heavy and light CDR3 sequences were mapped to publicly available CDR3 amino acid sequence datasets (Aguzzi et al., 2020; Croote et al., 2018; Hoh et al., 2016) with a 90% identity using the usearch tool (Version 11.0.667). The usearch performs a local alignment and we required a coverage of at least 70% of the query sequences.

#### *HLA genotyping, epitope prediction, and T cell activation assay*

Genomic DNA extracted from the peripheral blood of six CF patient samples was typed for HLA-DRB1, -DRB345, -DQA1, -DQB1, -DPB1, and -DPA1 alleles by 1x resolution typing NGS (HistoGenetics, Ossining, NY USA). We compared the observed allele frequencies using the following database: <http://allelefrequencies.net/hla.asp>, (Gonzalez-Galarza et al., 2020). The alignment and comparison of the amino acid sequences of HLA-DQA1\*05:01:01 with the DQA1\*01:01:01 allele and other DQA1\* alleles of the six individuals were performed using <https://www.ebi.ac.uk/ipd/imgt/hla/alignment/>, (Barker et al., 2023). In silico peptide binding predictions of Ara-h2 to the HLA-DQA/B combinations DQA1\*05:02/DQB1\*02:01:01 and DQA1\*05:02/DQB1\*03:01:01 were performed using NetMHCII-pan-4.0; see (Jensen et al., 2018). Prediction values are given in nM IC50, calculated as a % rank to a set of 1,000,000 random natural peptides, and indicated as strong ( $\leq 2\%$ ) or weak ( $\leq 10\%$ ) binding peptides. The following parameters were used for the algorithm: threshold: -99.9; threshold for strong binder (% Rank): 2; Threshold for weak binder (% Rank): 10. The sequence of Ara-h2 used was (Uniprot-ID: Q6PSU2): MAKLTILVA LALFLAAHA SARQQWELQG DRRCQSQLER ANLRPCEQHL MQKIQRDEDS YGRDPYSPSQ DPYSPSQDPD RRDYSPSPY DRRGAGSSQH QERCCNELNE FENNQRMCCE ALQQIMENQS DRLQGRQQEQ QFKRELRNLP QQCGLRAPQR CDLEVESGGR DRY.

#### *Sequencing of the CFTR gene in patients with peanut allergy*

DNA from PBMCs of an anonymised collection of 90 confirmed patients with allergies and of four pwCF was extracted using the Qiagen DNeasy Blood & Tissue kit. We then followed the *Advanta CFTR NGS Library Preparation on the LP48.48 IFC with Juno* protocol (FLUIDIGM, PN 101-627- C2 PROTOCOL, Fluidigm Corporation, South San Francisco, California, United States) to create suitable libraries for subsequent amplicon sequencing. We aimed to perform a target enrichment of *CFTR* variants from each of the 27 exons and select intronic regions. In short, we first pre-amplified the genomic (g)DNA as the concentration after

790 gDNA purification was < 30 ng/μl. The pre-amplified gDNA samples (n=96 samples, 90 samples from pa-  
tients with allergies, 4 samples from PwCF as positive controls, and 2 H<sub>2</sub>O samples as assay negative con-  
795 trols) were then indexed in a 96-well plate using targeted DNA sequencing barcodes, mixed with sample  
pre-mix (PCR water, TSP master mix, TSP sample loading reagent, TSP DNA polymerase), and were loaded  
into a flow cell containing integrated flow circuits (IFCs), together with the *CFTR* assay mix (loading agent  
and *CFTR* NGS assays pools) and the non-assay mix (loading agent with *CFTR* NGS assay pools). The JUNO  
scripts were run according to the manufacturer's instructions and the pooled samples were cleaned up  
following the protocol. This process was run twice for 48 samples per round as the IFCs used contained 48  
wells, and we obtained pools of 48 uniquely barcoded samples twice, i.e. 96 samples in total. Next, the  
sequencing adapter was added to the purified pooled libraries, another PCR was run as detailed by the  
800 manufacturer, and the PCR products were cleaned up. The PCR products after the first PCR round (PCR1)  
and after the second PCR round (PCR2) were then quality controlled on a Qubit Fluorometer (Thermo  
Fisher) following the manufacturer's instructions. We obtained the following concentrations: for library 1,  
after PCR1: 3.96 ng/μl; for library 2, after PCR1: 2.47 ng/μl; for library 1, after PCR2: 10.9 ng/μl; for library  
2, after PCR2: 9.15 ng/μl. We then looked at the samples with the TapeStation (Agilent) on an Agilent High  
805 Sensitivity DNA chip and observed the expected right shift of the electropherogram with a peak between  
300-400 bp. The library preparation was performed at the Genomic Diversity Centre, ETH Zurich. For Illu-  
mina sequencing, the two libraries were pooled and paired end 150 bp sequencing was performed on an  
Illumina MiSeq system at the Functional Genomics Centre Zurich (FGCZ). The quality of the raw data was  
assessed with FastQC 0.11.9. The reads were subsequently filtered with fastp 0.20.0 for minimum length  
810 of 18bp and mapped to the GRCh38 reference genome with Bowtie2 2.4.1. Marking of duplicates was  
performed with Picard tools 2.22.8 and sorting and indexing of the alignment was performed with  
samtools 1.11. Qualimap 2.2.1, SAMstat 1.5.1 and Picard tools 2.22.8 were used for assessing the quality  
of the mapping. Single nucleotide variants were called using GATK "HaplotypeCaller" pipeline (4.2.0) first  
for each patient data separately and then joint genotyping across the whole cohort was performed with  
815 GATK "CombineGVCFs" and "GenotypeGVCFs" pipelines (4.2.0), as recommended by the GATK Best Prac-  
tices. Vcftools 0.1.16 and bcftools 1.9 were used to calculate allele frequency and proportion of heterozy-  
gosity for each variant across the 90 Peanut-allergic patients, excluding the 4 CF control patients. Variants  
of low coverage (DP < 200) and detected outside of the *CFTR* gene were discarded with SnpSift 4.3. The  
identified sequence variants within the *CFTR* gene were then functionally annotated using SNPeff 4.3. Ad-  
820 ditional annotations were added with ClinVar (downloaded on the 07.03.2022) for a better understanding  
of the general clinical significance, with gnomad v3.1.1 allele frequencies using the pipeline available at  
[https://github.com/KalinNonchev/gnomAD\\_DB](https://github.com/KalinNonchev/gnomAD_DB) and with data from CFTR2 database (<https://cftr2.org/>,  
29.04.2022). The graphs were created in *Python* 3.8.3.

### *Online health survey to correlate health status with CFTR genotype*

825 We performed a fully anonymised data collection and data interpretation to correlate the *CFTR* genotype with health. The aim of the questionnaire was to identify whether the presence or absence of mutations in the *CFTR* gene were associated with allergies, in particular in heterozygous mutation carriers. We first designed a survey using REDCap (see **Supplemental File 13** for survey design and survey questions), ensuring that no clear link to allergies was visible to prevent bias. The online health survey was then announced on online platforms of CF patient organisations, notably Mukoviszidose e.V. Everybody whose *CFTR* genotype was known or was the parent of a child with CF, was amenable to participate. Data was analysed in R, similarly to previous accounts (Emmenegger, De Cecco, et al., 2023).

### *Registry-based and diagnostic code-based inference of prevalence of food allergy in mutation carriers and in pwCF*

835 PwCF (those confirmed by genetic sequencing of the *CFTR* gene) were identified using the Yale Paediatric CF Foundation's Patient Registry. This is a voluntary registry that captures the demographics and health information of pwCF who receive care in CF Foundation-accredited care centres in the United States. In parallel, *CFTR* mutation carriers (those with International Classification of Diseases, Tenth Revision (ICD-10) codes of Z14.1) were identified using Yale New Haven Health System's electronic medical records. Only *CFTR* mutation carriers with at least one clinic encounter at Yale from March 2018 to March 2023 who were between the ages of 0-18 at the time of that visit were included in the analysis. Data was analysed in R, similarly to previous accounts (Emmenegger, De Cecco, et al., 2023).

### *High-throughput microELISA of pwCF and controls using antigen panel*

845 The ELISA was conducted as detailed above (see Exploratory high-throughput screening), with the following modifications. (1) Instead of the unselected patient cohort, PwCF were selected from the biobank and random patient controls and healthy blood donor controls were included. (2) Instead of merely coating with nAra 2, an antigen panel was used (see list of antigens).

### *Indirect ELISA for IgG, IgG1, IgG2, IgG3, IgG4, and IgA with plasma samples*

850 Indirect ELISA was conducted in 384-well high-binding plates (PerkinElmer SpectraPlate) as previously shown. In short, plates were coated with 0.5 ug/ml nAra-h2 in sterile PBS and incubated at 37 °C for 1 h, followed by 3 washes with PBS 0.1% Tween-20 using Biotek EI406 and by blocking with 40 µL 5% milk in PBS-T for 1.5 h. Serum samples were diluted in sample buffer (1% milk in PBS 0.1% Tween-20) and a serial dilution was carried out (volume: 20 µL/well). After the sample incubation for 2 h at RT, the wells were washed five times with wash buffer and the presence of IgG, IgG1, IgG2, IgG3, IgG4, IgA, antibodies directed against nAra-h2 was detected using a panel of secondary antibodies, validated previously (Emmenegger et al., 2022) and referred to in **Key Resource Table**. The incubation of the secondary antibodies

for one hour at RT was followed by three washes with PBS 0.1% Tween-20, the addition of TMB, an incubation of five minutes at RT, and the addition of 0.5 M H<sub>2</sub>SO<sub>4</sub>. The plates were centrifuged after all dispensing steps, except for the addition of TMB. The absorbance at 450 nm was measured in a plate reader  
860 (Perkin Elmer, EnVision).

#### *ISAC and ImmunoCAP*

IgG4 and IgE ISAC (Thermo Fisher Science) were carried out according to the manufacturer's guidelines, using an array of 112 allergens from 48 sources, including food (oral), pollen (airway), venom, and contact allergens (Matricardi et al., 2016). ImmunoCAP (Thermo Fisher Science) assays were performed according  
865 to the manufacturer's guidelines.

#### *Rapid Extracellular Antigen Profiling*

REAP was performed as previously described (Wang, Dai, et al., 2021). In brief, IgG was purified from 50  $\mu$ l serum by 20  $\mu$ l protein G magnetic resin (Lytic Solutions) and quantified by NanoDrop 8000 Spectrophotometer (Thermo Fisher Scientific). Purified IgG was incubated with 10 OD induced empty yeast (same  
870 strain as the one used in library but displaying empty vector) for 3 hours in 4 °C with shaking to remove unspecific yeast binding IgG, and yeast-depleted IgG was filtered with 96-well 0.45- $\mu$ m filter plates (Thomas Scientific). 10  $\mu$ g yeast-depleted IgG was incubated with 10 OD induced library yeast in 100  $\mu$ l PBE (PBS with 0.5% BSA and 0.5 mM EDTA) with shaking for 1 hr at 4 °C, followed by 30 mins incubation with 1:100 biotin anti-human IgG Fc antibody (clone HP6017, BioLegend) in 100  $\mu$ l PBE and 30 mins incubation with 1:20 Streptavidin MicroBeads (Miltenyi Biotec) in 100  $\mu$ l PBE. Streptavidin MicroBeads captured yeast was positively selected by Multi-96 Columns (Miltenyi Biotec) placed in MultiMACS M96 Separator (Miltenyi Biotec). Selected yeast were recovered in 1 ml SDO -Ura at 30 °C for 24 hrs. DNA was extracted from yeast libraries using ZymoPrep-96 Yeast Plasmid Miniprep kits (Zymo Research). Purified plasmids were amplified and indexed (2 rounds Phusion PCR, 24 cycles/round), gel purified, and sequenced  
880 on an Illumina MiSeq Instrument with MiSeq Reagent Kit v3 (150-cycle). REAP score was calculated as previously described (Wang, Dai, et al., 2021). In brief, it is the log<sub>2</sub>-fold change between the frequency of each protein in pre and post selection library, i.e. before and after addition of the patient-derived purified IgG antibodies.

#### *Competitive Western Blotting*

885 Strips for competitive Western Blotting were prepared as follows: 100 ng nAra-h2 and 100 ng BSA were loaded into the same lane of NU-PAGE 4-12% Bis-Tris gel (ThermoFisher), run at 80 V for 10 min and then at 120 V for 60 min in MES running buffer. Gels were transferred onto PVDF membranes with a dry transfer system (iBlot 2 Gel Transfer Device, ThermoFisher) at 20 V for 7 min, cut into strips, and the membranes were blocked with 5% SureBlock (Lubio Science) for 1 hour at RT. Next, 1:100 dilutions of plasma samples  
890 were prepared in 1% SureBlock in PBS 0.1% Tween-20 (sample buffer) and matched samples were (i) mixed

with soluble 1 µg/ml nAra-h2 (competition) or (ii) without addition and incubated for two hours at 4° C in Eppendorf tubes. The pre-incubated plasma samples (both (i) and (ii)) were then added to the PVDF membrane strips and incubated overnight at 4° C. The following day, membranes were washed four times with PBS 0.1% Tween-20 and incubated for one hour with HRP-linked anti-human IgG antibody (Peroxidase AffiniPure Goat Anti-Human IgG, Fcγ Fragment Specific, Jackson, 109-035-098, at 1:5,000 dilution in sample buffer) at RT. Following another four-times wash with wash buffer, Immobilon Crescendo HRP Substrate (Merck Millipore) was added to the strips and the membranes were immediately imaged on the Fusion SOLO S imaging system (Vilber).

#### *Purification of IgG fraction from patient plasma*

10 mL 1:2 pre-diluted patient plasma (i.e. 5 mL undiluted plasma) collected during the isolation of peripheral blood mononuclear cells (PBMCs) (Emmenegger, De Cecco, et al., 2021) was first filtered through a sterile 0.4 µm filter. 2.5 mL Protein G Sepharose 4 fast flow (17-0618-05) was loaded onto a liquid chromatography column (Luer Lock, non-jacketed, I.D. x L 1.0 cm x 10 cm, Sigma), washed with 15-20 mL dH<sub>2</sub>O, and then equilibrated with 15-20 mL PBS. The pre-filtered plasma samples were then slowly loaded to the column, and the column was washed with 15 mL PBS. The tubing of the ÄKTA was then washed with dH<sub>2</sub>O and equilibrated with 0.1 M Glycine PH 2.7. The IgG fraction bound to Protein G on the column was then eluted using 20 mL 0.1 M Glycine pH 2.5 at a flow rate of 2.5 mL/min. The eluates were collected in tubes containing 50 µL 1 M TRIS pH 8, pooled, and dialysed against PBS overnight at 4 °C, resulting in 20 mL purified IgG per sample. We thus recovered 14-23 mg total IgG per patient, at total IgG concentrations between 0.7-1.15 mg/mL. The total IgG concentration in undiluted plasma is approximately four times higher, as the undiluted plasma (input: 5 mL) was diluted four times (output: 20 mL).

#### *Affinity determination using microfluidics diffusional sizing*

Alexa Fluor 647 N-hydroxysuccinimide ester (in DMSO, 4 equivalents) was added to nAra-h2 (typically 3.88 nmol, 1 equivalent) in 0.1 M NaHCO<sub>3</sub> (pH = 8.2). The mixture was incubated for 1 h at ca. 20 °C, protected from light, and subsequently purified by size exclusion chromatography on a Superdex 200 increase column connected to an ÄKTA pure instrument (GE Healthcare, UK) with a flow rate of 0.5 mL/min and PBS as eluent buffer, to yield labelled nAra-h2 (1.37 nmol).

Labelled HLA was incubated with varying fraction of purified IgGs derived from patient plasma and diluted in PBS 0.1% Tween-20 (Sigma Aldrich, US). The samples were incubated at RT for 30 minutes and the hydrodynamic radius, Rh, was determined by microfluidic antibody affinity profiling on a Fluidity One W (Fluidic Analytics).

The dissociation constant,  $K_d = \frac{[Ab][H]}{[AbH]}$ , and the antibody binding sites concentration,  $[Ab]_{tot}$ , was determined through Bayesian inference, as previously described (Fiedler et al., 2021; Schneider et al., 2022,

2023). [Ab] and [R] are the equilibrium concentrations of antibody binding sites and nAra-h2, respectively, and [AbR] is the concentration of bound nAra-h2. The data were analysed by Bayesian inference, according to the following equations. Following correction of fluorescence intensities for plasma autofluorescence, the fraction,  $f_d$  of nAra-h2 to diffuse into the distal channel is defined by (Linse et al., 2020)

$$f_d = \frac{[AbR](1 - \rho_b) + ([R]_0 - [AbR])(1 - \rho_f)}{[R]_0}$$

930

where  $[R]_0$  is the total concentration of nAra-h2, and  $\rho_b$  and  $\rho_f$  are the fractions of bound and free nAra-h2 to diffuse into the distal channel, respectively. By solving the binding equation, we obtain the following expression for [AbR]

935

$$[AbR] = \frac{\alpha[Ab]_{tot} + [R]_0 + K_D - \sqrt{(\alpha[Ab]_{tot} + [R]_0 + K_D)^2 - 4\alpha[Ab]_{tot}[R]_0}}{2}$$

940

where  $\alpha$  is the fraction of plasma used in the measurement and  $[Ab]_{tot}$  is the total concentration of antibody binding sites in the sample.  $K_d$  and  $[Ab]_{tot}$  were thus determined through Bayesian inference, with  $\rho_b$  and  $\rho_f$  as additional parameters to be inferred. The prior was considered to be flat in logarithmic space for  $K_d$  and  $[Ab]_{tot}$ , flat in linear space for  $\rho_b$  and  $\rho_f$ . The likelihood function was considered to be Gaussian, with a standard deviation obtained through triplicate measurements.

#### *IFN- $\gamma$ measurement in human plasma*

Measurement of IFN- $\gamma$  was performed according to the manufacturer's guidelines using the IFN gamma Human ELISA Kit, High Sensitivity (Invitrogen, BMS228-2HS). In short, 1:2 pre-diluted plasma samples (with PBS) were assayed in duplicates. The resulting OD was then interpolated with GraphPad Prism using a Sigmoidal, 4PL, X is log(Concentration) function. The resulting interpolated concentration was multiplied by four to account for the 1:2 predilution and the 1:2 dilution performed during the assay, and the concentration in pg/ml was calculated.

950

#### *Allergic sensitisation and challenge of Cftr<sup>-/-</sup> and wild type mice*

Transgenic CFTR<sup>-/+</sup> (B6.129P2-KOCftrtm1UNC; CF) mice were originally purchased from the Jackson Laboratory and bred with C57BL6 mice in the Yale animal facilities for over 5 generations. Cohoused, age (16-20 week-old mice) - and sex-matched Cftr<sup>-/-</sup> and Cftr<sup>+/+</sup> (WT) were sensitized to peanut allergen weekly for 6

weeks as previously described (Gertie et al., 2022). Briefly, a freshly prepared 5 mg ground blanched peanut was dissolved in 200  $\mu$ l of 0.2 mol sodium bicarbonate buffer with or without 10  $\mu$ g of cholera toxin  
955 (List Biologicals, Campbell, Calif) as an adjuvant and delivered by oral gavage. Blood was collected from the retro-orbital plexus from day 14 every 2 weeks until the endpoint, and plasma was isolated and preserved at -80 °C. After the 6 weeks of sensitization, mice were permitted to rest for 10 days before the challenge. Mice received no food for 4 hours before being challenged intraperitoneally with 500  $\mu$ g crude peanut extract (Greer, Cambridge, Mass) in 200  $\mu$ L PBS. Rectal temperature was measured every 15  
960 minutes for up to 60 minutes, clinical symptoms scores were assessed every 15 minutes (as described in **Fig. 9D**), and sera were collected for analysis 1 hour after the challenge. All procedures were performed in compliance with relevant laws and institutional guidelines and were approved by Yale University Institutional Animal Care and Use Committee.

#### *Cytokine measurement in murine plasma*

965 The cytokine concentrations in the plasma of *Cftr*<sup>-/-</sup> and WT mice were assessed with a U-plex Meso-Scale Multiplex immunoassay platform following the manufacturer's instructions (*Meso Scale Diagnostics, LLC*). The following cytokines were simultaneously evaluated in 50  $\mu$ l of plasma: IFN- $\gamma$ , IL-10 and IL-4.

#### *IgG and IgE ELISA with murine plasma*

Serum samples were analysed by ELISA to measure peanut-specific antibodies. Briefly, 20  $\mu$ g/mL of crude  
970 peanut extract (Greer Laboratories) in carbonate-bicarbonate buffer (pH 9.6) was coated on 96-well Maxisorp plates (Thermo Fisher Scientific) overnight at 4 °C. Plates were blocked with 1% bovine serum albumin (BSA) in PBS at RT for 1-hour followed by the addition of serially diluted serum samples with a 2-hour incubation at 37 °C. Peanut-specific antibodies of each isotype were detected with horseradish peroxidase (HRP)-conjugated goat anti-mouse IgE and, goat anti-mouse IgG, goat anti-mouse IgG1 (Polyclonal, Southern Biotech) antibodies at 37 °C. Antigen (Ag)-specific antibodies of each isotype were obtained from  
975 pooled sera from repeatedly immunized mice and used as standards, represented as arbitrary units (AU)/mL, for Ag-specific ELISAs.

#### *Identification of homologous sequences*

A protein BLAST was conducted using the BLOSUM-62 matrix on UniProt (Altschul et al., 1997; Schäffer et  
980 al., 2001). We have applied an E-Threshold of 0.01 without any further filtering, gapped=yes, and queried the UniProtKB reference proteomes plus Swiss-Prot (uniprotkb\_refprotswissprot, 64,899,224 sequences; 25,452,742,688 total letters). The sequence of Ara-h2 used was (Uniprot-ID: Q6PSU2): MAKLTILVALALFLLAAHA SARQQWELQG DRRCQSQLER ANLRPCEQHL MQKIQRDEDS YGRDPYSPSQ DPYSPSQDPD RRD-PYSPSPY DRRGAGSSQH QERCCNELNE FENNQRMCCE ALQQIMENQS DRLQGRQQEQ QFKRELRNLP  
985 QQCGLRAPQR CDLEVESGGR DRY.



## **Quantification and statistical analysis**

### *Exploratory data analysis*

PCA and UMAP were conducted as shown before (Emmenegger, De Cecco, et al., 2023). In short, PCA was carried out using the default implementation in the R stats package (prcomp, <https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/prcomp>) and data was visualized using the factoextra package (<https://cran.r-project.org/web/packages/factoextra/index.html>). For UMAP, the following configuration parameters from the umap package in R (<https://CRAN.R-project.org/package=umap>) were used, all of which are default, except for the metric where cosine was sometimes used instead of Euclidean due to the binary nature of the data (n\_neighbors: 15, n\_components: 2, metric: cosine, n\_epochs: 200, in-put: data, init: spectral, min\_dist: 0.1, set\_op\_mix\_ratio: 1, local\_connectivity: 1, bandwidth: 1, alpha: 1, gamma: 1, negative\_sample\_rate: 5, a: NA, b: NA, spread: 1, random\_state: NA, transform\_state: NA, knn: NA, knn\_repeats: 1, verbose: FALSE, umap\_learn\_args: NA). UMAP data was plotted using ggplot2 in R. Hierarchical clustering was conducted using the stats::hclust (<https://www.rdocumentation.org/packages/fastcluster/versions/1.2.3/topics/hclust>) library and dendrograms were visualized with functions of dendextend (<https://cran.r-project.org/web/packages/dendextend/vignettes/dendextend.html>), circlize ([https://jokergoo.github.io/circlize\\_book/book/](https://jokergoo.github.io/circlize_book/book/)), and factoextra. The method for hierarchical clustering was 'binary' when using the absence or presence of an ICD-10 code as feature or 'ward.D2' when using p(EC<sub>50</sub>) values.

### *Statistical data analysis*

Using a reactivity threshold of p(EC<sub>50</sub>) [sample dilution]  $\geq 2$  to define the seropositive samples ('hits' or 'seropositives') exceeding the threshold and the seronegative samples ('non-hits' or 'seronegatives') with p(EC<sub>50</sub>) values  $< 2$ , we aimed to correlate ICD-10 codes, active ingredients of prescribed drugs, and sex (male/female) with seropositivity. To this end, we first aggregated the dataset to obtain medical data occurring at least 5 times (i.e. in  $> 3.75\%$ ) in the seropositive group and then conducted  $\chi^2$  statistical testing, at  $\alpha$ -level 0.01 and applying Bonferroni correction for multiple comparisons.  $\chi^2$  analysis was performed using the stats.chi2\_contingency function of the SciPy package (<https://www.scipy.org/>) in Python.

We investigated associations between the p(EC<sub>50</sub>) values and medical (available ICD-10 codes, active ingredients of prescribed medication) or demographic (sex, age) data, without binarization of p(EC<sub>50</sub>) values into hits and non-hits. The analysis included 588 ICD-10 codes assigned to at least 100 patients and 384 active ingredients ('medications') appearing at least 100 times, sex, and age. Situated in a Bayesian framework, we applied multivariable regression with logit-transformed EC<sub>50</sub> values (i.e. -logit(EC<sub>50</sub>)) as outcome and with binary (0,1) predictor variables in the R package rstanarm (Goodrich et al., 2020), as shown before (Emmenegger, De Cecco, et al., 2023; Emmenegger, Emmenegger, et al., 2023; Emmenegger & Emmenegger,

ger, 2023), following the code set up by Julien Riou (Lamparter et al., 2022). Since regression using unin-  
1020 formative priors on regression coefficients (Normal(0,10)) led to the expected noise, given the number of  
parameters included in the analysis, we employed regularization techniques (Bayesian LASSO and regular-  
ized horseshoe priors, see (Gelman et al., 2013; Pironen & Vehtari, 2017)). We then depicted the associa-  
tions between ICD-10 codes, medication, or demographic features and the respective  $-\text{logit}(EC_{50})$  values  
as odds ratio with 95% credible interval. Statistical testing of age in hits ('seropositives) and non-hits ('ser-  
1025 onegatives') was performed with Mood's median test using the Python package `scipy.stats.median_test`  
(<https://www.scipy.org/>) or using Mann-Whitney U/Wilcoxon rank sum test at  $\alpha$ -level 0.01. Statistical tests  
(and respective post-hoc corrections, e.g. when accounting for multiple comparisons) were always speci-  
fied in the text and in the figure legend. To enhance visualisation, asterisks (\*) were occasionally used in  
the figures. They denote significance in the following manner:  $p < 0.01$ : \*,  $<0.001$ :\*\*,  $<0.0001$ :\*\*\*,  
1030  $<0.00001$ :\*\*\*\*.

## Acknowledgments

We thank all the hospital patients and the healthy individuals included in this study for their contribution to this research project. We acknowledge the help of the Research Ward of the Clinical Trials Center of the University Hospital Zurich as well as of the Research Data Service Center of the USZ. The Yale Pediatric CF Center is acknowledged for the banking of serum samples, and Eugenie Chung for coordinating the shipping of biological material between institutions. We are also very grateful to all patients and families who consented to the study. ISAC 112 test for IgG4 RUO and IgE has kindly been performed by Service laboratory, Thermo Fisher Scientific, Alleröd, Denmark, Bjarne Kristensen. The library preparation for the *CFTR* sequencing using the FLUIDIGM kit was performed in collaboration with the Genetic Diversity Centre (GDC), ETH Zurich, with the great and kind help of Dr. Roberta Benedetto (Fluidigm AG). We thank the entire high-throughput screening team, University Hospital Zurich, for help with this project, namely Leyla Batkitar, Magdalena Bialkowska, Lisa Caflisch, Berre Doğançay, Julie Domange, Jingjing Guo, Marigona Imeri, Lidia Madrigal, Lorène Mottier, Rea Müller, Antonella Rosati, Dezirae Schneider, and Anne Wiedmer. Dr. Pål Johansen and Muriel Träxler (Dermatology, University Hospital Zurich) are acknowledged for help with ImmunoCAP, and Dr. Thomas M. Kündig for inspiring discussions. We thank Rita Moos from the University Hospital Zurich for help with IgG purification and Raphael Schoen (Zurich, Switzerland) for help with figure design. The team of Mabyon AG, Schlieren, particularly Natascha Wuillemin and Jürgen Pahla, have been helpful throughout these studies and have contributed with access to samples and knowledge – we are thankful for all the help! We acknowledge Dr. Jacobo Sarabia del Castillo and Dr. Stefano Vavassori (both Cellerys AG, Schlieren, Switzerland) for help with Fluorospot assays. Warm thanks to Dr. Cezmi Akdis and Dr. Mübeccel Akdis for hosting us at the SIAF in Davos and for all the enriching interactions with their group members. All authors wish to thank their groups and collaborators for support. Lastly, heartfelt thanks to a wonderful wife and most beloved friend and partner, Dr. Vishalini Emmenegger (formerly Bio Engineering Laboratory, ETH Zurich) for daily discussions, care, love, and for being infinitely inspiring beyond life and death.

## Funding

Grants of the Swiss Personalized Health Network (Driver Grant 2017DRI17), of the Swiss National Science Foundation (SNF; grant #179040), of the Innovation Fund of the University Hospital Zurich (INOV00096), of the NOMIS Foundation, the Schwyzer Winiker Stiftung, and the Baugarten Stiftung (coordinated by the USZ Foundation, USZF27101) were awarded to AA and ME.

## Author contributions

Study design and overall idea: AA, ME. Planned and conducted the high-throughput antibody profiling experiments shown in **Figs. 1-2** and **S1**: AA, ME. Analysed the data from the high-throughput antibody profiling experiments shown in **Figs. 1-2** and **S1**: ME, with contributions of AC. Analysed the adult CF data shown in **Figs. 3** and **S1** and performed the comparisons: ME. Conducted the antibody isotyping and IgG subclass experiments with ELISA and visualized the ImmunoCAP experiments shown in **Fig. 4A-D**: ME. Planned and conducted the competitive Western Blot experiments shown in **Fig. 4E**: AA, ME. Conducted, analysed, and visualized the Microfluidic Affinity Antibody Profiling experiments (**Fig. 4F**): MMS, GM, with help from TJK. Carried out the ELISA-based experiments, analysed the data, and visualized the ISAC data shown in **Figs. 5** and **S2**: ME, with critical input from AA. Planned, conducted and analysed the Rapid extracellular antigen profiling data shown in **Figs. 6** and **S3**: YD, AMR, with analytical help from ME. Conducted the protein blast (not shown in a figure): ME. Conducted the BCR repertoire sequencing shown in **Figs. 7, S4, S5, S6**: CZ. Analysed the data from the BCR repertoire sequencing shown in **Figs. 7, S4, S5, S6**: FJN, HR, with help from CZ and ME. Planned, prepared, conducted, and analysed the HLA analysis (not shown in a figure): CZ, RM, MS, with help from ME. Planned, prepared, conducted, and analysed the T cell activation assays (**Fig. 7D**): RM, MS, ME. Had the idea to analyze *CFTR* in patients with allergies shown in **Fig. 8A-C**: AA. Prepared PCMBs and a DNA library of patients with peanut allergy for experiments shown in **Fig. 8A-C**: CZ, ME. Planned, prepared, and conducted the *CFTR* sequencing shown in **Fig. 8A-C**: ME, CZ. Analysed the data from the *CFTR* sequencing shown in **Fig. 8A-C**: NZ, with help from EMB and ME. Established a *CFTR* online questionnaire and analysed the data (**Fig. 8E-F**): ME, with help from KH and AA. Conducted a registry-based food allergy and *CFTR* study in Yale (not shown in a figure): SL, HN, with analytical help from ME. Planned, prepared, and conducted the peanut sensitisation in *Cftr* KO mouse model and related experiments (**Figs. 9** and **S7**): EMB, LRH, RG, MEE, SCE, with analytical help from ME. Conducted and analysed the IFN- $\gamma$  experiments shown in **Fig. S7D**: ME. Wrote the first version of the manuscript and conducted most analyses and created the figures: ME. Revised the manuscript and figures and suggested critical corrections: AA. Contributed to and critically revised the entire manuscript: all authors.

## Competing interests

AA is a member of the clinical and scientific advisory board of Fluidic Analytics and member of the board of directors of Mabyon AG and AB2Bio AG. AA has formerly been a scientific advisor to Mabyon AG. TPJK is a member of the board of directors of Fluidic Analytics. RM and MS are employed by Cellerys AG. All authors declare no competing interests.

## References

- 1095 Abbas, A. K., Trotta, E., Simeonov, D. R., Marson, A., & Bluestone, J. A. (2018). Revisiting IL-2: Biology and therapeutic prospects. *Science Immunology*, 3(25). <https://doi.org/10.1126/SCIIMMUNOL.AAT1482>
- Abdi, H., & Williams, L. J. (2010). Principal component analysis. *Wiley Interdisciplinary Reviews: Computational Statistics*, 2(4), 433–459. <https://doi.org/10.1002/WICS.101>
- 1100 Adriaanse, M. P. M., Vreugdenhil, A. C. E., Groeneweg, M., Brüggewirth, H. T., Castelijns, S. J. A. M., van der Ent, C. K., Voorter, C. E. M., & Tilanus, M. G. J. (2014). HLA frequencies and associations in cystic fibrosis. *Tissue Antigens*, 83(1), 27–31. <https://doi.org/10.1111/TAN.12265>
- Agache, I., Sampath, V., Aguilera, J., Akdis, C. A., Akdis, M., Barry, M., Bouagnon, A., Chinthrajah, S., Collins, W., Dulitzki, C., Erny, B., Gomez, J., Goshua, A., Jutel, M., Kizer, K. W., Kline, O., LaBeaud, A. D., Pali-Schöll, I., Perrett, K. P., ... Nadeau, K. C. (2022). Climate change and global health: A call to more research and more action. *Allergy*, 77(5), 1389–1407. <https://doi.org/10.1111/ALL.15229>
- 1105 Agaronyan, K., Sharma, L., Vaidyanathan, B., Glenn, K., Yu, S., Annicelli, C., Wigger, T. D., Penningroth, M. R., Hunter, R. C., Dela Cruz, C. S., & Medzhitov, R. (2022). Tissue remodeling by an opportunistic pathogen triggers allergic inflammation. *Immunity*, 55(5), 895-911.e10. <https://doi.org/10.1016/J.IMMUNI.2022.04.001>
- Aguzzi, A., Wullemmin, N., Sonati, T., & Bieli, D. (2020). *ANTI-ALLERGEN ANTIBODIES* (Patent 20200199207).
- 1110 Akdis, C. A. (2021). Does the epithelial barrier hypothesis explain the increase in allergy, autoimmunity and other chronic conditions? *Nature Reviews Immunology* 2021 21:11, 21(11), 739–751. <https://doi.org/10.1038/s41577-021-00538-7>
- Akdis, C. A., & Nadeau, K. C. (2022). Human and planetary health on fire. *Nature Reviews Immunology* 2022 22:11, 22(11), 651–652. <https://doi.org/10.1038/s41577-022-00776-3>
- 1115 Akdis, M., & Akdis, C. A. (2014). Mechanisms of allergen-specific immunotherapy: multiple suppressor factors at work in immune tolerance to allergens. *The Journal of Allergy and Clinical Immunology*, 133(3), 621–631. <https://doi.org/10.1016/J.JACI.2013.12.1088>
- 1120 Albon, D., Zhang, L., Patrie, J., Jones, M., Li, Z. G., Noonan, E., & Borish, L. (2023). Association between Cystic Fibrosis exacerbations, lung function, T2 inflammation and microbiological colonization. *Allergy, Asthma and Clinical Immunology*, 19(1), 1–10. <https://doi.org/10.1186/S13223-023-00760-Z/FIGURES/5>

- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, 25(17), 3389–3402. <https://doi.org/10.1093/NAR/25.17.3389>
- 1125 Aron, Y., Polla, B. S., Bienvenu, T., Dall’Ava, J., Dusser, D., & Hubert, D. (2012). HLA Class II Polymorphism in Cystic Fibrosis. *https://Doi.Org/10.1164/Ajrccm.159.5.9807046*, 159(5 I), 1464–1468. <https://doi.org/10.1164/AJRCCM.159.5.9807046>
- Barker, D. J., Maccari, G., Georgiou, X., Cooper, M. A., Flicek, P., Robinson, J., & Marsh, S. G. E. (2023). The IPD-IMGT/HLA Database. *Nucleic Acids Research*, 51. <https://doi.org/10.1093/nar/gkac1011>
- 1130 Bell, S. C., Mall, M. A., Gutierrez, H., Macek, M., Madge, S., Davies, J. C., Burgel, P. R., Tullis, E., Castaños, C., Castellani, C., Byrnes, C. A., Cathcart, F., Chotirmall, S. H., Cosgriff, R., Eichler, I., Fajac, I., Goss, C. H., Drevinek, P., Farrell, P. M., ... Ratjen, F. (2020). The future of cystic fibrosis care: a global perspective. *The Lancet Respiratory Medicine*, 8(1), 65–124. [https://doi.org/10.1016/S2213-2600\(19\)30337-6](https://doi.org/10.1016/S2213-2600(19)30337-6)
- 1135 Berin, M. C. (2012). Mucosal antibodies in the regulation of tolerance and allergy to foods. *Seminars in Immunopathology*, 34(5), 633. <https://doi.org/10.1007/S00281-012-0325-9>
- Blutspendedienst. (2021a). *Admission criteria for blood donors*. <https://www.blutspendezurich.ch/spender/ich-will-spenden/>
- Blutspendedienst. (2021b). *Detailed inclusion and exclusion criteria for blood donors*. [https://www.blutspendezurich.ch/fileadmin/pdf/Formulare/Spender/Fragebogen\\_01.02.2020/d-Neuspender-V16.pdf](https://www.blutspendezurich.ch/fileadmin/pdf/Formulare/Spender/Fragebogen_01.02.2020/d-Neuspender-V16.pdf)
- 1140
- Boehncke, W. H., Loeliger, C., Kuehnl, P., Kalbacher, H., Böhm, B. O., & Gall, H. (1998). Identification of HLA-DR and -DQ alleles conferring susceptibility to pollen allergy and pollen associated food allergy. *Clinical & Experimental Allergy*, 28(4), 434–441. <https://doi.org/10.1046/J.1365-2222.1998.00246.X>
- 1145 Bojanowski, C. M., Lu, S., & Kolls, J. K. (2021). Mucosal Immunity in Cystic Fibrosis. *The Journal of Immunology*, 207(12), 2901–2912. <https://doi.org/10.4049/JIMMUNOL.2100424>
- Branum, A. M., & Lukacs, S. L. (2009). Food allergy among children in the United States. *Pediatrics*, 124(6), 1549–1555. <https://doi.org/10.1542/PEDS.2009-1210>
- Bruscia, E. M., & Bonfield, T. L. (2016). Innate and Adaptive Immunity in Cystic Fibrosis. *Clinics in Chest Medicine*, 37(1), 17–29. <https://doi.org/10.1016/j.ccm.2015.11.010>
- 1150
- Bruscia, E. M., & Bonfield, T. L. (2022). Update on Innate and Adaptive Immunity in Cystic Fibrosis. *Clinics in Chest Medicine*, 43(4), 603–615. <https://doi.org/10.1016/J.CCM.2022.06.004>

- Ca, F. C. (2001). *Anti-inflammatory cytokines in asthma and allergy: interleukin-10, interleukin-12, interferon-g*. <https://doi.org/10.1080/09629350120054518>
- 1155 Chen, G., Shrock, E. L., Li, M. Z., Spergel, J. M., Nadeau, K. C., Pongracic, J. A., Umetsu, D. T., Rachid, R., MacGinnitie, A. J., Phipatanakul, W., Schneider, L., Oettgen, H. C., & Elledge, S. J. (2021). High-resolution epitope mapping by AllerScan reveals relationships between IgE and IgG repertoires during peanut oral immunotherapy. *Cell Reports Medicine*, 2(10), 100410. <https://doi.org/10.1016/J.XCRM.2021.100410>
- 1160 Cianferoni, A., & Spergel, J. M. (2009). Food allergy: review, classification and diagnosis. *Allergol Int*, 58(4), 457–466. <https://doi.org/10.2332/allergolint.09-RAI-0138>
- Collaco, A., Jakab, R., Hegan, P., Mooseker, M., & Ameen, N. (2010).  $\alpha$ -AP-2 directs myosin VI-dependent endocytosis of cystic fibrosis transmembrane conductance regulator chloride channels in the intestine. *Journal of Biological Chemistry*, 285(22), 17177–17187. <https://doi.org/10.1074/jbc.M110.127613>
- 1165 Collins, A. M., & Watson, C. T. (2018). Immunoglobulin light chain gene rearrangements, receptor editing and the development of a self-tolerant antibody repertoire. *Frontiers in Immunology*, 9(OCT), 2249. <https://doi.org/10.3389/FIMMU.2018.02249/FULL>
- Cosgriff, R., Ahern, S., Bell, S. C., Brownlee, K., Burgel, P. R., Byrnes, C., Corvol, H., Cheng, S. Y., Elbert, A., 1170 Faro, A., Goss, C. H., Gulmans, V., Marshall, B. C., McKone, E., Middleton, P. G., Ruseckaite, R., Stephenson, A. L., & Carr, S. B. (2020). A multinational report to characterise SARS-CoV-2 infection in people with cystic fibrosis. *Journal of Cystic Fibrosis*, 19(3), 355–358. <https://doi.org/10.1016/J.JCF.2020.04.012>
- Couroux, P., Ipsen, H., Stage, B. S., Damkjær, J. T., Steffensen, M. A., Salapatek, A. M., Lund, K., & Würtzen, 1175 P. A. (2019). A birch sublingual allergy immunotherapy tablet reduces rhinoconjunctivitis symptoms when exposed to birch and oak and induces IgG4 to allergens from all trees in the birch homologous group. *Allergy*, 74(2), 361–369. <https://doi.org/10.1111/ALL.13606>
- Croote, D., Darmanis, S., Nadeau, K. C., & Quake, S. R. (2018). High-affinity allergen-specific human antibodies cloned from single IgE B cell transcriptomes. *Science*, 362(6420), 1306–1309. 1180 <https://doi.org/10.1126/science.aau2599>
- Cutting, G. R. (2015). Cystic fibrosis genetics: From molecular understanding to clinical application. *Nature Reviews Genetics*, 16(1), 45–56. <https://doi.org/10.1038/nrg3849>

- Davidson, T. M., Murphy, C., Mitchell, M., Smith, C., & Light, M. (1995). Management of chronic sinusitis in cystic fibrosis. *The Laryngoscope*, 105(4), 354–358. <https://doi.org/10.1288/00005537-199504000-00002>
- 1185
- De Lisle, R. C. (2014). Disrupted tight junctions in the small intestine of cystic fibrosis mice. *Cell and Tissue Research*, 355(1), 131. <https://doi.org/10.1007/S00441-013-1734-3>
- De Lisle, R. C., & Borowitz, D. (2013). The cystic fibrosis intestine. *Cold Spring Harbor Perspectives in Medicine*, 3(9).
- 1190
- Dorsey, J., & Gonska, T. (2017). Bacterial overgrowth, dysbiosis, inflammation, and dysmotility in the Cystic Fibrosis intestine. *Journal of Cystic Fibrosis*, 16, S14–S23.
- Dreskin, S. C., Tripputi, M. T., Aubrey, M. T., Mustafa, S. S., Atkins, D., Leo, H. L., Song, B., Schlichting, D., Talwar, H., Wang, Q., & Freed, B. M. (2010). Peanut-allergic subjects and their peanut-tolerant siblings have large differences in peanut-specific IgG that are independent of HLA class II. *Clinical Immunology*, 137(3), 366–373. <https://doi.org/10.1016/J.CLIM.2010.08.009>
- 1195
- Durham, S. R., & Shamji, M. H. (2022). Allergen immunotherapy: past, present and future. *Nature Reviews Immunology* 2022 23:5, 23(5), 317–328. <https://doi.org/10.1038/s41577-022-00786-1>
- Elborn, J. S. (2016). Cystic fibrosis. *Lancet*, 388(10059), 2519–2531. [https://doi.org/10.1016/S0140-6736\(16\)00576-6](https://doi.org/10.1016/S0140-6736(16)00576-6)
- 1200
- El-Khouly, F., Lewis, S. A., Pons, L., Burks, A. W., & Hourihane, J. O. B. (2007). IgG and IgE avidity characteristics of peanut allergic individuals. *Pediatric Allergy and Immunology*, 18(7), 607–613. <https://doi.org/10.1111/J.1399-3038.2007.00542.X>
- Emmenegger, M., De Cecco, E., Hruska-Plochan, M., Eninger, T., Schneider, M. M., Barth, M., Tantardini, E., Rossi, P. de, Bacioglu, M., Langston, R. G., Kaganovich, A., Bengoa-Vergniory, N., Gonzalez-Guerra, A., Avar, M., Heinzer, D., Reimann, R., Häsler, L. M., Herling, T. W., Matharu, N. S., ... Aguzzi, A. (2021). LAG3 is not expressed in human and murine neurons and does not modulate  $\alpha$ -synucleinopathies. *EMBO Molecular Medicine*, 13(9), e14745. <https://doi.org/10.15252/EMMM.202114745>
- 1205
- Emmenegger, M., De Cecco, E., Lamparter, D., Jacquat, R. P. B., Riou, J., Menges, D., Ballouz, T., Ebner, D., Schneider, M. M., Morales, I. C., Doğançay, B., Guo, J., Wiedmer, A., Domange, J., Imeri, M., Moos, R., Zografou, C., Batkitar, L., Madrigal, L., ... Aguzzi, A. (2023). Continuous population-level monitoring of SARS-CoV-2 seroprevalence in a large European metropolitan region. *iScience*, 105928. <https://doi.org/10.1016/j.isci.2023.105928>
- 1210



- Emmenegger, M., & Emmenegger, V. (2023). *Code repository for Antiphospholipid antibodies are enriched post-acute COVID-19 but do not modulate the thrombotic risk*. Zenodo. <https://doi.org/10.5281/zenodo.10051978>
- 1215
- Emmenegger, M., Emmenegger, V., Shambat, S. M., Scheier, T. C., Gomez-Mejia, A., Chang, C. C., Wendel-Garcia, P. D., Buehler, P. K., Buettner, T., Roggenbuck, D., Brugger, S. D., & Frauenknecht, K. B. M. (2023). Antiphospholipid antibodies are enriched post-acute COVID-19 but do not modulate the thrombotic risk. *Clinical Immunology*, 257, 109845. <https://doi.org/10.1016/J.CLIM.2023.109845>
- 1220
- Emmenegger, M., Fiedler, S., Brugger, S. D., Devenish, S. R. A., Morgunov, A. S., Ilsley, A., Ricci, F., Malik, A. Y., Scheier, T., Batkitar, L., Madrigal, L., Rossi, M., Meisl, G., Lynn, A. K., Saleh, L., von Eckardstein, A., Knowles, T. P. J., & Aguzzi, A. (2022). Both COVID-19 infection and vaccination induce high-affinity cross-clade responses to SARS-CoV-2 variants. *iScience*, 25(8), 104766. <https://doi.org/10.1016/j.isci.2022.104766>
- 1225
- Emmenegger, M., Kumar, S. S., Emmenegger, V., Malinauskas, T., Buettner, T., Rose, L., Schierack, P., Sprinzl, M. F., Sommer, C. J., Lackner, K. J., Aguzzi, A., Roggenbuck, D., & Frauenknecht, K. B. M. (2021). Anti-prothrombin autoantibodies enriched after infection with SARS-CoV-2 and influenced by strength of antibody response against SARS-CoV-2 proteins. *PLOS Pathogens*, 17(12), e1010118. <https://doi.org/10.1371/journal.ppat.1010118>
- 1230
- Emmenegger, M., Worth, R., Fiedler, S., Devenish, S. R. A., Knowles, T. P. J., & Aguzzi, A. (2023). Protocol to determine antibody affinity and concentration in complex solutions using microfluidic antibody affinity profiling. *STAR Protocols*, 4(1), 102095. <https://doi.org/10.1016/J.XPRO.2023.102095>
- Ferreira, D. M., Darrieux, M., Silva, D. A., Leite, L. C. C., Ferreira, J. M. C., Ho, P. L., Miyaji, E. N., & Oliveira, M. L. S. (2009). Characterization of protective mucosal and systemic immune responses elicited by pneumococcal surface protein PspA and PspC nasal vaccines against a respiratory pneumococcal challenge in mice. *Clinical and Vaccine Immunology*, 16(5), 636–645. [https://doi.org/10.1128/CVI.00395-08/SUPPL\\_FILE/CVI\\_395\\_08\\_SUPPL\\_L\\_FIG\\_\\_LEGEND.DOC](https://doi.org/10.1128/CVI.00395-08/SUPPL_FILE/CVI_395_08_SUPPL_L_FIG__LEGEND.DOC)
- 1235
- Fiedler, S., Piziorska, M. A., Denninger, V., Morgunov, A. S., Ilsley, A., Malik, A. Y., Schneider, M. M., Devenish, S. R. A., Meisl, G., Kosmoliaptsis, V., Aguzzi, A., Fiegler, H., & Knowles, T. P. J. (2021). Antibody Affinity Governs the Inhibition of SARS-CoV-2 Spike/ACE2 Binding in Patient Serum. *ACS Infectious Diseases*, 7(8), 2362–2369. [https://doi.org/10.1021/ACSINFECDIS.1C00047/SUPPL\\_FILE/ID1C00047\\_SI\\_001.PDF](https://doi.org/10.1021/ACSINFECDIS.1C00047/SUPPL_FILE/ID1C00047_SI_001.PDF)
- 1240
- Filkins, L. M., & O’Toole, G. A. (2015). Cystic Fibrosis Lung Infections: Polymicrobial, Complex, and Hard to Treat. *PLOS Pathogens*, 11(12), e1005258. <https://doi.org/10.1371/journal.ppat.1005258>

- 1245 Florsheim, E. B., Sullivan, Z. A., Khoury-Hanold, W., & Medzhitov, R. (2021). Food allergy as a biological food quality control system. *Cell*, *184*(6), 1440–1454. <https://doi.org/10.1016/J.CELL.2020.12.007>
- Frossard, C. P., Hauser, C., & Eigenmann, P. A. (2004). Antigen-specific secretory IgA antibodies in the gut are decreased in a mouse model of food allergy. *Journal of Allergy and Clinical Immunology*, *114*(2), 377–382. <https://doi.org/10.1016/j.jaci.2004.03.040>
- 1250 Garcia, M. A. S., Yang, N., & Quinton, P. M. (2009). Normal mouse intestinal mucus release requires cystic fibrosis transmembrane regulator–dependent bicarbonate secretion. *The Journal of Clinical Investigation*, *119*(11), 3497–3497. <https://doi.org/10.1172/JCI38662C1>
- Garg, M., & Ooi, C. Y. (2017). The Enigmatic Gut in Cystic Fibrosis: Linking Inflammation, Dysbiosis, and the Increased Risk of Malignancy. *Current Gastroenterology Reports*, *19*(2), 1–8. <https://doi.org/10.1007/s11894-017-0546-0>
- 1255 Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). *Bayesian Data Analysis* (Third Edit). CRC Press, Taylor & Francis Group.
- Gertie, J. A., Zhang, B., Liu, E. G., Hoyt, L. R., Yin, X., Xu, L., Long, L. L., Soldatenko, A., Gowthaman, U., Williams, A., & Eisenbarth, S. C. (2022). Oral anaphylaxis to peanut in a mouse model is associated with gut permeability but not with Tlr4 or Dock8 mutations. *Journal of Allergy and Clinical Immunology*, *149*(1), 262–274. <https://doi.org/10.1016/J.JACI.2021.05.015>
- 1260 Gibbens, D. T., Gilsanz, V., Boechat, M. I., Dufer, D., Carison, M. E., & Wang, C. I. (1988). Osteoporosis in cystic fibrosis. *The Journal of Pediatrics*, *113*(2), 295–300. [https://doi.org/10.1016/S0022-3476\(88\)80268-3](https://doi.org/10.1016/S0022-3476(88)80268-3)
- 1265 Gibson-Corley, K. N., Meyerholz, D. K., & Engelhardt, J. F. (2016). Pancreatic pathophysiology in cystic fibrosis. *Journal of Pathology*, *238*(2), 311–320. <https://doi.org/10.1002/path.4634>
- Gonzalez-Galarza, F. F., McCabe, A., Santos, E. J. M. Dos, Jones, J., Takeshita, L., Ortega-Rivera, N. D., Cid-Pavon, G. M. D., Ramsbottom, K., Ghattaoraya, G., Alfirevic, A., Middleton, D., & Jones, A. R. (2020). Allele frequency net database (AFND) 2020 update: gold-standard data classification, open access genotype data and new query tools. *Nucleic Acids Research*, *48*(D1), D783–D788. <https://doi.org/10.1093/NAR/GKZ1029>
- 1270 Goodrich, B., Gabry, J., Ali, I., & Brilleman, S. (2020). *rstanarm: Bayesian applied regression modeling via Stan*. <https://mc-stan.org/rstanarm>
- Gorrieri, G., Scudieri, P., Caci, E., Schiavon, M., Tomati, V., Sirci, F., Napolitano, F., Carrella, D., Gianotti, A., Musante, I., Favia, M., Casavola, V., Guerra, L., Rea, F., Ravazzolo, R., Di Bernardo, D., & Galietta, L. J.
- 1275

- V. (2016). Goblet Cell Hyperplasia Requires High Bicarbonate Transport To Support Mucin Release. *Scientific Reports* 2016 6:1, 6(1), 1–15. <https://doi.org/10.1038/srep36016>
- Grimes, D. A., & Schulz, K. F. (2002). Bias and causal associations in observational research. *The Lancet*, 359(9302), 248–252. [https://doi.org/10.1016/S0140-6736\(02\)07451-2](https://doi.org/10.1016/S0140-6736(02)07451-2)
- 1280 Gustafsson, J. K., Ermund, A., Ambort, D., Johansson, M. E. V., Nilsson, H. E., Thorell, K., Hebert, H., Sjövall, H., & Hansson, G. C. (2012). Bicarbonate and functional CFTR channel are required for proper mucin secretion and link cystic fibrosis with its mucus phenotype. *The Journal of Experimental Medicine*, 209(7), 1263–1272. <https://doi.org/10.1084/JEM.20120562>
- Gustafsson, J. K., & Johansson, M. E. V. (2022). The role of goblet cells and mucus in intestinal homeostasis. 1285 *Nature Reviews Gastroenterology and Hepatology*, 19(12), 785–803. <https://doi.org/10.1038/S41575-022-00675-X>
- Hartl, D., Gaggar, A., Bruscia, E., Hector, A., Marcos, V., Jung, A., Greene, C., McElvaney, G., Mall, M., & Döring, G. (2012). Innate immunity in cystic fibrosis lung disease. *Journal of Cystic Fibrosis*, 11(5), 363–382. <https://doi.org/10.1016/j.jcf.2012.07.003>
- 1290 Hecker, T. M., & Aris, R. M. (2004). Management of Osteoporosis in Adults with Cystic Fibrosis. *Drugs*, 64(2), 133–147. <https://doi.org/10.2165/00003495-200464020-00002>
- Hill, M. O. (1973). Diversity and Evenness: A Unifying Notation and Its Consequences. *Ecology*, 54(2), 427–432. <https://doi.org/10.2307/1934352>
- Hoh, R. A., Joshi, S. A., Liu, Y., Wang, C., Roskin, K. M., Lee, J. Y., Pham, T., Looney, T. J., Jackson, K. J. L., 1295 Dixit, V. P., King, J., Lyu, S. C., Jenks, J., Hamilton, R. G., Nadeau, K. C., & Boyd, S. D. (2016). Single B-cell deconvolution of peanut-specific antibody responses in allergic patients. *Journal of Allergy and Clinical Immunology*, 137(1), 157–167. <https://doi.org/10.1016/J.JACI.2015.05.029>
- Hong, X., Hao, K., Ladd-Acosta, C., Hansen, K. D., Tsai, H. J., Liu, X., Xu, X., Thornton, T. A., Caruso, D., Keet, 1300 C. A., Sun, Y., Wang, G., Luo, W., Kumar, R., Fuleihan, R., Singh, A. M., Kim, J. S., Story, R. E., Gupta, R. S., ... Wang, X. (2015). Genome-wide association study identifies peanut allergy-specific loci and evidence of epigenetic mediation in US children. *Nature Communications* 2015 6:1, 6(1), 1–12. <https://doi.org/10.1038/ncomms7304>
- Howell, W. M., Turner, S. J., Hourihane, J. O. B., Dean, T. P., & Warner, J. O. (1998). HLA class II DRB1, DQB1 1305 and DPB1 genotypic associations with peanut allergy: evidence from a family-based and case-control study. *Clinical & Experimental Allergy*, 28(2), 156–162. <https://doi.org/10.1046/J.1365-2222.1998.00224.X>

- Husby, S., Oxelius, V.-A., Teisner, B., Jensenius, J. C., & Svehag, S.-E. (1985). Humoral immunity to dietary antigens in healthy adults. Occurrence, isotype and IgG subclass distribution of serum antibodies to protein antigens. *International Archives of Allergy and Applied Immunology*, 77(4), 416–422.  
1310 <https://doi.org/10.1159/000233819>
- Ioannidis, J. P. A. (2016). Why Most Clinical Research Is Not Useful. *PLoS Medicine*, 13(6).  
<https://doi.org/10.1371/JOURNAL.PMED.1002049>
- Irrgang, P., Gerling, J., Kocher, K., Lapuente, D., Steininger, P., Habenicht, K., Wytopil, M., Beileke, S., Schäfer, S., Zhong, J., Ssebyatika, G., Krey, T., Falcone, V., Schüle, C., Peter, A. S., Nganou-Makamdop, K., Hengel, H., Held, J., Bogdan, C., ... Tenbusch, M. (2023). Class switch toward noninflammatory, spike-specific IgG4 antibodies after repeated SARS-CoV-2 mRNA vaccination. *Science Immunology*, 8(79), eade2798. <https://doi.org/10.1126/SCIIMMUNOL.ADE2798>  
1315
- Jakob, T., Forstenlechner, P., Matricardi, P., & Kleine-Tebbe, J. (2015). Molecular allergy diagnostics using multiplex assays: methodological and practical considerations for use in research and clinical routine: Part 21 of the Series Molecular Allergology. *Allergo Journal International*, 24(8), 320–332.  
1320 <https://doi.org/10.1007/S40629-015-0087-8/TABLES/4>
- Jensen, K. K., Andreatta, M., Marcatili, P., Buus, S., Greenbaum, J. A., Yan, Z., Sette, A., Peters, B., & Nielsen, M. (2018). Improved methods for predicting peptide binding affinity to MHC class II molecules. *Immunology*, 154(3), 394–406. <https://doi.org/10.1111/IMM.12889>
- Jiménez-Saiz, R., Bruton, K., Koenig, J. F. E., Wasserman, S., & Jordana, M. (2018). The IgE memory reservoir in food allergy. *Journal of Allergy and Clinical Immunology*, 142(5), 1441–1443.  
1325 <https://doi.org/10.1016/J.JACI.2018.08.029>
- Jiménez-Saiz, R., Chu, D. K., Mandur, T. S., Walker, T. D., Gordon, M. E., Chaudhary, R., Koenig, J., Saliba, S., Galipeau, H. J., Utley, A., King, I. L., Lee, K., Ettinger, R., Wasserman, S., Kolbeck, R., & Jordana, M. (2017). Lifelong memory responses perpetuate humoral TH2 immunity and anaphylaxis in food allergy. *Journal of Allergy and Clinical Immunology*, 140(6), 1604-1615.e5.  
1330 <https://doi.org/10.1016/J.JACI.2017.01.018>
- Kayani, K., Mohammed, R., & Mohiaddin, H. (2018). Cystic fibrosis-related diabetes. *Frontiers in Endocrinology*, 9(FEB). <https://doi.org/10.3389/fendo.2018.00020>
- Kiyotani, K., Mai, T. H., Yamaguchi, R., Yew, P. Y., Kulis, M., Orgel, K., Imoto, S., Miyano, S., Burks, A. W., & Nakamura, Y. (2018). Characterization of the B-cell receptor repertoires in peanut allergic subjects undergoing oral immunotherapy. *Journal of Human Genetics*, 63(2), 239–248.  
1335 <https://doi.org/10.1038/s10038-017-0364-0>

- 1340 Kotz, D., Simpson, C. R., & Sheikh, A. (2011). Incidence, prevalence, and trends of general practitioner–recorded diagnosis of peanut allergy in England, 2001 to 2005. *Journal of Allergy and Clinical Immunology*, *127*(3), 623–630.e1. <https://doi.org/10.1016/J.JACI.2010.11.021>
- Lamparter, D., Jacquat, R. P. B., Riou, J., Menges, D., Ballouz, T., & Emmenegger, M. (2022). *Code repository for SARS-CoV-2 seroprevalence study*. <https://doi.org/10.5281/ZENODO.7454292>
- 1345 Laserson, U., Vigneault, F., Gadala-Maria, D., Yaari, G., Uduman, M., Vander Heiden, J. A., Kelton, W., Jung, S. T., Liu, Y., Laserson, J., Chari, R., Lee, J. H., Bachelet, I., Hickey, B., Lieberman-Aiden, E., Hanczaruk, B., Simen, B. B., Egholm, M., Koller, D., ... Church, G. M. (2014). High-resolution antibody dynamics of vaccine-induced immune responses. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(13), 4928–4933. <https://doi.org/10.1073/PNAS.1323862111/-/DCSUPPLEMENTAL>
- 1350 Leviatan, S., Vogl, T., Klompus, S., Kalka, I. N., Weinberger, A., & Segal, E. (2022). Allergenic food protein consumption is associated with systemic IgG antibody responses in non-allergic individuals. *Immunity*, *55*(12), 2454–2469.e6. <https://doi.org/10.1016/J.IMMUNI.2022.11.004>
- Linnemann, R. W., Yadav, R., Zhang, C., Sarr, D., Rada, B., & Stecenko, A. A. (2022). Serum anti-PAD4 autoantibodies are present in cystic fibrosis children and increase with age and lung disease severity. *Autoimmunity*, *55*(2), 109–117. <https://doi.org/10.1080/08916934.2021.2021193>
- 1360 Linse, S., Scheidt, T., Bernfur, K., Vendruscolo, M., Dobson, C. M., Cohen, S. I. A., Sileikis, E., Lundqvist, M., Qian, F., O'Malley, T., Bussiere, T., Weinreb, P. H., Xu, C. K., Meisl, G., Devenish, S. R. A., Knowles, T. P. J., & Hansson, O. (2020). Kinetic fingerprints differentiate the mechanisms of action of anti-A $\beta$  antibodies. *Nature Structural & Molecular Biology*, *27*(12), 1125–1133. <https://doi.org/10.1038/s41594-020-0505-6>
- 1365 Liu, E. G., Zhang, B., Martin, V., Anthonypillai, J., Kraft, M., Grishin, A., Grishina, G., Catanzaro, J. R., Chinthrajah, S., Sindher, T., Manohar, M., Quake, A. Z., Nadeau, K., Burks, A. W., Kim, E. H., Kulis, M. D., Henning, A. K., Jones, S. M., Leung, D. Y. M., ... Eisenbarth, S. C. (2022). Food-specific immunoglobulin A does not correlate with natural tolerance to peanut or egg allergens. *Science Translational Medicine*, *14*(671), eabq0599. <https://doi.org/10.1126/scitranslmed.abq0599>
- Liu, J., Walker, N. M., Ootani, A., Strubberg, A. M., & Clarke, L. L. (2015). Defective goblet cell exocytosis contributes to murine cystic fibrosis–associated intestinal disease. *The Journal of Clinical Investigation*, *125*(3), 1056–1068. <https://doi.org/10.1172/JCI73193>
- 1370 Lloyd, C. M., & Snelgrove, R. J. (2018). Type 2 immunity: Expanding our view. *Science Immunology*, *3*(25). <https://doi.org/10.1126/SCIIMMUNOL.AAT1604>

- Lockhart, A., Reed, A., Castro, T. R. de, Herman, C., Canesso, M. C. C., & Mucida, D. (2023). Dietary protein shapes the profile and repertoire of intestinal CD4+ T cells. *BioRxiv*, 2023.04.11.536475. <https://doi.org/10.1101/2023.04.11.536475>
- 1375 Losa, M., Emmenegger, M., Rossi, P. De, Schuerch, P. M., Serdiuk, T., Pengo, N., Capron, D., Bieli, D., Rupp, N. J., Carta, M. C., Frontzek, K. J., Lysenko, V., Reimann, R., Lakkaraju, A. K., Nuvolone, M., Westermarck, G. T., Nilsson, K. P., Polymenidou, M., Theocharides, A. P., ... Aguzzi, A. (2023). The ASC inflammasome adapter controls the severity of inflammation-related AA amyloidosis. *BioRxiv*, 2021.05.01.442282. <https://doi.org/10.1101/2021.05.01.442282>
- 1380 Lucarelli, S., Quattrucci, S., Zingoni, A. M., Frediani, T., Diamanti, S., Quintieri, F., Barbato, M., Cardi, E., & Antonelli, M. (1994). Food allergy in cystic fibrosis. *Minerva Pediatrica*, 46(12), 543–548.
- Lyczak, J. B., Cannon, C. L., & Pier, G. B. (2002). Lung infections associated with cystic fibrosis. *Clinical Microbiology Reviews*, 15(2), 194–222. <https://doi.org/10.1128/CMR.15.2.194-222.2002>
- 1385 Madore, A. M., Vaillancourt, V. T., Asai, Y., Alizadehfar, R., Ben-Shoshan, M., Michel, D. L., Kozyrskyj, A. L., Becker, A., Chan-Yeung, M., Clarke, A. E., Hull, P., Daley, D., Sandford, A. J., & Laprise, C. (2013). HLA-DQB1\*02 and DQB1\*06:03P are associated with peanut allergy. *European Journal of Human Genetics* 2013 21:10, 21(10), 1181–1184. <https://doi.org/10.1038/ejhg.2013.13>
- 1390 Magalhães, A. D., Emmenegger, M., Cecco, E. De, Carta, M., Frontzek, K., Chincisan, A., Hornemann, S., & Aguzzi, A. (2021). Large-scale seroepidemiology identifies a nephro-vascular syndrome associated with autoimmune reactivity to tau. *MedRxiv*, 2021.11.24.21266833. <https://doi.org/10.1101/2021.11.24.21266833>
- Maina, A., Mureithi, M., Kiiru, J., & Revathi, G. (2022). Systemic and Mucosal Concentrations of Nine Cytokines Among Individuals with *Neisseria gonorrhoeae* infection in Nairobi Kenya. *Open Research Africa*, 5, 12. <https://doi.org/10.12688/AASOPENRES.13351.1>
- 1395 Matricardi, P. M., Kleine-Tebbe, J., Hoffmann, H. J., Valenta, R., Hilger, C., Hofmaier, S., Aalberse, R. C., Agache, I., Asero, R., Ballmer-Weber, B., Barber, D., Beyer, K., Biedermann, T., Bilò, M. B., Blank, S., Bohle, B., Bosshard, P. P., Breiteneder, H., Brough, H. A., ... Ollert, M. (2016). EAACI Molecular Allergy User's Guide. *Pediatric Allergy and Immunology*, 27(S23), 1–250. <https://doi.org/https://doi.org/10.1111/pai.12563>
- 1400 Mattila, P., Joenväärä, S., Renkonen, J., Toppila-Salmi, S., & Renkonen, R. (2011). Allergy as an epithelial barrier disease. *Clinical and Translational Allergy*, 1(1), 1–8. <https://doi.org/10.1186/2045-7022-1-5/TABLES/3>

- Mazzini, E., Massimiliano, L., Penna, G., & Rescigno, M. (2014). Oral Tolerance Can Be Established via Gap Junction Transfer of Fed Antigens from CX3CR1+ Macrophages to CD103+ Dendritic Cells. *Immunity*, 40(2), 248–261. <https://doi.org/10.1016/j.immuni.2013.12.012>
- 1405 McDole, J. R., Wheeler, L. W., McDonald, K. G., Wang, B., Konjufca, V., Knoop, K. A., Newberry, R. D., & Miller, M. J. (2012). Goblet cells deliver luminal antigen to CD103+ dendritic cells in the small intestine. *Nature* 2012 483:7389, 483(7389), 345–349. <https://doi.org/10.1038/nature10863>
- McKendry, R. T., Kwok, M., Hemmings, O., James, L. K., & Santos, A. F. (2021). Allergen-specific IgG show distinct patterns in persistent and transient food allergy. *Pediatric Allergy and Immunology*, 32(7), 1508–1518. <https://doi.org/10.1111/PAI.13567>
- 1410 Miho, E., Yermanos, A., Weber, C. R., Berger, C. T., Reddy, S. T., & Greiff, V. (2018). Computational strategies for dissecting the high-dimensional complexity of adaptive immune repertoires. *Frontiers in Immunology*, 9(FEB), 224. <https://doi.org/10.3389/FIMMU.2018.00224/BIBTEX>
- Miller, A. C., Comellas, A. P., Hornick, D. B., Stoltz, D. A., Cavanaugh, J. E., Gerke, A. K., Welsh, M. J., Zabner, J., & Polgreen, P. M. (2020). Cystic fibrosis carriers are at increased risk for a wide range of cystic fibrosis-related conditions. *Proceedings of the National Academy of Sciences of the United States of America*, 117(3), 1621–1627.
- Moran, T. P., & Burks, A. W. (2015). Is Clinical Tolerance Possible after Allergen Immunotherapy? *Current Allergy and Asthma Reports*, 15(5), 23. <https://doi.org/10.1007/S11882-015-0523-3>
- 1420 Mueller, G. A., Gosavi, R. A., Pomés, A., Wünschmann, S., Moon, A. F., London, R. E., & Pedersen, L. C. (2011). Ara h 2: crystal structure and IgE binding distinguish two subpopulations of peanut allergic patients by epitope diversity. *Allergy*, 66(7), 878–885. <https://doi.org/10.1111/j.1398-9995.2010.02532.x>
- Nichols, D. P., & Chmiel, J. F. (2015). Inflammation and its genesis in cystic fibrosis. *Pediatric Pulmonology*, 50, S39–S56. <https://doi.org/10.1002/PPUL.23242>
- 1425 Noh, G., & Lee, S. S. (2009). A Pilot Study of Interferon- $\gamma$ -Induced Specific Oral Tolerance Induction (ISOTI) for Immunoglobulin E-Mediated Anaphylactic Food Allergy. <https://Home.Liebertpub.Com/Jir>, 29(10), 667–675. <https://doi.org/10.1089/JIR.2009.0001>
- Nwaru, B. I., Hickstein, L., Panesar, S. S., Roberts, G., Muraro, A., & Sheikh, A. (2014). Prevalence of common food allergies in Europe: a systematic review and meta-analysis. *Allergy*, 69(8), 992–1007. <https://doi.org/10.1111/ALL.12423>
- 1430

- Okafor, S., Kelly, K. M., & Halderman, A. A. (2020). Management of Sinusitis in the Cystic Fibrosis Patient. *Immunology and Allergy Clinics*, 40(2), 371–383. <https://doi.org/10.1016/J.IAC.2019.12.008>
- 1435 Ooi, C. Y., & Durie, P. R. (2016). Cystic fibrosis from the gastroenterologist’s perspective. *Nature Reviews Gastroenterology & Hepatology* 2016 13:3, 13(3), 175–185. <https://doi.org/10.1038/nrgastro.2015.226>
- 1440 Orengo, J. M., Radin, A. R., Kamat, V., Badithe, A., Ben, L. H., Bennett, B. L., Zhong, S., Birchard, D., Limnander, A., Rafique, A., Bautista, J., Kostic, A., Newell, D., Duan, X., Franklin, M. C., Olson, W., Huang, T., Gandhi, N. A., Lipsich, L., ... Yancopoulos, G. D. (2018). Treating cat allergy with monoclonal IgG antibodies that bind allergen and prevent IgE engagement. *Nature Communications* 2018 9:1, 9(1), 1–15. <https://doi.org/10.1038/s41467-018-03636-8>
- 1445 Ota, M., Hoehn, K. B., Ota, T., Aranda, C. J., Friedman, S., Braga, W. F., Malbari, A., Kleinstein, S. H., Sicherer, S. H., & Lafaille, M. A. C. de. (2023). The memory of pathogenic IgE is contained within CD23+IgG1+ memory B cells poised to switch to IgE in food allergy. *BioRxiv*, 2023.01.25.525506. <https://doi.org/10.1101/2023.01.25.525506>
- Pabst, O., & Mowat, A. M. (2012). Oral tolerance to food protein. *Mucosal Immunology* 2012 5:3, 5(3), 232–239. <https://doi.org/10.1038/mi.2012.4>
- 1450 Paolucci, M., Wuillemin, N., Homère, V., Bieli, D., Köhli, A., Ballmer-Weber, B., Waeckerle-Men, Y., Pengo, N., Kündig, T. M., Sonati, T., & Johansen, P. (2023). Targeting Ara h 2 with human-derived monoclonal antibodies prevents peanut-induced anaphylaxis in mice. *Allergy*, 78(6), 1605–1614. <https://doi.org/10.1111/ALL.15659>
- 1455 Patil, S. U., Ogunniyi, A. O., Calatroni, A., Tadigotla, V. R., Ruitter, B., Ma, A., Moon, J., Love, J. C., & Shreffler, W. G. (2015). Peanut oral immunotherapy transiently expands circulating Ara h 2–specific B cells with a homologous repertoire in unrelated subjects. *Journal of Allergy and Clinical Immunology*, 136(1), 125–134.e12. <https://doi.org/10.1016/J.JACI.2015.03.026>
- Perkin, M. R., Logan, K., Tseng, A., Raji, B., Ayis, S., Peacock, J., Brough, H., Marrs, T., Radulovic, S., Craven, J., Flohr, C., & Lack, G. (2016). Randomized Trial of Introduction of Allergenic Foods in Breast-Fed Infants. *New England Journal of Medicine*, 374(18), 1733–1743. <https://doi.org/10.1056/NEJMoa1514210>
- 1460 Piironen, J., & Vehtari, A. (2017). Sparsity information and regularization in the horseshoe and other shrinkage priors. *Electronic Journal of Statistics*, 11(2), 5018–5051. <https://doi.org/10.1214/17-EJS1337SI>
- Pillai, S. (2023). Is it bad, is it good, or is IgG4 just misunderstood? *Science Immunology*, 8(81). <https://doi.org/10.1126/SCIIMMUNOL.ADG7327>



- Polgreen, P. M., Brown, G. D., Hornick, D. B., Ahmad, F., London, B., Stoltz, D. A., & Comellas, A. P. (2018).  
1465 CFTR Heterozygotes Are at Increased Risk of Respiratory Infections: A Population-Based Study. *Open  
Forum Infectious Diseases*, 5(11).
- Price, C. E., & O'Toole, G. A. (2021). The Gut-Lung Axis in Cystic Fibrosis. *Journal of Bacteriology*, 203(20).  
<https://doi.org/10.1128/JB.00311-21>
- Ramsey, B., & Richardson, M. A. (1992). Impact of sinusitis in cystic fibrosis. *Journal of Allergy and Clinical  
1470 Immunology*, 90(3), 547–552. [https://doi.org/10.1016/0091-6749\(92\)90183-3](https://doi.org/10.1016/0091-6749(92)90183-3)
- Roesch, E. A., Nichols, D. P., & Chmiel, J. F. (2018). Inflammation in cystic fibrosis: An update. *Pediatric  
Pulmonology*, 53(S3), S30–S50. <https://doi.org/10.1002/PPUL.24129>
- Rogosch, T., Kerzel, S., Sikula, L., Gentil, K., Liebetrueth, M., Schlingmann, K.-P., Maier, R. F., & Zemlin, M.  
(2010). Plasma Cells and Nonplasma B Cells Express Differing IgE Repertoires in Allergic Sensitization.  
1475 *The Journal of Immunology*, 184(9), 4947–4954. <https://doi.org/10.4049/JIMMUNOL.0900859>
- Rowe, S. M., Miller, S., & Sorscher, E. J. (2005). Cystic fibrosis. *N Engl J Med*, 352(19), 1992–2001.  
<https://doi.org/10.1056/NEJMra043184>
- Rubelt, F., Busse, C. E., Bukhari, S. A. C., Bürckert, J. P., Mariotti-Ferrandiz, E., Cowell, L. G., Watson, C. T.,  
Marthandan, N., Faison, W. J., Hershberg, U., Laserson, U., Corrie, B. D., Davis, M. M., Peters, B.,  
1480 Lefranc, M. P., Scott, J. K., Breden, F., Luning Prak, E. T., & Kleinstein, S. H. (2017). Adaptive Immune  
Receptor Repertoire Community recommendations for sharing immune-repertoire sequencing data.  
*Nature Immunology* 2017 18:12, 18(12), 1274–1278. <https://doi.org/10.1038/ni.3873>
- Schäffer, A. A., Aravind, L., Madden, T. L., Shavirin, S., Spouge, J. L., Wolf, Y. I., Koonin, E. V., & Altschul, S.  
F. (2001). Improving the accuracy of PSI-BLAST protein database searches with composition-based  
1485 statistics and other refinements. *Nucleic Acids Research*, 29(14), 2994–3005.  
<https://doi.org/10.1093/NAR/29.14.2994>
- Schneider, M. M., Emmenegger, M., Xu, C. K., Condado Morales, I., Meisl, G., Turelli, P., Zografou, C., Zim-  
mermann, M. R., Frey, B. M., Fiedler, S., Denninger, V., Jacquat, R. P., Madrigal, L., Ilsley, A., Kos-  
moliaptsis, V., Fiegler, H., Trono, D., Knowles, T. P., & Aguzzi, A. (2022). Microfluidic characterisation  
1490 reveals broad range of SARS-CoV-2 antibody affinity in human plasma. *Life Science Alliance*, 5(2),  
e202101270. <https://doi.org/10.26508/lsa.202101270>
- Schneider, M. M., Scheidt, T., Priddey, A. J., Xu, C. K., Hu, M., Meisl, G., Devenish, S. R. A., Dobson, C. M.,  
Kosmoliaptsis, V., & Knowles, T. P. J. (2023). Microfluidic antibody affinity profiling of alloantibody-  
HLA interactions in human serum. *Biosensors and Bioelectronics*, 228, 115196.  
1495 <https://doi.org/10.1016/J.BIOS.2023.115196>

- Senatore, A., Frontzek, K., Emmenegger, M., Chincisan, A., Losa, M., Reimann, R., Horny, G., Guo, J., Fels, S., Sorce, S., Zhu, C., George, N., Ewert, S., Pietzonka, T., Hornemann, S., & Aguzzi, A. (2020). Protective anti-prion antibodies in human immunoglobulin repertoires. *EMBO Molecular Medicine*, *12*(9). <https://doi.org/10.15252/emmm.202012739>
- 1500 Shamji, M. H., Valenta, R., Jardetzky, T., Verhasselt, V., Stephen, |, Durham, R., & Würtzen, P. A. (2021). The role of allergen-specific IgE, IgG and IgA in allergic disease. *Allergy*, *00*, 1–15. <https://doi.org/10.1111/all.14908>
- Sheikh, S., Britt, R. D., Ryan-Wenger, N. A., Khan, A. Q., Lewis, B. W., Gushue, C., Ozuna, H., Jaganathan, D., McCoy, K., & Kopp, B. T. (2023). Impact of elexacaftor–tezacaftor–ivacaftor on bacterial colonization and inflammatory responses in cystic fibrosis. *Pediatric Pulmonology*, *58*(3), 825–833. <https://doi.org/10.1002/PPUL.26261>
- 1505 Shreffler, W. G., Charlop-Powers, Z., & Sicherer, S. H. (2006). Lack of association of HLA class II alleles with peanut allergy. *Annals of Allergy, Asthma & Immunology*, *96*(6), 865–869. [https://doi.org/10.1016/S1081-1206\(10\)61351-8](https://doi.org/10.1016/S1081-1206(10)61351-8)
- 1510 Sicherer, S. H., & Sampson, H. A. (2018). Food allergy: A review and update on epidemiology, pathogenesis, diagnosis, prevention, and management. *J Allergy Clin Immunol*, *141*(1), 41–58. <https://doi.org/10.1016/j.jaci.2017.11.003>
- Smith, N., Goncalves, P., Charbit, B., Grzelak, L., Beretta, M., Planchais, C., Bruel, T., Rouilly, V., Bondet, V., Hadjadj, J., Yatim, N., Pere, H., Merklings, S. H., Ghozlane, A., Kernéis, S., Rieux-Laucat, F., Terrier, B., Schwartz, O., Mouquet, H., ... Di Santo, J. P. (2021). Distinct systemic and mucosal immune responses during acute SARS-CoV-2 infection. *Nature Immunology* *2021* *22:11*, *22*(11), 1428–1439. <https://doi.org/10.1038/s41590-021-01028-7>
- 1515 Snouwaert, J. N., Brigman, K. K., Latour, A. M., Malouf, N. N., Boucher, R. C., Smithies, O., & Koller, B. H. (1992). An Animal Model for Cystic Fibrosis Made by Gene Targeting. *Science*, *257*(5073), 1083–1088. <https://doi.org/10.1126/SCIENCE.257.5073.1083>
- 1520 Soller, L., Ben-Shoshan, M., Harrington, D. W., Fragapane, J., Joseph, L., Pierre, Y. S., Godefroy, S. B., La Vieille, S., Elliott, S. J., & Clarke, A. E. (2012). Overall prevalence of self-reported food allergy in Canada. *Journal of Allergy and Clinical Immunology*, *130*(4), 986–988. <https://doi.org/10.1016/j.jaci.2012.06.029>
- 1525 Spolidoro, G. C. I., Amara, Y. T., Ali, M. M., Nyassi, S., Lisik, D., Ioannidou, A., Rovner, G., Khaleva, E., Venter, C., van Ree, R., Worm, M., Vlieg-Boerstra, B., Sheikh, A., Muraro, A., Roberts, G., & Nwaru, B. I. (2023).

- Frequency of food allergy in Europe: An updated systematic review and meta-analysis. *Allergy*, 78(2), 351–368. <https://doi.org/10.1111/ALL.15560>
- 1530 Sun, Y., Hu, B., Stanley, G., Harris, Z. M., Gautam, S., Homer, R., Koff, J. L., & Rajagopalan, G. (2023). IFN- $\gamma$  Is Protective in Cytokine Release Syndrome-associated Extrapulmonary Acute Lung Injury. *American Journal of Respiratory Cell and Molecular Biology*, 68(1), 75–89. [https://doi.org/10.1165/RCMB.2022-01170C/SUPPL\\_FILE/DISCLOSURES.PDF](https://doi.org/10.1165/RCMB.2022-01170C/SUPPL_FILE/DISCLOSURES.PDF)
- 1535 Tay, S. S., Clark, A. T., Deighton, J., King, Y., & Ewan, P. W. (2007). Patterns of immunoglobulin G responses to egg and peanut allergens are distinct: ovalbumin-specific immunoglobulin responses are ubiquitous, but peanut-specific immunoglobulin responses are up-regulated in peanut allergy. *Clinical & Experimental Allergy*, 37(10), 1512–1518. <https://doi.org/10.1111/J.1365-2222.2007.02802.X>
- 1540 Tiringer, K., Treis, A., Fucik, P., Gona, M., Gruber, S., Renner, S., Dehlink, E., Nachbaur, E., Horak, F., Jaksch, P., Döring, G., Cramer, R., Jung, A., Roehat, M. K., Hörmann, M., Spittler, A., Klepetko, W., Akdis, C. A., Szépfalusi, Z., ... Eiwegger, T. (2013). A Th17- and Th2-skewed Cytokine Profile in Cystic Fibrosis Lungs Represents a Potential Risk Factor for *Pseudomonas aeruginosa* Infection. <https://doi.org/10.1164/Rccm.201206-1150OC>, 187(6), 621–629. <https://doi.org/10.1164/RCCM.201206-1150OC>
- Tordesillas, L., & Berin, M. C. (2018). Mechanisms of oral tolerance. *Clinical Reviews in Allergy & Immunology*, 55(2), 107. <https://doi.org/10.1007/S12016-018-8680-5>
- 1545 Tordesillas, L., Berin, M. C., & Sampson, H. A. (2017). Immunology of Food Allergy. *Immunity*, 47(1), 32–50. <https://doi.org/10.1016/j.immuni.2017.07.004>
- 1550 Vander Heiden, J. A., Stathopoulos, P., Zhou, J. Q., Chen, L., Gilbert, T. J., Bolen, C. R., Barohn, R. J., Dimachkie, M. M., Ciafaloni, E., Broering, T. J., Vigneault, F., Nowak, R. J., Kleinstein, S. H., & O'Connor, K. C. (2017). Dysregulation of B Cell Repertoire Formation in Myasthenia Gravis Patients Revealed through Deep Sequencing. *The Journal of Immunology*, 198(4), 1460–1473. <https://doi.org/10.4049/JIMMUNOL.1601415/-/DCSUPPLEMENTAL>
- 1555 Vander Heiden, J. A., Yaari, G., Uduman, M., Stern, J. N. H., O'Connor, K. C., Hafler, D. A., Vigneault, F., & Kleinstein, S. H. (2014). pRESTO: a toolkit for processing high-throughput sequencing raw reads of lymphocyte receptor repertoires. *Bioinformatics*, 30(13), 1930–1932. <https://doi.org/10.1093/BIOINFORMATICS/BTU138>
- Walker, N. M., Liu, J., Young, S. M., Woode, R. A., & Clarke, L. L. (2022). Goblet cell hyperplasia is not epithelial-autonomous in the Cftr knockout intestine. *American Journal of Physiology. Gastrointestinal and Liver Physiology*, 322(2), G282–G293. <https://doi.org/10.1152/AJPGI.00290.2021>

- 1560 Wang, E. Y., Dai, Y., Rosen, C. E., Schmitt, M. M., Dong, M. X., Ferré, E. M. N., Liu, F., Yang, Y., Gonzalez-Hernandez, J. A., Meffre, E., Hinchcliffe, M., Koumpouras, F., Lionakis, M. S., & Ring, A. M. (2021). REAP: A platform to identify autoantibodies that target the human exoproteome. *BioRxiv*, 2021.02.11.430703. <https://doi.org/10.1101/2021.02.11.430703>
- 1565 Wang, E. Y., Mao, T., Klein, J., Dai, Y., Huck, J. D., Jaycox, J. R., Liu, F., Zhou, T., Israelow, B., Wong, P., Coppi, A., Lucas, C., Silva, J., Oh, J. E., Song, E., Perotti, E. S., Zheng, N. S., Fischer, S., Campbell, M., ... Ring, A. M. (2021). Diverse functional autoantibodies in patients with COVID-19. *Nature*, *595*(7866), 283–288. <https://doi.org/10.1038/s41586-021-03631-y>
- Weiner, H. L., da Cunha, A. P., Quintana, F., & Wu, H. (2011). Oral tolerance. *Immunol. Rev.*, *241*(1), 241–259. <https://doi.org/10.1111/j.1600-065x.2011.01017.x>
- 1570 Wood, R. A., Togias, A., Sicherer, S. H., Shreffler, W. G., Kim, E. H., Jones, S. M., Leung, D. Y. M., Vickery, B. P., Bird, J. A., Spergel, J. M., Iqbal, A., Olsson, J., Ligueros-Saylan, M., Uddin, A., Calatroni, A., Huckabee, C. M., Rogers, N. H., Yovetich, N., Dantzer, J., ... Chinthrajah, R. S. (2024). Omalizumab for the Treatment of Multiple Food Allergies. <https://doi.org/10.1056/NEJMoa2312382>, *390*(10), 889–899. <https://doi.org/10.1056/NEJMoa2312382>
- 1575 Yadav, R., Linnemann, R. W., Kahlenberg, J. M., Bridges, L. S., Stecenko, A. A., & Rada, B. (2020). IgA autoantibodies directed against self DNA are elevated in cystic fibrosis and associated with more severe lung dysfunction. *Autoimmunity*, *53*(8), 476–484. <https://doi.org/10.1080/08916934.2020.1839890>

## Figures and Tables

1580 **Table 1.** Population characteristics for different analyses performed in this study.

Assay	Characteristics	Non-hits	Hits	Total	
<i>High-throughput screening</i>	Individuals, number	24,409	127	24,536	
	Median age (IQR), years	54 (49-68)	45 (32-58.5)	54 (49-68)	
	Sex, female %	48	42	48	
	Sex, male %	52	58	52	
Assay	Characteristics	Non-hits	Hits	Total	
<i>CF profiling</i>	Individuals, number	88	31	119	
	Median age (IQR), years	33 (25-40)	31.5 (25.25-37.25)	32 (25-40)	
	Sex, female %	49	46	47	
	Sex, male %	51	54	53	
Assay	Characteristics	CF	PCF	PC	HC
<i>Secondary screen (ELISA)</i>	Individuals, number	177	76	319	358
	Median age (IQR), years	34 (26-42)	13 (9-17)	54 (39-67)	44 (31-55)
	Sex, female %	51	45	48	60
	Sex, male %	49	55	52	40

**Table 2.** Number and prevalence of select gastrointestinal, autoimmune, and inflammatory diseases recorded in individuals included in the primary HTS. The prevalence was calculated by dividing the number of occurrences of patients with the ICD-10 code by the total number of patients screened.

Disease	ICD-10	Number of occurrences	Prevalence in %
<i>Cystic fibrosis</i>	<i>E84</i>	169	0.6
<i>Gastritis and duodenitis</i>	<i>K29</i>	906	3.3
<i>Functional dyspepsia</i>	<i>K30</i>	88	0.3
<i>Crohn's disease</i>	<i>K50</i>	217	0.8
<i>Ulcerative colitis</i>	<i>K51</i>	220	0.8
<i>Other and unspecified noninfective gastroenteritis and colitis</i>	<i>K52</i>	541	2.0
<i>Irritable bowel syndrome</i>	<i>K58</i>	113	0.4
<i>Intestinal malabsorption</i>	<i>K90</i>	102	0.4
<i>Systemic lupus erythematosus</i>	<i>M32</i>	181	0.7
<i>Connective tissue disease</i>	<i>M35</i>	594	2.2
<i>Type I diabetes mellitus</i>	<i>E10</i>	437	1.6

<i>Type II diabetes mellitus</i>	<i>E11</i>	<i>2440</i>	<i>9.0</i>
<i>Asthma</i>	<i>J45</i>	<i>504</i>	<i>1.9</i>
<i>Other chronic obstructive pulmonary disease</i>	<i>J44</i>	<i>949</i>	<i>3.5</i>
<i>Rheumatoid arthritis</i>	<i>M05 and M06</i>	<i>798</i>	<i>2.9</i>
<i>Other disorders involving the immune mechanism</i>	<i>D89</i>	<i>43</i>	<i>0.2</i>

1585

**Table 3.** Population characteristics for online survey investigating differences in the prevalence of allergy between CFTR mutation carriers and pwCF. The questionnaire was provided in German language and announced by a German CF association.

	<i>Wildtype</i>	<i>Mutation carrier</i>	<i>Cystic fibrosis</i>	<i>Overall</i>
<i>Individuals, number</i>	<i>4</i>	<i>91</i>	<i>56</i>	<i>151</i>
<i>Median age (IQR), years</i>	<i>24.5 (14.0-33.5)</i>	<i>40.0 (34.0-45.0)</i>	<i>13.5 (6.0-31.0)</i>	<i>36.0 (16.5-43.0)</i>
<i>Sex, female %</i>	<i>75</i>	<i>63.7</i>	<i>58.9</i>	<i>62.3</i>
<i>Sex, male %</i>	<i>25</i>	<i>36.3</i>	<i>41.1</i>	<i>37.7</i>
<i>Food allergy %</i>	<i>0</i>	<i>13.2</i>	<i>7.1</i>	<i>10.6</i>

1590

**Table 4.** Population characteristics for registry-based survey investigating differences in the prevalence of allergy between CFTR mutation carriers and pwCF. The survey was conducted in Yale, New Haven, United States of America.

	<i>Mutation carrier</i>	<i>Cystic fibrosis</i>	<i>Overall</i>
<i>Individuals, number</i>	<i>56</i>	<i>59</i>	<i>115</i>
<i>Median age (IQR), years</i>	<i>3.0 (1.8-4.0)</i>	<i>13.0 (8.0-18.0)</i>	<i>5.0 (2.0-15.0)</i>
<i>Sex, female %</i>	<i>44.6</i>	<i>52.5</i>	<i>48.7</i>
<i>Sex, male %</i>	<i>55.4</i>	<i>47.5</i>	<i>51.3</i>
<i>Food allergy %</i>	<i>7.1</i>	<i>8.5</i>	<i>7.8</i>

## 1595 Supplementary Figures

**Fig. S1. Additional illustrations pertaining to Fig. 1 and Fig. 2.A.** **A.** The average OD plotted against the respective  $p(EC_{50})$  values, for all samples with  $p(EC_{50})$  value  $\geq 1.7$ . **B.** qq-plot showing distribution of  $p(EC_{50})$  values for nAra-h2 of all patient samples. **C.** Scatter dot plots visualizing PrP vs. nAra-h2 and CGRP vs. nAra-h2. Histograms display the distribution of values. In blue colour: seropositives in the screen against nAra-h2, shown within the distribution of PrP and CGRP. **D.** Sample replicability in pwCF, some of which are seropositive and others seronegative.

**Fig. S2. ISAC data displaying raw values, pertaining to Fig. 5.** **A.** Raw values of the IgG4 assay are shown. **B.** Raw values of the IgE assay are shown.

**Fig. S3. Additional hit characterization of REAP experiment, pertaining to Fig. 6.** **A.** Number of hits, i.e. REAP score  $\geq 2$ , for all conditions on the entire dataset. **B.** Number of hits, i.e. REAP score  $\geq 2$ , for all conditions on the conformational proteins. **C.** Number of hits, i.e. REAP score  $\geq 2$ , for all conditions on the protein epitopes. **A-C.** The red dotted line indicates the mean value. One-way ANOVA was performed to compare between all conditions. No statistically significant difference in the positivity rate was observed between either of the groups.

**Fig. S4. Additional illustrations pertaining to BCR repertoire sequencing shown in Fig. 7.** **A.** Number of sequences analysed. **B.** Sequences pertaining to respective immunoglobulin isotype, illustrated as fractions. Pairwise t-test indicated no significant differences between the groups. **C.** Distribution of heavy and light chain, illustrated as fractions. Pairwise t-test indicated no significant differences between the groups. **D.** VH family usage by isotype. Wilcoxon rank sum test was performed.

**Fig. S5. Additional illustrations pertaining to BCR repertoire sequencing shown in Fig. 7.** **A.** Evenness inferred from the Hill diversity index calculated over a range of diversity orders ( $q$ ) ranging from 0 to 4, in equally spaced increments, for each subset. **B.** Mutation rate by VH for different isotypes. Wilcoxon rank sum test was performed.

**Fig. S6. Additional illustrations pertaining to BCR repertoire sequencing shown in Fig. 7.** No relevant convergence in sequences of known binders to peanuts were found using the data derived from BCR repertoire sequencing.

**Fig. S7. Additional illustrations pertaining to peanut sensitisation in mice shown in Fig. 9.** **A.** Shown are the distributions of the optical densities (ODs) irrespective of the plasma dilution after two weeks, five weeks, and post-challenge. **B.** Distributions of body temperature measurements post challenge among the different conditions. **C.** Same as in (B) but with anaphylactic score as readout instead of body temperature. Ps from Wilcoxon rank sum test with Holms correction for multiple comparisons. **D.** Determination of IFN- $\gamma$  plasma concentrations in eight anti-Ara-h2-seropositive pwCF, three patients with systemic lupus erythematosus, 13 patients with mixed disease history, 10 COVID-19 patients, and six healthy controls. Distributional differences were assessed using Wilcoxon rank sum test, with CF as reference group for comparisons. No significant difference was observed before and after Holm's correction for multiple comparisons. The assay was conducted in technical duplicates.

## Supplemental File

**Supplemental File 01.**  $X^2$  statistics for ICD-10 codes.

1635 **Supplemental File 02.**  $X^2$  statistics for ICD-10 codes after removal of patients diagnosed with CF.

**Supplemental File 03.** Odds ratios for all ICD-10 codes that entered the Bayesian regression model.

**Supplemental File 04.**  $X^2$  statistics for active ingredients of medications.

**Supplemental File 05.** Odds ratios for all active ingredients of medications that entered the Bayesian regression model.

1640 **Supplemental File 06.** Statistical comparisons of cohort screened for antibodies against multiple food proteins and controls.

**Supplemental File 07.** Data for all samples and proteins tested with ISAC.

**Supplemental File 08.** Data from BLAST experiment using BLOSUM-62 matrix.

**Supplemental File 09.** Data for all samples and proteins tested with REAP.

1645 **Supplemental File 10.** HLA analysis.

**Supplemental File 11.** HLA sequence alignment.

**Supplemental File 12.** HLA binding prediction.

**Supplemental File 13.** Online survey codebook.

1650