

Advances in methods for characterizing dietary patterns: A scoping review

Authors: Joy M. Hutchinson¹, Amanda Raffoul², Alexandra Pepetone¹, Lesley Andrade¹, Tabitha E. Williams¹, Sarah A. McNaughton³, Rebecca M. Leech⁴, Jill Reedy⁵, Marissa M. Shams-White^{6,7}, Jennifer E. Vena⁸, Kevin W. Dodd⁹, Lisa M. Bodnar¹⁰, Benoît Lamarche¹¹, Michael P. Wallace¹², Megan Deitchler¹³, Sanaa Hussain¹, Sharon I. Kirkpatrick¹

1. School of Public Health Sciences, University of Waterloo, Waterloo, ON, Canada
2. Department of Nutritional Sciences, University of Toronto, Toronto, ON, Canada
3. Health and Well-Being Centre for Research Innovation, School of Human Movement and Nutrition Sciences, University of Queensland, St. Lucia, QLD, Australia
4. Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, Deakin University, Victoria, Geelong, Australia
5. Office of Dietary Supplements, National Institutes of Health, Bethesda, MD, USA
6. Population Science Department, American Cancer Society, Washington DC, USA
7. Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, MD, USA
8. Alberta's Tomorrow Project, Alberta Health Services, Edmonton, AB, Canada
9. Division of Cancer Prevention, National Cancer Institute, Bethesda, MD, USA
10. School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA
11. Centre Nutrition, santé et société (NUTRISS), Institut sur la nutrition et les aliments fonctionnels (INAF), Université Laval, Québec City, QC, Canada
12. Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, ON, Canada
13. Intake – Center for Dietary Assessment, FHI Solutions, Washington, DC, USA

Abstract

There is a growing focus on better understanding the complexity of dietary patterns and how they relate to health and other factors. Approaches that have not traditionally been applied to characterize dietary patterns, such as machine learning algorithms and latent class analysis methods, may offer opportunities to measure and characterize dietary patterns in greater depth than previously considered. However, there has not been a formal examination of how this wide range of approaches has been applied to characterize dietary patterns. This scoping review synthesized literature from 2005-2022 applying methods not traditionally used to characterize dietary patterns, referred to as novel methods. MEDLINE, CINAHL, and Scopus were searched using keywords including machine learning, latent class analysis, and least absolute shrinkage and selection operator (LASSO). Of 5274 records identified, 24 met the inclusion criteria. Twelve of 24 articles were published since 2020. Studies were conducted across 17 countries. Nine studies used approaches that have applications in machine learning to identify dietary patterns. Fourteen studies assessed associations between dietary patterns that were characterized using novel methods and health outcomes, including cancer, cardiovascular disease, and asthma. There was wide variation in the methods applied to characterize dietary patterns and in how these methods were described. The extension of reporting guidelines and quality appraisal tools relevant to nutrition research to consider specific features of novel methods may facilitate complete and consistent reporting and enable evidence synthesis to inform policies and programs aimed at supporting healthy dietary patterns.

Introduction

Dietary intake is among the top risk factors for chronic diseases.^{1,2} Research examining dietary intake has historically focused on single foods, nutrients, or other dietary constituents.³ As the focus of public health nutrition shifted from the prevention of deficiency to include a focus on the prevention of chronic diseases, research likewise shifted towards the examination of dietary patterns, aiming to capture how foods and beverages are consumed in real life.³⁻⁵ Humans typically do not consume foods or nutrients on their own, but in the context of a broader dietary pattern.^{3,4} Accordingly, food-based dietary guidelines are now typically focused on patterns of intake rather than single dietary components.⁶ It is likely the synergistic and antagonistic relationships among the multiple foods, beverages, and other dietary components that humans consume that influence health rather than individual components.⁴ In addition to this multidimensionality, dietary patterns are dynamic, changing from meal to meal, day to day and across the life course.^{4,7} Further, dietary patterns are shaped by culture, social position, and other contextual factors.^{8,9} However, incorporating the domains of multidimensionality, dynamism, and contextual factors into dietary patterns analysis is a difficult task.

Traditional approaches to identify dietary patterns, including ‘a priori’ and ‘a posteriori’ approaches, are useful for understanding overall dietary patterns or diet quality of populations and population subgroups.¹⁰ For example, ‘a priori’ methods like the Healthy Eating Index-2020 or the Healthy Eating Food Index-2019 are generally investigator driven,^{11,12} consider multiple components as inputs, such as fruits and vegetables and whole grains, but typically compress the multidimensional construct of total dietary patterns, condensing inputs to a single unidimensional score reflecting overall diet quality.^{13,14} ‘A posteriori’ approaches are data-driven and have also been widely used to identify dietary patterns. Commonly applied data-driven approaches include clustering methods (e.g., k-means, Ward’s method), principal component

analysis, and factor analysis, providing opportunities to identify dietary patterns through statistical modelling or clustering algorithms rather than relying on researcher hypotheses.¹⁵ These approaches compress dietary components to key food groupings typically expressed as single scores.^{10,16} By reducing the dimensionality of dietary patterns, these methods are limited in their ability to explain the wide variation in dietary intakes.⁴ Methods employed to traditionally characterize dietary patterns using ‘a priori’ and ‘a posteriori’ approaches thus address multidimensionality to some extent, but do not allow for explorations of dietary patterns in their totality because they miss potential synergistic or antagonistic associations among dietary components.^{4,14,17}

Novel methods that have not traditionally been used to identify dietary patterns, such as probabilistic graphical modelling, latent class analysis, and machine learning algorithms (e.g., random forest, neural networks), may capture complexities like dietary synergy. There is no clear delineation between traditional and novel methods, and specifically defining what is novel is challenging given it naturally implies an evolution of methods. Nonetheless, there is a growing interest among nutrition researchers in the application of methods that have not typically been used to capture dietary complexity, with these methods often centered in machine learning.¹⁸ To date, there have been perspectives and narrative reviews on the application of machine learning in nutrition,^{19–21} and a recent systematic review of studies that applied machine learning approaches to assess food consumption.²² However, there has not been an assessment of studies applying novel methods to characterize dietary patterns. Given the rapid adoption of these methods within the field of health,^{23–26} it is increasingly important for researchers to have a basic understanding of available methods and how they are being applied in the field. This will facilitate the synthesis of evidence from a range of methodological inputs to inform food-based dietary guidelines and other policies and programs that promote health. The objective of this

scoping review was therefore to describe the use of novel methods not traditionally used to characterize dietary patterns in the published literature.

Methods

The review was conducted in accordance with the JBI Manual for Evidence Synthesis,²⁷ which was developed using the Arksey and O'Malley framework.²⁸ Reporting follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR).²⁹

Defining novel methods

The novel methods considered were based on a preliminary search of the literature and the expertise of the research team and included systems methods (e.g., agent-based modelling, system dynamics), least absolute shrinkage and selection operator (LASSO), machine learning algorithms, copulas, and data-driven statistical modelling approaches (e.g., treelet transformations, principal balances and coordinates). Novel methods could also include those that have been used previously in nutrition research if applied in new ways to characterize dietary patterns (e.g., linear programming used to model a modified dietary pattern rather than to test scenarios). Methods that were not considered to be novel were those that have been applied to assess dietary patterns in numerous studies and have been considered by prior reviews and commentaries,^{2,10,30} including regression, 'a priori' approaches such as investigator-driven indices, and routinely used data-driven approaches, including factor analysis and cluster analysis.^{10,31}

Identifying relevant studies

Articles were eligible for inclusion if they were: a primary research article; focused on dietary intake as an exposure or outcome, including examination of dietary patterns (i.e., multiple dietary components in combination rather than single nutrients, foods, or other dietary components); used at least one or more novel methods to characterize dietary patterns as described above; were published in English; and focused on humans. Ineligible studies included those focused on individual foods or human milk rather than dietary patterns, and commentaries and reviews.

Searches of three research databases, MEDLINE (via PubMed), the Cumulative Index to Nursing and Allied Health Literature (CINAHL), and Scopus, were conducted in March 2022. These health-focused, specialized, and multidisciplinary databases were selected based on consultation with a research librarian (JS) to ensure a range of possibly relevant study types were included. The search strategies were developed in consultation with the research librarian using keywords and subject headings to capture diet-related constructs (e.g., dietary intake, patterns, recommendations, feeding behaviour, food habits) and novel methods to characterize dietary patterns (e.g., machine learning, network science, system dynamics model). No date limits were applied to the searches, and articles were included until the end of the search in March 2022. The search strategies for MEDLINE, CINAHL, and Scopus are available in **Supplemental File 1**.

Study selection

Two independent reviewers (two of AP, AR, SH, SIK) screened each record at the title and abstract and full-text screening stages using Covidence,³² with one consistent reviewer (AP) participating throughout the entire process. At the title and abstract screening stage, an initial pilot screening (25 records) generated 100% agreement (AR and AP) and 92% agreement (AP and SH). A second pilot screening (100 records) generated 91% agreement (AR and AP) and 93% agreement (AP and SH). When applicable, discrepancies were discussed by

reviewers and if needed deferred to a third reviewer (SIK) for decision. Following pilot screening, the reviewers independently reviewed the remaining articles (96% agreement, Kappa = 0.83).

The reviewers were intentionally liberal during the title and abstract screening stage because of the breadth of possible novel methods. This required iteratively revisiting the inclusion criteria. For example, reduced rank regression was initially considered to be novel but was found to be prevalent in the literature based on title and abstract screening and was excluded during full-text review. Further, articles that used ‘a posteriori’ methods to identify dietary intake but did not specify the exact method in the title or abstract were included for full-text review.

Pilot screening of full-text reviews (50 records) generated 82% agreement (AR and AP) and 96% agreement (AP and SIK); after discrepancies were discussed, two reviewers independently screened the remaining full-text articles (93% agreement, Kappa = 0.60). The high agreement between reviewers but relatively low Cohen’s Kappa is described as Cohen’s paradox, with a larger number of studies excluded than included.^{33–35}

Data extraction

Data extraction was completed by TEW and JMH using a pre-specified Excel template, with all extracted data subsequently verified by LA. Data extraction fields (**Supplemental File 2**) included information pertaining to authorship, study title, journal, year of publication, funding source, contextual details (e.g., study location), sample size, and participant characteristics (e.g., age). Details relating to study methods (e.g., analysis input variables, measurement of dietary intake, analytic approaches) and results (e.g., findings related to dietary patterns and if applicable, health risk and outcomes) were also extracted.

Results

Summary of search

A total of 5274 unique articles were identified after removing duplicates. Of these, 436 were identified as potentially relevant based on the title and abstract review and underwent full-text screening (**Figure 1**). Studies excluded during full-text screening included those that did not include methods defined as novel, those that did not focus on dietary patterns, commentaries, narrative reviews, systematic reviews, studies that were not published in English, studies that were not conducted with humans, and theses/dissertations. A final pool of 24 articles describing 24 unique studies met the inclusion criteria.

Characteristics of included studies

Across the 24 included studies, data from 17 countries were represented (**Table 1**). Half of the studies were published between 2005 and 2019,^{36–47} and the remaining 12 were published between 2020 and March 2022.^{48–59} Three studies used data from subsets of the European Prospective Investigation into Cancer and Nutrition,^{36,45,46} two studies used waves of data from the National Health and Nutrition Examination Survey,^{54,58} and two studies used data from the ELSA-Brasil cohort study (**Table 2**).^{42,57} Sample sizes ranged from 250 to over 73,000 participants. Nineteen studies were conducted using data from cohort or cross-sectional studies and five studies applied a case-control design.

The majority (n=15) of studies used food frequency questionnaires to assess dietary intake.^{36,37,39–43,45,47–49,52,55–57} Six studies used 24-hour recalls,^{46,51,53,54,58,59} two studies used food records/diaries,^{44,50} and one study used a food frequency questionnaire and a 24-hour recall.³⁸ Among the studies using 24-hour recalls and records/diaries, one used data from a single recall that was combined with data from a food frequency questionnaire.³⁸ The remaining studies,

including records or recalls, averaged or combined data from two or more days of intake. Apart from averaging recalls or records, none of the included studies applied substantial efforts to mitigate measurement error present in dietary intake data. Several studies noted potential misreporting as a limitation, and only five studies specifically noted that findings may have been influenced by measurement error present in self-reported dietary assessment instruments.^{40,43,51,52,59}

Novel methods applied to identify dietary patterns

The type of methods used and how they were implemented to identify dietary patterns varied widely (**Table 3**). Nine studies applied approaches that have applications in machine learning, including classification models, neural networks, and probabilistic graphical models (**Table 4**).^{36,38,44–46,49,53,56,59} The earliest study included in this review was published in 2005 and applied neural networks to characterize dietary patterns.³⁸ Fifteen studies applied other novel methods, including latent class analysis, mutual information, and treelet transform.^{37,39–43,47,48,50–52,54,55,57,58} Two studies identified dietary patterns using more than one novel method.^{44,53} Five studies included comparisons of different novel methods, though these were typically versions of the same model.^{44,45,47,49,53} For example, Solans et al. compared three models for compositional data analysis and reported that the best-performing model incorporated both investigator- and data-driven methods.⁴⁷

In twelve studies, two to eight distinct dietary patterns, such as the ‘prudent’ pattern or ‘Western’ pattern, were identified using methods such as latent class analysis, treelet transform, random forest with classification tree analysis, and multivariate finite mixture models.^{36–38,41–43,48,50,51,54,57,58} Six studies applied network methods, including probabilistic graphical models and mutual information, to identify networks of dietary patterns among populations.^{45,46,49,52,55,59}

Dynamism, or how dietary patterns vary across time, was incorporated into four studies' characterization or analysis of dietary patterns. Three studies incorporated stratification by meals to consider dynamism.^{44,46,59} In two studies using graphical models, separate networks were created for each meal to provide insights into how patterns of intake vary throughout the day.^{46,59} Hearty and Gibney used decision trees and neural networks and ran models by meals based on 62 food groups to predict diet quality.⁴⁴ Additionally, one study considered dynamism by using ANOVA and chi-square tests to descriptively show how a variety of characteristics were associated with stable or changing dietary patterns characterized using latent class analysis.⁴³

Fourteen studies examined relationships between dietary patterns characterized using novel methods and variables indicative of health risk or outcomes, such as periodontitis, cardiovascular disease, and metabolic syndrome (Table 3).^{36–40,48–56} Six studies included longitudinal analysis of the relationship between dietary patterns and health outcomes.^{36,38,40,50,52,53} Most studies that examined health risk or outcomes first identified dietary patterns using a novel method and then investigated relationships with health outcomes using regression models.^{36–40,48,50,51,54} In contrast, some studies incorporated variables indicative of health outcomes or risk directly into the machine learning models.^{49,56} For example, Zhao et al.⁵⁶ applied Bayesian kernel machine regression, a machine learning model designed to incorporate high dimensional data, to jointly model the relationship between several dietary components and cardiovascular disease risk. Similarly, a paper by Hoang et al.⁴⁹ included health variables within mixed graphical models, though directionality of diet-health relationships could not be ascertained given the cross-sectional nature of the data. In two case-control studies, dietary patterns were identified using mutual information to estimate dietary pattern networks, with stratification by health outcomes.^{52,55}

Nineteen studies considered sociodemographic characteristics, such as sex, age, race/ethnicity, education, and income.^{36,37,39–43,45,48–58} In one case, sociodemographic characteristics were included in models used to characterize dietary patterns.⁴⁹ Two studies stratified by sociodemographic characteristics, examining dietary patterns by sex⁴⁵ or age groups.⁴² Studies that used case-control designs typically considered sociodemographic characteristics through matching.^{52,55} In the remaining studies that considered sociodemographic characteristics, these were incorporated in regression models to explore how dietary patterns characterized using novel methods were associated with health and other characteristics.

Two studies included comparisons of novel methods and traditional statistical approaches.^{36,57} While there was overlap between the patterns identified through novel and traditional approaches in these studies, the patterns identified differed across approaches. For instance, Biesbroek et al.³⁶ found that dietary patterns identified through reduced rank regression were more strongly associated with coronary artery disease compared to those identified through random forest with classification tree analysis.

Discussion

The application of novel methods to dietary pattern research is rapidly expanding, with the aim of better understanding their complexity and how they are related to health and other factors. Many studies used methods that characterize distinct dietary patterns based on the population being studied, such as the ‘prudent’ pattern or the ‘Western’ pattern. Most studies used cross-sectional data, limiting opportunities to examine the effect of dietary patterns on health.

Methods newly being applied in this field offer promising capacity to better understand the totality of dietary patterns and synergistic relationships among dietary components when compared with traditional approaches that do not assume synergy.^{4,14} Given the large variation in how dietary patterns were characterized using novel methods, multidimensionality and potential synergistic relationships between dietary components were considered and presented in a range of ways, from latent classes to networks. Several studies incorporated dynamism into their consideration of dietary patterns, though in most cases this was through stratification, for example, by meal, rather than through direct use of novel methods.^{43,44,46,59} In these cases, it was a combination of input variables, stratification by time, and the novel method that enabled explorations of dynamism.

The methods highlighted have a range of strengths and limitations for the characterization of dietary patterns. Methods that focused on the classification of distinct patterns allowed for the assessment of relationships between these patterns and health outcomes or other indicators of interest but explored the interrelationships between dietary components to a lesser degree.^{36–38,41–43,48,50,51,54,57,58} Other methods were designed to better consider synergistic relationships among dietary components, such as compositional data analysis, mutual information, or probabilistic graphical models, but required further analyses, such as the development of a score, to assess relationships with health outcomes.^{45–47,49,52,55,59}

There are trade-offs between novel and traditional methods that should be considered when contemplating the most appropriate methods for a given study. Though potential benefits such as a greater ability to discern multidimensionality may be desirable, these must be weighed against the implications for interpretability and computational costs. The application of novel methods may not always yield insights beyond those gained from traditional approaches. For example, Biesbroek et al.³⁶ found that random forest models did not outperform reduced rank regression when examining associations of dietary patterns with coronary artery disease.

Conversely, a study that was not included in this review because it first identified dietary patterns using a traditional method—principal component analysis—found that machine learning algorithms were better able to classify the identified dietary patterns according to cardiometabolic risk compared to traditional approaches.⁶⁰

Several sociodemographic characteristics are indicators of systemic health inequity and have been shown to be associated with dietary patterns among populations.^{61–63} The degree to which studies incorporated sociodemographic characteristics into their consideration of dietary patterns or relationships between dietary patterns and health varied, with adjusted regression models applied after dietary patterns were characterized as the most common approach. Consistent with nutrition research more broadly,^{63–65} there was little consideration of possible interactions among sociodemographic characteristics in relation to dietary patterns. Methods particularly suited to pattern recognition and complexity could be leveraged to simultaneously explore potential joint relationships among facets of social identity and dietary patterns⁶⁶ and advance our understanding of how broader systems of oppression and intersecting characteristics contribute to dietary patterns.⁶²

Beyond the inclusion of sociodemographic characteristics in models, considering equity from the beginning of study design is a critical consideration given potential bias in data and algorithms that can have immense implications for those who already experience inequities, such as structural racism.^{67–69} The included studies did not explicitly discuss the incorporation of equity into study design, and many conducted secondary analyses of existing datasets. The use of directed acyclic graphs has been identified as a potential solution to mitigate some possible issues with bias through careful model design⁶⁷ and has been applied in other domains of nutrition research using novel methods.¹⁴ Engaging individuals with lived experience and the integration of interdisciplinary teams with broad expertise that can combine content knowledge with data-driven approaches can help to mitigate potential bias in algorithms.⁶⁶

The level of description of methods varied and it was sometimes challenging to decipher the specifics of how novel methods were applied. Although the Strengthening the Reporting of Observational Studies in Epidemiology—Nutritional Epidemiology (STROBE-nut) reporting guidelines provide guidance for transparently reporting nutritional epidemiology and dietary assessment research,⁷⁰ it was not designed specifically for the methods used in the studies considered in this review and the ways in which they are being applied in dietary patterns research. Other reporting guidelines, such as the Consolidated Standards of Reporting Trials (CONSORT), have been extended to consider the application of artificial intelligence (AI).⁷¹ Motivations related to the extension of CONSORT included inadequate reporting of studies using AI and the lack of full consideration of potential sources of bias specific to AI within existing reporting guidelines.⁷¹ Relevant items added to CONSORT-AI pertain to the role of AI in the study, the nature of the data used in AI systems, and how humans interacted with AI systems, for example.⁷¹ The extension of reporting guidelines such as STROBE-nut to consider applications of AI, including machine learning, and other methods that are becoming more commonly used, may facilitate consistent and complete reporting and improved comparability of studies. Reporting guidelines should continue to emphasize strategies applied to mitigate measurement error in dietary intake data,⁷⁰ as studies using novel methods are not immune to the effects of error on findings.⁷² Along with reporting guidelines, the development of tailored quality appraisal tools may facilitate synthesis of high-quality evidence to inform recommendations about dietary patterns and health.

This review provides a snapshot of a rapidly evolving field,^{73,74} with the involvement of an interdisciplinary team of researchers lending to a robust consideration of emerging methods in dietary patterns research. While prior reviews have provided perspectives on the potential applications of machine learning within the field of nutrition,^{19–21} this review considered dietary patterns in particular, as well as considering approaches beyond machine learning that have not

been traditionally used in this area, broadening the scope compared to prior reviews.^{22,75} The search terms were informed by preliminary searching, though it is unlikely that all relevant articles applying novel methods to characterize dietary patterns were captured. This is partially driven by the wide range of descriptors used for these methods and the lack of reporting standards. As well, determining whether a method is novel is somewhat subjective. Methods such as factor analysis and principal component analysis once revolutionized dietary pattern analysis, providing data-driven approaches to identify patterns.¹⁵ Now, they are widely applied and recognized as limited in their capabilities to capture complexity compared to some newer approaches. Further, the search terms skewed toward multidimensionality versus dynamism, potentially overlooking some studies focusing on variation of dietary patterns over time or across eating occasions. Nonetheless, this review documents an acceleration of the application of a range of novel methods to dietary patterns research and captures a broad scope of methods being used to characterize these patterns, highlighting the need for researchers to develop the lexicon and knowledge needed to interpret the emerging literature.

Conclusion

The findings of this review indicate a strong motivation to apply novel methods, including but not limited to machine learning, to improve understanding of dietary patterns and how they relate to health and other factors. The application of these methods may help us to learn about complex relationships that may not be possible to discern through traditional approaches. However, these methods may not be suitable for every question and do not necessarily overcome the limitations of more traditional approaches.

Given the proliferation of these methods, it is becoming increasingly worthwhile for nutrition researchers to have at least a basic understanding of novel methods such as machine

learning and latent class analysis, so they can interpret the results of emerging studies. The development and implementation of reporting guidelines and quality appraisal mechanisms for studies that apply novel methods may improve the capacity for synthesis of evidence generated to inform strategies that promote improved population health and well-being.

Acknowledgements: We thank research librarian Jackie Stapleton (JS) of the University of Waterloo for support with the search strategy.

Conflict of Interest: RML is a statistical editor for the British Journal of Nutrition. Other authors have none to declare.

Funding: This review was funded by the Canadian Institutes of Health Research, a University of Waterloo Research Incentive Fund award, an Ontario Ministry of Research and Innovation Early Researcher Award held by SIK, and Microsoft AI for Good. RML is funded by a National Health and Medical Research Council Emerging Leadership Fellowship (APP1175250). LMB was funded by the National Institutes of Health (R01 HD102313, MPI Bodnar LM, Naimi AI).

Contributions: SIK conceived of the review and planned it with the co-authors; AR, AP, SH, SIK conducted the search and screening; JMH and TEW conducted extraction; LA conducted verification; JMH led the first draft of the manuscript with support from AP to write the methods; all co-authors provided critical input to the manuscript and all co-authors read and approved the final manuscript.

References

1. Afshin A, Sur PJ, Fay KA, et al. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*. 2019;393(10184):1958-1972. doi:10.1016/S0140-6736(19)30041-8
2. English LK, Ard JD, Bailey RL, et al. Evaluation of dietary patterns and all-cause mortality: A systematic review. *JAMA network open*. 2021;4(8). doi:10.1001/JAMANETWORKOPEN.2021.22277
3. Mozaffarian D, Rosenberg I, Uauy R. History of modern nutrition science—implications for current research, dietary guidelines, and food policy. *BMJ*. 2018;361. doi:10.1136/BMJ.K2392
4. Reedy J, Subar AF, George SM, Krebs-Smith SM. Extending methods in dietary patterns research. *Nutrients*. 2018;10(5). doi:10.3390/NU10050571
5. Schulz CA, Oluwagbemigun K, Nöthlings U. Advances in dietary pattern analysis in nutritional epidemiology. *Eur J Nutr*. 2021;60(8):4115-4130. doi:10.1007/s00394-021-02545-9
6. Herforth A, Arimond M, Álvarez-Sánchez C, Coates J, Christianson K, Muehlhoff E. A global review of food-based dietary guidelines. *Advances in Nutrition*. 2019;10(4):590. doi:10.1093/ADVANCES/NMY130
7. National Academies of Sciences, Engineering, and Medicine; Health and Medicine Division; Food and Nutrition Board; Committee to Review the Process to Update the Dietary Guidelines for Americans. *Redesigning the Process for Establishing the Dietary Guidelines for Americans*. National Academies Press; 2017. doi:10.17226/24883
8. Imamura F, Micha R, Khatibzadeh S, et al. Dietary quality among men and women in 187 countries in 1990 and 2010: a systematic assessment. *The Lancet Global Health*. 2015;3(3):e132-e142. doi:10.1016/S2214-109X(14)70381-X
9. Delormier T, Frohlich K, Potvin L. Food and eating as social practice – understanding eating patterns as social phenomena and implications for public health. *Sociology of Health & Illness*. 2009;31(2):215-228. doi:10.1111/j.1467-9566.2008.01128.x
10. Ocké MC. Evaluation of methodologies for assessing the overall diet: dietary quality scores and dietary pattern analysis. *Proceedings of the Nutrition Society*. 2013;72(2):191-199. doi:10.1017/S0029665113000013
11. Shams-White MM, Pannucci TE, Lerman JL, et al. Healthy Eating Index-2020: Review and update process to reflect the Dietary Guidelines for Americans, 2020-2025. *J Acad Nutr Diet*. 2023;123(9):1280-1288. doi:10.1016/j.jand.2023.05.015
12. Brassard D, Munene LAE, Pierre SS, et al. Development of the Healthy Eating Food Index (HEFI)-2019 measuring adherence to Canada’s Food Guide 2019 recommendations on healthy food choices. *Applied Physiology, Nutrition, and Metabolism*. 2022;47(5):595-610. doi:10.1139/APNM-2021-0415

13. Kirkpatrick SI, Reedy J, Krebs-Smith SM, et al. Applications of the Healthy Eating Index for surveillance, epidemiology, and intervention research: Considerations and caveats. *Journal of the Academy of Nutrition and Dietetics*. 2018;118(9):1603. doi:10.1016/J.JAND.2018.05.020
14. Bodnar LM, Cartus AR, Kirkpatrick SI, et al. Machine learning as a strategy to account for dietary synergy: an illustration based on dietary intake and adverse pregnancy outcomes. *The American Journal of Clinical Nutrition*. 2020;111(6):1235-1243. doi:10.1093/AJCN/NQAA027
15. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Current Opinion in Lipidology*. 2002;13(1):3-9. doi:10.1097/00041433-200202000-00002
16. Michels KB, Schulze MB. Can dietary patterns help us detect diet-disease associations? *Nutr Res Rev*. 2005;18(2):241-248. doi:10.1079/NRR2005107
17. Reedy J, Krebs-Smith SM, Hammond RA, Hennessy E. Advancing the science of dietary patterns research to leverage a complex systems approach. *Journal of the Academy of Nutrition and Dietetics*. 2017;117(7):1019-1022. doi:10.1016/J.JAND.2017.03.008
18. National Academies of Sciences, Engineering, and Medicine; Health and Medicine Division; Food and Nutrition Board; Alice Vorosmarti and Joe Alper, Rapporteurs. *The Role of Advanced Computation, Predictive Technologies, and Big Data Analytics in Food and Nutrition Research*. National Academies Press; 2024. Accessed May 10, 2024. <https://www.nationalacademies.org/our-work/the-role-of-advanced-computation-predictive-technologies-and-big-data-analytics-in-research-related-to-food-and-nutrition-a-workshop-series>
19. Kirk D, Kok E, Tufano M, Tekinerdogan B, Feskens EJM, Camps G. Machine learning in nutrition research. *Advances in Nutrition*. 2022;13(6):2573-2589. doi:10.1093/ADVANCES/NMAC103
20. Morgenstern JD, Rosella LC, Costa AP, Souza RJ de, Anderson LN. Perspective: Big data and machine learning could help advance nutritional epidemiology. *Advances in Nutrition*. 2021;12(3):621-631. doi:10.1093/ADVANCES/NMAA183
21. Côté M, Lamarche B. Artificial intelligence in nutrition research: perspectives on current and future applications. *Appl Physiol Nutr Metab*. Published online 2021:1-8. doi:10.1139/apnm-2021-0448
22. Oliveira Chaves L, Gomes Domingos AL, Louzada Fernandes D, Ribeiro Cerqueira F, Siqueira-Batista R, Bressan J. Applicability of machine learning techniques in food intake assessment: A systematic review. *Critical Reviews in Food Science and Nutrition*. 2023;63(7):902-919. doi:10.1080/10408398.2021.1956425
23. Le Glaz A, Haralambous Y, Kim-Dufor DH, et al. Machine learning and natural language processing in mental health: Systematic review. *J Med Internet Res*. 2021;23(5):e15708. doi:10.2196/15708
24. Payedimarri AB, Concina D, Portinale L, et al. Prediction Models for Public Health Containment Measures on COVID-19 Using Artificial Intelligence and Machine Learning: A

- Systematic Review. *Int J Environ Res Public Health*. 2021;18(9):4499.
doi:10.3390/ijerph18094499
25. Sidey-Gibbons JAM, Sidey-Gibbons CJ. Machine learning in medicine: a practical introduction. *BMC Medical Research Methodology*. 2019;19(1):64. doi:10.1186/s12874-019-0681-4
 26. Morgenstern JD, Buajitti E, O'Neill M, et al. Predicting population health with machine learning: a scoping review. *BMJ Open*. 2020;10(10):e037860. doi:10.1136/BMJOPEN-2020-037860
 27. Chapter 11: Scoping reviews. In: *JBI Manual for Evidence Synthesis*. JBI; 2020.
doi:10.46658/JBIMES-20-12
 28. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology*. 2005;8(1):19-32.
doi:10.1080/1364557032000119616
 29. Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med*. 2018;169(7):467-473. doi:10.7326/M18-0850
 30. Newby PK, Tucker KL. Empirically derived eating patterns using factor or cluster analysis: A review. *Nutrition Reviews*. 2004;62(5):177-203. doi:10.1111/j.1753-4887.2004.tb00040.x
 31. Krebs-Smith SM, Subar AF, Reedy J. Examining dietary patterns in relation to chronic disease: Matching measures and methods to questions of interest. *Circulation*. 2015;132(9):790-793. doi:10.1161/CIRCULATIONAHA.115.018010
 32. Covidence - Better systematic review management. Covidence. Accessed September 12, 2023. <https://www.covidence.org/>
 33. Gwet KL. Computing inter-rater reliability and its variance in the presence of high agreement. *British Journal of Mathematical and Statistical Psychology*. 2008;61(1):29-48.
doi:10.1348/000711006X126600
 34. Cicchetti DV, Feinstein AR. High agreement but low kappa: II. Resolving the paradoxes. *Journal of Clinical Epidemiology*. 1990;43(6):551-558. doi:10.1016/0895-4356(90)90159-M
 35. Belur J, Tompson L, Thornton A, Simon M. Interrater reliability in systematic review methodology: Exploring variation in coder decision-making. *Sociological Methods & Research*. 2021;50(2):837-865. doi:10.1177/0049124118799372
 36. Biesbroek S, van der A DL, Brosens MCC, et al. Identifying cardiovascular risk factor-related dietary patterns with reduced rank regression and random forest in the EPIC-NL cohort. *Am J Clin Nutr*. 2015;102(1):146-154. doi:10.3945/ajcn.114.092288
 37. Fonseca MJ, Gaio R, Lopes C, Santos AC. Association between dietary patterns and metabolic syndrome in a sample of Portuguese adults. *Nutr J*. 2012;11:64.
doi:10.1186/1475-2891-11-64

38. Jamiołkowski J, Szpak A, Pawłowska D. Dietary habits of men from Podlasie region of Poland in the years 1987-1998 analysed with self-organizing neural networks. *Rocz Akad Med Białymst.* 2005;50 Suppl 1:220-224.
39. Oliveira A, Rodríguez-Artalejo F, Gaio R, Santos AC, Ramos E, Lopes C. Major habitual dietary patterns are associated with acute myocardial infarction and cardiovascular risk markers in a southern European population. *J Am Diet Assoc.* 2011;111(2):241-250. doi:10.1016/j.jada.2010.10.042
40. Wu Y, Sánchez BN, Goodrich JM, et al. Dietary exposures, epigenetics and pubertal tempo. *Environ Epigenet.* 2019;5(1):dvz002. doi:10.1093/eep/dvz002
41. Affret A, Severi G, Dow C, et al. Socio-economic factors associated with a healthy diet: results from the E3N study. *Public Health Nutr.* 2017;20(9):1574-1583. doi:10.1017/S1368980017000222
42. Bezerra IN, Bahamonde NMSG, Marchioni DML, et al. Generational differences in dietary pattern among Brazilian adults born between 1934 and 1975: a latent class analysis. *Public Health Nutr.* 2018;21(16):2929-2940. doi:10.1017/S136898001800191X
43. Harrington JM, Dahly DL, Fitzgerald AP, Gilthorpe MS, Perry IJ. Capturing changes in dietary patterns among older adults: a latent class analysis of an ageing Irish cohort. *Public Health Nutr.* 2014;17(12):2674-2686. doi:10.1017/S1368980014000111
44. Hearty AP, Gibney MJ. Analysis of meal patterns with the use of supervised data mining techniques--artificial neural networks and decision trees. *Am J Clin Nutr.* 2008;88(6):1632-1642. doi:10.3945/ajcn.2008.26619
45. Iqbal K, Buijsse B, Wirth J, Schulze MB, Floegel A, Boeing H. Gaussian graphical models identify networks of dietary intake in a German adult population. *The Journal of Nutrition.* 2016;146(3):646-652. doi:10.3945/JN.115.221135
46. Schwedhelm C, Knüppel S, Schwingshackl L, Boeing H, Iqbal K. Meal and habitual dietary networks identified through Semiparametric Gaussian Copula Graphical Models in a German adult population. *PLoS One.* 2018;13(8):e0202936. doi:10.1371/journal.pone.0202936
47. Solans M, Coenders G, Marcos-Gragera R, et al. Compositional analysis of dietary patterns. *Stat Methods Med Res.* 2019;28(9):2834-2847. doi:10.1177/0962280218790110
48. Dalmartello M, Decarli A, Ferraroni M, et al. Dietary patterns and oral and pharyngeal cancer using latent class analysis. *Int J Cancer.* 2020;147(3):719-727. doi:10.1002/ijc.32769
49. Hoang T, Lee J, Kim J. Network analysis of demographics, dietary intake, and comorbidity interactions. *Nutrients.* 2021;13(10):3563. doi:10.3390/NU13103563/S1
50. Hose AJ, Pagani G, Karvonen AM, et al. Excessive unbalanced meat consumption in the first year of life increases asthma risk in the PASTURE and LUKAS2 birth cohorts. *Front Immunol.* 2021;12:651709. doi:10.3389/fimmu.2021.651709

51. Madeira T, Severo M, Oliveira A, Gorjão Clara J, Lopes C. The association between dietary patterns and nutritional status in community-dwelling older adults-the PEN-3S study. *Eur J Clin Nutr.* 2021;75(3):521-530. doi:10.1038/s41430-020-00745-w
52. Samieri C, Sonawane AR, Lefèvre-Arbogast S, Helmer C, Grodstein F, Glass K. Using network science tools to identify novel diet patterns in prodromal dementia. *Neurology.* 2020;94(19):e2014-e2025. doi:10.1212/WNL.00000000000009399
53. Shang X, Li Y, Xu H, et al. Leading dietary determinants identified using machine learning techniques and a healthy diet score for changes in cardiometabolic risk factors in children: a longitudinal analysis. *Nutr J.* 2020;19(1):105. doi:10.1186/s12937-020-00611-2
54. Wright DM, McKenna G, Nugent A, Winning L, Linden GJ, Woodside JV. Association between diet and periodontitis: a cross-sectional study of 10,000 NHANES participants. *Am J Clin Nutr.* 2020;112(6):1485-1491. doi:10.1093/ajcn/nqaa266
55. Xia Y, Zhao Z, Zhang S, et al. Complex dietary topologies in non-alcoholic fatty liver disease: A network science analysis. *Front Nutr.* 2020;7:579086. doi:10.3389/fnut.2020.579086
56. Zhao Y, Naumova EN, Bobb JF, Claus Henn B, Singh GM. Joint associations of multiple dietary components with cardiovascular disease risk: A machine-learning approach. *Am J Epidemiol.* 2021;190(7):1353-1365. doi:10.1093/aje/kwab004
57. de Almeida Alves M, Molina MDCB, da Fonseca M de JM, Lotufo PA, Benseñor IM, Marchioni DML. Different statistical methods identify similar population-specific dietary patterns: an analysis of Longitudinal Study of Adult Health (ELSA-Brasil). *Br J Nutr.* 2022;128(11):2249-2257. doi:10.1017/S0007114522000253
58. Farmer N, Lee LJ, Powell-Wiley TM, Wallen GR. Cooking frequency and perception of diet among US adults are associated with US healthy and healthy Mediterranean-style dietary related classes: A latent class profile analysis. *Nutrients.* 2020;12(11):3268. doi:10.3390/nu12113268
59. Schwedhelm C, Lipsky LM, Shearrer GE, et al. Using food network analysis to understand meal patterns in pregnant women with high and low diet quality. *International Journal of Behavioral Nutrition and Physical Activity.* 2021;18(1):101. doi:10.1186/s12966-021-01172-1
60. Panaretos D, Koloverou E, Dimopoulos AC, et al. A comparison of statistical and machine-learning techniques in evaluating the association between dietary patterns and 10-year cardiometabolic risk (2002-2012): the ATTICA study. *The British Journal of Nutrition.* 2018;120(3):326-334. doi:10.1017/S0007114518001150
61. Hanson KL, Connor LM. Food insecurity and dietary quality in US adults and children: A systematic review. *Am J Clin Nutr.* 2014;100(2):684-692. doi:10.3945/ajcn.114.084525
62. Doan N, Olstad DL, Vanderlee L, Hammond D, Wallace M, Kirkpatrick SI. Investigating the intersections of racial identity and perceived income adequacy in relation to dietary quality among adults in Canada. *The Journal of Nutrition.* 2022;13(152):67S-75S. doi:10.1093/JN/NXAC076

63. Hiza HAB, Casavale KO, Guenther PM, Davis CA. Diet quality of Americans differs by age, sex, race/ethnicity, income, and education level. *Journal of the Academy of Nutrition and Dietetics*. 2013;113(2):297-306. doi:10.1016/J.JAND.2012.08.011
64. Olstad DL, Nejatnamini S, Victorino C, Kirkpatrick SI, Minaker LM, McLaren L. Trends in socioeconomic inequities in diet quality between 2004 and 2015 among a nationally representative sample of children in Canada. *The Journal of Nutrition*. 2021;151(12):3781-3794. doi:10.1093/JN/NXAB297
65. Brassard D, Munene LAE, Pierre SS, et al. Evaluation of the Healthy Eating Food Index (HEFI)-2019 measuring adherence to Canada's Food Guide 2019 recommendations on healthy food choices. *Applied Physiology, Nutrition, and Metabolism*. 2022;47(5). doi:10.1139/APNM-2021-0416
66. Wang A, Ramaswamy VV, Russakovsky O. Towards intersectionality in machine learning: Including more identities, handling underrepresentation, and performing evaluation. In: *Proceedings of the 2022 ACM Conference on Fairness, Accountability, and Transparency*. FAccT '22. Association for Computing Machinery; 2022:336-349. doi:10.1145/3531146.3533101
67. Robinson WR, Renson A, Naimi AI. Teaching yourself about structural racism will improve your machine learning. *Biostatistics*. 2020;21(2):339-344. doi:10.1093/biostatistics/kxz040
68. Rojas JC, Fahrenbach J, Makhni S, et al. Framework for integrating equity into machine learning models: A case study. *Chest*. 2022;161(6):1621-1627. doi:10.1016/j.chest.2022.02.001
69. Rajkomar A, Hardt M, Howell MD, Corrado G, Chin MH. Ensuring fairness in machine learning to advance health equity. *Annals of Internal Medicine*. 2018;169(12):866-872. doi:10.7326/M18-1990
70. Lachat C, Hawwash D, Ocké MC, et al. Strengthening the Reporting of Observational Studies in Epidemiology-Nutritional Epidemiology (STROBE-nut): An extension of the STROBE statement. *PLOS Medicine*. 2016;13(6). doi:10.1371/JOURNAL.PMED.1002036
71. Liu X, Cruz Rivera S, Moher D, et al. Reporting guidelines for clinical trial reports for interventions involving artificial intelligence: the CONSORT-AI extension. *Nature Medicine*. 2020;26(9):1364-1374. doi:10.1038/s41591-020-1034-x
72. Spicker D, Nazemi A, Hutchinson J, et al. Challenges for predictive modeling with neural network techniques using error-prone dietary intake data. Published online November 15, 2023. doi:10.48550/arXiv.2311.09338
73. Lampignano L, Tatoli R, Donghia R, et al. Nutritional patterns as machine learning predictors of liver health in a population of elderly subjects. *Nutr Metab Cardiovasc Dis*. 2023;33(11):2233-2241. doi:10.1016/j.numecd.2023.07.009
74. Slurink IA, Corpeleijn E, Bakker SJ, et al. Dairy consumption and incident prediabetes: prospective associations and network models in the large population-based Lifelines Study. *Am J Clin Nutr*. 2023;118(6):1077-1090. doi:10.1016/j.ajcnut.2023.10.002

75. Kirk D, Catal C, Tekinerdogan B. Precision nutrition: A systematic literature review. *Computers in Biology and Medicine*. 2021;133:104365. doi:10.1016/j.compbiomed.2021.104365
76. Breiman L. Random Forests. *Machine Learning*. 2001;45(1):5-32. doi:10.1023/A:1010933404324
77. Song Y yan, Lu Y. Decision tree methods: applications for classification and prediction. *Shanghai Arch Psychiatry*. 2015;27(2):130-135. doi:10.11919/j.issn.1002-0829.215044
78. Grossi E, Buscema M. Introduction to artificial neural networks. *European Journal of Gastroenterology & Hepatology*. 2007;19(12):1046. doi:10.1097/MEG.0b013e3282f198a0
79. Bianchi D, Calogero R, Tirozzi B. Kohonen neural networks and genetic classification. *Mathematical and Computer Modelling*. 2007;45(1):34-60. doi:10.1016/j.mcm.2006.04.004
80. Drton M, Maathuis MH. Structure learning in graphical modeling. <https://doi.org/10.1146/annurev-statistics-060116-053803>. 2017;4:365-393. doi:10.1146/ANNUREV-STATISTICS-060116-053803
81. Altenbuchinger M, Weihs A, Quackenbush J, Grabe HJ, Zacharias HU. Gaussian and Mixed Graphical Models as (multi-)omics data analysis tools. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms*. 2020;1863(6):194418. doi:10.1016/J.BBAGRM.2019.194418
82. Koller D, Friedman N. *Probabilistic Graphical Models: Principles and Techniques*. MIT Press; 2009.
83. Liu H, Han F, Yuan M, Lafferty J, Wasserman L. High-dimensional semiparametric Gaussian copula graphical models. *The Annals of Statistics*. 2012;40(4):2293-2326.
84. Freund Y, Schapire RE. A decision-theoretic generalization of on-line learning and an application to boosting. *Journal of Computer and System Sciences*. 1997;55(1):119-139. doi:10.1006/jcss.1997.1504
85. Bobb JF, Valeri L, Claus Henn B, et al. Bayesian kernel machine regression for estimating the health effects of multi-pollutant mixtures. *Biostatistics*. 2015;16(3):493-508. doi:10.1093/biostatistics/kxu058
86. Aitchison J. The statistical analysis of compositional data. *Journal of the Royal Statistical Society Series B (Methodological)*. 1982;44(2):139-177.
87. Rodrigues LA, Daunis-i-Estadella P, Figueras G, Henestrosa S. Flying in compositional morphospaces: evolution of limb proportions in flying vertebrates. In: *Compositional Data Analysis: Theory and Applications*. ; 2011:235-254. doi:10.1002/9781119976462.ch17
88. Zhang CH, Huang J. The sparsity and bias of the Lasso selection in high-dimensional linear regression. *The Annals of Statistics*. 2008;36(4):1567-1594.

89. Oliveira A, Lopes C, Torres D, Ramos E, Severo M. Application of a latent transition model to estimate the usual prevalence of dietary patterns. *Nutrients*. 2021;13(1):133. doi:10.3390/nu13010133
90. Fahey MT, Thane CW, Bramwell GD, Coward WA. Conditional Gaussian mixture modelling for dietary pattern analysis. *Journal of the Royal Statistical Society Series A (Statistics in Society)*. 2007;170(1):149-166.
91. Cover T, Thomas J. *Elements of Information Theory*. Wiley; 2005. Accessed August 23, 2023. <https://books-scholarsportal-info.proxy.lib.uwaterloo.ca/en/read?id=/ebooks/ebooks2/pda/2011-12-01/1/13568.9780471748816#page=1>
92. Gorst-Rasmussen A, Dahm C, Dethlefsen C, Scheike T, Overvad K. Exploring dietary patterns by using the treelet transform. *Am J Epidemiol*. 2011;173:1097-1104. doi:10.1093/aje/kwr060
93. Lee AB, Nadler B, Wasserman L. Treelets—An adaptive multi-scale basis for sparse unordered data. *The Annals of Applied Statistics*. 2008;2(2):435-471. doi:10.1214/07-AOAS137




Figure 1. PRISMA diagram illustrating the screening process for a scoping review exploring innovative methods for the analysis of dietary intake data and characterization of dietary patterns.

Table 1: Study characteristics across included studies applying novel methods to characterize dietary patterns

Characteristic	Number of articles
Country of study population [§]	
Austria	1
Brazil	2
China	2
Finland	1
France	3
Germany	3
Italy	1
Mexico	1
Netherlands	1
Northern Ireland	1
Poland	1
Portugal	3
Republic of Ireland	2
South Korea	1
Spain	1
Switzerland	1
USA	3
Year published	
2005-2009	2
2010-2014	3
2015-2019	7
2020-2022	12
Included health risk or outcomes	
Health risk/outcome (cross-sectional)	8
Health outcome (longitudinal)	6
No	10

[§]Some studies included more than one country

Table 2: Characteristics of studies (n=24) identified in a scoping review of novel analytic methods to characterize dietary patterns

Authors, date, country(s)	Study aims	n	Study design and data source	Participant characteristics	Diet assessment method(s)
Affret et al., 2017, France ⁴¹	To characterize the relationship of dietary patterns with socioeconomic factors	73 031	Cohort; E3N cohort ^a	Adults in France, females only, mean age 52.9 y	FFQ ^b
Bezerra et al., 2018, Brazil ⁴²	To examine how dietary patterns differ among generations in a Brazilian cohort	15 069	Cross-sectional analysis of cohort; ELSA-Brasil cohort ^c	Adults in Brazil, aged 35-74 y	FFQ
Biesbroek et al., 2015, Netherlands ³⁶	To compare if two methods (traditional vs. novel) derived differential dietary patterns and associations with health risk	34644	Cohort; EPIC ^d Prospect and EPIC MORGEN	Adults in the Netherlands, male and female, aged 20-70 y	FFQ
Dalmartello et al., 2020, Italy ⁴⁸	To identify dietary patterns associated with oropharyngeal cancer	946 cases, 2492 controls	Case-control; Unnamed study	Adults in Italy, admitted to major hospitals, 19-82 y	FFQ
De Almeida et al., 2022, Brazil ⁵⁷	To investigate how a variety of approaches identify dietary patterns in the same dataset	12 816	Cohort; ELSA-Brasil cohort	Adults in Brazil, aged 35-74 y	FFQ
Farmer et al., 2020, USA ⁵⁸	To explore how latent class analysis groups individuals into dietary patterns	11 481	Repeat cross-sectional; NHANES ^e 2007-2010	Adults in the USA, male and female, ≥19 y	Up to two 24HDRs ^f (per day)

Fonseca et al., 2012, Portugal ³⁷	To investigate the association between dietary patterns and metabolic syndrome	2167	Cross-sectional analysis of cohort study; EPIPorto cohort	Adults in Portugal, male and female, age not specified	FFQ
Harrington et al., 2014, Republic of Ireland ⁴³	To investigate changes in dietary patterns across 10 years among Irish adults	925 (baseline) 320 (follow-up)	Cohort; Cork and Kerry Diabetes and Heart Disease Study	Adults in the Republic of Ireland, 50-69 y	FFQ
Hearty & Gibney, 2008, Republic of Ireland & Northern Ireland ⁴⁴	To compare the ability of two supervised machine learning techniques to predict diet quality based on dietary patterns	1379	Cross-sectional; North-South Ireland Food Consumption Survey	Adults in Northern Ireland and the Republic of Ireland, aged 18-64 y	7-day food diary
Hoang et al., 2021, South Korea ⁴⁹	To describe interactions among dietary components, demographic characteristics, and chronic disease risk factors	7423	Cross-sectional analysis of cohort study; Cancer Screening Examination Cohort	Adults in South Korea, 30-70 y	FFQ
Hose et al., 2021, Austria, Finland, France, Germany, Switzerland ⁵⁰	To explore how feeding patterns across the first year of life relate to asthma in school-aged children	1246	Cohort; PASTURE ^g Birth Cohort and LUKAS2 ^h Cohort	Children in 5 European countries, birth cohort	Food diaries (weekly & monthly), dietary screener (monthly)
Iqbal et al., 2016, Germany ⁴⁵	To identify dietary patterns among a German adult population	27 120	Cross-sectional analysis of cohort; EPIC Potsdam	Adults in Germany, male and female, aged 35-65 y	FFQ

Jamiolkowski et al., 2005, Poland ³⁸	To identify dietary patterns and their associations with markers of health status	556	Cohort; Unnamed study	Adult males in Poland, age not specified	FFQ + 24HDR "24-hour consumption questionnaire"
Madeira et al., 2021, Portugal ⁵¹	To investigate associations between dietary patterns, malnutrition, and body mass index among older adults	849	Cross-sectional; PEN-3S Study ⁱ	Adults in Portugal, ≥65 y	Two 24HDR (per day)
Oliveira et al., 2011, Portugal ³⁹	To assess associations between a posteriori dietary patterns and health outcomes	3016 *820 cases, 2196 controls	Case-control; Unnamed study	Adults in Portugal, male and female, aged ≥ 18 years	FFQ
Samieri et al., 2020, France ⁵²	To model dietary patterns prior to the onset of dementia	1522	Case-control; 3-City Study	Adults in France, ≥65 y	FFQ
Schwedhelm et al., 2018, Germany ⁴⁶	To describe dietary networks using machine learning methods across meals and usual patterns of eating	814	Cross-sectional analysis of cohort; EPIC Potsdam	Adults in Germany, male and female, average age 65.5 y	Up to three 24HDR (per day)
Schwedhelm et al., 2021, USA ⁵⁹	To compare dietary patterns by meals among pregnant women with high and low diet quality	365	Cohort; Pregnancy Eating Attributes Study	Women receiving prenatal care in the USA, 18-44 y	Up to three 24HDRs, pooled
Shang et al., 2020, China ⁵³	To characterize top dietary components that are associated with changes in cardiometabolic risk	5676	Cohort; Unnamed study	Children in China, 6-13 y	Three 24HDRs, assessed at baseline and one year. Average of recalls

Solans et al., 2019, Spain ⁴⁷	factors among children To compare three compositional data analysis approaches to identify dietary patterns	3471	Case-control; Multi-Case Control Spain	Adults in Spain, male and female, aged 20-85 y (controls only)	FFQ
Wright et al., 2020, USA ⁵⁴	To examine how dietary patterns are associated with periodontitis	10 010	Repeat cross-sectional; NHANES 2009-2014	Adults in the USA, male and female, ≥30 y	Up to two 24HDRs Average of recalls
Wu et al., 2019, Mexico ⁴⁰	To explore how maternal and adolescent diet are associated with pubertal tempo	250 (baseline) 222 (follow-up)	Cohort; ELEMENT cohort ⁱ	Children in Mexico enrolled in a birth cohort	FFQ
Xia et al., 2020, China ⁵⁵	To examine how dietary patterns differ between individuals with non-alcoholic fatty liver disease and controls	4086 *split 50/50 between cases and controls	Case-control; Tianjin Chronic Low-grade Systemic Inflammation and Health Cohort Study	Individuals in China	FFQ
Zhao et al., 2021, USA ⁵⁶	To explore the relationship between dietary components, dietary patterns, and cardiovascular disease	1928	Cross-sectional analysis of cohort; CARDIA cohort ^k	Adults in the USA, 18-30 y	FFQ

^aEtude Epidémiologique auprès de femmes de la Mutuelle Générale de l'Education Nationale

^bFood frequency questionnaire

^cBrazilian Longitudinal Study of Adult Health

^dEuropean Prospective Investigation into Cancer and Nutrition

^eNational Health and Nutrition Examination Survey

^f24 hour dietary recall

^gProtection Against Allergy: Study in Rural Environments

^hFinnish, rural-suburban birth cohort

ⁱPortuguese Elderly and Nutritional Status Surveillance System

^jEarly Life Exposure in Mexico to Environmental Toxicants

^kCoronary Artery Risk Development in Young Adults

Table 3: Description of dietary patterns (n=24) identified in a scoping review of novel analytic methods to characterize dietary patterns

<i>Studies that characterized dietary patterns to assess health outcomes/risk</i>					
Authors, date, country(s)	Novel method(s) to analyze diet	Dietary input variables	Characterization of dietary patterns	Consideration of health outcomes/risk	Consideration of sociodemographic characteristics[§]
Biesbroek et al., 2015, Netherlands ³⁶	Random forest with classification tree analysis	Food items were grouped into 35 food groups based on Dutch standard food groups, represented as % of total energy.	7 patterns were derived to explain cardiovascular disease risk factors: Prudent-like 1, Prudent-like 2, Traditional-like 1, Traditional-like 2, Traditional-like 3, Western-like 1, Western-like 2.	Investigated the association between dietary patterns and first occurrence of coronary artery disease, stroke, death from any cause through hazard ratios	All hazard ratios were adjusted for age and sex and one model additionally considered smoking status and education.
Dalmartello et al., 2020, Italy ⁴⁸	Latent class analysis	Items from the FFQ were grouped into 25 food groups and daily intake (g/d) was estimated using standard portion sizes.	Latent class analysis was used to identify distinct dietary patterns: Western, Prudent, Lower consumers- combination, Higher consumers- combination.	Investigated associations of dietary patterns with risk of oropharyngeal cancer using odds ratios.	Not stated how cases and controls were matched. Odds ratios were adjusted for age, sex, education, smoking status.
Fonseca et al., 2012, Portugal ³⁷	Multivariate finite mixture models	Items from the FFQ were grouped into 14 food groups based on similar nutritional characteristics.	Identified four dietary patterns among males and females. Males: healthy, fish, red meat and alcohol, intermediate. Females: healthy, low fruit and vegetable, red meat and alcohol, in transition to fast food.	Associations between metabolic syndrome and dietary patterns were assessed using crude and adjusted odds ratios	Identified dietary patterns by sex. Odds ratios were adjusted for age, education, smoking status.
Hoang et al., 2021, South Korea ⁴⁹	Gaussian graphical models and	16 food groups, assessed as g/d, were included in graphical models.	A network of intake was identified among the 16 food groups, with edges representing conditional	Graphical models also included comorbidity factors, such as	Sociodemographic characteristics including age, sex, marital status,

	mixed graphical models	These food groups were also included in a graphical model as a single dietary score.	dependencies between food groups. Food groups that were not connected by an edge were conditionally independent.	blood pressure or total cholesterol, to identify associations with dietary components.	education, and income were included in graphical models.
Hose et al., 2021, Austria, Finland, France, Germany, Switzerland ⁵⁰	Latent class analysis	Included 17 food items from 19-item screener that assessed time of introduction (screeners repeated at 2, 12, 18, 24, 36, 48, 60, 72 months of age) and frequency of consumption.	Four food introduction patterns were identified. Latent class 1 was characterized by daily fruit, vegetable, and milk consumption; Latent class 2 was characterized by daily fruit, vegetable, and meat and cereal consumption, but less frequent milk consumption; Latent class 3 was characterized by slower introduction to foods, a diverse range of food items, but less than daily fruit and vegetable consumption; Latent class 4 was similar to Latent class 3 but with quicker introduction to foods.	Linear or logistic/multinomial regression was used to estimate associations between latent class analysis food introduction patterns, possible determinants, and asthma outcomes.	Regression analyses adjusted for sex.
Jamiolkowski et al., 2005, Poland ³⁸	Kohonen neural networks	41-item FFQ was included as input.	Three dietary patterns were identified: 1- high frequency of margarine, fried vegetable oil, low consumption of butter, lard, bacon, most frequent consumption of poultry, fresh fish, fruit; 2- high frequency butter, lard,	Impact of dietary habits on blood serum lipids assessed using Kruskal-Wallis procedure.	No.

Madeira et al., 2021, Portugal ⁵¹	Latent class transition model	Items were grouped from 71 food items to 27 food groups.	giblets, smoked 'second-rate' meats, sweets, sugary beverages; 3- low frequency butter, margarine, milk, cheese. Latent class transition models identified two dietary patterns, labelled as a protein-foods pattern and a Mediterranean pattern.	Linear regression was used to examine relationships between identified dietary patterns and nutritional status and BMI	Regression models adjusted for sex, age, education.
Oliveira et al., 2011, Portugal ³⁹	Multivariate and univariate finite fixture models	Items from an 82-item FFQ were grouped into 14 food groups (g/d) based on nutritional similarities.	Distinct dietary patterns were identified among males and females using multivariate and univariate fixture models. Males: Healthy, Fish, Red meat and alcohol, intermediate intake. Females: Healthy, Low fruit and vegetables, red meat and alcohol, in transition to fast food.	Logistic regression was used to examine the association between dietary patterns and acute myocardial infarction (with and without confounders)	Identified dietary patterns by sex. Multivariate models included age, education, smoking status as confounders.
Samieri et al., 2020, France ⁵²	Mutual information	148-item FFQ grouped into 62 dietary variables.	A mutual information matrix was computed to estimate the network relationships between dietary components. Networks of intake were estimated among cases and controls. The case network showed one mainly connected network of foods, whereas the	Two mutual information networks were developed and compared – one for dementia cases and one for controls. A third network representing the differences	Nested case-control. Matched cases and controls by age, sex, and education.

			controls network was represented by several disconnected subnetworks.	between the cases' and controls' dietary patterns was developed.	
Shang et al., 2020, China ⁵³	Random forest, gradient boost machine	Foods were categorized into 26 food groups (% energy/d)	The components from baseline dietary intake served as inputs into machine learning algorithms to determine the most important components for changes in cardiometabolic disease risk. The most important components were then summed to create a Healthy Diet Score.	Changes in cardiometabolic risk were evaluated by subtracting the results at baseline from those at follow-up. The Healthy Diet Score was then tested in an external dataset.	Regression models adjusted for age, sex, household income, parental education.
Wright et al., 2020, USA ⁵⁴	Treelet transform	Food and Nutrient Database for Dietary Studies food groups were used as inputs.	Eight treelet components (dietary patterns) were identified that explained substantial variation. 1: diets high in salads, moderate vegetables, poultry seafood, fruit, water, tea. 2: loaded on milk, ready-to-eat cereal, bananas. 3: loaded on bread, luncheon meat, table fat, coffee, sugar. 4: loaded on fruit juice, fruit-flavoured drinks. 5: low intake bread, meat, table fats, high intake coffee and sugar.	Quantile regression was used to estimate associations between dietary patterns and periodontitis.	Adjusted models for sex, ethnicity, age, education, smoking status.

Wu et al., 2019, Mexico ⁴⁰	LASSO	From a 116-item FFQ among adolescents and 104-item FFQ among pregnant women, 21 dietary components were identified as potential methyl-donor rich foods for inclusion in LASSO models.	6: loaded on beef, potatoes, beans, corn. 7: loaded on candy, crackers, pretzels, low-calorie carbonated soft drinks. 8: loaded on pancakes, eggs, French toast. Seven food items were retained from LASSO models for the maternal diet and eight from the adolescent diet. An Epigenetics Associated Diet Score was created for methyl donor-rich foods, with weights for dietary components obtained from LASSO.	Time-to-event analysis with interval-censored regression models was used to investigate the relationship between Epigenetics Associated Diet Scores and pubertal timing.	Final models adjusted for household socioeconomic status (included material wellbeing, highest education level in household).
Xia et al., 2020, China ⁵⁵	Mutual information	100-item FFQ condensed to 25 food groups (g/d) based on similar characteristics.	Dietary patterns were identified using mutual information, identifying five primary dietary components in the cases and six in the controls.	Separate networks were developed for non-alcoholic fatty liver disease cases and controls and were compared.	Case-control. Used propensity score matching to match cases and controls. This process considered sex, age, education, income, smoking status, employment status. Cases that could not be matched closely enough were not included in analyses.

Zhao et al., 2021, USA ⁵⁶	Bayesian kernel machine regression	FFQ items were assigned to 166 food groups and were further grouped to 12 dietary factors based on evidence of associations with cardiovascular disease.	Bayesian kernel machine regression simultaneously modelled the relationship between the 12 dietary components of interest and cardiovascular disease risk, including the change in each dietary component and disease risk and the dose-response relationship of each dietary component.	Bayesian kernel machine regression was used to jointly consider the 12 dietary components and cardiovascular disease risk.	Balanced recruitment by sociodemographic characteristics. 10-year CVD calculation considers age, sex, race/ethnicity, smoking status. Additionally adjusted regression models for income and education.
--------------------------------------	------------------------------------	--	--	--	---

Studies that characterized dietary patterns (no health outcomes/risk)

Authors, date, country(s)	Analytic method(s)	Dietary input variables	Characterization of dietary patterns	Consideration of sociodemographic characteristics
Affret et al., 2017, France ⁴¹	Latent Class Analysis	Created 57 food groups from the 208 food items available in the FFQ.	Used Latent Class Analysis to obtain distinct dietary patterns from these groups, deriving a 'Healthy' and 'Less Healthy' pattern.	Used logistic regression to assess the individual and socioeconomic factors associated with 'healthy diet'. These factors included education, employment status, income, number of children, marital status, place of residence, size of city, deprivation index, geographical location of birth.
Bezerra et al., 2018, Brazil ⁴²	Latent Class Analysis	A 114-item FFQ was condensed to create 13 mutually exclusive groups based on nutritional content and level of processing (cereals and tubers and roots; skimmed milk and dairy products; white meats and fish; fruit;	Latent class models identified dietary patterns for the whole sample and each generation. Traditionalists: mixed pattern, restricted pattern, prudent pattern. Baby Boomers: mixed pattern, processed pattern, prudent pattern.	Assessed how dietary patterns differed by three birth cohorts (age).

		vegetables; legumes and nuts; whole milk and dairy products; red meats; processed meats; ultra-processed products; soft drinks and industrialized fruit drinks; sweets; coffee).	Generation X: mixed pattern, processed pattern, prudent pattern.	
De Almeida et al., 2022, Brazil ⁵⁷	Treelet transform	114-item FFQ condensed to 25 food groups based on nutritional similarity.	Used three analytic approaches to derive dietary patterns (treelet transform, factor analysis, reduced rank regression). Treelet transform identified three patterns: convenience, prudent, and rice and beans. Factor analysis identified three patterns: Convenience, Brazilian traditional, and prudent. Reduced rank regression identified three patterns but only retained one, the convenience pattern.	Assessed how dietary patterns differed by sociodemographic characteristics: age, sex, smoking status, education, race
Farmer et al., 2020, USA ⁵⁸	Latent class profile analysis	11 out of 12 components from the Healthy Eating Index-2010 served as inputs (1. total vegetable, 2. greens/beans, 3. total fruit, 4. whole	Distinct classes for total day intake and dinner intake were identified. Classes included (total day intake): Standard American Diet, Standard American Diet with low sodium, Healthy US,	Multinomial logistic regression used to assess how latent classes differed by sociodemographic characteristics, including age, sex/gender, race/ethnicity, marital status, education, employment status, poverty/income ratio, number of people in household.

		grains, 5. dairy, 6. total protein foods, 7. seafood and plant proteins, 8. fatty acid ratio, 9. sodium, 10. refined grains, 11. solid fats, alcohol and added sugars).	Healthy US with high vegetable, Healthy US low sodium.	
Harrington et al., 2014, Republic of Ireland ⁴³	Latent class analysis	Food items were aggregated from a 159-item and 167-item FFQ into 23 mutually exclusive food groups (g/d, ml/d)	Latent class analysis identified three dietary patterns: Western, Healthy, and low-energy.	Assessed relationships between latent classes and sociodemographic characteristics, including smoking status, education, marital status.
Hearty & Gibney, 2008, Republic of Ireland & Northern Ireland ⁴⁴	Artificial neural networks and decision trees	Food codes were aggregated to 62 food groups.	Food groups were provided to neural networks and decision trees to identify dietary patterns predictive of diet quality quintile by meal.	No.
Iqbal et al., 2016, Germany ⁴⁵	Gaussian graphical models (GGM); confirmed with Semiparametric Gaussian Copula graphical models	Items from the 148-item FFQ were grouped into 49 groupings and were log-transformed.	GGMs were applied to derive dietary patterns among males and females. Edges represented conditional dependencies between food groups. Food groups that were not connected by an edge were conditionally independent.	Identified sex-specific graphical networks.
Schwedhelm et al., 2018, Germany ⁴⁶	Semiparametric Gaussian Copula	Food codes were condensed to 39 food groups.	Semiparametric Gaussian Copula graphical models derived meal-specific networks. For example, the	No.

	graphical models		breakfast network was characterized by one primary network with five communities and included foods such as breakfast cereals, margarine, and eggs. Edges represented conditional dependencies between food groups. Food groups that were not connected by an edge were conditionally independent.	
Schwedhelm et al., 2021, USA ⁵⁹	Gaussian graphical model	Foods were grouped into 40 food groups based on the Food and Nutrient Database for Dietary Studies.	Diet quality was assessed using the Healthy Eating Index 2015, and participants were split into tertiles of HEI 2015 scores, to compare dietary patterns by meal of women in the low and high HEI tertile generated using Gaussian graphical models. Edges represented conditional dependencies between food groups. Food groups that were not connected by an edge were conditionally independent.	No.
Solans et al., 2019, Spain ⁴⁷	3 Compositional data analysis approaches: compositional PC analysis,	9 items (vegetables, fruit, legumes, seafood, cereals, meat, dairy, monounsaturated fats, and saturated	Three variations of compositional data analysis were used to explain the variance among the dietary	No.

balances,
principal
balances

fats) from the 140-
item FFQ were
included as inputs.

components and identify
dietary patterns.

§We considered sociodemographic characteristics that are related to social position or are indicators of equity including age, sex, gender, race/ethnicity, marital status, education, employment status, smoking status as examples. We did not include physical activity, BMI, or alcohol consumption.

Table 4: Novel methods applied to identify dietary patterns across included studies

Method of analysis	Description	References
Methods with applications in machine learning		
Classification models	Algorithms that can be used for prediction and classification. ^{76,77}	
Random forest	The random forest is generated from a wide array of trees that are randomly sampled and averaged. Random forest models can additionally be used on regressors. ⁷⁶	(Biesbroek et al. 2015; Shang et al. 2020)
Decision tree	Trees include nodes (variables) and branches, representing possible outcomes from a node. This forms an 'inverted tree', beginning with a root node and branching to many possible end nodes. ⁷⁷	(Hearty and Gibney 2008)
Neural network	The structure of this algorithm was based on the human brain and is well-suited to complex, non-linear problems. ⁷⁸	
Artificial neural network	The algorithm starts with input nodes, which communicate, forming hidden layers of connections within the network, and provide learned output. ⁷⁸	(Hearty and Gibney 2008)
Kohonen neural network	A neural network comprised of a single layer. Data enter through input nodes and are organized to a simple structure of output nodes. ⁷⁹	(Jamiołkowski, Szpak, and Pawłowska 2005)
Probabilistic graphical models	Algorithm that can be used to determine joint relationships between nodes, forming a graphical network, where connections between nodes are represented by edges. These connections may be full- or partial-order correlations. ⁸⁰⁻⁸²	
Gaussian graphical models	Probabilistic graphical model where data follows a Gaussian distribution. ⁸¹	(Hoang, Lee, and Kim 2021; Iqbal et al. 2016; Schwedhelm et al. 2021)
Mixed graphical models	Probabilistic graphical model where data is mixed (i.e., Gaussian and categorical). ⁸¹	(Hoang, Lee, and Kim 2021)
Semiparametric Gaussian Copula graphical models	Probabilistic graphical model where the data is not required to be normally distributed. ⁸³	(Iqbal et al. 2016; Schwedhelm et al. 2018)
Other machine learning approaches		
Gradient boost machine	Machine learning algorithm that applies a boosting ensemble method to decision trees. This allows a weak learning algorithm to provide predictions through weighted averages of weak predictions. ^{53,84}	(Shang et al. 2020)

Bayesian kernel machine regression	Machine learning algorithm that uses a kernel function to consider the effects of multiple exposures, covariates, and confounders on an outcome. ⁸⁵	(Zhao et al. 2021)
Compositional data analysis	Method that identifies the relative magnitude of differences between variables to determine relative importance. ^{86,87}	(Solans et al. 2019)
Least Absolute Shrinkage and Selection Operator (LASSO)	A method that selects the most important variables and reduces the number of variables in the model. ⁸⁸	(Wu et al. 2019)
Multivariate finite mixture models		
Latent Class Analysis	A method that creates mutually exclusive groups (classes) of participants based on similar latent characteristics. ^{41,42}	(Affret et al. 2017; Bezerra et al. 2018; Dalmartello et al. 2020; Harrington et al. 2014; Hose et al. 2021)
Latent class transition model	Allows for transition between identified groups. ^{51,89}	(Madeira et al. 2021)
Latent class profile analysis	Latent class analysis when continuous variables are used. ⁵⁸	(Farmer et al. 2020)
Multivariate finite mixture models	Identifies mutually exclusive groupings where cluster shape is not limited. ⁹⁰	(Fonseca et al. 2012; Oliveira et al. 2011)
Mutual information	This method creates networks, estimating the amount that uncertainty is reduced in one variable by knowing another. ⁹¹	(Samieri et al. 2020; Xia et al. 2020)
Treelet transform	Data-driven approach that reduces dimensionality and is a hybrid between hierarchical clustering and principal component analysis. ^{92,93}	(de Almeida Alves et al. 2022; Wright et al. 2020)

*Studies may have used more than one novel approach to characterize dietary patterns.