

## Sleep and long COVID: Preexisting sleep issues and the risk of PASC in a large general population using 3 different model definitions

Stuart F. Quan, M.D.<sup>a,i</sup>  
Matthew D. Weaver, Ph.D.<sup>a,i</sup>  
Mark É. Czeisler, Ph.D.<sup>b,c,d</sup>  
Laura K. Barger, Ph.D.<sup>a,i</sup>  
Lauren A. Booker, Ph.D.<sup>d,e</sup>  
Mark E. Howard, M.B.B.S, Ph.D.<sup>d,f,g</sup>  
Melinda L. Jackson, Ph.D.<sup>c,d</sup>  
Rashon I. Lane, Ph.D.<sup>a</sup>  
Christine F. McDonald, M.B.B.S., Ph.D.<sup>d,g,h,j</sup>  
Anna Ridgers, M.B.B.S.<sup>d,h,g</sup>  
Rebecca Robbins, Ph.D.<sup>a,i</sup>  
Prerna Varma, Ph.D.<sup>c</sup>  
Joshua F. Wiley, Ph.D.<sup>c</sup>  
Shantha M.W. Rajaratnam, Ph.D.<sup>a,c,d,i</sup>  
Charles A. Czeisler, M.D., Ph.D.<sup>a,i</sup>

<sup>a</sup>Division of Sleep and Circadian Disorders, Brigham and Women's Hospital, Boston, MA; <sup>b</sup>Francis Weld Peabody Society, Harvard Medical School, Boston, MA; <sup>c</sup>School of Psychological Sciences, Turner Institute for Brain and Mental Health, Monash University, Melbourne, Australia; <sup>d</sup>Institute for Breathing and Sleep, Austin Health, Heidelberg, Victoria, Australia; <sup>e</sup>University Department of Rural Health, La Trobe Rural Health School, La Trobe University, Bendigo, Victoria, Australia; <sup>f</sup>Turner Institute for Brain and Mental Health, Monash University, Melbourne, Australia; <sup>g</sup>Department of Medicine, The University of Melbourne, Melbourne, Victoria, Australia; <sup>h</sup>Department of Respiratory and Sleep Medicine, Austin Health, Heidelberg, Victoria, Australia; <sup>i</sup>Division of Sleep Medicine, Harvard Medical School, Boston, MA; <sup>j</sup>Faculty of Medicine, Monash University, Melbourne Australia

Running Head: Sleep and Long COVID

Correspondent: Stuart F. Quan, M.D.  
Division of Sleep and Circadian Disorders  
Brigham and Women's Hospital  
221 Longwood Ave.  
Boston, MA 02115  
Email: [Stuart.Quan@hms.harvard.edu](mailto:Stuart.Quan@hms.harvard.edu)

Author Approval: All authors have seen and approved the manuscript.

Conflicts of Interest: MDW reports institutional support from the US Centers for Disease Control and Prevention, National Institutes of Occupational Safety and Health, and Delta Airlines; as well as consulting fees from the Fred Hutchinson Cancer Center and the University of Pittsburgh. LKB reports institutional support from the US Centers for Disease Control and Prevention, National Institutes of

## Sleep and Long COVID

Occupational Safety and Health, Delta Airlines, and the Puget Sound Pilots; as well as honorariums from the National Institutes of Occupational Safety and Health, University of Arizona and University of British Columbia. M EC reported personal fees from Vanda Pharmaceuticals Inc., research grants or gifts to Monash University from WHOOP, Inc., Hopelab, Inc., CDC Foundation, and the Centers for Disease Control and Prevention. SMWR reported receiving grants and personal fees from Cooperative Research Centre for Alertness, Safety, and Productivity, receiving grants and institutional consultancy fees from Teva Pharma Australia and institutional consultancy fees from Vanda Pharmaceuticals, Circadian Therapeutics, BHP Billiton, and Herbert Smith Freehills. SFQ has served as a consultant for Best Doctors, Bryte Foundation, Jazz Pharmaceuticals, Apnimed, and Whispersom. RR reports personal fees from SleepCycle AB; Rituals Cosmetics BV; Sonesta Hotels International, LLC; Ouraring Ltd; AdventHealth; and With Deep, LLC. CAC serves as the incumbent of an endowed professorship provided to Harvard Medical School by Cephalon, Inc. and reports institutional support for a Quality Improvement Initiative from Delta Airlines and Puget Sound Pilots; education support to Harvard Medical School Division of Sleep Medicine and support to Brigham and Women’s Hospital from: Jazz Pharmaceuticals PLC, Inc, Philips Respironics, Inc., Optum, and ResMed, Inc.; research support to Brigham and Women’s Hospital from Axome Therapeutics, Inc., Dayzz Ltd., Peter Brown and Margaret Hamburg, Regeneron Pharmaceuticals, Sanofi SA, Casey Feldman Foundation, Summus, Inc., Takeda Pharmaceutical Co., LTD, Abbaszadeh Foundation, CDC Foundation; educational funding to the Sleep and Health Education Program of the Harvard Medical School Division of Sleep Medicine from ResMed, Inc., Teva Pharmaceuticals Industries, Ltd., and Vanda Pharmaceuticals; personal royalty payments on sales of the Actiwatch-2 and Actiwatch-Spectrum devices from Philips Respironics, Inc; personal consulting fees from Axome, Inc., Bryte Foundation, With Deep, Inc. and Vanda Pharmaceuticals; honoraria from the Associated Professional Sleep Societies, LLC for the Thomas Roth Lecture of Excellence at SLEEP 2022, from the Massachusetts Medical Society for a New England Journal of Medicine Perspective article, from the National Council for Mental Wellbeing, from the National Sleep Foundation for serving as chair of the Sleep Timing and Variability Consensus Panel, for lecture fees from Teva Pharma Australia PTY Ltd. and Emory University, and for serving as an advisory board member for the Institute of Digital Media and Child Development, the Klarman Family Foundation, and the UK Biotechnology and Biological Sciences Research Council. CAC has received personal fees for serving as an expert witness on a number of civil matters, criminal matters, and arbitration cases, including those involving the following commercial and government entities: Amtrak; Bombardier, Inc.; C&J Energy Services; Dallas Police Association; Delta Airlines/Comair; Enterprise Rent-A-Car; FedEx; Greyhound Lines, Inc./Motor Coach Industries/FirstGroup America; PAR Electrical Contractors, Inc.; Puget Sound Pilots; and the San Francisco Sheriff’s Department; Schlumberger Technology Corp.; Union Pacific Railroad; United Parcel Service; Vanda Pharmaceuticals. CAC has received travel support from the Stanley Ho Medical Development Foundation for travel to Macao and Hong Kong; equity interest in Vanda Pharmaceuticals, With Deep, Inc, and Signos, Inc.; and institutional educational gifts to Brigham and Women’s Hospital from Johnson & Johnson, Mary Ann and Stanley Snider via Combined Jewish Philanthropies, Alexandra Drane, DR Capital, Harmony Biosciences, LLC, San Francisco Bar Pilots, Whoop, Inc., Harmony Biosciences LLC, Eisai Co., LTD, Idorsia Pharmaceuticals LTD, Sleep Number Corp., Apnimed, Inc., Avadel Pharmaceuticals, Bryte Foundation, f.lux Software, LLC, Stuart F. and Diana L. Quan Charitable Fund. Dr Czeisler’s interests were reviewed and are managed by the Brigham and Women’s Hospital and Mass General Brigham in accordance with their conflict-of-interest policies. No other disclosures were reported.

## Sleep and Long COVID

Preprint Information: A preprint of this manuscript has posted on medRxiv:

Funding Information: This work was supported by the Centers for Disease Control and Prevention. Dr. M. Czeisler was supported by an Australian-American Fulbright Fellowship, with funding from The Kinghorn Foundation. The salary of Drs. Barger, Czeisler, Robbins and Weaver were supported, in part, by NIOSH R01 OH011773 and NHLBI R56 HL151637. Dr. Robbins also was supported in part by NHLBI K01 HL150339.

### Acknowledgments:

Concept and Design: SFQ

Data collection: MDW, MÉC, MEH

Data analysis and interpretation: SFQ, MEH, CFM, MLJ, MEH

Drafting of the manuscript: SFQ

Critical feedback and revision of manuscript: SFQ, MDW, MÉC, LKB, LAB, MEH, MLJ, RL, CFM, AR, RR, PV, SMWR, CAC

Tables: 5

Figures: 0

Word Count: Text: 3873

Abstract: 246

Brief Summary: 91

## Sleep and Long COVID

### **Abstract**

*Study Objectives:* Insomnia, poor sleep quality and extremes of sleep duration are associated with COVID-19 infection. This study assessed whether these factors are related to Post-Acute Sequelae of SARS-CoV-2 infection (PASC).

*Methods:* Cross-sectional survey of a general population of 24,803 U.S. adults to determine the association of insomnia, poor sleep quality and sleep duration with PASC.

*Results:* Prevalence rates of PASC among previously COVID-19 infected participants for three definitions of PASC were COPE (21.9%), NICE (38.9%) and RECOVER PASC Score (15.3%). PASC was associated with insomnia in all 3 models in fully adjusted models with adjusted odds ratios (aORs) and 95% confidence intervals (CI) ranging from 1.30 (95% CI: 1.11-1.52,  $p \leq 0.05$ , PASC Score) to 1.52 (95% CI: 1.34-1.71,  $p \leq 0.001$ , (NICE). Poor sleep quality was related to PASC in all models with aORs ranging from 1.77 (95% CI: 1.60-1.97,  $p \leq 0.001$ , NICE) to 2.00 (95% CI: 1.77-2.26,  $p \leq 0.001$ , COPE). Sleep <6 hours was associated with PASC with aORs between 1.59 (95% CI: 1.40-1.80,  $p \leq 0.001$ , PASC Score) to 1.70 (95% CI: 1.53-1.89,  $p \leq 0.001$ , COPE). Sleep  $\geq 9$  hours was not associated with PASC in any model. Although vaccination with COVID-19 booster decreased the likelihood of developing PASC, it did not attenuate associations between insomnia, poor sleep quality and short sleep duration with PASC in any of the models.

*Conclusions:* Insomnia, poor sleep quality and short sleep duration are potential risk factors for PASC. Interventions to improve sleep may decrease the development of PASC.

### **Key Words**

Long COVID, PASC, insomnia, poor sleep quality, sleep duration, Post-Acute Sequelae of SARS-CoV-2 infection

## Sleep and Long COVID

### Brief Summary

*Current Knowledge/Study Rationale:* Insomnia, poor sleep quality, and extremes of sleep duration have been associated with a higher likelihood of COVID-19 infection. However, evidence implicating an association with the development of Post-Acute Sequelae of SARS-CoV-2 infection (PASC) is scant.

*Study Impact:* Results indicate that insomnia, poor sleep quality and sleep duration  $\leq 6$  hours are associated with an increase in the prevalence of PASC among persons who have previously had a COVID-19 infection. The findings provide support for employing interventions to improve sleep as a means to decrease the development of PASC.

## Sleep and Long COVID

### Introduction

As the COVID-19 pandemic has evolved into global endemic status, the emergence of persistent and/or relapsing-remitting physical, cognitive and mental health symptoms commonly known as “long COVID” or more formally as Post-Acute Sequelae of SARS-CoV-2 infection (PASC) has become an increasingly important public health concern.<sup>1-4</sup> Risk factors for the development of PASC include greater severity of initial COVID-19 illness, multiple infections, female sex and preexisting health conditions.<sup>2,5</sup> Recently, obstructive sleep apnea (OSA) has been demonstrated in two large studies to be associated with the development of PASC.<sup>6,7</sup> Although poor sleep quality and short sleep duration are associated with an increased risk of acute COVID-19 infection,<sup>8,9</sup> it is uncertain whether they also are related to a higher likelihood of PASC.

Estimates of the prevalence of PASC are imprecise, ranging from 7.5 to 41%.<sup>10</sup> In part, this imprecision is related to the lack of a generally accepted definition of this syndrome. Several definitions have been proposed that include the use of various combinations of duration of time post infection, number of symptoms and a scoring system that weights some symptoms more than others.<sup>6,11-14</sup> However, there are few data that have prospectively tested or compared definitions to each other, control groups and long-term outcomes.

In this study, we aimed to determine in a large general population cohort whether preexisting insomnia, poor sleep quality, and short or long sleep duration are associated with the development of PASC using 3 different model definitions of the syndrome. In addition, we compared the prevalence of PASC among these models.

## Sleep and Long COVID

### Methods

#### *Study Design and Participants*

From March 10 to October 15, 2022, the COPE Initiative<sup>15</sup> administered five successive waves of surveys focused on accumulating data on the prevalence and sequelae of COVID-19 infection in the United States (U.S.). Dates of administration were Wave 1 (March 10-30, 2022), Wave 2 (April 4-May 1, 2022), Wave 3 (May 4-June 2, 2022), Wave 4 (August 1-18, 2022), Wave 5 (September 26-October 15, 2022). Each wave consisted of approximately 5000 unique participants who were recruited to create samples that approximated population estimates for age, sex, race, and ethnicity based on the 2020 U.S. census. Surveys were conducted online by Qualtrics, LLC (Provo, Utah, and Seattle, Washington, U.S.), using their network of participant pools with varying recruitment methodologies. Informed consent was obtained electronically. The study was approved by the Monash University Human Research Ethics Committee (Study #24036).

#### *Survey Items*

Participants self-reported demographic, anthropometric, and socioeconomic information including age, race, ethnicity, sex, height and weight, education level, employment status and household income. In addition, they provided information on several current and past medical conditions by answering the question: “Have you ever been diagnosed with any of the following conditions?” In addition to insomnia, opportunity was provided to endorse obstructive sleep apnea (OSA), high blood pressure, cardiovascular disease (e.g., heart attack, stroke, angina), gastrointestinal disorder (e.g., acid reflux, ulcers, indigestion), cancer, chronic kidney disease, liver disease, sickle cell

## Sleep and Long COVID

disease, chronic obstructive pulmonary disease, and asthma. Possible responses to each condition were “Never”, “Yes I have in the past, but don’t have it now”, “Yes I have, but I do not regularly take medications or receiving treatment”, and “Yes I have, and I am regularly taking medications or receiving treatment”.

The following two questions pertaining to sleep quality were asked of the participants:

- a. “Thinking about the past month, to what extent has poor sleep troubled you in general?”. Possible responses were “Not at all”, “A little”, “Somewhat”, “Much” and “Very Much”.
- b. From the Pittsburgh Sleep Quality Index,<sup>16</sup> “During the past month, how would you rate your sleep quality overall?” Possible responses were “Very good”, “Fairly good”, “Fairly bad” and “Very bad”.

Sleep duration was assessed using a question from the Pittsburgh Sleep Quality Index.<sup>16</sup> Responses were rounded to the nearest hour; those <3 hours or >12 hours were excluded as improbable estimates (N=987). In addition, sleep duration was stratified as short sleep ( $\leq 6$  hours), recommended sleep (7-8 hours) and long sleep ( $\geq 9$  hours).

Symptoms of OSA were obtained from responses to the Pittsburgh Sleep Quality Index and included items related to roommate or bedpartner reported “loud snoring” and “long pauses between breaths while you sleep”.<sup>16</sup> In addition, sleepiness was assessed from the following item in the questionnaire: “During the past month, how often have you had trouble staying awake while driving, eating meals, or engaging in social activity”. Possible responses to all three items were “not during the past month”, “less than once a week”, “once or twice a week” or “three or more times a week”. As done in our previous studies, participants were considered to have symptoms of OSA if they had either of the



## Sleep and Long COVID

following combination of symptoms: 1) snoring “three or more times a week” and witnessed apnea or sleepiness “once or twice a week”; 2) witnessed apnea and sleepiness “once or twice a week”.<sup>6,17</sup>

Each survey contained identical items related to COVID-19 infection status and the number of COVID-19 vaccinations participants had received. Ascertainment of past COVID-19 infection was obtained using responses to the following questions related to COVID-19 testing or the presence of loss of taste or smell:

1. “Have you ever tested positive?”
2. “Despite never testing positive, are you confident that you have had COVID-19?”
3. “Despite never testing positive, have you received a clinical diagnosis of COVID-19?”
4. “Have you experienced a problem with decreased sense of smell or taste at any point since January 2020?”

COVID-19 vaccination status was ascertained by asking “How many COVID-19 vaccine doses have you received? (If you have had two doses of one brand and one of another, please select three)”. Participants were allowed to respond from 0 to 4.

### *Statistical Analyses*

Exposures: Based on our previous analyses in this cohort, we defined a positive history of COVID-19 infection as an affirmative response to having tested positive for COVID-19, a clinical diagnosis of COVID-19, or loss of taste or smell.<sup>6,9,17</sup> Participants were classified as having OSA if they affirmed currently having the condition whether treated or not or if they had two or more symptoms of OSA.<sup>6,9,17</sup> Insomnia was present if endorsed by participants irrespective of treatment status. Poor sleep quality was defined

## Sleep and Long COVID

as being troubled by poor sleep “Much” or “Very Much” or rating their sleep quality as “Fairly Bad” or “Very Bad”.<sup>9</sup>

Vaccination status was dichotomized as Boosted (>2 vaccinations) or Not Boosted ( $\leq 2$  vaccinations). Comorbid medical conditions were defined as currently having the condition whether treated or untreated. The effect of comorbid medical conditions was evaluated by summing the number of conditions reported by the participant (minimum value 0, maximum value 9).<sup>6,9,17</sup> Body mass index (BMI) was calculated using self-reported height and weight as kg/m<sup>2</sup>. Socioeconomic covariates were dichotomized as follows: employment (retired vs. not retired), education (high school or less vs. some college) and income in U.S. Dollars (<\$50,000 vs  $\geq$ \$50,000).

Outcomes: Based on previous analyses of this cohort, PASC was defined as the presence of 3 or more symptoms commonly associated with PASC for at least 3 months after their COVID-19 infection (COPE).<sup>6</sup> For comparison, 2 other definitions of PASC were also used: 1) a PASC Score  $\geq 12$  modified from the RECOVER cohort proposal,<sup>13</sup> and 2) one or more symptoms commonly associated with PASC for at least 3 months after their COVID-19 infection as suggested by the United Kingdom’s National Institute for Health and Care Excellence (NICE).<sup>12</sup> Because the COPE survey did not contain a few items comparable to those in the RECOVER cohort, we substituted symptoms contained in the COPE survey which were correlated<sup>13</sup> or similar as follows: sore throat for thirst, sleep problems for sexual function, headache for dizziness, and other aches for abnormal movements.

Analytics: Summary data for continuous or ordinal variables are reported as their respective means and standard deviations (SD) and for categorical variables as their percentages. Comparisons of co-morbid medical, demographic, and social characteristic

## Sleep and Long COVID

variables stratified by presence or absence of PASC were performed using Student's unpaired t-test for continuous or ordinal variables and  $\chi^2$  for categorical variables.

Multivariable modelling using logistic regression was utilized to determine whether insomnia, poor sleep quality and sleep duration were associated with the COPE, RECOVER and NICE models of PASC. For each definition, a baseline model was constructed using only insomnia, poor sleep quality and sleep duration stratified as short ( $\leq 6$  hours), recommended (7-8 hours),<sup>18</sup> and long ( $\geq 9$  hours) sleep. We then developed increasingly complex models by sequentially including demographic factors, comorbidities, boosted vaccination status, socioeconomic factors, and other sleep conditions. Inasmuch as the sleep conditions evaluated in the models may have been correlated, diagnostic tests for collinearity were performed; the variance inflation factor was less than 10 and the tolerance was greater than 0.1 for all variables. However, because poor sleep quality can be considered a component of insomnia, models for insomnia were not adjusted for poor sleep quality. Conversely, models for poor sleep quality were not adjusted by insomnia. Results of the logistic regression models are presented as unadjusted or adjusted odds ratios (aOR) and their 95% confidence intervals (95% CI).

To determine whether our definition of COVID-19 infection status influenced our results, we performed sensitivity analyses with stricter (i.e., using COVID-19 infection as a positive test or loss of taste or smell) and broader (i.e., our original definition plus assumed positive for COVID-19 without a positive test or clinical diagnosis as an indicator of a past COVID-19 infection).

All analyses were conducted using IBM SPSS version 28 (Armonk, NY). A  $p < 0.05$  was considered statistically significant.

## Sleep and Long COVID

### Results

There were 24803 participants with evaluable data from the five COPE initiative surveys from which a positive history of COVID-19 infection (see Methods) was reported in 10324 (41.6%), which represented the analytic sample. Table 1 presents the demographic, anthropometric, co-morbid medical and social characteristics of these COVID-19 positive participants in the COPE cohort stratified by PASC status for the 3 models of PASC analyzed in this study. Characteristics of participants positive for PASC were generally similar for all 3 models. Those who were PASC positive were younger and more likely to be Hispanic, not retired, and have a higher income. They were less likely to have received a COVID-19 booster vaccination (COPE: 24.2% vs. 29.3%; NICE: 23.7% vs. 31.0%; PASC Score: 24.5% vs. 28.8%; all  $p < 0.01$ ). In terms of sleep, 3452 (45.6%) participants reported having slept the recommended 7-8 hours, 2973 (36.9%) screened positive for poor sleep quality, 1671 (20.7%) screened positive for insomnia, and 1697 (21.0%) screened positive for OSA. Compared with participants who did not experience PASC, those with PASC had a higher prevalence of OSA, insomnia, and poor sleep quality. In addition, they were more likely to report a short sleep duration whereas there was a slight tendency for those with PASC to report a long sleep duration.

The prevalence of PASC as defined in the COPE, NICE, and PASC score models is shown in Table 1. The most restrictive model was the PASC score (15.3%) followed by the COPE (21.9%) and NICE (38.9%) models.

In Table 2 are logistic regression analyses describing among participants with a positive history of COVID-19 infection the association between insomnia and PASC in all 3 PASC models at baseline and subsequently in partially and fully adjusted models. The

## Sleep and Long COVID

aORs in the fully adjusted models were comparable with narrow confidence intervals (CI) ranging from 1.30 (95% CI: 1.11-1.52,  $p \leq 0.05$ , PASC Score) to 1.52 (95% CI: 1.34-1.71,  $p \leq 0.001$ , (NICE). In none of the models did boosted vaccination affect the relationship between insomnia and PASC (insomnia x boosted vaccination aORs were not significant, data not shown).

Table 3 describes the association of poor sleep quality with PASC among participants with a positive history of COVID-19 infection. For all 3 models, poor sleep quality was related to an increased probability of PASC; aORs at baseline were comparable ranging from 2.36 (95% CI: 2.18-2.56,  $p \leq 0.001$ , NICE) to 2.63 (95% CI: 2.39-2.90,  $p \leq 0.001$ , COPE). Subsequent adjustment for demographic and socioeconomic factors, and sleep conditions as well as comorbidities attenuated these associations, but all remained significant in fully adjusted models with aORs ranging from 1.77 (95% CI: 1.60-1.97,  $p \leq 0.001$ , NICE) to 2.00 (95% CI: 1.77-2.26,  $p \leq 0.001$ , COPE). In none of the models did boosted vaccination affect the relationship between poor sleep quality and PASC (poor sleep quality x boosted vaccination aORs were not significant, data not shown).

In Table 4 are shown the relationships between the full distribution of hourly sleep duration (3 to 12 hours) with 3 models of PASC among participants with a positive history of COVID-19 infection. For all 3 models, the unadjusted aORs suggested that both short and long sleep durations were associated with PASC in comparison to the recommended 7-8 hours of sleep. For sleep durations less than 7-8 hours, these findings were attenuated, but remained significant. However, for sleep durations greater than 7-8 hours, the association with PASC was no longer present with the solitary exception of 11 hours in the PASC Score model.

## Sleep and Long COVID

Shown in Table 5 are the associations of short sleep ( $\leq 6$  hours) and long sleep ( $\geq 9$  hours) with PASC among participants with a positive history of COVID-19 infection. In comparison to recommended hours of sleep (7-8 hours;  $n = 3452$ ), short sleep ( $n = 3284$ ) was associated with a greater likelihood of PASC. The aORs at baseline were comparable for all 3 models and ranged from 1.59 (95% CI: 1.40-1.80,  $p \leq 0.001$ , PASC Score) to 1.70 (95% CI: 1.53-1.89,  $p \leq 0.001$ , COPE). These associations were attenuated but remained significant in fully adjusted models with aORs of 1.17 (95% CI: 1.01-1.36,  $p \leq 0.05$ , PASC Score), 1.23 (95% CI: 1.08-1.41,  $p \leq 0.001$ , COPE), and 1.23 (95% CI: 1.10-1.37,  $p \leq 0.001$ , NICE). Long sleep ( $n = 826$ ) also was associated with a higher probability of PASC at baseline in all 3 models with aORs ranging from 1.33 (95% CI: 1.12-1.58,  $p \leq 0.001$ , COPE) to 1.50 (95% CI: 1.24-1.82,  $p \leq 0.001$ , PASC Score). However, in contrast to short sleep, adjustment for demographic and socioeconomic factors, and sleep conditions as well as comorbidities negated these associations in fully adjusted models. The associations between short and long sleep, and all 3 models of PASC were not mitigated by having received a COVID-19 booster (sleep duration x boosted vaccination aORs were not significant, data not shown).

In sensitivity analyses, using a stricter definition of COVID-19 that required a positive test or loss of taste or smell, the positive associations between insomnia, poor sleep quality and short sleep duration remained (data not shown). Similarly, using a less stringent definition of COVID-19 that allowed for a positive diagnosis for any unconfirmed report of infection did not alter these relationships (data not shown).

## Discussion

In this study, the prevalence of PASC among participants with a positive history of COVID-19 infection was estimated in a large U.S. general population cohort using three

## Sleep and Long COVID

different models of PASC and found to range between 15.3% and 38.9%. In all three models, insomnia, poor sleep quality and sleep shorter than recommended levels were associated with PASC after adjustment for demographic, anthropometric, and socioeconomic factors, and comorbid medical conditions. In contrast, sleep longer than recommended was not related to PASC.

Although it is generally acknowledged that a substantial proportion of persons who develop acute COVID-19 infection have a multitude of lingering symptoms, there is no agreement on a unifying definition of PASC. Such a definition is important for public health case identification as well as for guiding future research. However, several definitions have been proposed using different requirements for number of symptoms and time elapsed after acute infection. In the current study, we estimated the prevalence of PASC using a definition proposed by NICE in the United Kingdom<sup>12</sup> and another modified from the RECOVER cohort in the U.S.<sup>13</sup> and compared them to one we developed for the COPE Initiative.<sup>6</sup> We found that among participants with a positive history of COVID-19 infection, prevalence estimates of PASC were 21.9%, 38.9% and 15.3% using the COPE, NICE, and RECOVER models, respectively. In contrast, prevalence rates of 21.21% and 8.68% were observed using the NICE and PASC Score definitions in a survey of the adult population of Mexico.<sup>19</sup> Analysis of the Behavioral Risk Factor Surveillance System (BRFSS) found a 21.7% prevalence using the NICE definition.<sup>20</sup> The prevalence of PASC in the RECOVER cohort was 23%,<sup>13</sup> but a much higher rate of 61% was noted in a study from the International COVID Sleep Study II (ICOSSII) group using the World Health Organization definition of at least one symptom lasting at least 3 months.<sup>21</sup> Notably, a meta-analysis of a heterogeneous array of 33 studies found an overall prevalence of 37% 30 days after infection with a 95% confidence interval between 26 and

## Sleep and Long COVID

49%.<sup>22</sup> The explanation for the considerable heterogeneity in PASC prevalence rates is not entirely clear, but investigations differed in both methods used to ascertain COVID-19 positivity and definition of PASC as well as in the population evaluated. Most studies utilized self-report of both COVID-19 infection and PASC associated symptoms,<sup>6,20,21,23,24</sup> but in some, objective documentation of infection was required resulting in lower prevalence rates.<sup>13,19</sup> To our knowledge, this is the first study in the U.S. to estimate prevalence rates of PASC using different definitions in the same cohort. However, the most appropriate definition to ultimately use may depend on which one best predicts clinical outcomes or responses to therapy.

We observed that among participants with a positive history of COVID-19 infection, both insomnia and poor sleep quality were strongly associated with PASC in all three of the models evaluated. With respect to insomnia, our results are consistent with data from a longitudinal study of 2,759 Italians in which higher pre-COVID-19 insomnia severity index scores were found to predict greater risk of individual symptoms of PASC 1 to 3 months after COVID-19 infection.<sup>25</sup> Our findings extend this previous report by demonstrating in a larger cohort that insomnia is associated with several global definitions of PASC. Additionally, our results related to poor sleep quality are supported by data from the aforementioned Italian cohort study which showed that higher scores on the Pittsburgh Sleep Quality Index predicted a greater risk for developing symptoms associated with PASC.<sup>25</sup> Furthermore, in a study of 1,581 persons from the United Kingdom who were surveyed concerning their sleep quality one month before their COVID-19 infection, persons who reported average to very poor sleep quality were 2.4-3.5 times more likely to self-report having “Long COVID”.<sup>24</sup> Our study extends these



## Sleep and Long COVID

previous observations by using a more precise definition of poor sleep quality as well as a more formal definitions of PASC.

Sleep duration  $\leq 6$  hours per night in comparison to recommended levels of 7-8 hours was found to be associated with PASC in all 3 models. Our findings replicate those noted in analyses of the BRFSS dataset in which  $< 7$  hours of sleep were also associated with higher odds of PASC.<sup>20</sup> However, they differ from a report from the the International COVID Sleep Study II (ICOSSII) group who noted increased likelihood of PASC in those sleeping  $< 6$  hours per night only among individuals who had pre-existing medical co-morbidities; no association was observed in the absence of medical co-morbidities.<sup>21</sup> In contrast to the strong association between short sleep duration and PASC, no such linkage was observed in our study for long sleep duration. Thus, our results with respect to long sleep duration are consistent with reports from the BRFSS and ICOSSII studies which also did not find increased odds of PASC with longer than recommended sleep duration.<sup>20,21</sup>

We observed that for all 3 PASC models, the prevalence of having received a COVID-19 booster vaccination was lower in persons with PASC. This finding is consistent with a meta-analysis of 18 studies<sup>26</sup> as well as with a recent large retrospective matched cohort study demonstrating that COVID-19 vaccination decreased the risk of developing PASC.<sup>27</sup> Previous results from the COPE initiative indicated that receiving a COVID-19 booster vaccination reduced the risk of COVID-19 hospitalization attributable to OSA.<sup>17</sup> In contrast, we did not find that boosted vaccination for COVID-19 affected the odds of developing PASC related to insomnia, poor sleep quality or short sleep duration.

There are several mechanisms that could explain the association between insomnia, poor sleep quality and short sleep duration, and PASC. Insomnia and most

## Sleep and Long COVID

likely poor sleep quality are characterized by a state of stress and hyperarousal.<sup>28</sup> This can lead to alterations in the hypothalamic-pituitary-adrenal axis related to inflammation; in a meta-analysis, sleep disturbance was associated with higher levels of C Reactive Protein and Interleukin-6.<sup>29</sup> Reduced sleep can result in an increase in transcriptional pathways and greater cellular production of inflammatory cytokines from monocytes.<sup>30</sup> Furthermore, it has been suggested that inflammatory processes are important in the pathogenesis of the dysautonomia, endothelial dysregulation, and neurocognitive impairments observed with PASC.<sup>31</sup> Sleep disturbances also have been implicated in dysfunction of the immune system. Chronic sleep loss and disturbed sleep promote not only a proinflammatory state, but also affect the number and function of immune regulating cells<sup>32,33</sup> and prolong shedding of SARS-CoV-2 after infection.<sup>34</sup> Both factors may promote viral persistence in tissues which is hypothesized as a mechanism resulting in PASC.<sup>31</sup>

Our findings that sleep disruption is associated with a greater likelihood of PASC suggest interventions improving sleep may mitigate or reduce the impact of PASC on overall public health. For example, information pertaining to the benefits of healthy sleep could be disseminated to persons who develop COVID-19 infection. Inasmuch as the prevalence of PASC in our study as well as others ranges between 26% and 49%,<sup>22</sup> even a small reduction in the incidence of PASC could have profound public health importance.

There are several limitations to the interpretation of our findings. First, sleep disturbances can occur *de novo* after acute COVID-19 infection.<sup>35,36</sup> Our analyses are cross-sectional, and therefore a conclusion that insomnia, poor sleep quality and short sleep duration are risk factors for the development of PASC should be interpreted cautiously. Second, sleep duration was self-reported. Self-reported sleep has historically

## Sleep and Long COVID

overestimated sleep duration compared with the gold-standard objective measurement by polysomnography.<sup>37</sup> However, inasmuch as both 3 and 4 hours of sleep in this study were associated with PASC, it is unlikely that any overestimates of sleep duration would have been sufficiently extreme to alter our findings of an association between short sleep duration and PASC. Third, the sample could be susceptible to misclassification bias with respect to history of COVID-19 infection, as infection with COVID-19 was based on self-report and included dyssomnia and dysgeusia. However, in sensitivity analyses our findings were robust when incorporating less and more stringent criteria. Finally, the COPE survey did not include all of the symptoms included in the RECOVER PASC score, which had not been published at the time the COPE survey was developed and initially administered. Nevertheless, the findings observed with our modification of the PASC Score were consistent with the COPE and NICE definitions suggesting that they were similarly representative of participants with PASC.

In summary, the prevalence of PASC ranged between 15.3% and 38.9% in persons who previously had a COVID-19 infection. Insomnia, poor sleep quality, and short sleep duration are associated with higher odds of PASC. Boosted vaccination for COVID-19 did not affect these observations. These findings emphasize the importance of sleep disturbance in COVID-19 infection and PASC and suggest that interventions to improve sleep may reduce the incidence of PASC.

## Sleep and Long COVID

### References

1. Healey Q, Sheikh A, Daines L, Vasileiou E. Symptoms and signs of long COVID: A rapid review and meta-analysis. *J Glob Health*. 2022;12:05014.
2. Yong SJ. Long COVID or post-COVID-19 syndrome: putative pathophysiology, risk factors, and treatments. *Infect Dis (Lond)*. 2021;53:737-754.
3. Marshall M. The lasting misery of coronavirus long-haulers. *Nature*. 2020;585:339-341.
4. Nasserie T, Hittle M, Goodman SN. Assessment of the Frequency and Variety of Persistent Symptoms Among Patients With COVID-19: A Systematic Review. *JAMA Netw Open*. 2021;4:e2111417.
5. Tsampasian V, Elghazaly H, Chattopadhyay R, et al. Risk Factors Associated With Post-COVID-19 Condition: A Systematic Review and Meta-analysis. *JAMA Intern Med*. 2023;183:566-580.
6. Quan SF, Weaver MD, Czeisler MÉ, et al. Association of Obstructive Sleep Apnea with Post-Acute Sequelae of SARS-CoV-2 Infection. *Am J Med*. 2024;137:529-37.
7. Mandel HL, Colleen G, Abedian S, et al. Risk of post-acute sequelae of SARS-CoV-2 infection associated with pre-coronavirus disease obstructive sleep apnea diagnoses: an electronic health record-based analysis from the RECOVER initiative. *Sleep*. 2023;46:zsad126.
8. Shafiee A, Jafarabady K, Rajai S, Mohammadi I, Mozhgani S-H. Sleep disturbance increases the risk of severity and acquisition of COVID-19: a systematic review and meta-analysis. *Eur J Med Res*. 2023;28:442.
9. Quan SF, Weaver MD, Czeisler MÉ, et al. Insomnia, Poor Sleep Quality and Sleep Duration, and Risk for COVID-19 Infection and Hospitalization. *Am J Med*. 2023;136:780-788.e5.
10. Nittas V, Gao M, West EA, et al. Long COVID Through a Public Health Lens: An Umbrella Review. *Public Health Rev*. 2022;43:1604501.
11. A clinical case definition of post COVID-19 condition by a Delphi consensus, 6 October 2021. [https://www.who.int/publications-detail-redirect/WHO-2019-nCoV-Post\\_COVID-19\\_condition-Clinical\\_case\\_definition-2021.1](https://www.who.int/publications-detail-redirect/WHO-2019-nCoV-Post_COVID-19_condition-Clinical_case_definition-2021.1).
12. 1 Identification | COVID-19 rapid guideline: managing the long-term effects of COVID-19 | Guidance | NICE. <https://www.nice.org.uk/guidance/ng188/chapter/1-Identification#case-definition>.
13. Thaweethai T, Jolley SE, Karlson EW, et al. Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection. *JAMA*. 2023;329:1934-1946.

## Sleep and Long COVID

14. Fineberg HV, Brown L, Worku T, Goldowitz I, eds. A Long COVID Definition: A Chronic, Systemic Disease State with Profound Consequences. Washington, D.C.: National Academies Press; 2024.
15. The COPE Initiative. The COPE Initiative. <https://www.thecopeinitiative.org>.
16. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 1989;28:193-213.
17. Quan SF, Weaver MD, Czeisler MÉ, et al. Associations between obstructive sleep apnea and COVID-19 infection and hospitalization among US adults. *J Clin Sleep Med.* 2023;19:1303-1311.
18. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's updated sleep duration recommendations: final report. *Sleep Health.* 2015;1:233-243.
19. Bello-Chavolla OY, Fermín-Martínez CA, Ramírez-García D, et al. Prevalence and determinants of post-acute sequelae after SARS-CoV-2 infection (Long COVID) among adults in Mexico during 2022: a retrospective analysis of nationally representative data. *Lancet Reg Health Am.* 2024;30:100688.
20. Hejazian SS, Sadr AV, Shahjouei S, Vemuri A, Abedi V, Zand R. Prevalence and Determinants of Long-Term Post-COVID Conditions in the United States: 2022 Behavioral Risk Factor Surveillance System. *Am J Med.* February 2024:S0002-9343(24)00090-1.
21. Berezin L, Waseem R, Merikanto I, et al. Habitual short sleepers with pre-existing medical conditions are at higher risk of Long COVID. *J Clin Sleep Med.* 2024;20:111-119.
22. Chen C, Hauptert SR, Zimmermann L, Shi X, Fritsche LG, Mukherjee B. Global Prevalence of Post-Coronavirus Disease 2019 (COVID-19) Condition or Long COVID: A Meta-Analysis and Systematic Review. *J Infect Dis.* 2022;226:1593-1607.
23. Chen S-J, Morin CM, Ivers H, et al. The association of insomnia with long COVID: An international collaborative study (ICOSS-II). *Sleep Med.* 2023;112:216-222.
24. Paul E, Fancourt D. Health behaviours the month prior to COVID-19 infection and the development of self-reported long COVID and specific long COVID symptoms: a longitudinal analysis of 1581 UK adults. *BMC Public Health.* 2022;22:1716.
25. Salfi F, Amicucci G, Corigliano D, et al. Poor sleep quality, insomnia, and short sleep duration before infection predict long-term symptoms after COVID-19. *Brain Behav Immun.* 2023;112:140-151.
26. Gao P, Liu J, Liu M. Effect of COVID-19 Vaccines on Reducing the Risk of Long COVID in the Real World: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health.* 2022;19:12422.

## Sleep and Long COVID

27. Malden DE, Liu I-LA, Qian L, et al. Post-COVID conditions following COVID-19 vaccination: a retrospective matched cohort study of patients with SARS-CoV-2 infection. *Nat Commun.* 2024;15:4101.
28. Bonnet MH, Arand DL. Hyperarousal and insomnia: state of the science. *Sleep Med Rev.* 2010;14:9-15.
29. Irwin MR, Olmstead R, Carroll JE. Sleep Disturbance, Sleep Duration, and Inflammation: A Systematic Review and Meta-Analysis of Cohort Studies and Experimental Sleep Deprivation. *Biol Psychiatry.* 2016;80:40-52.
30. Irwin MR, Opp MR. Sleep Health: Reciprocal Regulation of Sleep and Innate Immunity. *Neuropsychopharmacology.* 2017;42:129-155.
31. Sherif ZA, Gomez CR, Connors TJ, Henrich TJ, Reeves WB, RECOVER Mechanistic Pathway Task Force. Pathogenic mechanisms of post-acute sequelae of SARS-CoV-2 infection (PASC). *Elife.* 2023;12:e86002.
32. Besedovsky L, Lange T, Haack M. The Sleep-Immune Crosstalk in Health and Disease. *Physiol Rev.* 2019;99:1325-1380.
33. Yin K, Peluso MJ, Luo X, et al. Long COVID manifests with T cell dysregulation, inflammation, and an uncoordinated adaptive immune response to SARS-CoV-2. *Nat Immunol.* 2024;25:218-25.
34. Lin YN, Zhou LN, Liu ZR, et al. Short Sleep Duration is Associated with Prolonged Virus Shedding in SARS-CoV-2 Omicron-Infected Patients. *Nat Sci Sleep.* 2023;15:547-554.
35. Tański W, Tomasiewicz A, Jankowska-Polańska B. Sleep Disturbances as a Consequence of Long COVID-19: Insights from Actigraphy and Clinimetric Examinations-An Uncontrolled Prospective Observational Pilot Study. *J Clin Med.* 2024;13:839.
36. Batool-Anwar S, Fashanu OS, Vassell C, Quan SF. Long-Term Effects of COVID-19 on Sleep Patterns. *Preprints.* 2023;2023050836.
37. Silva GE, Goodwin JL, Sherrill DL, et al. Relationship Between Reported and Measured Sleep Times. *J Clin Sleep Med.* 2007;03:622-630.

## Sleep and Long COVID

Table 1: Associations Between PASC Status and Insomnia, Poor Sleep Quality, Sleep Duration, Co-morbid Medical, Demographic and Social Characteristics

	COPE <sup>‡</sup> (≥3 Symptoms)				NICE <sup>#</sup> (≥1 Symptom)				PASC <sup>†</sup> Score (≥12)				Overall	
	PASC Negative		PASC Positive		PASC Negative		PASC Positive		PASC Negative		PASC Positive			
Prevalence (N/%)	8067/78.1		2257/21.9		6310/61.1		4014/38.9		8747/84.7		1577/15.3		10324	
	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>
Age (y)	41.9	16.6	37.4*	13.6	43.1	17.0	37.6*	13.9	41.6	16.5	37.0*	13.1	40.9	16.1
Body Mass Index (kg/m <sup>2</sup> )	27.6	6.2	27.5	6.6	27.5	6.1	27.7	6.6	27.6	6.2	27.7	6.8	27.6	6.3
No. Comorbidities	1.6	2.5	3.7*	3.5	1.4	2.2	3.3*	3.4	1.8	2.6	3.8*	3.5	2.1	2.9
	<i>N</i>	<i>%</i>	<i>N</i>	<i>%</i>	<i>N</i>	<i>%</i>	<i>N</i>	<i>%</i>	<i>N</i>	<i>%</i>	<i>N</i>	<i>%</i>	<i>N</i>	<i>%</i>
Sex														
Male	3951	49.5	1056	47.6	3121	49.9	1886†	47.7	4251	49.1	756	48.7	5007	49.1
Female	4034	50.5	1164	52.4	3129	50.1	2069	52.3	4403	50.9	795	51.3	5198	50.9
Race/Ethnicity														
White	4763	59.0	1269	56.2*	3792	60.1	2240*	55.8	5140	58.8	892†	56.6	6032	58.4
Black	805	10.0	251	11.1	609	9.7	447	11.1	866	9.9	190	12.0	1056	10.2
Asian	429	5.3	84	3.7	355	5.6	158	3.9	464	5.3	49	3.1	513	5.0
Hispanic	1728	21.4	536	23.7	1278	20.3	986	24.6	1900	21.7	364	23.1	2264	21.9
Other	342	4.2	117	5.2	276	4.4	183	4.6	377	4.3	82	5.2	459	4.4
Employment														
Retired	1230	15.2	152	6.7*	1079	17.1	303	7.5*	1272	14.5	110	7.0*	1382	13.4
Not Retired	6837	84.8	2105	93.3	5231	82.9	3711	92.5	7475	85.5	1467	93.0	8942	86.6
Education														
High School or Less	2129	26.4	541	24.0†	1661	26.3	1009	25.1	2290	26.2	380	24.1	2670	25.9
Some College	5938	73.6	1716	76.0	4649	73.7	3005	74.9	6457	73.8	1197	75.9	7654	74.1
Income (Yearly)														

## Sleep and Long COVID

< \$50,000	329 2	42. 2	844	38.2 *	257 6	42. 4	1560	39.6 *	353 5	41. 8	601	38.7 †	413 6	41. 3
≥ \$50,000	450 6	57. 8	136 7	61.8	349 6	57. 6	2377	60.4	492 2	58. 2	951	61.3	587 3	58. 7
Vaccination Boosted														
No (≤2 Vaccinations)	552 4	70. 7	168 3	75.8 *	420 6	69. 0	3001	76.3 *	602 8	71. 2	1179	75.5 *	720 7	71. 9
Yes (>2 Vaccinations)	228 5	29. 3	538	24.2	189 0	31. 0	933	23.7	244 1	28. 8	382	24.5	282 3	28. 1
Obstructive Sleep Apnea														
Yes	169 7	21. 0	1131	50.1 *	107 1	17. 0	1757	43.8 *	2011	23. 0	817	51.8 *	282 8	72. 6
No	637 0	79. 0	1126	49.9 *	523 9	83. 0	2257	56.2	673 6	77. 0	760	48.2	749 6	27. 4
Insomnia														
Yes	167 1	20. 7	991	43.9 *	109 0	17. 3	1573	39.2 *	196 8	22. 5	695	44.1 *	266 3	25. 8
No	639 5	79. 3	126 6	56.1	522 0	82. 7	2441	60.8	677 9	77. 5	882	55.9	766 1	74. 2
Poor Sleep Quality														
Yes	297 3	36. 9	136 7	60.6 *	214 0	33. 9	2200	52.8 *	338 6	38. 7	954	60.5 *	434 0	42. 0
No	509 4	63. 1	890	39.4	417 0	66. 1	1814	45.2	536 1	61. 3	623	39.5	598 4	58. 0
Sleep Duration (Hourly)														
3	234	3.1	142	7.1*	160	2.7	216	6.1*	281	3.4	95	6.8*	376	3.9
4	532	7.0	250	12.5	380	6.3	402	11.3	607	7.4	175	12.5	782	8.2
5	955	12. 6	341	17.0	724	12. 0	572	16.1	106 6	13. 0	230	16.5	129 6	13. 5
6	156 3	20. 7	370	18.5	125 0	20. 8	683	19.2	168 3	20. 6	250	17.9	193 3	20. 2
7	164 7	21. 8	295	14.7	136 7	22. 7	575	16.2	174 7	21. 4	195	14.0	194 2	20. 3
8	180 5	23. 9	388	19.4	148 3	24. 7	710	20.0	191 3	23. 4	280	20.1	219 3	22. 9
9	433	5.7	90	4.5	355	5.9	168	4.7	460	5.6	63	4.5	523	5.5
10	244	3.2	75	3.7	186	3.1	133	3.7	256	3.1	63	4.5	319	3.3
11	39	0.5	16	0.8	31	0.5	24	0.7	42	0.5	13	0.9	55	0.6
12	110	1.5	36	1.8	75	1.2	71	2.0	115	1.4	31	2.2	146	1.5
Sleep Duration (Categorical)														
≤6 Hours	328 4	43. 4	1103	55.1 *	251 4	41. 8	1873	52.7 *	363 7	44. 5	750	53.8 *	438 7	45. 9
7-8 Hours	345 2	45. 6	683	34.1	285 0	47. 4	1285	36.2	366 0	44. 8	475	32.1	413 5	43. 2



## Sleep and Long COVID

≥9 Hours	826	10.9	217	10.8	647	10.8	396	11.1	873	10.7	170	12.2	1043	10.9
----------	-----	------	-----	------	-----	------	-----	------	-----	------	-----	------	------	------

† $P < 0.05$ , \* $P < 0.01$  vs. corresponding placebo Group

‡COPE: The COVID-19 Outbreak Public Evaluation Initiative

#NICE: National Institute for Health and Care Excellence

¶PASC: Post-Acute Sequelae of SARS-CoV-2 infection

## Sleep and Long COVID

Table 2: Association of Self-reported Insomnia with Three Models of PASC

Model	COPE <sup>a</sup> (≥3 Symptoms)		NICE <sup>b</sup> (≥ 1 Symptom)		PASC <sup>c</sup> Score (≥12)	
	<i>aOR</i>	95% CI	<i>aOR</i>	95% CI	<i>aOR</i>	95% CI
Baseline	2.94 <sup>§</sup>	2.71-3.31	3.58 <sup>§</sup>	3.22-3.73	3.58 <sup>§</sup>	3.23-3.96
+Demographics*	2.89 <sup>§</sup>	2.61-3.20	3.20 <sup>§</sup>	2.99-3.45	3.21 <sup>§</sup>	2.89-3.57
+Comorbidities†	1.75 <sup>§</sup>	1.55-1.98	1.79 <sup>§</sup>	1.63-1.96	1.63 <sup>§</sup>	1.43-1.86
+Socioeconomic‡	1.74 <sup>§</sup>	1.54-1.97	1.78 <sup>§</sup>	1.62-1.95	1.61 <sup>§</sup>	1.42-1.84
+Sleep Factors§	1.46 <sup>§</sup>	1.28-1.68	1.52 <sup>§</sup>	1.34-1.71	1.30 <sup>§</sup>	1.11-1.52

<sup>a</sup>COPE: The COVID-19 Outbreak Public Evaluation Initiative

<sup>b</sup>NICE: National Institute for Health and Care Excellence

<sup>c</sup>PASC: Post-Acute Sequelae of SARS-CoV-2 infection

Models adjusted as follows (see text for definitions):

\*Age, Sex, Race

†BMI, # of the following conditions: diabetes, asthma, sickle cell disease, cardiovascular disease, hypertension, cancer, chronic kidney disease, liver disease, chronic obstructive pulmonary disease, boosted vaccination status

‡Education, Income, Employment

§Sleep Duration, Obstructive Sleep Apnea

§p<0.001

## Sleep and Long COVID

Table 3: Association of Poor Sleep Quality with Three Models of PASC

Model	COPE <sup>a</sup> (≥3 Symptoms)		NICE <sup>b</sup> (≥ 1 Symptom)		PASC <sup>c</sup> Score (≥12)	
	aOR	95% CI	aOR	95% CI	aOR	95% CI
Baseline	2.63 <sup>§</sup>	2.39-2.90	2.36 <sup>§</sup>	2.18-2.56	2.42 <sup>§</sup>	2.17-2.71
+Demographics*	2.56 <sup>§</sup>	2.32-2.82	2.27 <sup>§</sup>	2.09-2.47	2.36 <sup>§</sup>	2.11-2.64
+Comorbidities†	2.13 <sup>§</sup>	1.91-2.38	1.92 <sup>§</sup>	1.75-2.11	1.99 <sup>§</sup>	1.75-2.25
+Socioeconomic‡	2.11 <sup>§</sup>	1.89-2.37	1.92 <sup>§</sup>	1.75-2.12	1.98 <sup>§</sup>	1.75-2.25
+Sleep Factors§	2.00 <sup>§</sup>	1.77-2.26	1.77 <sup>§</sup>	1.60-1.97	1.89 <sup>§</sup>	1.64-2.18

<sup>a</sup>COPE: The COVID-19 Outbreak Public Evaluation Initiative

<sup>b</sup>NICE: National Institute for Health and Care Excellence

<sup>c</sup>PASC: Post-Acute Sequelae of SARS-CoV-2 infection

Models adjusted as follows (see text for definitions):

\*Age, Sex, Race

†BMI, # of the following conditions: diabetes, asthma, sickle cell disease, cardiovascular disease, hypertension, cancer, chronic kidney disease, liver disease, chronic obstructive pulmonary disease, boosted vaccination status

‡Education, Income, Employment

§Sleep Duration, Obstructive Sleep Apnea

<sup>§</sup>p<0.001

## Sleep and Long COVID

Table 4: Odds ratio (adjusted) for association of self-reported hourly sleep duration\* with 3 models of PASC

		Unadjusted		Fully Adjusted†		Unadjusted		Fully Adjusted†		Unadjusted		Fully Adjusted†	
		COPE <sup>a</sup> (≥3 Symptoms)				NICE <sup>b</sup> (≥ 1 Symptom)				PASC <sup>c</sup> Score (≥12)			
Sleep Duration, Hours*	N	aOR	95% CI	aOR	95% CI	aOR	95% CI	aOR	95% CI	aOR	95% CI	aOR	95% CI
3	376	3.07 <sup>§</sup>	2.45-3.84	1.56 <sup>&amp;</sup>	1.18-2.05	2.99 <sup>§</sup>	2.42-3.71	1.49 <sup>&amp;</sup>	1.15-1.94	2.61 <sup>§</sup>	2.03-3.35	1.40 <sup>§</sup>	1.04-1.89
4	782	2.38 <sup>§</sup>	2.00-2.82	1.57 <sup>§</sup>	1.28-1.92	2.35 <sup>§</sup>	2.01-2.74	1.57 <sup>§</sup>	1.30-1.89	2.22 <sup>§</sup>	1.83-2.70	1.55 <sup>§</sup>	1.24-1.95
5	1296	1.81 <sup>§</sup>	1.56-2.09	1.23 <sup>§</sup>	1.03-1.48	1.75 <sup>§</sup>	1.54-1.99	1.25 <sup>&amp;</sup>	1.07-1.46	1.66 <sup>§</sup>	1.40-1.97	1.15	0.94-1.41
6	1933	1.20 <sup>§</sup>	1.04-1.38	1.05	0.89-1.24	1.21 <sup>§</sup>	1.08-1.36	1.09	0.95-1.25	1.15	0.97-1.35	0.99	0.82-1.20
7-8	4135	Reference		Reference		Reference		Reference		Reference		Reference	
9	523	1.05	0.83-1.34	0.96	0.71-1.28	1.05	0.86-1.28	0.92	0.72-1.17	1.06	0.80-1.40	0.92	0.66-1.30
10	319	1.55 <sup>§</sup>	1.18-2.04	1.15	0.83-1.61	1.59 <sup>§</sup>	1.26-2.00	1.20	0.90-1.59	1.90 <sup>§</sup>	1.42-2.54	1.34	0.94-1.92
11	55	2.07 <sup>§</sup>	1.15-3.73	2.05	0.97-4.34	1.72 <sup>§</sup>	1.00-2.94	1.56	0.76-3.19	2.39 <sup>&amp;</sup>	1.27-4.48	2.45 <sup>§</sup>	1.13-5.31
12	146	1.65 <sup>&amp;</sup>	1.12-2.43	1.15	0.73-1.82	2.10 <sup>§</sup>	1.51-2.92	1.47	0.98-2.19	2.08 <sup>§</sup>	1.38-3.12	1.53	0.95-2.46

<sup>a</sup>COPE: The COVID-19 Outbreak Public Evaluation Initiative

<sup>b</sup>NICE: National Institute for Health and Care Excellence

<sup>c</sup>PASC: Post-Acute Sequelae of SARS-CoV-2 infection

\*Self-reported sleep duration (3 to 12 hours); Reference is 7-8 hours

†Fully adjusted model includes Age, Sex, Race, BMI, insomnia, # of comorbidities, Education, Income and Employment, Insomnia, Obstructive Sleep Apnea, Poor Sleep Quality

<sup>§</sup>p<0.05

<sup>&</sup>p<0.01

<sup>§</sup>p<0.001

## Sleep and Long COVID

Table 5: Association of Self-reported Sleep Duration with Three Models of PASC

Model	COPE <sup>a</sup> (≥3 Symptoms)		NICE <sup>b</sup> (≥ 1 Symptom)		PASC Score <sup>c</sup> (≥12)	
	aOR	95% CI	aOR	95% CI	aOR	95% CI
<b>SLEEP ≤6 HOURS<sup>#</sup></b>						
Unadjusted	1.70 <sup>§</sup>	1.53-1.89	1.62 <sup>§</sup>	1.51-1.81	1.59 <sup>§</sup>	1.40-1.80
+Demographics <sup>*</sup>	1.66 <sup>§</sup>	1.49-1.85	1.61 <sup>§</sup>	1.47-1.76	1.56 <sup>§</sup>	1.37-1.76
+Comorbidities <sup>†</sup>	1.52 <sup>§</sup>	1.35-1.72	1.48 <sup>§</sup>	1.33-1.64	1.41 <sup>§</sup>	1.23-1.63
+Socioeconomic <sup>‡</sup>	1.55 <sup>§</sup>	1.37-1.75	1.51 <sup>§</sup>	1.36-1.68	1.43 <sup>§</sup>	1.24-1.65
+Sleep Factors <sup>§</sup>	1.23 <sup>§</sup>	1.08-1.41	1.23 <sup>§</sup>	1.10-1.37	1.17 <sup>§</sup>	1.01-1.36
<b>SLEEP ≥9 HOURS<sup>#</sup></b>						
Unadjusted	1.33 <sup>§</sup>	1.12-1.58	1.36 <sup>§</sup>	1.18-1.56	1.50 <sup>§</sup>	1.24-1.82
+Demographics <sup>¶</sup>	1.20 <sup>¶</sup>	1.01-1.43	1.22 <sup>&amp;</sup>	1.05-1.41	1.35 <sup>&amp;</sup>	1.11-1.64
+Comorbidities <sup>†</sup>	1.05	0.86-1.28	1.05	0.89-1.24	1.18	0.95-1.48
+Socioeconomic <sup>‡</sup>	1.12	0.92-1.38	1.11	0.94-1.32	1.25	0.99-1.56
+Sleep Factors <sup>§</sup>	1.12	0.91-1.37	1.12	0.94-1.33	1.24	0.99-1.55

<sup>a</sup>COPE: The COVID-19 Outbreak Public Evaluation Initiative

<sup>b</sup>NICE: National Institute for Health and Care Excellence

<sup>c</sup>PASC: Post-Acute Sequelae of SARS-CoV-2 infection

Models adjusted as follows (see text for definitions):

<sup>#</sup>Referent is 7-8 Hours

<sup>\*</sup>Age, Sex, Race

<sup>†</sup>BMI, # of the following conditions: diabetes, asthma, sickle cell disease, cardiovascular disease, hypertension, cancer, chronic kidney disease, liver disease, chronic obstructive pulmonary disease, boosted vaccination status

<sup>‡</sup>Education, Income, Employment

<sup>§</sup>Insomnia, Obstructive Sleep Apnea, Sleep Quality

<sup>¶</sup>p≤0.05; <sup>&</sup>p≤0.01, <sup>§</sup>p<0.001