

Original article

Risk Factors and Outcomes of Pulmonary Hemorrhage in Preterm Infants born before 32 weeks.

Gieng Thi My Tran², Nhat Phan Minh Nguyen^{1,4}, Nguyen Phuoc Long⁶, Duc Ninh Nguyen⁷,
Thu-Tinh Nguyen^{1,3,5*}.

Affiliations

¹Department of Pediatrics, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, 700000, Vietnam. E-mail: nhatnguyen@ump.edu.vn

²Department of Neonatology, Thien Hanh Hospital, Daklak Province, 630000, Vietnam. E-mail: mygiengtranthi@gmail.com

³Neonatal Intensive Care Unit, Children's Hospital 2, Ho Chi Minh City, 700000, Vietnam.

⁴Neonatal Department, Children's Hospital 2, Ho Chi Minh City, 700000, Vietnam.

⁵University Medical Center, Ho Chi Minh City, 70000, Vietnam.

⁶Department of Pharmacology and Pharmacogenomics Research Center, Inje University College of Medicine, Busan 47392, Republic of Korea. E-mail: phuoclong@inje.ac.kr

⁷Section for Comparative Pediatrics and Nutrition, Department of Veterinary and Animal Sciences, University of Copenhagen, Denmark. E-mail: dnn@sund.ku.dk

***Corresponding Author:**

Tinh-Thu Nguyen, MD, PhD;

Department of Pediatrics,

University of Medicine and Pharmacy at Ho Chi Minh City,

217 Hong Bang Street, District 5, Ho Chi Minh City, 700000, Vietnam.

Tel: +84-937-911-277

E-mail: tinhnguyen@ump.edu.vn

ORCID: 0000-0002-8198-3170

Category of study: Pediatric Research

Impact

1. High incidence and mortality of pulmonary hemorrhage in preterm infants < 32 weeks' gestational age with respiratory distress in a Neonatal Intensive Care Unit in Vietnam.
2. Pulmonary hemorrhage should be considered in the clinical deterioration of preterm infants given invasive ventilation in the first 2-4 days of life.
3. Independent risk factors for pulmonary hemorrhage: intubation in the first 24 hours, coagulation disorders, and transfusion of blood products in the first 3 days of life.
4. Urgent need to seek diagnostic criteria for intraventricular hemorrhage as soon as pulmonary hemorrhage occurs.

1 **Abstract**

2 **Background:** Pulmonary hemorrhage (PH) is a catastrophic event associated with significant
3 morbidity and mortality among preterm infants. Understanding PH risk factors in preterm
4 newborns, particularly those in low-to-middle-income countries like Vietnam, remains
5 limited. This study aimed to investigate the risk factors and short-term outcomes of PH in
6 very preterm infants.

7 **Methods:** We conducted an observational study of newborns aged < 72 hours with
8 gestational age < 32 weeks, admitted to our unit from April 1, 2018 to March 31, 2019.

9 **Results:** Of 118 recruited newborns, 28 (23.7%) had PH. The logistic regression analysis
10 showed that intubation within 24 first hours, blood transfusion, and coagulation disorders
11 within the first 3 days were risk factors for PH (aOR = 4.594, 95% CI = 1.200-17.593; aOR =
12 5.394, 95% CI = 1.243-23.395 and aOR = 7.242 95% CI = 1.838-28.542, respectively). Intra-
13 ventricular hemorrhage (IVH) and mortality rates were higher in patients with PH compared
14 to those without ($p < 0.001$). The length of invasive ventilation was longer in the PH group
15 ($p < 0.001$).

16 **Conclusion:** We have identified intubation, blood transfusion, and coagulation disorders
17 shortly after birth as risk factors for PH in very preterm infants, which was associated with
18 increased mortality and occurrence of IVH.

Keywords: blood coagulation disorders, blood transfusion, intratracheal intubation,
premature, pulmonary hemorrhage.

19

20 **Introduction**

21 Pulmonary hemorrhage (PH) is a life-threatening condition with blood extravasation into the
22 alveoli, causing significant morbidity and mortality in hospitalized preterm infants. It occurs
23 in up to 1.2% of total live births¹, with a particularly high incidence in preterm infants (up to
24 5%)¹. PH currently has no preventive treatment and is most commonly diagnosed during the
25 first few days of life; its mortality rates could rise to 50%, especially in extremely preterm
26 infants². Its diagnostic criteria include the aspiration of secretions from the trachea or
27 endotracheal tube^{3,4}. The severity can vary from a mild, self-limited disorder to a massive,
28 deteriorating, and end-stage syndrome. Upon PH diagnosis in preterm infants, invasive
29 ventilation and prolonged intensive care are often required, further elevating the risks of other
30 significant morbidity and mortality^{3,4,3}. In preterm infants, it is often manifested from
31 pulmonary edema and hemodynamically significant patent ductus arteriosus (hsPDA)^{3,4}.
32 Asphyxia, prematurity, intrauterine growth restriction, early-onset sepsis, patent ductus
33 arteriosus, and surfactant replacement have been shown as major risk factors of PH in several
34 geographical settings²⁻⁵. Awareness of PH risk factors may help clinicians realize the early
35 symptoms and develop prompt strategies for PH management.

36 The already reported risk factors of PH in preterm newborns are variable across different
37 settings. In Vietnam, PH incidence and associated factors are unknown. Therefore, this study
38 aimed to explore the perinatal associated factors and short-term outcomes of PH in very
39 preterm newborns in a neonatal intensive care unit (NICU) in Vietnam, which will provide
40 future benefits for early triage and the management of critically ill patients.

41 **Materials and Methods**

42 **Study design**

This retrospective observational study was conducted at the NICU of the Children's Hospital
1 (a level IV NICU) from April 1st, 2018 to March 31st, 2019 in Ho Chi Minh City, Vietnam.

The study protocol was reviewed and approved by the Institutional Review Board of Children's Hospital 1 (IORG0007285, FWA00009748, Approval number: 2784/QĐ-BVND1, On October 16th, 2019). Written informed consent was waived by the Institutional Review Board because this was a retrospective study. There was no patient contact, and some of the patients were deceased.

43 The inclusion criteria include: (i) gestational age at birth < 32 weeks, (ii) being admitted to
44 the NICU within 72 hours following the birth, and (iii) absence of congenital anomalies,
45 including congenital heart disease other than patent ductus arteriosus (PDA), structural
46 anomalies of the gastrointestinal tract, and other anomalies requiring surgery. Patients
47 developing PH prior to admission to our center were also excluded.

48 PH diagnosis was performed according to the international standard with hemorrhagic
49 secretions aspirated from the trachea or endotracheal tube, accompanied by clinical
50 deterioration, which included the requirement of intubation and ventilation or the need for an
51 increased fraction of inspired oxygen (FiO₂) by at least 10% ³. A chest X-ray was also taken
52 to confirm diffuse pulmonary infiltrates.

53 Information during the pregnancy and hospitalization was collected retrospectively from the
54 medical notes. Maternal history including antenatal corticosteroids, complications of this
55 pregnancy (gestational diabetes, pre-eclampsia/ eclampsia, hypertension, urinary infection),
56 intrapartum fever. Neonatal baseline details including gender, gestational age (calculated
57 from the first day of the last menstrual period or prenatal obstetric ultrasonography in the first
58 trimester), birth weight, mode of delivery, small for gestational age (SGA < 10 percentile on
59 Fenton chart), intubation within first 24h of age, treatment with surfactant, multiple doses of
60 surfactant prior the onset of PH, coagulation problems within the first 3 days, blood
61 transfusion within the first 3 days (prior the occurrence of PH), platelet count < 100000/mm³,
62 hsPDA. Echocardiography was used to diagnose PDA by a cardiologist from day 3 – 7 and

63 repeated when needed. The hsPDA was managed by consensus among neonatologists and
64 cardiologists, and oral ibuprofen or intravenous acetaminophen was given to close the PDA.
65 Surgery was performed in cases of contraindications to ibuprofen/ acetaminophen or failed 2
66 courses of ibuprofen/acetaminophen therapy.

67 The study's primary outcomes include the incidence of PH. Secondary outcomes include PH
68 risk factors, the rate of mortality, IVH, retinopathy of prematurity (ROP) and
69 bronchopulmonary dysplasia (BPD), length of invasive ventilation, and NICU stay in patients
70 with vs. without PH. IVH was identified by cranial ultrasound and classified based on Papile
71 criteria. ROP was screened by an ophthalmologist when infants were 4 weeks of age or 31
72 weeks of postmenstrual age, whichever came last. BPD was defined as the requirement for
73 oxygen supplementation at 36 weeks postmenstrual age or at 28 days old.

74 *Statistical analysis*

75 The data were imported into Excel. Descriptive statistical analysis was used to describe the
76 characteristics of preterm newborns and their mothers. The normally distributed results are
77 reported as the mean and standard deviation (SD); the remaining results are reported as the
78 median, interquartile range (IQR), or percentage. Chi-squared test, Mann-Whitney test,
79 Fisher's test, Student's *t*-test, and logistic regression model were used for statistical analysis.
80 Multiple logistic regression was applied to classify patients into the binary state of PH
81 (Yes/No). Since the sample size was relatively small ($N = 118$) and the PH class ratio was
82 imbalanced ($N_{\text{Yes}} = 28$, $N_{\text{No}} = 90$), a ℓ_2 -penalization was applied to the logistic regression
83 model to alleviate the potential bias in classification and coefficient estimation⁶. A nested *k*-
84 fold cross-validation procedure was conducted to assess the performance of the classification
85 model. The outer loop ($k = 5$) is for splitting training (70%) and test (30%) data, while the
86 inner loop ($k = 10$) is for hyperparameters tuning. The generalization performance of the
87 model was obtained from the nested cross-validation procedure by taking the average for

88 each metric over all the outer-loop folds. The area under the curve (AUC) of the Receiver
89 Operating Characteristic curve was used as a metric to assess model performance. Moreover,
90 the ridge logistic regression was interpreted, and adjusted odds ratios (aORs) with their
91 corresponding 95% confidence intervals (CIs) were computed based on regression
92 coefficients and standard error. Specifically, a ridge logistic regression model was first
93 trained on the whole dataset using a 10-fold cross-validation procedure to select the best-
94 performing hyperparameters. Subsequently, using the optimal set of hyperparameters, we
95 fitted the ℓ_2 -penalized logistic regression model on the whole dataset to estimate the
96 regression coefficients for each clinical covariate. Model training and validation were
97 conducted using the *caret* package (version 6.0-94)⁷ in R version 4.3.2. Before the analyses,
98 categorical variables were dummy encoded, while continuous variables were centered and
99 scaled.

100 **Results**

101 *Baseline characteristics*

102 From April 1st, 2018 to March 31st, 2019, there were 171 preterm newborns with gestational
103 age < 32 weeks admitted to the NICU of Children Hospital 1. One had severe congenital
104 heart disease, one had hydrocephalus, five cases had congenital gastrointestinal anomalies,
105 and 46 were admitted after 72 hours of age. There were a total of 118 patients enrolled in the
106 study: 28/118 (23.7%) patients were diagnosed with PH, and 21/28 (75%) patients in the PH
107 group were extremely preterm newborns. The male/ female ratio was 0.96. Other baseline
108 parameters comparing patients with vs. without PH are depicted in Table 1.

109 *Clinical manifestations*

110 The median time for the onset of PH is 47 (32.5-73) hours. The earliest and latest cases in the
111 PH group happened at 28 hours and 144 hours, respectively. It took 38 (26-48.5) hours from
112 the first dose of surfactant to PH manifestation.

113 The main clinical characteristics of the enrolled patients are also described in Table 1. The
114 analysis of neonatal factors showed that newborns with PH had lower gestational age and
115 birth weight ($p < 0.001$). Additionally, infants in the PH group were more likely to be
116 intubated in the first 24 hours, treated with surfactant, had been transfused, and had
117 coagulation problems.

118 Risk factors for PH in this cohort were analyzed via a logistic regression model. The model
119 exhibited considerable performance in classifying patients who underwent PH and those who
120 did not [AUC and standard deviation (SD) from the nested cross-validation = 0.863 ± 0.065]
121 (Figure 1). The aORs of clinical covariates and their corresponding 95% CIs were presented
122 in Figure 2 and Table 2. Three covariates were significantly associated with PH. In detail,
123 patients who had coagulation dysfunction in the first 3 days increased the odds of acquiring
124 PH with an aOR = 7.242 (95% CI = 1.838-28.542). In the same vein, blood transfusion prior
125 to PH in the first 3 days (aOR = 5.394, 95% CI = 1.243-23.395) and intubation in the first 24
126 hours (aOR = 4.594, 95% CI = 1.200-17.593) were positively associated with undergoing
127 PH.

128 Short-term outcomes between PH and no-PH group are described in Table 3. Newborns
129 who had PH showed a higher mortality rate (64.3% vs 25.6%, $p = 0.001$). Grade II to IV
130 intraventricular hemorrhage was more frequent in the group with PH (32.1% vs 4.4%,
131 $p = 0.001$). Patients in the PH group needed a longer time of invasive ventilation than those in
132 the no-PH group.

133 **Discussion**

134 PH is thought to occur in up to 4% to 12% of VLBW infants⁴. Similarly, the incidence of PH
135 was 54/1350 (4%) in a prospective cohort study by Gezmu et al., which was conducted in a
136 tertiary-level hospital in Botswana that included 1350 newborns enrolled during a 2-year
137 period⁸. The population of that study had a mean (SD) birth weight was 2154 (± 997.5) grams

138 and a gestational age was 34.3 (± 4.7) weeks. However, the incidence of PH in our center
139 accounts for 23.7% among newborns with gestational age at birth below 32 weeks. The mean
140 gestational age of our patients is lower than those of Gezmu's study (25.5 weeks vs. 34.3
141 weeks), and it may affect the rate of PH.

142 The mean age of onset of PH varied between 40 - 72 hours^{9,10}. More than 80% of cases of PH
143 onset within 3 days of life with a median of 40 hours¹⁰. In 2013, a study by Yen TA et al.
144 demonstrated that the mean age of infants who have birth weight < 1500gr with PH
145 occurrence was 2.4 ± 1.3 days¹¹. Additionally, the study of Wang et al. revealed that the
146 median age of PH occurrence was 3 (2-4,5) days in extremely low birth weight (ELBW)
147 infants³. Similarly, the mean time for the onset of PH was 47 hours after birth in our study.

148 In our study, we assessed risk factors for PH by multivariate analysis and identified that
149 intubation in the first 24 hours, blood transfusion prior to PH in the first 3 days, and
150 coagulation problems in the first 3 days were independently associated factors for PH.
151 Similar to our result, Ferreira et al. also found that preterm infants who needed intubation in
152 the delivery room were associated with PH occurrence⁴. As can be seen in our study design,
153 Children's Hospital 1 has a level IV NICU, where obtaining patients' referrals from nearby
154 regional hospitals and intubation may be indicated conditionally for airway control while
155 transferring. The need for intubation and positive pressure ventilation can cause excessive
156 alveolar distension and lead to stress damage to the alveolar capillaries, which may activate
157 the genesis of PH¹².

158 In the multiple logistic regression, we found there is an association between PH and usage of
159 blood products prior to PH in the first 3 days of life. Transfusion of blood products is thought
160 to increase pulmonary capillary pressure, leading to stress injury of the capillary wall⁴.
161 Additionally, a clinical disorder which is known as transfusion-related acute lung injury
162 (TRALI) can be detected in patients who experience a great deal of stress, which may impact

163 the occurrence of PH¹³. TRALI begins with increasing the levels of interleukin-8 and
164 interleukin-6, which results in neutrophil recruitment to the pulmonary vasculature, and
165 pulmonary edema can eventually happen. The onset of TRALI can be during or from 6–72
166 hours after a transfusion.

167 Coagulation dysfunction in the first 3 days of life also showed an association with PH in a
168 multivariate logistic regression model. In 2021, a study by Jing Li et al. found that PDA,
169 coagulopathy, and 10-minute Apgar Score were risk factors for PH¹⁴. Coagulation
170 dysfunction in preterm infants can be related to hypoxia due to respiratory distress syndrome
171 (RDS) and sepsis. On the other hand, preterm infants may need surfactant replacement in the
172 case of RDS, and surfactant was reported to have an impact on the impairment of coagulation
173 in-vitro¹⁵.

174 Prior studies reported the association of PDA with the occurrence of PH^{1,14}. From 2 days of
175 age, the reduction of pulmonary vascular resistance leads to increased left-to-right shunting
176 through PDA and pulmonary blood flow and results in elevating the pressure state of the
177 pulmonary vessels, which may compromise cardiac function with an increased risk of PH.
178 However, we do not reveal this association when analyzed in a multivariate logistic
179 regression model. It can be explained that the preterm infants less than 28 weeks in our center
180 had been given prophylactic treatment with acetaminophen and fluid restriction to reduce the
181 risk of hspDA.

182 Several studies showed that surfactant replacement may increase the risk of PH^{2,4,16}, whereas
183 other studies reported no impact of surfactant replacement^{3,17}. In the current study, our data
184 showed that PH was associated with surfactant replacement within 24 hours of age and
185 repeated surfactant therapy. It is reasonable to postulate that the illness of infants who need
186 surfactant are clinically severe and more likely to have PH than those who do not need

187 surfactant. However, multivariate logistic regression analysis revealed no interactive effects
188 of surfactant and PH. This is in agreement with some of the previous reports^{17,18}.

189 For short-term outcomes of PH, we found a strong association between PH and mortality
190 rate, similar to previous studies^{2,5,17}. Another significant outcome observed in this study was
191 IVH. PH and IVH were reported as a result of hemodynamic instability and coagulation
192 disorders⁸. The study by Wang TT et al. found that the IVH rate of the group of extremely
193 low birth weight infants (ELBWIs) with PH was significantly higher than those of the group
194 of ELBWIs without PH³. In our study, the rate of grade II-IV IVH was significantly higher in
195 infants with PH than those without PH (32.1% and 4.4%, respectively, $P < 0.001$). On the
196 other hand, IVH can also be found in 24.4% of extremely preterm infants with hsPDA, but
197 there has been no evidence that the presence of a hsPDA is associated with a higher
198 probability of IVH. As mentioned above, the incidence of hsPDA was 48.3% among our
199 patients, and it was higher in the PH group compared to the no-PH group but not statistically
200 significant.

201 Supportive management of PH was positive ventilation, blood transfusion, and vasopressor
202 administration¹⁹. In our study, the duration of invasive ventilation and nCPAP support was
203 significantly longer in the PH group than in the no-PH group ($p=0.001$). However, there were
204 no significant differences in NICU stay between surviving infants in the PH and no-PH
205 groups ($p=0.6$).

206 Our unit is a level IV NICU that primarily cares for out-born infants transferred from
207 neonatal units in the Obstetric Hospitals in Ho Chi Minh City and the surrounding regions.

208 This was a retrospective study in a single center, which may not be able to highlight all the
209 associated factors of PH in very preterm newborns due to the limited data and small sample
210 size. However, analyzing the risk factors of PH will help physicians to understand better why
211 PH occurs and how to prevent it.

212 **Conclusion**

213 PH is a severe and dreadful complication that has a common onset at 2-4 days of age in the
214 population of newborns with <32 weeks, particularly in extremely preterm infants. The risk
215 of PH increases with decreasing gestational age of live-born infants, especially those who
216 have been intubated for mechanical ventilation on the first day of life, have coagulation
217 problems within the first 3 days of life, and have blood products transfusion within the first 3
218 days. IVH should be investigated as soon as the diagnosis of PH is confirmed.

219 **Acknowledgments**

We would like to acknowledge all participants who were enrolled in the study and their parents and families. We would like to acknowledge our supporting colleagues at the University of Medicine and Pharmacy at Ho Chi Minh City and Children's Hospital 1 in Ho Chi Minh City.

Funding

The authors received no financial support for the research, authorship, and publication of this article.

Author contributions

Conceptualization: TTN, GTMT; Data collection: GTMT, TTN; Formal analysis: GTMT, TTN, NPMN, NPL, DNN; Methodology: TTN, GTMT, NPMN, NPL, DNN; Supervision: TTN; Writing draft manuscript: GTMT, TTN, NPMN, NPL, DNN; Review and edit manuscript: GTMT, TTN, NPMN, NPL, DNN; Approve the final manuscript: all authors.

Conflicting Interests

The authors have no conflicts of interest to declare.

Data availability statement:

Data for this study are contained within patient medical records and are not publicly available due to privacy restrictions. The data are available from the corresponding author, Thu-Tinh Nguyen, upon reasonable request.

References

- 220 1. Zahr, R. A., Ashfaq, A. & Marron-Corwin, M. Neonatal Pulmonary Hemorrhage.
221 *Neoreviews* **13**, e302–e306 (2012).
- 222 2. Ahmad, K. A., Bennett, M. M., Ahmad, S. F., Clark, R. H. & Tolia, V. N. Morbidity
223 and mortality with early pulmonary haemorrhage in preterm neonates. *Arch Dis Child*
224 *Fetal Neonatal Ed* **104**, F63–F68 (2019).
- 225 3. Wang, T.-T., Zhou, M., Hu, X.-F. & Liu, J.-Q. Perinatal risk factors for pulmonary
226 hemorrhage in extremely low-birth-weight infants. *World Journal of Pediatrics* **16**,
227 299–304 (2020).
- 228 4. Ferreira, C. H., Carmona, F. & Martinez, F. E. Prevalence, risk factors and outcomes
229 associated with pulmonary hemorrhage in newborns. *J Pediatr (Rio J)* **90**, 316–322
230 (2014).
- 231 5. Cao, Z.-L. *et al.* [Pulmonary hemorrhage in very low birth weight infants: risk factors
232 and clinical outcome]. *Zhongguo Dang Dai Er Ke Za Zhi* **24**, 1117—1123 (2022).
- 233 6. Šinkovec, H., Heinze, G., Blagus, R. & Geroldinger, A. To tune or not to tune, a case
234 study of ridge logistic regression in small or sparse datasets. *BMC Med Res Methodol*
235 **21**, 199 (2021).
- 236 7. Kuhn, M. Building Predictive Models in *R* Using the **caret** Package. *J Stat Softw* **28**,
237 (2008).
- 238 8. Gezmu, A. M. *et al.* Pulmonary hemorrhage and associated risk factors among
239 newborns admitted to a tertiary level neonatal unit in Botswana. *Front Pediatr* **11**,
240 (2023).
- 241 9. Finlay, E. R. & Subhedar, N. V. Pulmonary haemorrhage in preterm infants. *Eur J*
242 *Pediatr* **159**, 870–871 (2000).

- 243 10. Tomaszewska, M. *et al.* Pulmonary Hemorrhage. *Arch Pediatr Adolesc Med* **153**, 715
244 (1999).
- 245 11. Yen, T.-A. *et al.* Short-term Outcome of Pulmonary Hemorrhage in Very-Low-Birth-
246 Weight Preterm Infants. *Pediatr Neonatol* **54**, 330–334 (2013).
- 247 12. Berger, T. M., Allred, E. N. & Van Marter, L. J. Antecedents of Clinically Significant
248 Pulmonary Hemorrhage Among Newborn Infants. *Journal of Perinatology* **20**, 295–
249 300 (2000).
- 250 13. Marik, P. E. & Corwin, H. L. Acute lung injury following blood transfusion:
251 Expanding the definition. *Crit Care Med* **36**, 3080–3084 (2008).
- 252 14. Li, J., Xia, H., Ye, L., Li, X. & Zhang, Z. Exploring prediction model and survival
253 strategies for pulmonary hemorrhage in premature infants: a single-center,
254 retrospective study. *Transl Pediatr* **10**, 1324–1332 (2021).
- 255 15. Strauss, T. *et al.* Surfactant impairs coagulation in-vitro: A risk factor for pulmonary
256 hemorrhage? *Thromb Res* **132**, 599–603 (2013).
- 257 16. Chen, Y. *et al.* Pulmonary hemorrhage in very low birthweight infants: Risk factors
258 and management. *Pediatrics International* **54**, 743–747 (2012).
- 259 17. Berger, T. M., Allred, E. N. & Van Marter, L. J. Antecedents of Clinically Significant
260 Pulmonary Hemorrhage Among Newborn Infants. *Journal of Perinatology* **20**, 295–
261 300 (2000).
- 262 18. De Carolis, M. P. *et al.* Pulmonary haemorrhage in infants with gestational age of less
263 than 30 weeks. *Eur J Pediatr* **157**, 1037–1038 (1998).
- 264 19. Barnes, M. E. *et al.* Pulmonary haemorrhage in neonates: Systematic review of
265 management. *Acta Paediatr* **111**, 236–244 (2022).
- 266
- 267

268 **Table 1:** Comparison of clinical characteristics between patients with and without pulmonary

Variables	aOR	95% Confidence Interval	P-value
Characteristics			
Gestational age (n (%) or median [25-75pt])	Pulmonary hemorrhage 0.435	No Pulmonary hemorrhage 0.148-1.279	p-value 0.130
Birth weight	n = 28 0.956	n = 90 0.319-2.867	0.937
Not given antenatal steroids	1.097	0.287-4.201	0.892
Gender (female)	15 (53.6)	45 (50.0)	0.741
Gestational age (weeks)	25.5 [25-27.8]	28 [27-30]	0.001*
Birth weight (g)	725 [657.5-1000]	1100 [947.5-1400]	0.001*
Small for gestational age	3 (10.7)	5 (5.6)	0.392
Not given antenatal steroids	22 (78.6)	51 (56.7)	0.046
Cervical incompetence	2 (7.1)	7 (7.8)	1.00
Intrapartum mother's fever	2 (7.1)	7 (7.8)	1.00
Intubation within first 24 hours	18 (64.3)	22 (24.4)	0.001
Surfactant therapy within first 24 hours	23 (82.1)	41 (45.6)	0.001
Given second surfactant dose	6 (21.4)	2 (2.2)	0.002
Blood transfusion prior PH within the first 3 days of life.	20 (71.4)	8 (8.9)	0.001
Coagulopathy within the first 3 days	22 (78.6)	8 (8.9)	0.001 [†]
Hemodynamically significant PDA	17 (60.7)	40 (44.4)	0.132 [†]
Platelet count < 100.000/mm ³ ‡	7 (25.0)	8 (8.9)	0.046

269 hemorrhage.

270 * *Mann-Whitney test* † *Chi-square test*, ‡: *Fisher's test*.

271 *Abbreviation: PDA (Patent Ductus Arteriosus), PH (Pulmonary Hemorrhage)*

Intubation within first 24 hours	4.594	1.200-17.593	0.026
Surfactant therapy within first 24 hours	0.890	0.217-3.650	0.871
Given second surfactant dose	1.250	0.195-8.005	0.813
Blood transfusion prior PH within the first 3 days of life	5.394	1.243-23.395	0.024
Coagulopathy within the first 3 days	7.242	1.838-28.542	0.005
Hemodynamically significant PDA	2.434	0.594-9.966	0.216
PDA prophylaxis	0.394	0.082-1.899	0.246
PDA ligation	0.452	0.108-1.887	0.276
Platelet count < 100.000/mm ³	0.790	0.166-3.758	0.767

272 **Table 2:** Adjusted odds ratio and 95% confidence intervals estimated from ridge logistic
273 regression model

274

275 **Table 3:** Short-term outcomes of pulmonary hemorrhage

Variables	Pulmonary hemorrhage (n = 28)	No Pulmonary hemorrhage (n = 90)	p-value
Death, n (%) [*]	18 (64.3)	23 (25.6)	0.001
Intraventricular hemorrhage grade II-IV, n (%) [†]	9 (32.1)	4 (4.4)	0.001
Moderate- severe BPD, n (%) [†]	5 (17.9)	11 (12.2)	0.528
Stage 3 ROP, n (%) [†]	2 (7.1)	7 (7.8)	1.00
Length of invasive ventilation (days) [‡]			
IMV	5.4 (1.12; 14.4)	0.5 (0; 4)	0.001
HFO	1.75 (1; 6.75)	0 (0; 1.56)	0.001
Length of using NCPAP (days) [‡]	2 (0; 25)	11 (3; 25)	0.016
NICU stay (days) [‡]	21 (3.75; 52)	24 (7; 50.25)	0.604


276 ^{*}Chi-square test, [†]Fisher's test, [‡]Mann-Whitney test.

277 Abbreviations: BPD (Bronchopulmonary Dysplasia), HFO (High-Frequency Oscillations),

278 IMV (Intermittent Mandatory Ventilation), NICU (Neonatal Intensive Care Unit), ROP


279 (Retinopathy of Prematurity).

280



281

282 **Figure 1.** The receiver operating characteristic curve of the ridge logistic regression model
283 classifies patients who acquired PH and those who did not. The total sample size was 118
284 patients ($N_{\text{Yes}} = 28$, $N_{\text{No}} = 90$).



285

286 **Figure 2.** Odds ratio of factors associated with pulmonary hemorrhage. Adjusted odds ratios

287 and corresponding confidence intervals estimated from the ridge logistic regression model.

288 The confidence level is 95%, and non-significant results are presented in blue.